

Ecological larval fish groups in Chabahar Bay in relation to day/night and monsoon variations

Nasiri H.R.¹; Rabhaniha M.^{2*}; Attaran-Fariman G.³;
Mousavi Golefid S.A.⁴

Received: April 2016

Accepted: May 2016

1-Chabahar Offshore Fisheries Research Center, Iranian Fisheries Science Research Institute, Agricultural Research, Education and Extension Organization, Chabahar, Iran.

2- Iranian Fisheries Science Research Institute, Agricultural Research, Education and Extension Organization, Tehran, Iran. P.O. Box 14155-775.

3- Chabahar Maritime University, Faculty of Marine Sciences, Department of Marine Biology, Daneshgah Avenue, 99717-56499, Chabahar, Iran.

4- International Sturgeon Research Institute, Iranian Fisheries Science Research Institute, Agricultural Research, Education and Extension Organization

* Corresponding author's Email: rab.mahnaz@gmail.com

Keywords: Larval fish, Ecological groups, Day/night, Monsoon

Introduction

Studies on larval fish abundance allow important inferences about the spawning grounds, reproduction season and migration pattern of fishes (Goulding, 1980; Pavlov, 1994). Larval densities can also be used to estimate the abundance index, which in turn can be correlated to the fisheries yield, and provide an alternative approach for measuring the size of the spawning stock (Smith and Richardson, 1977). Previous studies in this ecosystem have focused on the abundance of fish larva without consideration of daily fish larva variations (Thangaraja, 1987, 1989, 1991; Thangaraja and Al-Aisry, 2001; Rabhaniha *et al.*, 2014). Sanvicente-Añorve *et al.* (2000) reported that seasonality and day/night variations

seem to play an important role on larval fish abundance and composition. In this research we consider the monsoon as an important phenomenon, affecting the daily fish larval distribution.

Material and methods

Day and night samplings of fish larvae were done at 3 stations on the south-eastern coastal areas of Chabahar Bay in the middle of each season during 2013 (Fig. 1) using a plankton-sampler with a mesh size of 300 μ (Smith and Richardson, 1977). Fish larvae identification was based on Leis and Rennis (1983), Houde, *et al.* (1986), Leis and Transky (1989), Olivar, *et al.* (1999) and Richards (2006) and the larva were allocated to ecological groups.



Figure 1: The position of sampling stations in Chabahar Bay.

Data analysis

Fish larvae abundance was standardized to the number of larvae per 10 m² (Smith and Richardson, 1977). Then the fish larvae were separated into different ecological groups based on their spawning behavior pattern following Leis and Rennis (1983), Leis and Transky (1989).

Shannon–Wiener diversity index (H'), richness index (R) and evenness (j) were measured in night and day samples (Ludwig and Reynolds 1988).

Results and discussion

Totally 1163 fish larvae specimens belonging to 29 families with the average abundance of 647.24±30.87 (mean± standard deviation) larvae per 10 m² were collected. Clupeidae, Gobiidae and Blenniidae were the dominant families.

Day time samples included a total of 518 fish larvae specimens belonging to 23 fish families, with an average abundance of 282.12±29.38 larvae per 10 m². Families of Blenniidae,

Scombridae and Clupeidae with a total relative abundance of 65.17% were dominant.

The average Shannon, evenness and richness indices were calculated as 0.88±0.54, 0.66±0.34 and 1.0±0.67 respectively (Table 1). The family names with their relative larval abundance are listed in Table 2. The ratios of ecological groups with consideration of the monsoon period are shown in Fig. 2.

Night samples consisted of 650 fish larvae specimens belonging to 24 fish families, with an average abundance of 365.11±33.23 larvae per 10 m². The family names with their relative larval abundance are listed in table 2. Families of Clupeidae and Gobiidae with a total relative abundance of 65.17% were dominant. Larvae of the families Nomeidae, Paralichthyidae, Platycephalidae and Sphyraneidae were not recorded in night samples. Moreover larvae of the families Blenniidae and Scombridae which were dominant in day time samples,

significantly decreased showing that these families are diurnal or crepuscular. Brebbia and Zubir (2012) reported that new Blenniid larvae feed only on rotifers and other small zooplanktons, while the Scombrid larvae is zooplanktivorous in the early larval stages and starts eating fishes at later stages, whereas others shift early in the larval period. The shift to piscivorous feeding invariably results in an increase in predator growth rate (Juanes *et al.*, 2008). It would be logical to assume that predation in the daytime is more successful. The average

Shannon, evenness and richness indices were calculated as 0.94 ± 0.6 , 0.56 ± 0.3 and 1.36 ± 0.89 , respectively (Table 1). Fig. 3, shows the relative ecological groups in total, pre, and post monsoon at night.

The result of the ratio of day/night fish larvae sampling shows the fish larvae assemblage at night was more than that in the day time. Bonecker *et al.* (2009) found that larval fish densities in Mucuri estuary were significantly higher during the night compared to daylight sampling.

Table 1: The bio- factors of fish larva during sampling periods.

Sampling period	number	abundance	H'	J	R
Day	518	282.12±29.38	0.88±0.54	0.66±0.34	1.0±0.67
Night	650	365.11±33.23	0.94±0.6	0.56±0.3	1.36±0.89

Shannon–Wiener diversity index (H'), richness index (R) and evenness (j)

Table 2: Relative abundance of fish larva and ecological habitats during sampling periods.

family	habitat	spawning	Relative abundance day	Relative abundance night	Relative abundance pre-monsoon	Relative abundance post-monsoon
Apogonidae	coral reef	demersal	0.131	0.292	0.55	0.00
Blenniidae	demersal	demersal	31.242	1.777	4.18	21.60
Bothidae	demersal	pelagic	0	0.427	0.13	0.32
Callionymidae	demersal	pelagic	0.407	0.655	1.35	0.00
Carangidae	pelagic	pelagic	10.885	4.365	3.65	9.58
Clupeidae	pelagic	pelagic	14.833	27.361	47.86	4.30
Cynoglossidae	demersal	pelagic	0.898	0.998	1.40	0.65
Engraulidae	pelagic	pelagic	5.030	9.486	14.59	2.76
Gerridae	demersal	pelagic	0.122	0.889	0.13	0.84
Gobiidae	demersal	demersal	5.360	26.389	5.69	25.01
Lethrinidae	demersal	pelagic	0	0.407	0.41	0.11
Leiognathidae	demersal	pelagic	0.261	2.706	3.44	0.42
Lutjanidae	demersal	pelagic	0.253	0.817	1.41	0.00
Monacanthidae	demersal	demersal	0.430	0.115	0.00	0.42
Mugilidae	demersal	pelagic	1.383	0.182	1.55	0.13
Myctophidae	mesoplagic	pelagic	1.435	0.439	0.00	1.46
Nemipteridae	demersal	pelagic	0	2.254	2.45	0.47
Nomeidae	pelagic	pelagic	0.388	0	0.42	0.00
Paralichthyidae	demersal	pelagic	0.625	0	0.00	0.46
Platycephalidae	demersal	pelagic	0.131	0	0.14	0.00

Table 2 continued:

Polynemidae	demersal	pelagic	0	2.337	3.27	0.00
Pomacenteridae	coral reef	pelagic	0	0.790	0.63	0.32
Scomberidae	pelagic	pelagic	19.327	0.257	0.00	14.32
Scorpanidae	demersal	pelagic	0	0.105	0.00	0.10
Serranidae	demersal	pelagic	0.697	4.526	1.38	3.85
Solenostomidae	demersal	brooding eggs	0.208	0	0.00	0.15
Sparidae	demersal	pelagic	3.236	11.921	1.69	12.49
Sphyracidae	pelagic	pelagic	1.220	0	1.31	0.00
Triacanthidae	demersal	pelagic	1.500	0.507	2.18	0.10




Figure 2: The percent of ecological groups in the day time (a: total; b: pre-monsoon; c: post-monsoon).




Figure 3: The percent of ecological groups in the night time (a: total; b: pre-monsoon; c: post-monsoon).

In this research we found that during pre-monsoon, the fish larvae assemblage was mainly dominated by the larvae emerging from pelagic eggs (89.4%). During post-monsoon, this pattern changed to fish larvae with demersal eggs (47%) and reduction of pelagic eggs (52%). Malzahn and Boersma (2007) obtained the same results in the North Sea. By comparing Figs. 2 and 3, we notice that during post monsoon, the abundance of the demersal fish group increased, whereas that of the pelagic group decreased considerably. In a study on the fish larvae composition of the coastal waters of Bushehr in Persian Gulf, Rabbaniha *et al.* (2015) reported that the pelagic fish larvae were dominated during the warm period, but during the cold period the demersal groups were dominant, which was in agreement with our results. Our study indicated that Clupeid larvae were dominant in Chabahar Bay during pre-monsoon. This family is pelagic and a high abundance of its larvae was also reported from the Iranian coastal line of the Persian Gulf during the warm season (Vosoghi *et al.*, 2010) which is equivalent to pre-monsoon in our research. During post-monsoon, especially in the night time samples, clupeid abundance decreased to 1/10 of its pre-monsoon period, showing a sharp increase in the ratio of demersal to pelagic groups (Fig. 3c). This result corresponds with the clupeid reproduction cycle which occurs during the pre-monsoon period.

Acknowledgements

We thankfully acknowledge the cooperation of the Ecology Department of Chabahar Offshore Fisheries Research Center for their assistance in sampling.

References

- Bonecker, F. T., Castro, M. S. and Bonecker, C. T. 2009.** Larval fish assemblage in a tropical estuary in relation to tidal cycles, day/night and seasonal variations. *Pan-American Journal of Aquatic Sciences*, 4(2): 239-246.
- Brebbia, C.A. and Zubir, S.S. 2012.** Management of natural resources, sustainable development and ecological hazards .WIT press .659 P.
- Goulding, M., 1980.** The fishes and the forest: Explorations in Amazonian natural history. University of California Press, Los Angeles, 200P.
- Houde, E.D., Almatar, J.C. Leak, Down, C.E., 1986.** Ichthyoplankton abundance and diversity in the Western the Gulf. Kuwait Bulletin of Marine Science, No. 8, Kuwait Institute for Scientific Research (KISR), Kuwait, pp.107-393.
- Juanes, F., Buckel, J.A. and Scharf, F.S., 2008.** Feeding ecology of piscivorous fishes (pp. 267–283). Part 3: Fish as predators and prey. Handbook of fish biology and fisheries, Volume 1: Fish biology. 413P.

- Leis, J.M. and Rennis, D.S., 1983.** The larvae of Indo Pacific coral reef fishes. New South Wales University Press, Sydney. 269P.
- Leis, J.M. and Transky, T., 1989.** The larvae of Indo- Pacific shore fishes. New South Wales University Press, Sydney. 371P.
- Ludwig, J.A. and Reynolds, J.F., 1988.** Statistical ecology: A primer of methods and computing. Wiley Press, New York, New York. 337P.
- Malzahn, A.M. and Boersma, M., 2007.** Year-to-year variation in larval fish assemblages of the Southern North Sea. *Helgoland Marine Research*, 61, 117–126. DOI: 10.1007/s10152-007-0059-6.
- Olivar, M.P., Moser, H.G. and Beckley, L.E., 1999.** Lanternfish larvae from the Agulhas current (SW Indian Ocean). *SCI Scientia Marina Journal*, 63(2), 101-120.
- Pavlov, D.S., 1994,** The downstream migration of young fishes in rivers: mechanisms and distribution. *Folia Zoologica*, 43, 193-208.
- Rabbaniha, M., Mossavi Golesefid, A. and Owfi, F., 2014.** The effect of monsoon on fish larva assemblage changes in Gowatr Bay, North Oman Sea. *Iranian Journal of Fisheries Science*, 13(2), 427-436.
- Rabbaniha, M., Molinero, J.C., López-López, L., Javidpour, J., Primo, A.L., Owfi, F. and Sommer, U., 2015.** Habitat association of larval fish assemblages in the northern Persian Gulf. *Marine Pollution Bulletin*, 97(1-2), 105-110.
- Richards, W.J., 2006.** Early stages of Atlantic fishes, an identification guide for the Western Central North Atlantic, Two Volume Set. Taylor & Francis. 2640P.
- Sanvicente-Añorve, L., Flores-Coto, C. and Ciappa-Carrara, X., 2000.** Temporal and spatial scales of ichthyoplankton distribution in the southern Gulf of Mexico. *Estuarine, Coastal and Shelf Science*, 51, 463-475.
- Smith, P.E. and Richardson, S.L., 1977.** Standard techniques for pelagic fish eggs and larvae survey. FAO, Rome, Italy. 110P.
- Thangaraja, M., 1987.** Ichthyoplankton studies. A research brief 1987. Ministry of Agriculture and Fisheries, Marine Science and Fisheries Centre, Marine Ecology Section, Sultanate of Oman, 40P.
- Thangaraja, M., 1989.** Ichthyoplankton studies of the coastal waters off Oman. MSFC Research Report Number 89-2. Ministry of Agriculture and Fisheries, Marine Science and Fisheries Centre, Marine Ecology Section, Sultanate of Oman, 29P.
- Thangaraja, M., 1991.** Fish eggs, larvae and their ecology of the coastal waters off Muscat. MSFC Research Report Number 91-3. Ministry of Agriculture and Fisheries, Marine Science and Fisheries Centre, Marine Ecology Section, Sultanate of Oman, 38P.

- Thangaraja, M. and Al-Aisry, A., 2001.** Studies on the occurrence and abundance of fish eggs and larvae in the waters of Sultanate of Oman. In: Clareboudt M., Goddard S., Al-Oufi H. and McIlwain J. eds. Proc. 1st International Conference on Fisheries, Aquaculture and Environment in NW Indian Ocean, Sultan Qaboos University, Muscat, Sultanate of Oman. pp.13-36.
- Vosoghi, G.H., Fatemi, M.R., Jamili, S., Nikoyan, A.R. and Rabbaniha, M., 2010.** The fluctuation of coralline fish larvae & non coralline fish larvae abundance of Khark & Kharko (Persian Gulf). *Iranian Journal of Fisheries Sciences*, 4(3), 136-142.