Research Article Carbon dioxide anesthesia: A potential application to improve the air exposure duration of tilapia

Fan X.¹; Zhao M.M.¹; Guan W.L.¹; Liu T.T.; Chen D.W.^{1*}

Received: March 2018

Accepted: January 2019

Abstract

The behavioral and physiological responses of tilapia to air exposure after pure saturated carbon dioxide (CO₂) anesthesia were analyzed, and a potential application of live tilapia dehydrated transport was evaluated by comparing it with the tricaine methanesulfonate (MS-222) anesthesia and no anesthesia. Tilapia displayed about 1 min of hyperactivity, and the induction time of anesthetized was about 3 min during CO₂ anesthesia, both the behavioral analysis and plasma cortisol levels showed that tilapia were more sensitive and stressed to the CO₂ anesthesia than MS-222 anesthesia. During air exposure, both anesthesia groups had good sedative effects, and the survival time of unanesthetized group, CO₂ anesthesia group and MS-222 anesthesia group were 331, 489 and 528 min respectively, and the survival time and survival rates of the anesthetized tilapia were significantly increased. Considering the efficacy, ease of use, as well as safety, the results of this study showed that CO₂ was an ideal anesthetic to improve air exposure duration of tilapia, and carbon dioxide anesthesia of tilapia potentially can be translated into an industrial short-time air exposure procedure.

Keywords: Carbon dioxide, Anesthesia, Tricaine Methanesulfonate, Tilapia, Behavioral and physiological response, Air exposure duration

1-Department of Food Science, Guangxi University, 530004, Nanning, Guangxi, P.R.China. *Corresponding author's Email: chendw@gxu.edu.cn.

Introduction

In recent years. tilapia (mainly Oreochromis niloticus) has become one of the largest single cultured fishery species in China, with a production of 1.65 million metric tons in 2013, of which 0.86 million tons of the production were used in processing frozen tilapia product (Fisheries Bureau, Ministry of Agriculture of The People's Republic of China, 2013). Currently, most farms are far from fish processing plants, tilapia are netted and placed in tank with an agitator for aeration, mixing with oxygen, and transported for about 2-5 hours to the processing plant. During transportation, water quality, physical damage and physiological problems were of great impacts on the mortality of tilapia (Colt and Kroeger, 2013). Fish were subjected to intense handling stress and might be injured or even die during the transport and handling procedures (Conte, 2004; Harmon, 2009). Air exposure is unavoidable for fish processing, however, air exposure for a short period of time (less than 10 min) is a significant stressor to fish (White et al., 2008; Trushenski et al., 2010). Air exposure also cause physical damage, which influence post-release behavior and survival of fish (Cook et al., 2015). Although dehydrated (without large volume of water, only skin and gills were kept moist, and fish were exposed to air) transport or other handling of live tilapia procedure seems attractive, it is seldom used because of strong stress responses and high mortality. However, fish anesthetized properly could be transported without

water, provided the skin and gills were kept moist under low temperature (Peter, 2008).

Anesthetics, such tricaine as methanesulfonate (MS-222), carbon dioxide benzocaine $(CO_2),$ and phenoxyethanol have been used to sedate fish. lower metabolism and oxygen demand, and minimize stress responses and physical damages during handling procedures (Pramod et al., 2010; Erikson, 2011; Vandergoot et al., 2011; Cosenza, et al., 2014; Oberg et al., 2015). MS-222 is one of the most widely used anesthetic in fish, and it has high efficacy for anesthetizing fish (Carter et al., 2011). However, a 21-day of withdrawal period is required by the US FDA for use on fish for food, and use of **MS-222** should be restricted to Ictaluridae, Salmonidae, Esocidae and Percidae, and water temperature should not exceed 10°C. Therefore, use of MS-222 for anesthetizing food fish is often not practical. CO₂ is a food additive and widely used in the food industry, and it is also used as an anesthetic in fishery to immobilize fish before slaughter (Roth et al., 2006; Erikson, 2008; Vandergoot et al., 2011). Compared with other fish anesthetics, CO₂ is cheap and easy to use, in combination with the advantage that it leaves no toxic residues in the product. However, CO₂ anesthesia usually results in mild or strong stress response, so fish may be physically damaged or even die (Erikson, 2011; Seth et al., 2013).

The objective of this study was to evaluate the effects of air exposure after carbon dioxide anesthesia on behavioral and physiological responses of tilapia, by comparing it with the tricaine methanesulfonate anesthesia and no anesthesia. Results from this study have important implications for developing a short-time air exposure procedure for handling tilapia.

Materials and methods

Animals

All the experiments were performed according to Guidelines of the European Union Council (86/609/EU) for the use of laboratory animals, and ethical permit Ethics approved by the Animal Committee of Guangxi University. Live tilapia (Oreochromis niloticus) ranging from 470-550g (mean±SE: 510±21g) were purchased in a local farm and transported to the laboratory, where they were acclimated for two days before the experiment began in continuously aerated 50L tanks, with a stocking density of 20 fish per tank ($15\pm1^{\circ}$ C, pH 6.8-7.2, dissolved oxygen levels 5.9-7.3 mg L⁻¹).

Behavioral analysis

Behavior of the two fish groups, the CO₂ and MS-222 anesthesia tilapia, was recorded using a video camera and visually observed. The indicator of consciousness was assessed according to the method of Roth *et al.* (2003). Behavioral indicators at different stages of anesthesia are described in Table 1: (such as Stage 3: No swimming activity, problems of ventilation of operculum, total loss of equilibrium - fish turn over). A total of 16 tilapia were observed for each group. Each experiment was performed in quadruplicate.

respectively.			
Stage	Behavioral responses	Induction time (s)	
		CO ₂ anesthesia	MS-222 anesthesia
0 (Normal)	Normal Then appear agitated, exhibit jumping behavior.	0-30 30-90	0-10 10-30
1 (Light sedation)	Reduced swimming activity, had problems with equilibrium, but with normal ventilation of operculum.	117±23 ^{a,x}	34±5 ^{a,y}
2 (Narcosis)	Weak swimming activity, slow and long ventilation rate, equilibrium loss with efforts to right.	135±17 ^a .x	157±9 ^{b,y}
3 (Deep narcosis)	No swimming activity, problems of ventilation of operculum and total loss of equilibrium - fish turn over.	171±15 ^{b,x}	293±16 ^{c,y}

Table 1: Behavioral responses and induction time	(seconds) of tilapia exposure to CO ₂ and MS-222,
normostivaly	

^{*a*, *b*, *c*} Data in column with the same superscripts are not significantly different while data with different superscripts are significantly different (p<0.05). ^{*x*, *y*} Data between CO₂ and MS-222 anesthesia with different superscripts are significantly different (p<0.05).

Carbon dioxide anesthesia experiment Forty-five fish were randomly divided into three groups: Group 1 (CO₂ anesthesia group), CO₂saturated water (pH 5.60-5.65) was prepared via flushing with pure CO₂ gas during this procedure. Fish (n=16) were exposed to CO₂-saturated water with the density of 2 fish L^{-1} at $15\pm1^{\circ}C$. Each experiment was performed in quadruplicate.

Group 2 (MS-222 anesthesia group), Fish (n=16) were exposed to 100 mg L⁻¹ of MS-222 buffered with NaHCO₃ (200 mg L⁻¹) in aerated water, with the density of 2 fish L⁻¹ at 15±1°C. Each experiment was performed in quadruplicate. MS-222 anesthesia was used as a positive control.

As long as the fish of the two groups reached the third stage of anesthesia, they were immediately netted from the anesthetized tank, then placed in a temperature-controlled $(15\pm1^{\circ}C)$ box for air exposure experiment. The choice of doses, temperature and time used was based on our previous study (Fan *et al.*, 2014).

Group 3 (control/unanesthetized group), Control fish (n=13) were put in the basket directly without being anesthetized.

Air exposure experiment

During the air exposure experiment, fish were placed in a basket and then the basket was put in a constant-temperature box $(15\pm1^{\circ}C)$, and fish were sprayed constant-temperature water $(15\pm1^{\circ}C)$ every 10 min to kept moist of the skin and gills. Fish was considered to be dead when its respiratory movement ceased (Roth *et al.*, 2003). Fish survival time and survival rates at different times were the metric used to evaluate each treatment.

Determination of blood variables

Blood samples (0.5 mL sample⁻¹) were collected by using heparinised syringes as soon as fish were anesthetized before air exposure and after 450 min of air exposure (for the CO₂ and MS-222 anesthesia group), and after 300 min of air exposure (for the control group). Whole blood was spun and plasma was collected. Subsequently, a clinical blood analyzer (TBA-120FR, Toshiba medical systems Co., Japan) was used to analyze lactate (Lac) and glucose (Glu). Plasma cortisol analyzed with was the radioimmunoassay technique (DPC Immulite 2000, DPC Cirrus Co., USA). Each experiment was performed in triplicate. The of **MS-222** data anesthesia before air exposure in this study were also used as the baseline data.

Statistical analysis

SPSS statistics 17.0 (SPSS Inc., Chicago, USA) was used for analyzing the data. Analysis of Variance (ANOVA) was used to determine the significant differences (p<0.05) among the treatments.

Results

Behavioral responses

Initially after anesthetic exposure, tilapia in both groups swam around calmly, then after tens of seconds, they appeared agitated by exhibition jumping behavior (hyperactivity) and attempted to escape (Table 1). The hyperactivity was more intense in the CO_2 anesthesia experiment. After the anesthetics began to take effect, fish reached the second stage of light sedation (narcosis). In the third stage, fish were in deep narcosis, and they had reduced ventilation of operculum and exhibited total loss of equilibrium. At the third stage, tilapia were immediately netted from the anesthetized tank, and placed in a temperature-controlled box for the air exposure experiment.

In the CO₂ anesthesia experiment, although tilapia displayed a brief and violent struggle, there were markedly faster induction times compared with MS-222 anesthesia experiment (171 s vs. 293 s). During air exposure, there were no jumping behavior in both the CO₂ and MS-222 anesthesia group fish, while the control group fish were jumping for nearly 20 min, then they were exhausted.

Blood parameters and plasma cortisol

As shown in Figure 1, the CO_2 anesthesia experiment induced a significant increase in plasma cortisol levels as compared with MS-222 anesthesia experiment. Because MS-222 is a very mild anesthetic, and it's widely used for anesthetizing fish to get the baseline data of fish physiology (Carter et al., 2011), the data of MS-222 anesthesia before air exposure in this study were also used as the baseline data. After 300 min (for the control group) and 450 min (for the CO₂ and MS-222 anesthesia groups) of air exposure, the plasma cortisol levels in all groups increased up to 4-5 times higher than the baseline data. The plasma cortisol level in CO₂ anesthesia experiment was higher than those of the MS-222 anesthesia and control treatments. The blood plasma glucose level in CO_2 anesthesia treatment was significantly higher than that of the MS-222 anesthesia treatment, while lactate level in CO_2 anesthesia treatment was significantly lower than that of the MS-222 anesthesia treatment. After air exposure, as expected, the plasma glucose levels in all groups dropped sharply, while lactate levels rose rapidly at the same time.

Survival rates and survival time during air exposure

Survival rate of the control group after 280 min of air exposure was 100% (Table 2), and then began decreasing as air exposure time continued, with the bulk of mortalities occurring at 280-400 min of air exposure. No mortality was observed in the CO₂ and MS-222 anesthesia groups within 450 min of air exposure. Most of the fish died between 480-500 min (for CO₂ anesthesia group) 480-600 min (for **MS-222** and anesthesia group) of air exposure. The survival time of the control group, CO₂ anesthesia group and MS-222 anesthesia group were 331, 489 and 528 min, respectively. Survival rates and survival time in the control group were significantly lower than in the CO₂ and MS-222 anesthesia groups. The MS-222 anesthesia group had the highest survival rate and survival time.



Figure 1: Plasma cortisol and blood parameters in tilapia at time of 0 min (before air exposure) and after 300 min (for the control group) and 450 min (for the CO₂ and MS-222 anesthesia groups) of air exposure.

Air exposure time	Survival rate (%)			
(min)	CO ₂ anesthesia group	MS-222 anesthesia group	Control group	
280	100	100	100	
300	100	100	80	
350	100	100	40	
400	100	100	0	
450	100	100	0	
480	80	100	0	
500	10	70	0	
550	0	40	0	
600	0	0	0	
Survival time (min)	489 ± 9^{a}	528±32 ^b	331±25°	

Table 2: Survival rates at different times and survival time of the three groups of tilapia during air exposure (N=10).

a, b, c Data in row with different superscripts are significantly different (p < 0.05).

Discussion

Behavioral analysis showed that tilapia were more sensitive and stressed to CO₂ anesthesia than MS-222. MS-222 is generally considered to be effective for anesthetizing fish, and it has little side effect on stress when properly used (Carter et al., 2011). MS-222 is absorbed rapidly through gills and has direct actions on the central nervous system, and exerts it sedative effect by preventing generation and conduction of nervous impulse (Carter et al., 2011). In this study, tilapia displayed a very short and weak struggling behavior, and then came into the light sedation stage during MS-222 anesthesia, which proved that MS-222 was effective for anaesthetizing tilapia. The effectiveness of CO_2 is due to causing hypercapnia, which can depress the central nervous system, resulting in the loss of consciousness and voluntary motor action (Sanderson and Hubert, 2007). When anesthetizing fish with CO₂, there is a hypothesis that supplemental oxygen (i.e. hyperoxic carbon dioxide anesthesia) may reduce stress during CO₂ anesthesia (Kohler et al., 1999). However, some studies (Sandblom et al., 2013) have examined the physiological responses to hyperoxic carbon dioxide anesthesia in fish, and found that the increase in plasma cortisol was similar with the pure CO_2 anesthesia. which suggested that supplemental oxygen did not markedly reduce stress. Therefore, pure saturated CO_2 was deliberately chosen to anesthetize the fish rapidly for this study. In this study, tilapia displayed about 1 min of hyperactivity, with an induction time of about 3 min during CO₂ anesthesia. These results were similar to those of using CO₂ anesthesia of Arctic char (Sandblom et al., 2013; Seth et al., 2013), common snook (Centropomus undecimalis), and Florida pompano (Trachinotus carolinus) (Oberg et al., 2015).

Plasma cortisol level is a sensitive indicator of primary stress response (Bonga, 1997). Therefore, measurements of cortisol are common in the assessments of stress response in fish (Foo and Lam, 1993; Sandblom *et al.*, 2013; Seth *et al.*, 2013). In this study, plasma cortisol levels increased significantly after CO₂ anesthesia, and the cortisol level in CO₂ anesthesia treatment was twice that of the MS-222 anesthesia, which means that CO₂ anesthesia treatment tilapia specimens were more stressed as compared with MS-222 anesthesia treatment. Plasma cortisol levels coincided with the behavioral analysis results. Stress during anesthesia process may be unavoidable. However, compared with other more severe handling stress, for example, the cortisol level was over 100 ng mL⁻¹ when tilapia were confined (Vijayan et al., 1997). Therefore, CO_2 anesthesia in the present study led to a medium stress response as compared with the very mild MS-222 anesthesia.

An increase in plasma glucose level is a secondary typical stress response resulting from activation of glycogenolysis and/or glyconeogenesis by the action of cortisol (Polakof et al., 2012). As usual, elevated plasma cortisol level is bounded with the increase of plasma glucose and lactate levels. In the case of our CO₂ and MS-222 anesthesia experiment, the relationship between changes in the plasma cortisol and plasma glucose was quite obvious. CO2 anesthesia induced a strong release of cortisol, and the maximum plasma cortisol level corresponded to the maximum plasma glucose. However, lactate level in CO₂ anesthesia treatment was significantly lower compared with that in the MS-222 anesthesia treatment; perhaps as the induction time of CO₂ anesthesia was shorter than MS-222 anesthesia (3 min vs. 5 min), and so the accumulation of lactate was less.

As expected, air exposure had great affects on tilapia. Plasma cortisol levels increased sharply after air exposure, so air exposure was a severe stressor to tilapia. During air exposure, there was oxygen available and aerobic no metabolism was shifted to anaerobic metabolism. Anaerobic glycolysis provides energy by breakdown of glucose without the need of oxygen, and lactate is the by-product of anaerobic metabolism (Polakof et al., 2012). Therefore, the plasma glucose levels dropped and lactate levels increased sharply after air exposure. After 300 min (for the control group) and 450 min (for the CO₂ and MS-222 anesthesia groups) of air exposure, the plasma lactate levels in all the groups accumulated to 18.5-18.9 mmol L⁻¹, and energy depletion made fish difficult to stay alive.

Two remarkable phenomena were observed that both the survival rates and survival time of the anesthetized tilapia were significantly increased during air exposure. Anesthesia could sedate fish and lower the metabolism (Carter et al., 2011). For example, at the beginning of air exposure, the control fish were jumping until exhaustion, while anesthetized fish were sedate. Therefore, anesthesia might have prolonged the survival time of tilapia during air exposure. In this study, both the MS-222 anesthesia and the CO₂ anesthesia had exactly good effects, and survival time of their groups was 197 min and 158 min longer than that of the control group, respectively. It would be convenient for the tilapia processing plant if the tilapia were sedate when it is exposed to air for

a short time (about 100-400 min) during the transport and processing procedure. In the present study both anesthesia groups were under good sedative effects that prolonged the survival time during air exposure. Therefore, it seemed that anesthesia would be suitable for improving the air exposure duration of tilapia and would be useful in the tilapia processing plant.

Considering the efficacy, ease of use, as well as safety, CO_2 was an ideal anesthetic for tilapia. Pure saturated CO_2 anesthesia only led to a medium stress response of tilapia. CO_2 anesthetized tilapia was under good sedative effects during air exposure, the air exposure duration was improved and the survival time was prolonged to 489 min. Therefore, a short-time air exposure procedure could be practical when tilapia was properly anesthetized by CO_2 .

Acknowledgments

This work was funded by "National Natural Science Foundation of China (No. 31660448)", "Guangxi Natural Science Foundation (No. 2016GXNSFAA380290 and 2016GXNSFEA380003)", "The Nanning City Science and Technology Promotion Project (No. 20142306)" and "BaGui Scholars Program of Food Biotechnology".

References

Bonga, S.E.W., 1997. The stress response in fish. *Physiological Reviews*, 77(3), 591-625. DOI:10.1152/physrev.1997.77.3.591

- Carter, K.M., Woodley, C.M. and Brown, R.S., 2011. A review of tricaine methanesulfonate for anesthesia of fish. *Reviews in Fish Biology and Fisheries*, 21, 51-59. DOI:10.1007/s11160-010-9188-0.
- **Colt, J. and Kroeger, E.L., 2013.** Impact of aeration and alkalinity on the water quality and product quality of transported tilapia - A simulation study. *Aquacultural Engineering*, 55, 46-58.

DOI:10.1016/j.aquaeng.2013.03.002.

- Conte, F.S., 2004. Stress and the welfare of cultured fish. *Applied Animal Behaviour Science*, 86(3-4), 205-223. DOI:10.1016/j.applanim.2004.02.00 3.
- Cook, K.V., Lennox, R.J., Hinch, S.G. and Cooke, S.J., 2015. Fish out of water: How much air is too much? *Fisheries*, 40(9), 452-461. DOI:10.1080/03632415.2015.10745 70.
- Claudiano, Cosenza, **G.R. G.S.** Marcusso, **P.F.** Eto. **S.F.** Manrique, W.G., Loureiro, B.A., M.T., Salvador, Shimada, R., Moraes, J.R.E. and Moraes, F.R., 2014. Influence of glyceryl guaiacolate ether on anesthetics in tilapia compared to benzocaine and eugenol. Revista MVZ Córdoba, 19(1). 3944-3953. DOI:10.21897/rmvz.114.
- Erikson, U., 2008. Live chilling and carbon dioxide sedation at slaughter of farmed Atlantic salmon: a description of a number of commercial case studies. *Journal of Applied Aquaculture*, 20(1), 38-61. DOI:10.1080/10454430802022078.
- Erikson, U., 2011. Assessment of different stunning methods and

91 Fan et al., Carbon dioxide anesthesia: A potential application to improve the air exposure duration ...

recovery of farmed Atlantic salmon (*Salmo salar*), isoeugenol, nitrogen and three levels of carbon dioxide. *Animal Welfare*, 20(**3**), 365-375.

- Fan, X., Lu, Y.Q., Chen, M.S., Yan, M.Y. and Chen, D.W., 2014. Effect of MS-222 anaesthesia on the transport of live tilapia without water. *Food Science and Technology*, 39, 174-179, in Chinese.
- Fisheries Bureau, Ministry of Agriculture of The People's Republic of China. 2013. China Fishery Statistical Yearbook 2013. China Agriculture Press, Peking, China, in Chinese.
- Foo, J.T.W. and Lam, T.J., 1993. Serum cortisol response to handling stress and the effect of cortisol implantation on testosterone level in the tilapia, *Oreochromis mossambicus*. *Aquaculture*, 115(1-2), 145-158. DOI:10.1016/0044-8486(93)90365-6.
- Harmon, T.S., 2009. Methods for reducing stressors and maintaining water quality associated with live fish transport in tanks: a review of the basics. *Reviews in Aquaculture*, 1(1), 58-66. DOI:10.1111/j.1753-5131.2008.01003.x.
- Kohler, I., Meier, R., Busato, A., Neiger-Aeschbacher, G. and Schatzmann, U., 1999. Is carbon dioxide (CO2) a useful short acting anaesthetic for small laboratory animals? *Laboratory Animals*, 33(2), 155-161.

DOI:10.1258/002367799780578390.

Oberg, E.W., Perez, K.O. and Fuiman, L.A., 2015. Carbon dioxide is an effective anesthetic for multiple marine fish species. *Fisheries* *Research*, 165, 22-27. DOI:10.1016/j.fishres.2014.12.019.

- Peter, C., 2008. W02008097108 -Waterless storage and transport of live aquatic animals. *Patentscope*, W0/2008/097108, PAT -AU2008213166.
- Polakof, S., Panserat, S., Soengas, J.L. and Moon, T.W., 2012. Glucose metabolism in fish: a review. Journal of Comparative Physiology - B: Biochemical, Systemic and Environmental Physiology, 182(8), 1015-1045. DOI:10.1007/s00360-012-0658-7.
- Pramod, P.K., Ramachandran, A., Sajeevan, T.P., Thampy, S. and Pai, S.S., 2010. Comparative efficacy of MS-222 and benzocaine as anaesthetics under simulated transport conditions of a tropical ornamental fish Puntius filamentosus (Valenciennes). Aquaculture 309-314. Research, 41(2), DOI:10.1111/j.1365-2109.2009.02333.x.
- Roth, B., Imsland, A., Moeller, D. and Slinde, E., 2003. Effect of electric field strength and current duration on stunning and injuries in market-sized Atlantic salmon held in seawater. *North American Journal of Aquaculture*, 65(1), 8-13. DOI:10.1577/1548-

8454(2003)065<0008:eoefsa>2.0.co; 2.

Roth, B., Slinde, E. and Robb, D.H.F., 2006. Field evaluation of live chilling with CO₂ on stunning Atlantic salmon (*Salmo salar*) and the subsequent effect on quality. *Aquaculture Research*, 37(8), 799-804. DOI:10.1111/j.1365-2109.2006.01495.x. Sandblom, E., Seth, H., Sundh, H., Sundell, K., Axelsson, M. and Kiessling, A., 2013. Stress responses in Arctic char (*Salvelinus alpinus* L.) during hyperoxic carbon dioxide immobilization relevant to aquaculture. *Aquaculture*, 414-415, 254-259. DOI:10.1016/j.aquaculture.2013.07.

047.

- Sanderson, T.B. and Hubert, W.A., 2007. Assessment of gaseous CO₂ and AQUI-S as anesthetics when surgically implanting radio transmitters into cutthroat trout. *North American Journal of Fisheries Management*, 27(4), 1053-1057. DOI:10.1577/m06-215.1
- Seth, H., Axelsson, M., Sundh, H., Sundell, K., Kiessling, A. and Sandblom, E., 2013. Physiological responses and welfare implications of hypothermia rapid and immobilisation with high levels of CO₂ at two temperatures in Arctic char (Salvelinus alpinus). Aquaculture, 402-403, 146-151. DOI:10.1016/j.aquaculture.2013.04. 004.
- Trushenski, J., Schwarz, M., Takeuchi, R., Delbos, B. and Sampaio, L.A., 2010. Physiological responses of cobia *Rachycentron canadum* following exposure to low

water and air exposure stress challenges. *Aquaculture*, 307(1-2), 173-177.

DOI:10.1016/j.aquaculture.2010.07. 015.

- Vandergoot, C.S., Murchie, K.J., Cooke, S.J., Dettmers, J.M., Bergstedt, R.A. and Fielder, D.G., 2011. Evaluation of two forms of electroanesthesia and carbon dioxide for short-term anesthesia in Walleye. North American Journal of Fisheries Management, 31(5), 914-922. DOI:10.1080/02755947.2011.62971 7.
- Vijayan, M.M., Pereira, C., Grau,
 E.G. and Iwama, G.K., 1997.
 Metabolic responses associated with confinement stress in tilapia: the role of cortisol. *Comparative Biochemistry and Physiology Part C: Pharmacology, Toxicology and Endocrinology*, 116(1), 89-95.
 DOI:10.1016/s0742-8413(96)00124-7.
- White, A.J., Schreer, J.F. and Cooke, **Behavioral** 2008. and S., physiological responses of the congeneric largemouth (Micropterus salmoides) and smallmouth bass (M. dolomieu) to various exercise and air Fisheries exposure durations. 9-16. Research, 89(1). DOI:10.1016/j.fishres.2007.08.008.