

Production of artificial diets for female broodstock of western white shrimp (*Litopenaeus vannamei*) and study on their singular effect

Ghorbani Vaghei R.^{*1}; Abolhasani M.H.²; Matinfar A.³; Dadgar Sh.³; Ghorbani R.⁴,

Received: January 2016

Accepted: August 2016

Abstract

Advantages of pelleted feeds in terms of storage, transportation and lower costs compared to natural fresh feed justified the replacement of artificial feeds instead of natural ones partially on maturation performances. This study comprised nine treatments (with three replications) including control treatment: polychaete worm (*Perinereis nuntia*), Squid and calf liver (natural feed); T₁: pellet feed with 50% crude protein and 8% crude lipid+ *Perinereis nuntia*, Squid and calf liver; T₂: pellet feed with 50% crude protein and 10% crude lipid+ *Perinereis nuntia*, Squid and calf liver; T₃: pellet feed with 40% crude protein and 10% crude lipid+ *Perinereis nuntia*, Squid and calf liver; T₄: pellet feed with 40% crude protein and 8% crude lipid+ *Perinereis nuntia*, Squid and calf liver; T₅: pellet feed with 50% crude protein and 10% crude lipid; T₆: pellet feed with 50% crude protein and 8% crude lipid; T₇: pellet feed with 40% crude protein and 10% crude lipid; T₈: pellet feed with 40% crude protein and 8% crude lipid. In treatments 1, 2, 3 and 4, 50% of the natural diets were removed and pellet foods used instead. The amount of pellet feeds that given daily, was 3.5% of shrimp broodstock biomass. In treatments 5, 6, 7 and 8, the amount of pellet feeds that given daily was 7.5% daily. Feeding was done 4 times a day. GSI in the control (3.23%), treatment 3 (3.20%) and treatment 6 (3.02%) were significantly higher than that in the other treatments ($p < 0.05$). Absolute fecundity in the control (29980 eggs) and treatment 3 (29683 eggs) was significantly higher than other treatments ($p < 0.05$). HSI reached the lowest level in treatment 8 (2.14%). But in treatments 3 and 5 was more than control, but did not any statistical significant ($p > 0.05$). Survival rate in the treatment 8 (26.66%) was significantly lower than other treatments ($p < 0.05$). Generally, In the treatment 3, with the elimination of 50% of the natural foods, and the use of pellet food instead of them, result of comparative indices of reproduction were in desirable proportions.

Keywords: Natural wet feed, Pellet feed, Sexual maturation, Western white shrimp female brood

1- Shrimp Research Center, Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research Education and Extension Organization (AREEO), Bushehr, Iran

2- Department of Natural Resources, Faculty of Agriculture, Islamic Azad University, Khorasgan Isfahan Branch, Esfahan, Iran.

3- Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research Education and Extension Organization (AREEO), Bushehr, Iran

4- Department of Natural Resources, Agriculture and Natural Resources, Gorgan, Iran

* Corresponding author's E-mail: ghorbani_v2@yahoo.com

Introduction

It has been reported that total replacement of natural feeds with artificial ones has not been achieved, yet (FAO, 2010). Several factors have made it impossible to use large amount of pellet feed for shrimp broodstock' maturation. Chimsung (2014) have reported that polychaete worm improves shrimp reproduction due to having high nutritional value, arachidonic acid and important reproduction hormones. In addition, it has been believed that polychaete worm is an excellent source for not only unsaturated fatty acids, but also sexual hormones similar to those of shrimp. Also, squid not only is a rich source of cholesterol, but also contains sexual steroids involved in vitellogenesis (Wouters *et al.*, 2005). As mentioned before, natural feeds for breeder feeding have high amounts of n-3 fatty acid. Also, fecundity rate of broodstock is correlated with n-3 fatty acid (Wouters *et al.*, 2005).

Amino acids, fatty acids, protein and lipid play an important role in shrimp breeder maturation and need more protein during the sexual maturation period (Coutteau, 2001). In addition, highly unsaturated fatty acids (HUFA) especially eicosapentaenoic acid and docosapentaenoic acid are important components of breeder diets and are found abundantly in ovary tissue (Teikwa and Mgaya, 2003). Diets deficient in n-3 HUFA adversely affect ovary development, fecundity and egg quality (Teikwa and Mgaya, 2003).

Western white shrimp broodstock with frequent spawning have higher protein content in their hepatopancreas and ovaries than those broodstock with weak spawning (Magalhaes *et al.*, 2012). Successful sexual maturation in western white shrimp can be obtained with freshly-frozen wet diet (with or without artificial diets). The researchers have emphasized that the weight of matured shrimp ovaries may increase 4-6 fold within about one week (Verstraete *et al.*, 1995). Divan *et al.* (2009) reported that pelleted feeds containing 50% protein and 10% lipid can be used in combination with fresh feeds. Alava *et al.* (1993) reported that diets lacking HUFA cause a decrease in gonado-somatic index (GSI). Consequently, it is believed that HUFAs are needed for sexual maturation. Pelleted feeds are important in many aspects, like ease of storage, transportation, feeding and low costs. The aim of the present study was to provide a suitable pelleted diet for the sexual maturation of female broodstock and determining their effects on sexual parameters.

Materials and methods

Broodstock' acclimation

A total number of 270 female broodstock with an average weight of 37 ± 2 g were sampled from Bandargah Research Station and kept in two 4-ton fiberglass tanks for one week to acclimatize prior to stocking in 300-L tanks. Eyestalk ablation was done a week after stocking in the 300-L tanks

filled with 150 L water. Feeding and sexual maturation in broodstock were checked every day for 5 weeks.

Treatments

This study was conducted with 9 treatments each with three replications containing 10 broodstock. Treatments included control: polychaete worm (*Perinereis nuntia*), Squid and calf liver (natural feed); T₁: pellet feed with 50% crude protein and 8% crude lipid+ *Perinereis nuntia*, Squid and calf liver; T₂: pellet feed with 50% crude protein and 10% crude lipid+ *Perinereis nuntia*, Squid and calf liver; T₃: pellet feed with 40% crude protein and 10% crude lipid+ *Perinereis nuntia*, Squid and calf liver; T₄: pellet feed with 40% crude protein and 8% crude lipid+ *Perinereis nuntia*, Squid and calf liver; T₅: pellet feed with 50% crude protein and 10% crude lipid; T₆: pellet feed with 50% crude protein and 8% crude lipid; T₇: pellet feed with 40% crude protein and 10% crude lipid; T₈: pellet feed with 40% crude protein and 8% crude lipid.

Feeding rate (natural feeds) was 25% of shrimp broodstock biomass daily (Brock and Main, 1994). Their composition is detailed (Tables 1 and 2). In treatments 1, 2, 3 and 4, 50% of the natural diets were removed and pellet foods used instead. The amount of pellet feeds that given daily, was 3.5% of shrimp broodstock biomass. In treatments 5, 6, 7 and 8, the amount of pellet feeds that given daily was 7.5% daily. Feeding was done 4 times a day.

Feed types

Wet feed

Polychaete worm (*Perinereis nuntia*), Squid and calf liver were used for the broodstock feeding and sexual maturation.

Pellet feed

Pellet feeds were manufactured with laboratory equipment. In addition, depending to treatment type, four pellet feeds were prepared.

Analyses

Shrimp length including carapace length and total length were measured by a digital calipers (Model TITAN), whereas, total wet weight was measured using a digital scale (Model WANT) at 0.001 g nearest (Conides *et al.*, 2008). The sexual stage, and weight of ovaries, oocyte count, fecundity and hepatopancreas weight in female broodstock were determined (Islam *et al.*, 2012). Also, eyestalk ablation and sexual development assessment of broodstock was performed according to Aelgamal (2015). Water temperature, dissolved oxygen, salinity and pH were measured daily. Data were analyzed using one-way ANOVA and the differences among the treatments were determined using Duncan test ($\alpha=0.05$). All statistical analyses were performed in SPSS 17.

fecundity of the control and treatment 3 was significantly higher ($p<0.05$) than that in the other treatments. Survival

rate in treatment 8 was significantly lower ($p<0.05$) than that in the other treatments.

Table 5: Mean (\pm SD) of growth and reproductive indices in different treatments.

| Features | Control | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
|----------------------|--------------------------------|--------------------------------|--------------------------------|-------------------------------|-------------------------------|-------------------------------|-------------------------------|--------------------------------|--------------------------------|
| Weight (g) | 4.77 \pm 39.06 ^a | 6.17 \pm 39.72 ^a | 3.34 \pm 39.14 ^a | 3.43 \pm 39.68 ^a | 3.45 \pm 38.34 ^a | 3.47 \pm 36.82 ^a | 2.43 \pm 37.61 ^a | 1.52 \pm 36.23 ^a | 2.83 \pm 37.6 ^a |
| Carapace length (cm) | 0.13 \pm 2.54 ^a | 0.21 \pm 2.56 ^a | 0.16 \pm 2.57 ^a | 0.12 \pm 2.58 ^a | 0.08 \pm 2.57 ^a | 0.12 \pm 2.55 ^a | 0.11 \pm 2.52 ^a | 0.11 \pm 2.52 ^a | 0.05 \pm 2.48 ^a |
| Body length (cm) | 0.78 \pm 15.47 ^{ab} | 1.19 \pm 15.33 ^{ab} | 0.64 \pm 15.58 ^{ab} | 1.08 \pm 16.05 ^a | 0.5 \pm 15.66 ^{ab} | 0.79 \pm 14.96 ^b | 0.77 \pm 14.82 ^b | 0.54 \pm 15.17 ^{ab} | 0.25 \pm 15.16 ^{ab} |

In each row, different letters show significant difference among the treatments ($p<0.05$).




Figure 1: Correlation between the treatments and frequency of various stages of maturation.




Figure 2: Effect of different treatments on GSI (different letters above the bars mean significant difference at $\alpha=0.05$).




Figure 3: Effect of different treatments on HSI (different letters above the bars mean significant difference at $\alpha=0.05$).




Figure 4: Effect of different treatments on fecundity (different letters above the bars mean significant difference at $\alpha = 0.05$).




Figure 5: Effect of different treatments on survival rate (different letters above the bars mean significant difference at $\alpha = 0.05$).

There were no significant differences ($p>0.05$) in survival rate and final weight of broodstock among the treatments. The broodstock' length in

the treatments 5 and 6 was significantly lower than that in the other treatments (Table 5).

Mean (\pm SD) of water physico-chemical parameters during the experiment are

presented in Table 6.

Table 6: Mean (\pm SD) of water physico-chemical parameters during the experiment.

| Parameters | Water temperature ($^{\circ}$ C) | Dissolved oxygen (mg/L) | Salinity (ppt) | pH |
|------------|-----------------------------------|-------------------------|----------------|-----------------|
| amount | 30 \pm 1 | 6.93 \pm 0.53 | 33 \pm 2 | 8.16 \pm 0.11 |

Discussion

At present, the nutritional requirements of brood shrimps are hard to supply in propagation centers. The main problems with fresh wet feeds (worm and squid) are their availability, high price, difficulty of transportation and washing prior to use. Thus, in this study 50% and 100% of natural wet feeds were replaced with pellet feeds to investigate the possibility of the replacement of pelleted feeds to wet feeds. Also, it has been reported that total replacement of natural feeds with artificial ones has not been achieved, yet (FAO, 2010). Several factors have made it impossible to use large amounts of pellet feeds for shrimp breeder maturation. Chimsung (2014) reported that polychaete worm, improves shrimp reproduction due to having high nutritional value, arachidonic acid and important reproduction hormones. In addition, it has been believed that polychaete worm is an excellent source of not only unsaturated fatty acids, but also sexual hormones similar to those of shrimp. Also, it was reported that squids contain high levels of protein, HUFA and cholesterol, which are necessary for successful growth and reproduction of shrimp (Chimsung, 2014). In addition, it has been reported that bloodworm, squid, mussel and clam meat along with

pelleted feeds should be used for shrimp feeding (Brock and Main, 1994). Divan *et al.* (2009) reported that pelleted feeds containing 50% protein and 10% lipid (PUFA) can be used in combination with fresh feeds. However, the rate of replacement of fresh feeds with pelleted feeds has not been suggested. Thus, in the present study, the effect of even 100% replacement was investigated on brood shrimp. Sangpradub *et al.*, (1994) investigated the effect of different feeding regimes on ovary maturation and spawning of black tiger shrimp (*Penaeus monodon*) and in line with the results of present study, found that the highest ovary maturation and spawning was related to the broodstock fed either wet feeds or wet feeds plus pelleted feeds.

As, mating occurs predominantly when female broodstock are in stage 4 of sexual maturation, low mating frequency means a few of the broodstock are in stage 4. Such a condition was observed in the present study considering that there were 10 broodstock in each 300-L tank. In the investigation of the breeder sexual maturation, stress delivered by tank siphoning and water replacement should be considered, particularly in small tanks.

In the case of the importance of dietary protein in shrimp broodstock reproduction, Wouters *et al.* (2001) reported that due to synthesis of new biotic materials during the sexual maturation, the animals need higher protein. Also, they reported that protein content of artificial feeds is about 50% depending on shrimp species and protein source. Wouters *et al.* (2004) found an increase in ovary protein content during oocyte development and reproduction. Requirement for dietary lipid depends on HUFA, phospholipids, sterols and energy supplementation. For a long time it has been known that the crustaceans have little ability to synthesis HUFA, phospholipids and sterols (Sangpradub *et al.*, 1994). Artificial feeds have seemingly lower EPA (20:5n-3) compared to natural feeds, thus, they have lower n-3-HUFA and DHA: EPA ratio. Diets for the shrimp brooders during sexual maturation should have high n-3/n-6 ratios. It seems that, they require dietary phospholipids (Sangpradub *et al.*, 1994). Verstracte *et al.* (1995) reported that successful sexual maturation of western white shrimp can be met using freshly-frozen marine organisms without formulated diets. The results of present study showed that, it is possible to eliminate 50% of natural diets and use pellet feed instead.

ACE (2003) reported that for western white shrimp propagation, water salinity, temperature and pH should be 32 ppt, 27-29°C and about 8

respectively and saturated dissolved oxygen of above 5 ppm is required.

Stickney (2000) recommended the following water properties for sexual maturation of tropical shrimp broodstock: salinity=27-36±0.5 ppt, temperature=27-29±0.2°C, pH=7.8±0.2, and dissolved oxygen 5±0.5 mg/L. Regarding the mentioned cases, in the present study the physic-chemical factors of water were in the right range.

Acknowledgements

This study was conducted with the support of Iranian National Science Foundation. We thank the foundation chief, Projects Evaluating Director and other staff. Also, we thank the General Manager, Research Vice-President, Planning and Support Vice-President, Aquaculture Section Head, Research Coordination Manager, Nutrition Group Manager and the other staff at the Iranian Fisheries Research Organization. Our thanks are also forwarded to the General Manager, Research Vice-President, Planning and Support Vice-President of the Iranian Shrimp Research Organization and the other staff at Bandargah Research Station.

References

- Aelgamal, A., 2015.** Ovarian development and eyestalk ablation in white leg shrimp (*Litopenaeus vannamei*). *Fish Consulting Group*, pp. 10-24.
- Alava, V.R., Kanazawa, A., Teshima, S.I. and Koshio, S., 1993.** Effect of

- dietary phospholipids and n-3 highly unsaturated fatty acids on ovarian development of Kuruma prawn. *Nippon Suisan Gakkaishi*, pp. 345-351.
- Automatic Content Extraction (ACE), 2003.** Tiger prawn (*Penaeus monodon*) and white legged shrimp (*Litopenaeus vannamei*). 18P.
- Brock, J.A. and Main, K.L., 1994.** A guide to the common problems and disease of culture *Penaeus vannamei*. The Oceanic Institute Makapuu Point. Thiland. 241P.
- Chimsung, N., 2014.** Maturation diets for black tiger shrimp (*Penaeus monodon*) broodstock: A review. Songklanakarin. *Journal of Science and Technology*, 36(3), 265-273,
- Conides, A., Glamuzina, B., Dulcic, J., Kapiris, K., Jug-Dujakovic, J. and Papaconstantinou, C., 2008.** Study of the reproduction of the Karamote shrimp *Penaeus (Melicertus) kerathurus* in Amvrakikos Gulf, Western Greece. *Acta Adriatica*, 49(2), 97-106.
- Coutteau, P., Pham, D., Pinon, E and Balcazar, Y., 2001.** Mixed maturation diets improve shrimp broodstock performance. *Global Aquaculture Alliance*, created 26.02.2015. 20P.
- Divan, A.D., Joseph, S. and Ayyappan, S., 2009.** Physiology of reproduction, breeding and culture of tiger shrimp *Penaeus monodon (Fabricus)*. Arendra Publishing House, 292P.
- FAO, 2010.** The pre-spawning process. Originated by Fisheries and Aquaculture Department. pp. 9-24.
- Islam, M.R., Sultana, N., Belal Hossain, M. and Mondal, S., 2012.** Estimation of fecundity and gonadosomatic index (GSI) of Gangetic whiting, *Sillaginopsis panijus* (Hamilton, 1822) from the Meghna River Estuary, Bangladesh. *World Applied Sciences Journal*, 17(10), pp. 1253-1260.
- Magalhaes, T., Mossolin, E.C. and Mantelatto, F.L., 2012.** Gonadosomatic and hepatosomatic indexes of the freshwater shrimp *Macrobrachium olfersii* (Decapoda, Palaemonidae) from Sao Sebastiao Island, Southeastern Brazil. *Pan-American Journal of Aquatic Sciences*, 7 (1), pp. 1-9.
- Sangpradub, S., Fast, A.W., Piyatiratorakul, S. and Menasveta, P., 1994.** Effects of different feeding regimes on ovarian maturation and spawning of pond-reared giant tiger prawn in Thiland. Biotec Publication Thiland, pp. 10-23.
- Stickney, R.R., 2000.** Encyclopedia of aquaculture. A Wiley-Interscience Publication. 1060P.
- Teikwa, E.D. and Mgaya, Y.D., 2003.** Abundance and reproductive biology of the penaeid prawns of Bagamoya coastal waters, Tanzania. Western Indian Ocean. *Journal of Marine Science*, pp. 117-126.
- Verstraete, P., Mora, B. and Lavens, P., 1995.** Maturation of *Penaeus*

vannamei by using dry pellets as a partial substitute of the natural diet. 76-78. Ed. I. (Eds) Lavens, P., Jaspers, E., Roelants. Larvi 95. European Aquaculture Society.

Wouters, R., Lavens, Nieto, J. and Sorgeloos, P., 2001. Penaeid shrimp broodstock nutrition: an updated

review on research and development. *Aquaculture*, 202, 1-21.

Wouters, R., Nieto, J. and Sorgeloos, P., 2005. A review of recent research on shrimp broodstock nutrition and artificial diets. Retrieved from <https://www.Feedofbroodstock.htm>., X.