

## ***In vitro* screening of *Sonneratia alba* extract against the oomycete fish pathogen, *Aphanomyces invadans***

**Afzali S.F.; Wong W.L.\***

Received: November 2016

Accepted: January 2017

Department of Biological Science, Faculty of Science, Universiti Tunku Abdul Rahman, Jalan Universiti, Bandar Barat, Kampar 31900, Perak, Malaysia.

\*Corresponding author's Email: wongwl@utar.edu.my

**Keywords:** *Aphanomyces invadans*, Antifungal, Herbal extract, *In vitro* screening, *Sonneratia alba*

### **Introduction**

Epizootic ulcerative syndrome (EUS), caused by the aquatic oomycete fish pathogen, *Aphanomyces invadans* (David and Kirk, 1997) is one of the OIE-listed diseases that leads to huge economic losses in the fish industry in the world. Traditional disease management strategies relying on chemotherapy continue to cause undesirable effects such as antibiotic resistance, environmental pollution and food security issues (Pandey *et al.*, 2012). In addition, there has been no vaccine available to prevent the outbreak of EUS according to the Office International des Epizooties (OIE) (2015). Therefore, many current researches focus on alternative strategies such as using plants for EUS

treatments (Fairweather, 1999; Campbell *et al.*, 2001; Chowdhury and Rahman, 2013; Alam *et al.*, 2014; Uthayakumar *et al.*, 2014; Kumar *et al.*, 2015; Yogeshwari *et al.*, 2015).

Generally, biological compounds are known to be easily biodegradable, environmentally friendly, inexpensive, renewable and locally available, and can be easily prepared (Ortuño *et al.*, 2002; Valladão *et al.*, 2015). More than 400,000 species of tropical plants are known to have medicinal properties (Citarasu, 2010). Biomedicinal plants such as mangrove plant extracts or their by-products contain several active compounds such as phenols, polyphenols, alkaloids, quinones, terpenoids, lectines, and polypeptides that have been shown to be effective

alternatives to traditional chemotherapies and vaccines for disease treatments and control (Citarasu *et al.*, 2002; Sivaram *et al.*, 2004; Kjer *et al.*, 2009; Citarasu, 2010; Harikrishnan *et al.*, 2011). A number of plants, such as *Mikania cordata* (Burm), *Rauwolfia tetraphylla* L., neem (*Azadirachta indica* A. Juss.), akand (*Calotropis gigantea* L.), turmeric (*Curcuma longa* L.), *Curcuma zedoaria* (Christm.), Indian sorrel (*Oxalis corniculata* L.) and *Nigella sativa* L. have been proved to enhance disease resistance of fish against EUS (Campbell *et al.*, 2001; Chowdhury and Rahman, 2013; Kumar *et al.*, 2015; Yogeshwari, *et al.*, 2015). However, most of these plants are required in relatively high amounts to show effectiveness in the treatment of EUS infected fish.

The present study is the first attempt in applying the mangrove plant, *Sonneratia alba* as a biomedicine against EUS causative Oomycete fungus. *Sonneratia alba* (Smith) is a mangrove plant belonging to the family Sonneratiaceae. It is known as mangrove apple or “perepat” in Malaysia (Duke and Jackes, 1987). Members of this family are rich in cyclitol, polyol, sucrose, glucose, fructose, condensed and hydrolysable tannins, minerals, nucleotides (Bandaranayake, 2002). Tannin is known for its antimicrobial activity and probably evolved in plants as a defense against microbial attack (Balasooriya *et al.*, 1982). In the present study, we

aimed to investigate the *in vitro* antifungal effects of the methanol extract of *S. alba* against *A. invadans* using the disc diffusion methods.

### Materials and methods

The leaves of *S. alba* were collected from Sungai Prai in Penang, Malaysia and identified according to Duke and Jackes (1987) (Fig. 1). The fresh leaves were washed with distilled water, blotted dried and freeze-dried using a freeze dryer (Martin Christ, Germany). The dried leaves were then powdered using a mechanical grinder and weighed. The methanol extraction of dried leaves was performed using a Soxhlet extraction device. The extract was distilled using a rotary evaporator (Büchi, Switzerland) and stored in air tight containers at -20°C until required. The agar disk diffusion technique was conducted according to Bailey (1983). Briefly, the *A. invadans* strain (NJM9701) (courtesy of Dr. Oidtmann B, U.K.) was inoculated onto a sterilized glucose peptone (GP) agar media (3 gL<sup>-1</sup> glucose, 1 gL<sup>-1</sup> peptone, 0.128 gL<sup>-1</sup> MgSO<sub>4</sub>.7H<sub>2</sub>O, 0.014 gL<sup>-1</sup> KH<sub>2</sub>PO<sub>4</sub>, 0.029 gL<sup>-1</sup> CaCl<sub>2</sub>.2H<sub>2</sub>O, 2.4 mgL<sup>-1</sup> FeCl<sub>3</sub>.6H<sub>2</sub>O, 1.8 mgL<sup>-1</sup> MnCl<sub>2</sub>.4H<sub>2</sub>O, 3.9 mgL<sup>-1</sup> CuSO<sub>4</sub>.5H<sub>2</sub>O, 0.4 mgL<sup>-1</sup> ZnSO<sub>4</sub>.7H<sub>2</sub>O, 12 gL<sup>-1</sup> technical agar, 10 mL L<sup>-1</sup> Penicillin) plate and incubated for five days at 25°C for preparation of agar discs. A set of three dilutions (100, 500 and 1000 ppm) of *S. alba* extract was prepared with distilled water.



**Figure 1:** *Sonneratia alba* mangrove plant tree can be morphologically characterized by rounded leaf tips (arrow), corolla width flattened fruits (circles), and flower with white and small petals (rectangle).

Distilled water was used as a positive control. The agar diffusion discs experiments were prepared by soaking the pre-inoculated agar discs (6 mm in diameter) individually in different concentrations of 100  $\mu$ l of *S. alba* extract. The discs were soaked in the extracts for one hour at 25°C. The treated discs were then washed three times in the distilled water and placed on the GP agar plates and incubated for seven days at 25°C. Mycelial growth of *A. invadans* was monitored daily for seven days. The antifungal activities of the extracts were evaluated by measuring the mycelial growth (mm) from the edge of the pre-inoculated discs on the GP plates. The boundary dilution without any visible growth was defined as the minimal inhibitory concentration (MIC) for *A. invadans* at the given *S. alba* extract concentrations. The filter paper disk diffusion method (Kohner *et al.*, 1994) was applied to compare the inhibition zone of *S. alba*

extracts with commercially available antifungal agents, namely malachite green, known to be the most effective drug against EUS, and formalin at their minimal inhibiting concentration for *A. invadans* mycelia growth. Sterile filter papers (Whatman, No. 4) with a diameter of 5 mm were impregnated with 100  $\mu$ L of each of the three dilutions of extract solution and placed on the GP agar plate including a pre-inoculated agar cube (10 mm) in the center which was incubated for seven days at 25°C to monitor the mycelia growth of *A. invadans*. All the experiments were performed in triplicate.

### Results and discussion

The methanol extract of *S. alba* effectively inhibited the mycelial growth of *A. invadans* at a minimum concentration of 1000 ppm for both the agar and filter paper diffusion experiments. In the agar diffusion test,


500 ppm of the extract inhibited the fungus mycelial growth up to 96 hours. The mycelial growth from the edge of the pre-inoculated *A. invadans* agar discs treated with *S. alba* extracts at concentrations of 100, 500 and 1000 ppm were 15, 8 and 0 mm respectively (Fig. 2). The results of the filter paper disc test showed that the *S. alba* extract at its minimal inhibitory concentration (1000 ppm) has similar qualitative inhibitory effects as malachite green at 1 ppm and formalin at 250 ppm (Fig. 3).

The antimicrobial activity of *S. alba* extract was previously reported by Saad *et al.* (2012) who suggested that this plant extract may possess therapeutic property against human pathogens and infectious diseases caused by *Escherichia coli*, *Staphylococcus aureus* and *Bacillus cereus*. However, the authors had noted that *S. alba* extract was not sensitive even at high concentration (10,000 ppm) against the


tested fungal strain, *Candida albicans*, and yeast, *Cryptococcus neoformans*.

Campbell *et al.* (2001) had tested several plant extracts and showed that high concentrations (5000-10000 ppm) of herbals such as turmeric (*Curcuma longa*), propolis resin, *Calophyllum inophyllum* L., *Eugenia caryophyllus* L., and *Ficus pumila* L. are effective in inhibiting the mycelial growth of *A. invadans in vitro*. However, the application of high concentration of herbals seems not to be practical due to the expenses may be imposed.

It is reported that neem, *A. indica*, extract (azadirachtin) may inhibit *A. invadans* sporulation at 1000 ppm *in vitro* (Campbell *et al.*, 2001), and possesses antifungal effect on EUS-affected fish (Harikrishnan *et al.*, 2005; Chowdhury and Rahman, 2013) as well.



**Figure 2:** Mycelial growth of pre-inoculated *Aphanomyces invadans* agar discs treated with different concentrations of *Sonneratia alba* extract and distilled water on a glucose peptone plate. a) Distilled water (positive control); b) *Sonneratia alba* extract of 500 ppm; c) *Sonneratia alba* extract of 100 ppm; d) *Sonneratia alba* extract of 1000 ppm.



**Figure 3:** Inhibition zone of mycelial growth of *Aphanomyces invadans* resulting from malachite green, formalin and different concentrations of *Sonneratia alba* extract on the filter paper discs. a) formalin at 250 ppm; b) *Sonneratia alba* extract at 1000 ppm; c) *Sonneratia alba* extract at 500 ppm; d) Malachite green at 1 ppm.

Campbell and his colleagues (2001), also tested other herbal extracts including *Psidium guajava*, *Phyllanthus urinaria*, *Phytolacca americana*, garlic clove, *Acalypha lanceolata*, *Acalypha indica*, tea tree oil and witch-hazel that did not show any inhibitory effects against *A. invadans* even at higher concentration (5000 – 1000000 ppm), and cannot be recommended as prophylactic measure against EUS. In another study, Fairweather (1999) found that administration of D-limonene, a natural compound found in orange peels with anti-cancer properties (Sun, 2007), is totally ineffective on *A. invadans* growth at all concentrations tested *in vitro*. Even though malachite green and formalin were reported to be able to inhibit the mycelial growth of *A. invadans* at 1 ppm and 250 ppm, respectively (Campbell *et al.*, 2001), these chemicals are toxic and malachite

green, for example, is forbidden by U.S. food and drug administration (FDA) in aquaculture (Khan *et al.*, 2011). The present study suggests that *S. alba* methanol extract with a concentration of 1000 ppm may be used as an alternative in controlling the invasive growth of the EUS causative agent, *A. invadans*. This can be exploited further in the development of potent antifungal compounds.

#### Acknowledgements

This research was supported by a research grant from the Universiti Tunku Abdul Rahman, Malaysia (UTAR RF. No. 6200/W48) awarded to Wong WL and Afzali SF, and by the Universiti Tunku Abdul Rahman post-doctoral research scholarship awarded to Afzali SF. The authors are indebted to Dr. Birgit Oidtmann (CEFAS, United

Kingdom) for her support in providing *A. invadans* strain (NJM9701).

## References

- Alam, M.N., Ahmed, G.U. and Chowdhury, M.B.R., 2014.** Performance of herbal extracts on diseased fish. *Bangladesh Journal of Veterinary Medicine*, 12, 225–230.
- Bailey, T.A., 1983.** Method for *in vitro* screening of aquatic fungicides. *Journal of Fish Diseases*, 6, 91–100.
- Balasoorya, S.J., Sotheeswaran, S. and Balasubramaniam, S., 1982.** Economically useful plants of Sri Lanka. Part IV. Screening of Sri Lanka plants for tannins. *Journal of the National Science Council of Sri Lanka*, 10, 213–219.
- Bandaranayake, W.M., 2002.** Bioactivities, bioactive compounds and chemical constituents of mangrove plants. *Wetlands Ecology and Management*, 10, 421–452.
- Campbell, R.E., Lilley, J.H., Taukhid, Panyawachira, V. and Kanchanakhan, S., 2001.** *In vitro* screening of novel treatments for *Aphanomyces invadans*. *Aquaculture Research*, 32, 223–233.
- Chowdhury, M.B.R. and Rahman, T., 2013.** Use of low-cost chemotherapeutic and medicinal plants against Thai silver barb (*Barbonymus gonionotus*). *Journal of Bangladesh Agricultural University*, 10, 385–390.
- Citarasu, T., Babu, M.M., Sekar, R.R.J. and Petermarian, M., 2002.** Developing Artemia enriched herbal diet for producing quality larvae in *Penaeus monodon*, Fabricius. *Asian Fisheries Science*, 15(1), 21–32.
- Citarasu, T., 2010.** Herbal biomedicines: a new opportunity for aquaculture industry. *Aquaculture International*, 18, 403–414.
- David, J.C. and Kirk, P.M. and IMI (International Mycological Institute), 1997.** Index of fungi, Vol. 6, Part 13 in Ainsworth and Bisby's dictionary of the fungi, 8<sup>th</sup> edition, CAB International, Wallingford, UK, 706.
- Duke, N.C. and Jackes, B.R., 1987.** A systematic revision of the mangrove genus *Sonneratia* (Sonneratiaceae) in Australasia. *Blumea-Biodiversity Evolution and Biogeography of Plants*, 32, 277–302.
- Fairweather, D.J., 1999.** Development of a bath challenge system to study component causes, and preventative treatments, of epizootic ulcerative syndrome (EUS) in snakehead fish (*Channa striata*). Master of Science Dissertation. University of Plymouth, UK.
- Harikrishnan, R., Balasundaram, C. and Bhuvaneshwari, R., 2005.** Restorative effect of *Azadirachta indica* aqueous leaf extract dip treatment on haematological parameter changes in *Cyprinus carpio* (L.) experimentally infected with *Aphanomyces invadans* fungus. *Journal of Applied Ichthyology*, 21(5), 410–413.
- Harikrishnan, R., Balasundaram, C. and Heo, M.S., 2011.** Impact of



- plant products on innate and adaptive immune system of cultured finfish and shellfish. *Aquaculture*, 317, 1–15.
- Khan, M.R., Rahman, M.M., Shamsuddin, M., Islam, M.R. and Rahman, M., 2011.** Present status of aqua-drugs and chemicals in Mymensingh district. *Journal of the Bangladesh Society for Agricultural Science and Technology*, 8(1-2), 169–174.
- Kjer, J., Wray, V., Edrada-Ebel, R., Ebel, R., Pretsch, A., Lin, W. and Proksch, P., 2009.** Xanalteric acids I and II and related phenolic compounds from an endophytic *Alternaria* sp. isolated from the mangrove plant *Sonneratia alba*. *Journal of Natural Products*, 72, 2053–2057.
- Kohner, P.C., Rosenblatt, J.E. and Cockerill, F.R., 1994.** Comparison of agar dilution, broth dilution and disk diffusion testing of ampicillin against *Haemophilus* spp. by using in-house and commercially prepared media. *Journal of Clinical Microbiology*, 32, 1594–96.
- Kumar, V., Suvra, R. and Debtanu, B., 2015.** Effect of *Mikania cordata* (Burm) B.L. Robins on non-specific immune response of *Catla catla* (Hamilton, 1822) against *Aphanomyces invadans*. *Fish Technology*, 52, 20–25.
- OIE, 2015.** Manual of diagnostic tests for aquatic animals. Available: <http://www.oie.int/fileadmin/Home/e> ng/Health\_standards/aahm/current/c\_hapitre\_aphanomyces\_invadans.pdf.
- Ortuño, J., Cuesta, A., Rodriguez, A., Esteban, M.A. and Meseguer, J., 2002.** Oral administration of yeast, *Saccharomyces cerevisiae*, enhances the cellular innate immune response of gilthead seabream (*Sparus aurata* L.). *Veterinary Immunology and Immunopathology*, 85, 41–50.
- Pandey, G., Sharma, M. and Mandloi, A.K., 2012.** Medicinal plants useful in fish diseases. *Plant Archives*, 12, 1–4.
- Saad, S., Taher, M., Susanti, D. Qaralleh, H. and Awang AFIB., 2012.** *In vitro* antimicrobial activity of mangrove plant *Sonneratia alba*. *Asian Pacific Journal of Tropical Biomedicine*, 2, 427–429.
- Sivaram, V., Babu, M.M., Immanuel, G., Murugadass, S., Citarasu, T. and Marian, M.P., 2004.** Growth and immune response of juvenile greasy groupers (*Epinephelus tauvina*) fed with herbal antibacterial active principle supplemented diets against *Vibrio harveyi* infections. *Aquaculture*, 237, 9–20.
- Sun, J., 2007.** D-Limonene: safety and clinical applications. *Alternative Medicine Review*, 12(3), 259–264.
- Uthayakumar, V., Chandirasekar, R., Sreedevi, P.R., Senthilkumar, D., Jayakumar, R. and Ramasubramanian, V., 2014.** Immunostimulatory effect and disease resistance induced by *Lawsonia inermis* against *Aphanomyces invadans* in striped

murrels (*Channa striatus*). *Malaya Journal of Biosciences*, 1, 231–241.

**Valladão, G.M.R., Gallani, S.U. and Pilarski, F., 2015.** Phytotherapy as an alternative for treating fish disease. *Journal of Veterinary Pharmacology Therapeutics*, 38, 417–428.

**Yogeshwari, G., Jagruthi, C., Anbazahan, S.M., Mari, L.S.S., Selvanathan, J., Arockiaraj, J., Dhayanithi, N.B., Ajithkumar, T.T., Balasundaram, C. and Ramasamy, H., 2015.** Herbal supplementation diet on immune response in *Labeo rohita* against *Aphanomyces invadans*. *Aquaculture*, 437, 351–359.