

Use of otolithic morphometrics and ultrastructure for comparing between three goatfish species (family: Mullidae) from the northern Red Sea, Hurghada, Egypt

Osman A.G.M.^{1*}; Farrag M.M.¹; Mehanna S.F.²; Osman Y.A.³

Received: December 2017

Accepted: July 2018

Abstract

This work highlights the role of otolithic morphometrics, shape indices and ultrastructure in the identification of three Mullidae species from the Red Sea. Differences in otolithic measurements were detectable in all three-goatfish species. The statistical analysis of otolithic morphometric parameters showed that otolithic measurements are good indicators of fish size. For all three species, the correlation between fish length and different otolithic variables was statistically significant. The coefficient of determination (r^2) ranged from 0.83 to 0.92 in the three species, being higher for *M. vanicolensis* in all cases. The otolith area of *M. vanicolensis* was most strongly related to fish length, with a high correlation between otolith area and fish length ($r^2=0.92$) being observed in this species. The mean values of the six examined shape indices of the otoliths were considerably different among the three species, and the high degree of differentiation of these indices among species makes them useful for other researchers who wish to use otoliths in fish identification and classification. Remarkable variations in the morphological characteristics of fish otoliths were recorded among the studied species via scanning electron microscopy, including variations in the ornamentation of the ostium, cauda, and column of the otolith. These differences in otolithic characteristics and morphology might be useful for fisheries, biologists, archaeologists and geologists in discriminating *Mulloidichthys flavolineatus*, *M. vanicolensis*, and *Parupeneus forsskali*. This work contributes to the bioecological knowledge regarding commercially important fishes and provides key information for studying the trophic ecology of fish-eating species and fishery management.

Keywords: Goatfish species, Otoliths, Morphometrics, Scanning electron microscope, Red Sea

1-Department of Zoology, Faculty of Science, Al-Azhar University (Assiut Branch), 71524 Assiut, Egypt.

2-National Institute of Oceanography and Fisheries, Suez, Egypt

3-National Institute of Oceanography and Fisheries, Hurghada, Egypt

*Corresponding author's Email: agosman@azhar.edu.eg

Introduction

Family Mullidae (Goatfishes) is widely distributed globally, especially in the Pacific and Indo-Pacific regions, representing important food-chain components in coastal ecosystems (Pavlov *et al.*, 2015). The family consists of six genera and approximately 62 species (Nelson, 2006). Goatfishes are one of the most economically and commercially important groups inhabiting the northern Egyptian Red Sea sector (Sabrah, 2015). Three common goatfish genera are present along the Egyptian Red Sea coast: *Mulloidichthys*, *Parupeneus* and *Upeneus* (Kuronuma and Abe, 1986).

Otoliths are calcareous structures found in the inner ear of fishes (Campana, 2004). There are three pairs of otoliths (sagittae, asterisci and lapilli) in the otic sacs (Popper and Lu, 2000), and they function as mechanoreceptors that are involved in balance and hearing (Popper *et al.*, 2005). In most species, the sagittal otoliths (the largest of the three otolith pairs) are most commonly used to estimate age and growth, movement and habitat, population structure, and trophic ecology (Campana and Casselman, 1993; Rooker *et al.*, 2008). Otoliths are often collected in routine fisheries investigations for age determination as well as from predator stomachs (Nielsen and Andersen, 2001), allowing precise information on the length, weight, age, and quantity of individual fish prey to be obtained in many cases. Furthermore, otoliths are generally considered to serve as

taxonomical and biological archives, as they reflect species' growth and development. Most importantly for the current study, otolithic morphology is species specific (Popper *et al.*, 2005). Otolithic morphology ranges from ellipsoidal to more complex shapes, with protrusions and invaginations (Campana, 2004), and has been used in many studies for species identification (Tuset *et al.*, 2006; Lord *et al.*, 2012; Bani *et al.*, 2013; Sadighzadeh *et al.*, 2014). However, for accurate species identification, a reference collection of otoliths from known species must be produced to generate sufficient comparative material to produce accurate descriptions of morphological characteristics.

To our knowledge, there is no available information concerning the identification of goatfish from the Egyptian Red Sea using otolithic morphometrics and/or ultrastructure. Therefore, this study had three objectives: 1) to determine the possibility of readily discriminating common goatfish species based on the shape and ultrastructure of their otoliths; 2) to quantify otolith length, width, and height relationships with respect to fish length; and 3) to identify characteristics and ultrastructural features allowing otoliths from common Mullidae species to be distinguished from each other.

Materials and methods

Sampling

Three goatfish species (Family: Mullidae) were collected (n=275) from the northern part of the Red Sea

Province at Hurghada (latitudes $27^{\circ} 10' \text{ N}$ – $27^{\circ} 33' \text{ N}$ and longitudes $33^{\circ} 70' \text{ E}$ – $33^{\circ} 85' \text{ E}$) (Fig. 1) using artisanal fishing gear (gill net and trammel net) from September 2015 to August 2016. These species included *Mulloidichthys flavolineatus* (n=152), *Mulloidichthys vanicolensis* (n=21), and *Parupeneus forsskali* (n=102). For all individuals,

total length (TL) was measured to the nearest millimeter and weight (W) to the nearest gram. The sagittal otoliths (Fig. 2) of the fish were removed, washed and dried and then stored in plastic vials until being examined and photographed.




Figure 1: Map showing the study area at Hurghada, Egypt.



Figure 2: Proximal view of the left sagitta of common goatfish species from the Red Sea, Egypt.

Morphometric and shape analysis of otoliths

For morphometric analysis, the otoliths from the left side of the fish were oriented with the inner side (sulcus acusticus) upwards and the rostrum to the right, for digitization using a stereomicroscope linked to a digital camera (Optica 2.1) (Fig. 2). Then, otolith length (OL, mm), otolith width (OWid, mm), otolith area (OA, mm²), and otolith perimeter (OP, mm) were measured using ImageJ software. Otolith weight (OW, mg) was measured using an AS220 kL⁻¹ model balance. The relationships between fish total length (TL) and otolith variables were estimated using the power equation

(otolith variables = aTL^b), followed by log transformation to estimate a and b via simple linear regression, in which a is the angular coefficient characterizing the otolith's growth rate and b is a constant specific to the species.

To describe otolith shape, six dimensionless shape factors (aspect ratio (AS), compactness (CO), form factor (FF), rectangularity (RE), roundness (RO), and ellipticity (EL)) were obtained by combining size parameters (Russ, 1990; Tuset *et al.*, 2003b; Pinkerton, 2015) (Table 1).

Table 1: Otoliths shape indices.

Index	Formula
Aspect ratio (AS)	(OWid / OL)
Compactness (CO)	(OP ² / OA)
Form factor (FF)	$4 \pi OA / OP^2$
Rectangularity (RE)	OA / (L * l)
Roundness (RO)	$4 OA / \pi L^2$
Ellipticity (EL)	(OL - OWid) / (OL + OWid)

OL, otolith length, OWid-otolith width, OA- otolith area, OP- otolith perimeter.

Scanning electron microscopy

The otoliths were physically cleaned by carefully removing any adhering tissue and debris without damaging the scale surface. Then, they were immersed in a solution of sodium hypochlorite for several minutes to soften adhering tissues for further cleaning. For scanning electron microscopy (SEM) examination, the otoliths were fixed on a specimen holder using sticker tape and coated with a 30-nm layer of gold. Electron micrographs were produced on a GAOL, GSMS 400 LV scanning electron microscope in back-scattering

mode and on a Stereo Scan Cambridge Mark 2A (15 KV) at the Assiut University Electron Microscope Center, Assiut, Egypt. The morphological descriptions of the otoliths were based on the terminology proposed by Tuset *et al.*, (2008) (Fig. 2).

Statistical analysis

A Kolmogorov-Smirnov test was used to check the normality of the data distributions and variance homogeneity. Statistical description of the weight, length and otolith size of goatfish from the Red Sea was conducted using SPSS.

A paired sample t-test was employed to compare the left and right otoliths (SPSS, Version 18). Summaries of the descriptive statistics for the otolith shape indices of common species of family Mullidae were performed using SPSS. A power equation was applied to estimate the interaction between fish length and otolithic measurements.

Results

The summarized results for the left and right otolithic measurements as well as

descriptive statistics (minimum, maximum, mean, standard error and standard deviation) and paired-t tests are shown in Table 2. An insignificant difference ($p>0.05$) in the weight, length, width, area, and perimeter of the otoliths was observed between the left and right sagittal pairs for each goatfish species (Table 2). Therefore, the left otoliths were selected for the recording of other measurements and statistics.

Table 2: Summary of descriptive statistics and paired t-test results for left and right sagitta otoliths of three goatfish (*Mulloidichthys flavolineatus*, *M. vanicolensis* and *Parupeneus forsskali*) from the Red Sea, Egypt.

	Min		Max		Mean		SE		SD		p value
	Right	Left	Right	Left	Right	Left	Right	Left	Right	Left	Right-Left
<i>M. flavolineatus</i>											
Weight	0.0013	0.0012	0.0086	0.0086	0.004	0.004	0.0001	0.0001	0.0012	0.0012	0.741
Length	2.38	2.4	4.77	4.76	3.62	3.61	0.03	0.03	0.37	0.37	0.687
Width	1.77	1.71	3.54	3.43	2.5	2.42	0.02	0.02	0.26	0.25	0.000
Area	2.85	2.54	11.12	10.28	6.3	5.88	0.1	0.1	1.28	1.2	0.001
Perimeter	3.21	6.55	16.21	14.98	11.1	10.58	0.15	0.12	1.8	1.45	0.002
<i>M. Vanicolensis</i>											
Weight	0.0035	0.0035	0.0146	0.0146	0.0077	0.0078	0.0007	0.0007	0.0033	0.0034	0.386
Length	3.54	3.59	4.89	4.87	4.37	4.35	0.1	0.09	0.44	0.4	0.219
Width	2.39	2.32	3.81	3.68	3.06	2.99	0.09	0.09	0.42	0.41	0.01
Area	6.1	5.97	12.13	10.85	9.16	8.75	0.42	0.38	1.9	1.73	0.000
Perimeter	11.11	11.03	17.58	18	14.84	14.49	0.48	0.43	2.21	1.96	0.127
<i>P. forsskali</i>											
Weight	0.0012	0.0013	0.0054	0.0055	0.0025	0.0025	0.0001	0.0001	0.0008	0.0009	0.827
Length	2.51	2.62	4.16	4.11	3.1	3.11	0.03	0.03	0.33	0.31	0.519
Width	1.61	1.58	2.88	2.84	2	1.96	0.02	0.02	0.24	0.24	0.000
Area	2.03	2.83	6.64	6.62	3.8	4.14	0.09	0.08	0.95	0.81	0.001
Perimeter	3.51	7.14	11.73	12.15	7.64	8.94	0.13	0.1	1.35	0.98	00.000

The morphometric measurements of the fish and otoliths are shown in Table 3, with statistical descriptions of each species. Fish total length (TL) ranged from 110–334 mm, 174–287 and 144–271 for *M. flavolineatus*, *M. vanicolensis* and *P. forsskali*, respectively. The weight of the fish ranged from 11.85–453.26 g, 52.8–297.9 g and 32.13–237.37 g for *M. flavolineatus*, *M. vanicolensis* and *P.*

forsskali, respectively. Individual maximum otolith length (OL) ranged from 4.1 mm (*P. forsskali*) to 4.8 mm (*M. vanicolensis*). The otoliths of *M. vanicolensis* were the largest, reaching up to 4.8 mm. The otoliths of this species also exhibited the greatest perimeter (ranging from 11.03 to 18). The otoliths of *P. forsskali* presented the smallest values for all examined parameters. Overall individual otolith

weight (OW) ranged from 0.005 (*P. forsskali*) to 0.01 g (*M. vanicolensis*). *M. vanicolensis* displayed the heaviest otoliths, with an average OW of 0.0077 ± 0.0007 g, as well as the widest otoliths and the greatest surface area.

The overall otolithic measurements of *M. flavolineatus* occupied an intermediate position between those of *M. vanicolensis* and *P. forsskali*.

Table 3: Statistical Descriptions of fish weight (FW), fish total length (TL), otolith weight (OW), otolith length (OL), otolith width (OWid), otolith perimeter (OP), and otolith area (OA) of three goatfish (*Mulloidichthys flavolineatus*, *M. vanicolensis* and *Parupeneus forsskali*) from the Red Sea, Egypt.

Species	Variables	Minimum	Maximum	Mean	SE	SD
<i>M. flavolineatus</i>	FW	11.85	453.26	96.64	5.23	64.47
	TL	110	334	207.4	2.75	33.92
	OW	0.0012	0.0086	0.0039	0.000098	0.00122
	OL	2.4	4.76	3.61	0.03	0.37
	OWid	1.71	3.43	2.42	0.02	0.25
	OP	6.55	14.98	10.58	0.12	1.45
	OA	3.54	10.28	5.92	0.09	1.12
	No.			152		
<i>M. vanicolensis</i>	FW	52.8	297.9	190.14	19.69	90.22
	TL	174	287	246.91	9.06	41.50
	OW	0.0035	0.015	0.008	0.00078	0.0034
	OL	3.59	4.87	4.35	0.089	0.40
	OWid	2.32	3.68	2.99	0.09	0.41
	OP	11.03	18	14.49	0.43	1.96
	OA	5.97	10.85	8.7493	0.38	1.73
	No.			21		
<i>P. forsskali</i>	FW	32.13	237.37	76.94	3.94	39.75
	TL	144	271	185.30	2.63	26.53
	OW	0.0013	0.006	0.0025	0.00009	0.0009
	OL	2.62	4.11	3.11	0.031	0.31
	OWid	1.58	2.84	1.96	0.024	0.24
	OP	7.14	12.15	8.94	0.097	0.98
	OA	2.83	6.62	4.12	0.08	0.81
	No.			102		

According to the power equation (otolith variables = aTL^b), all morphometric measurements of the three selected species' otoliths exhibited a good relationship with fish total length (Table 4). The maximum

length, weight, width, area and perimeter of the otoliths were linearly related to total fish length for the studied fish species. All regressions were highly significant, and the analysis of otolith morphometric parameters

versus TL indicated that the regression models explained nearly all of the variance that best fit TL. The coefficients of determination (r^2) ranged from 0.78 to 0.81 in *M. flavolineatus*, from 0.83 to 0.92 in *M. vanicolensis*, and from 0.78 to 0.85 in *P. forsskali*, being higher for *M. vanicolensis* in all cases. The otolith area of *M.*

vanicolensis was highly correlated with fish length ($r^2=0.92$) and was most strongly related to fish length. The statistical analysis of otolith morphometric parameters showed that the otolithic measurements were good indicators of fish size.

Table 4: Power relationships between fish length and otolith variables for the three-goatfish species (*Mulloidichthys flavolineatus*, *M. vanicolensis* and *Parupeneus forsskali*) from the Red Sea. OW, otolith weight; TL, total length; OL, otolith length; OWid, otolith width; OA, otolith area; OP, otolith perimeter.

<i>Mulloidichthys flavolineatus</i>		<i>Mulloidichthys vanicolensis</i>		<i>Parupeneus forsskali</i>	
OL=0.1501TL ^{0.597}	$r^2=0.84$	OL=0.299TL ^{0.486}	$r^2=0.85$	OL=0.0996TL ^{0.659}	$r^2=0.85$
OWid=0.1058TL ^{0.58^A}	$r^2=0.80$	OWid=0.058TL ^{0.716}	$r^2=0.83$	OWid=0.0346TL ^{0.773}	$r^2=0.83$
OW=8E-07TL ^{1.602}	$r^2=0.82$	OW=1E-08TL ^{2.431}	$r^2=0.87$	OW=2E-08TL ^{2.237}	$r^2=0.83$
OA=0.0237TL ^{1.033}	$r^2=0.78$	OA=0.0168TL ^{1.134}	$r^2=0.92$	OA=0.00647TL ^{1.236}	$r^2=0.83$
OP=0.1646TL ^{0.780}	$r^2=0.78$	OP=0.8502TL ^{0.717}	$r^2=0.85$	OP=0.2275TL ^{0.703}	$r^2=0.78$

The mean values of the six examined shape indices of the three species' otoliths are shown in Table 5. They were considerably different among the three species. The highest mean values of AS, CO, and RO were recorded in *M. vanicolensis*. The lowest mean values of AS and RO were observed in *P. forsskali*, whereas the lowest value of COM was recorded in *M. flavolineatus*. The highest FF was recorded in *M. flavolineatus* and the lowest in *M. vanicolensis*. All three species exhibited nearly the same value of RE. The highest ellipticity was observed in *P. forsskali* (Table 5). The Kolmogorov-Smirnov Z test confirmed a normal distribution ($p>0.05$; non-significant) for nearly all measurements (excluding the weight and width of the otoliths of *M. flavilineatus*). According to the present results, the aspect ratio

gradually increased as fish otolith length increased.

The relationship between otolith length and the aspect ratio was close to 1; thus, the otoliths of the three species were elongated (Fig. 3). Roundness and ellipticity values increased as otolith length increased. Form factor and rectangularity values decreased as otolith length decreased. Pearson's correlation coefficients between the examined shape factors and maximal otolith length are presented in Table 6, and all of the variables for *M. flavolineatus* were significantly correlated with otolith length. Nearly all of the variables for *M. vanicolensis* were also significantly correlated with otolith length (excluding AS and EL). For *M. vanicolensis*, only FF and RO were significantly correlated with otolith length.

Table 5: Minimum, maximum, mean, standard error, and standard deviation of otoliths six shape descriptors (AS, CO, FF, RO, RE, and EL) of three goatfish species (*Mulloidichthys flavolineatus*, *M. vanicolensis* and *Parupeneus forsskali*) from the Red Sea, Egypt.

	Variables	N	Minimum	Maximum	Mean	SE	SD
<i>M. flavolineatus</i>	AS	152	0.58	0.77	0.67	0.003	0.036
	CO	152	11.71	26.04	19.03	0.212	2.616
	FF	152	0.48	1.07	0.67	0.008	0.100
	RO	152	25.98	286.78	103.09	3.612	44.53
	RE	152	0.57	0.90	0.67	0.004	0.049
	EL	152	0.13	0.26	0.19	0.002	0.025
<i>M. vanicolensis</i>	AS	21	0.62	0.80	0.69	0.011	0.049
	CO	21	20.38	32.58	24.18	0.579	2.653
	FF	21	0.39	0.62	0.52	0.011	0.052
	RO	21	98.03	314.34	219.27	16.715	76.599
	RE	21	0.60	0.72	0.67	0.006	0.029
	EL	21	0.11	0.24	0.19	0.007	0.034
<i>P. forsskali</i>	AS	102	0.56	0.72	0.63	0.003	0.035
	CO	102	16.13	24.36	19.52	0.169	1.704
	FF	102	0.52	0.78	0.65	0.006	0.057
	RO	102	27.41	142.39	53.31	2.240	22.57
	RE	102	0.57	0.79	0.68	0.0042	0.042
	EL	102	0.16	0.28	0.23	0.003	0.026




Figure 3: Relationship between the shape indices (Aspect ratio, Compactness, Form factor, Rectangularity, Roundness and Ellipticity) of the left otolith and otolith length in three goatfish species (*Mulloidichthys flavolineatus*, *M. vanicolensis* and *Parupeneus forsskali*) from the Red Sea, Egypt.

Table 6: Correlation coefficients of Pearson r between shape indices of the left otoliths and otoliths length in the three-goatfish species (*Mulloidichthys flavolineatus*, *M. vanicolensis* and *Parupeneus forsskali*) from the Red Sea, Egypt.

Index	<i>M. flavolineatus</i>		<i>M. vanicolensis</i>		<i>M. forsskali</i>	
	r	p	r	p	r	p
AS	-0.243**	0.003	0.401	0.072	0.098	0.325
Com	0.513**	0.000	0.433	0.50	0.239*	0.015
FF	-0.517**	0.000	-0.480*	0.028	-0.235*	0.018
RO	0.953**	0.000	0.992**	0.000	0.972**	0.000
RE	-0.402**	0.000	-0.302	0.183	-0.432**	0.000
ELL	0.242**	0.003	-0.409	0.066	-0.095	0.341

Correlation is significant at the 0.01 level

Scanning electron microscopy analysis of goatfish otolith topography

Images of the otoliths of the three examined goatfish species are presented in Fig. 4. The sagittal shape was oval in *M. flavolineatus* (Fig. 4 A and D) and oblong in *M. vanicolensis* (Fig. 4 B and E) and *P. forsskali* (Fig. 4 C and F). The rostrum occupied the anterior margin in all three goatfish species. The posterior margin was rounded in all three species but was also equipped with projections in *M. vanicolensis* and *P. forsskali* (Fig. 4 B, C). All three sagittae exhibited scalloped or crenulated ventral and dorsal rims. The dorsal rim of *M. flavolineatus* was

lobed and sinuate (Fig. 4 A). In the case of *M. vanicolensis* and *P. forsskali*, the dorsal rim was sinuate and emarginated (Fig. 4 B, C). The ventral rim of *M. flavolineatus* presented regular lobes only in the most dorsal portion (Fig. 4 A). In *M. vanicolensis*, the ventral lobes occurred in a single row and were irregular, extending from the rostrum anteriorly to the post rostrum posteriorly (Fig. 4 B). *P. forsskali* exhibited many rows of irregular dense lobes covering the entire ventral rim, extending from the rostrum anteriorly to the post rostrum posteriorly (Fig. 4 C).




Figure 4: The scanning electron microscope photographs of the proximal (A, B, C) and distal (D, E, F) views of the left sagitta of the three-goatfish species from the Red Sea (A and D, *Mulloidichthys flavolineatus*; B and E, *Mulloidichthys vanicolensis*; C and F, *Parupeneus forsskali*).

The rostrum of *M. flavolineatus* was elongated, free of projections and did not exhibit an antirostrum (Fig. 4 A). In contrast, the rostra of *M. vanicolensis* and *P. forsskali* were nearly round, with projections and antirostra (Fig. 4 B, C). The proximal face of the sagittal otolith displayed a sulcus acusticus lined dorsally by crista superior and ventrally by crista inferior. The sulcus was ostial and homosulcoid. The crista superior of *M. flavolineatus* (Fig. 4 A) was smooth, while those of *M. vanicolensis* and *P. forsskali* were rough and crushed (Fig. 4 B, C). The sagittae presented depressions in dorsal and ventral areas, oriented parallel to the sulcus acusticus. These depressions were shallow in the case of *M. flavolineatus* (Fig. 4 A) and deep in the case of *M. vanicolensis* and *P. forsskali* (Fig. 4 B, C). The sulcus acusticus was open anteriorly and posteriorly and divided into the ostium, column, and cauda (Fig. 4). The ostium was provided with a pore in all three species. This pore was small and rounded in *M. flavolineatus* (Fig. 4 A), while it was large and irregularly shaped in both *M. vanicolensis* and *P. forsskali* (Fig. 4 B, C). The cauda of *M. flavolineatus* was funnel shaped (Fig. 4 A), while those of both *M. vanicolensis* and *P. forsskali* were crucible shaped and were initially straight, with curves then occurring distally toward the inner rim (Fig. 4 B, C). The column of the *M. flavolineatus* was long, narrow, and deep, with straight crista superior and inferior (Fig. 4 A). The columns of *M.*

vanicolensis and *P. forsskali* were short and shallow, with curved crista superior and inferior (Fig. 4 B, C).

Ornamentations in the sulcus acusticus

At low magnification, the surface appeared relatively smooth, but a remarkable difference was observed in the ornamentation of the ostium, cauda, and column.

Ostium

Two types of ornamentation were observed on the ostium of each species (Fig. 5). Fine ornamentation was observed toward the edge of the ostium, transforming into a more coarsely textured surface or a coarser crystalline structure toward the inner side of the ostium in both *M. flavolineatus* and *M. vanicolensis* (Fig. 5 A, B). In contrast, the ostium of *P. forsskali* exhibited nearly the same textured surface on the edge and the inner side (Fig. 5 C). In *M. flavolineatus*, the ornamentation of the edge of the ostium had a sand-like appearance, whereas that of the inner side of the ostium had the appearance of rectangular stones (Fig. 5 A). The ostium of *M. vanicolensis* presented fine sandy ornamentation externally and projecting lath-like crystals internally (Fig. 5 B). Both the outer and the inner projections of *P. forsskali* consisted of projecting lath-like crystals (Fig. 5 C).




Figure 5: Scanning electron microscope photographs of the ostium of the left sagitta of the three-goatfish species from the Red Sea (A, *Mulloidichthys flavolineatus*; B, *Mulloidichthys vanicolensis*; C, *Parupeneus forsskali*).

Cauda

Each species exhibited only one type of ornamentation on the cauda. *M. flavolineatus* presented sand-like crystals (Fig. 6 A), whereas *M.*

vanicolensis and *P. forsskali* presented the same type of ornamentation, in the form of projecting lath-like crystals (Fig. 6 B, C).




Figure 6: Scanning Electron Microscope photographs of the cauda of the left sagitta of the three-goatfish species from the Red Sea (A, *Mulloidichthys flavolineatus*; B, *Mulloidichthys vanicolensis*; C, *Parupeneus forsskali*).

Column

The column's ornamentation differed among the three species. The ornamentation of *M. flavolineatus* consisted of triangular and square stone-like shapes (Fig. 7 A), while that

of *M. vanicolensis* took the form of irregular projections. In the case of *P. forsskali*, the ornamentation was rectangular and stone-like (Fig. 7 B, C).




Figure 7: Scanning Electron Microscope photographs of the column of the left sagitta of the three-goatfish species from the Red Sea (A, *Mulloidichthys flavolineatus*; B, *Mulloidichthys vanicolensis*; C, *Parupeneus forsskali*).

Discussion

Goatfish represent one of the most economically important fisheries in Egyptian waters, in both the Red and Mediterranean Seas. The fishery status of any fish stock needs to be identified correctly. The fishes of this group (family Mullidae) are teleost fishes comprising many species with some shape similarities; hence, these species need to be clearly distinguished. Many studies have been performed on the

fisheries and biology of goatfishes in Egypt (Golani and Ritte, 1999; Sabrah, 2015), with little attention being paid to goatfish taxonomy and phylogeny. Recently, stock identification was achieved using otolithic features (Lombarte *et al.*, 2006; Mehanna *et al.*, 2016). The morphological characteristics of fish otoliths are the most widely used tools in species identification and comparative taxonomy of fishes, due to the large

size and inter-specific variability of fish otoliths (Battaglia *et al.*, 2010). To our knowledge, the otolithic shape indices and measurements of goatfish from the Red Sea have never been estimated previously. Thus, the present study may be the first to discriminate goatfish species using otolithic indices and their relationships with fish size and otolithic ultrastructure. In the present work, sagittal otoliths were employed because they are easily accessible structures and exhibit a distinctive degree of interspecific variation in their form, weight, and growth, in accordance with Nolf (1985). The morphometric parameters of sagittal otoliths have been used in earlier studies to identify species in other groups of teleosts (Tuset *et al.*, 2003a; Ponton, 2006; Short *et al.*, 2006; Tuset *et al.*, 2006). Our results suggest that differences in otolithic measurements are detectable in all three goatfish species examined in this work.

The results of the present study indicate no significant morphometric differences between the measurements of the left and right otoliths. Remarkable differences are not usually observed between left and right otolithic measurements (for many fish species) (Morat *et al.*, 2008; Jawad *et al.*, 2011a; Yilmaz *et al.*, 2014). This finding is in agreement with previous work (Valinassab *et al.*, 2012) on clupeids from the Persian Gulf. In another previous study, Hunt, (1992) studied eight species from the Atlantic Ocean, and no statistically significant difference was observed between the left and right otoliths. Similarly,

Harvey *et al.*, (2000) observed no significant differences between the left and right otoliths of 63 species collected from the Eastern North Pacific Ocean. However, significant differences between left and right otolithic variables have been recorded for some other fish species (Mérigot *et al.*, 2007).

The relationships between the morphological parameters (length, width, perimeter, and area) of the otoliths and fish body length (TL) in the three goatfish species were examined using power equations. The relationships between otolith size and body length (TL) have also been estimated in other fish species using linear functions (Morat *et al.*, 2008; Pavlov, 2016). Previous studies have focused mainly on the relationship between otolith length and width and fish size (Jawad *et al.*, 2011b). This work supplies additional information (compared with other similar studies) by considering six otolithic measurements (OL, OH, OW, OWid, OA, and OP). Thus, our findings are more reliable than those relying on a single equation, since the tip of the otolithic rostrum may be damaged, making it impossible to measure OL or OW (Jawad *et al.*, 2011b; Yilmaz *et al.*, 2014), influencing the reliability of subsequent calculations. Presenting the six models ($OL=aTL^b$, $OH=aTL^b$, $OW=aTL^b$, $OWid=aTL^b$, $OA=aTL^b$ and $OP=aTL^b$) for each species helps mitigate this potential problem. It is appropriate to use the functions indicated in this paper within the range of the fish size examined in the present

work. Further studies on fish size–otolithic variable relationships involving a larger sample size, a wider range of fish lengths, and different growth phases would help to support the results presented herein. The ability to identify species using morphometric parameters of the sagittal otoliths has been demonstrated for other groups of teleosts (Tuset *et al.*, 2003b; Ponton, 2006; Short *et al.*, 2006; Tuset *et al.*, 2006). The differences in the morphological characteristics of the sagittal otoliths among these species may be associated with the size of the skull (Bani *et al.*, 2013). Although relative skull size was not recorded in this study, personal observations indicate that the greatest head length belongs to *M. vanicolensis*, which also exhibits the longest otoliths.

The power relationship between sagittal otoliths and fish length was described by a negative allometric relationship (b less than 3). Similar results were previously recorded in the freckled goatfish, *Upeneus tragula* (Pavlov *et al.*, 2015). Several studies have shown a linear relationship between body length and otolith length in fish (Harvey *et al.*, 2000; Fossen *et al.*, 2003; Lychakov *et al.*, 2006). Linear otolith growth might represent a common relationship between otoliths and body length in the juvenile stage (Huang and Chiu, 1997). In the present study, power equations were applied and showed a strong correlation between the variables, indicating that otolithic measurements are good indicators of fish size. For all three species, the correlation between fish

length and different otolithic variables was statistically significant (0.005), with otolith length showing the highest (0.70–0.96) Pearson's correlation coefficient, compromised by high variability in otolith shape. This result could be explained by the fact that otolithic length is most sensitive to variations in the growth rate and most closely related to changes in fish metabolism (Pawson, 1990; Flecher, 1991). In the present study, the coefficient of determination (r^2) ranged from 0.83 to 0.92 in the three species, being higher for *M. vanicolensis* in all cases. The otolith area of *M. vanicolensis* was most strongly related to fish length, with a high correlation between otolith area and fish length ($r^2=0.92$) being observed in this species. Concerning findings of this study, the observed variability of otolithic shape encourages further research to verify the potential role of otolithic morphometric measurements in fish identification.

Otolithic shape indices such as the aspect ratio, compactness or circularity, form factor, roundness, rectangularity, and ellipticity have been described for many fish species (Tuset *et al.*, 2003b; Pavlov *et al.*, 2015), confirming identifications performed based on morphometric variables (Tuset *et al.*, 2003a; Tuset *et al.*, 2003b). The results of this study show that the shape indices differ significantly in the analyzed species. Such differences were observed for almost all shape indices of the three selected goatfish species. Among the shape indices, AS, COM, RO, and FF were found to be more

efficient than other factors in the studied species. The highest mean values of AS, COM, and RO were recorded in *M. vanicolensis*, whereas the lowest mean values of AS and RO were found in *P. forsskali*, although the lowest value of COM was recorded in *M. flavolineatus*. Thus, in the last species, otolith shape is more circular than in *M. vanicolensis*, indicating that all of the otoliths of *M. flavolineatus* grow equally in terms of length and height. The numeric values of the FF show that the sagittal shape is geometrically irregular in the three species. The highest FF was recorded in *M. flavolineatus*, and the lowest was observed in *M. vanicolensis*, which presented the most irregular otolithic shape among the studied species. The shape analyses revealed rectangular shapes in the otoliths of the three species, as they exhibited nearly the same

Although the study of shape indices is complicated, it can confirm identifications made based on the differentiation of fish species (Špiranec and Banek Zorica, 2010; Sadighzadeh *et al.*, 2012) and populations (Mérigot *et al.*, 2007; Duarte-Neto *et al.*, 2008; Pavlov, 2016). In the present study, the shape indices were compromised by their high differentiation; thus, the results of this study will be useful for other researchers in verifying the role of the otolith in fish identification and classification. These findings were supported by previous results (Tuset *et al.*, 2008). The obtained ellipticity values show that the shape of the three species is irregular, with the lowest

value being found in *M. vanicolensis* and the highest in *P. forsskali*. According to the Kolmogorov-Smirnov Z test, a normal distribution was confirmed for nearly all otolithic measurements in each of the examined fish species. Otolith area, perimeter and shape indices have been suggested as an easier means of discriminating stocks compared with other methods (Bolles and Begg, 2000; Tuset *et al.*, 2003b). The aspect ratio and ellipticity were found to be directionally proportional to otolith length, while the form factor (FF) and roundness (RD) were inversely proportional to this parameter. Such similarity in otolithic shape indices may be derived from the fact that all of the studied goatfish species occupy the same ecological niche. Fish occupying the same ecological niche show similarities in otolithic shape variables (Parmentier *et al.*, 2001). The correlation between shape factors and otolith length were strong and significant in the case of *M. flavolineatus*. Excluding AS and EI, all other shape factors in *M. forsskali* exhibited significant correlations with otolith length. In the case of *M. vanicolensis*, only FF and RO were significantly correlated with otolith length. The results of this study show that the shape indices significantly differ from species to species, although they indicate a similar otolithic pattern. These results correlate with those of Tuset *et al.* (2008), who postulated that otoliths are the most widely employed tool for the discrimination of fish species because of their form, weight,

growth, consistency and chemical composition.

The current study also compared otolithic shape among the three goatfish species using SEM, to observe variations in otolithic morphology. In *M. flavolineatus*, the sagittal shape was found to be oval, with regular lobes on the dorsal and ventral rims. In *M. vanicolensis* and *P. forsskali*, the otolithic shape is oblong, with irregularly lobed dorsal and ventral rims. Remarkable variations in the morphological characteristics of fish otoliths were recorded between the studied species, including variations in the rostrum, sulcus acusticus, ostium, column and cauda. Remarkable variations were also recorded in the ornamentation of the ostium, cauda, and column. These differences in otolithic characteristics might be important to fishery biologists, archaeologists and geologists, who can use them to distinguish *M. flavolineatus*, *M. vanicolensis* and *P. forsskali*. This work contributes to the bioecological knowledge regarding commercially important fishes and provides key information for studying the trophic ecology of fish-eating species and fisheries management.

References

- Bani, A., Poursaeid, S. and Tuset, V. M., 2013.** Comparative morphology of the sagittal otolith in three species of south Caspian gobies. *Journal of Fish Biology*, 82(4), 1321-1332.
- Battaglia, P., Malara, D., Romeo, T. and Andaloro, F., 2010.** Relationships between otolith size and fish size in some mesopelagic and bathypelagic species from the Mediterranean Sea (Strait of Messina, Italy). *Scientia Marina*, 74(3), 605-612.
- Bolles, K.L. and Begg, G.A., 2000.** Distinction between silver hake (*Merluccius bilinearis*) stocks in US waters of the northwest Atlantic based on whole otolith morphometrics. *Fishery Bulletin*, 98(3), 451-451.
- Campana, S.E., 2004.** Photographic atlas of fish otoliths of the Northwest Atlantic Ocean Canadian special publication of fisheries and aquatic sciences No. 133. NRC Research Press, 284P.
- Campana, S.E. and Casselman, J.M., 1993.** Stock discrimination using otolith shape analysis. *Canadian Journal of Fisheries and Aquatic Sciences*, 50(5), 1062.
- Duarte-Neto, P., Lessa, R., Stosic, B. and Morize, E., 2008.** The use of sagittal otoliths in discriminating stocks of common dolphinfish (*Coryphaena hippurus*) off northeastern Brazil using multishape descriptors. *ICES Journal of Marine Science: Journal du Conseil*, 65(7), 1144-1152.
- Flecher, W., 1991.** A test of the relationship between otolith weight and age for the pilchard *Sardinops neopilchardus*. *Canadian Journal of Fisheries and Aquatic Sciences*, 48(1), 35-38.
- Fossen, I., Albert, O.T. and Nilssen, E.M., 2003.** Improving the precision of ageing assessments for long rough dab by using digitised pictures and

- otolith measurements. *Fisheries Research*, 60(1), 53-64.
- Golani, D. and Ritte, U., 1999.** Genetic relationship in goatfishes (Mullidae: Perciformes) of the Red Sea and the Mediterranean, with remarks on Suez Canal migrants. *Scientia Marina*, 63(2), 129-135.
- Harvey, J.T., Loughlin, T.R., Perez, M.A. and Oxman, D.S., 2000.** Relationship between fish size and otolith length for 63 species of fishes from the eastern North Pacific Ocean. NOAA Technical Report NMFS 150, 36P.
- Huang, W.B. and Chiu, T.S., 1997.** Daily increments in otoliths and growth equation of black porgy, *Acanthopagrus schlegeli*, larvae. *Acta Zoologica Taiwanica*, 8(2), 121-131.
- Hunt, J., 1992.** Morphological characteristics of otoliths for selected fish in the Northwest Atlantic. *Journal of Northwest Atlantic Fishery Science*, 13, 63-75.
- Jawad, L., Al-Mamry, J. and Al-Busaidi, H., 2011a.** Relationship between fish length and otolith length and width in the lutjanid fish, *Lutjanus bengalensis* (Lutjanidae) collected from Muscat City coast on the Sea of Oman. *Journal of Black Sea/Mediterranean Environment*, 17(2).
- Jawad, L., Ambuali, A., Al-Mamry, J. and Al-Busaidi, H., 2011b.** Relationships between fish length and otolith length, width and weight of the Indian mackerel *Rastrelliger kanagurta* (Cuvier, 1817) collected from the Sea of Oman. *Croatian Journal of Fisheries*, 69(2), 51-61.
- Kuronuma, K. and Abe, Y., 1986.** Fishes of the Arabian Gulf. Kuwait Institute for Scientific Research, Kuwait, 356P.
- Lombarte, A., Rufino, M.M. and Sánchez, P., 2006.** Statolith identification of Mediterranean Octopodidae, Sepiidae, Loliginidae, Ommastrephidae and Enoploteuthidae based on warp analyses. *Journal of the Marine Biological Association of the United Kingdom*, 86(4), 767-771.
- Lord, C., Morat, F., Lecomte-Finiger, R. and Keith, P., 2012.** Otolith shape analysis for three Sicyopterus (Teleostei: Gobioidae: Sicydiinae) species from New Caledonia and Vanuatu. *Environmental Biology of Fishes*, 93(2), 209-222.
- Lychakov, D., Rebane, Y., Lombarte, A., Fuiman, L. and Takabayashi, A., 2006.** Fish otolith asymmetry: morphometry and modeling. *Hearing Research*, 219(1), 1-11.
- Mehanna, S., Jawad, L., Ahmed, Y., Abu El-Regal, M. and Dawood, D., 2016.** Relationships between fish size and otolith measurements for *Chlorurus sordidus* (Forsskal, 1775) and *Hipposcarus harid* (Forsskal, 1775) from the Red Sea coast of Egypt. *Journal of Applied Ichthyology*, 32(2), 356-358.
- Mérigot, B., Letourneur, Y. and Lecomte-Finiger, R., 2007.** Characterization of local populations of the common sole *Solea solea* (Pisces, Soleidae) in the NW Mediterranean through otolith

- morphometrics and shape analysis. *Marine Biology*, 151(3), 997-1008.
- Morat, F., Banaru, D., M rigot, B., Batjakas, I. E., Betoulle, S., Vignon, M., Lecomtefiniger, R. and Letourneur, Y., 2008.** Relationships between fish length and otolith length for nine teleost fish species from the Mediterranean basin, Kerguelen Islands, and Pacific Ocean. *Cybiurn*, 32(3), 265-269.
- Nelson, J., 2006.** Fishes of the World. 4th eds. John Wiley & Sons Inc., New Jersey, 601P
- Nielsen, J.R. and Andersen, M., 2001.** Feeding habits and density patterns of Greenland cod, *Gadus ogac* (Richardson 1836), at West Greenland compared to those of the coexisting Atlantic cod, *Gadus morhua* L. *Journal of Northwest Atlantic Fishery Science*, 29, 1–22.
- Nolf, D., 1985.** Otolithi piscium: handbook of paleoichthyology, v. 10. Stuttgart, New York: Gustav Fischer, 145P.
- Parmentier, E., Vandewalle, P. and Lagard re, F., 2001.** Morpho-anatomy of the otic region in carapid fishes: eco-morphological study of their otoliths. *Journal of Fish Biology*, 58(4), 1046-1061.
- Pavlov, D., 2016.** Differentiation of three species of the genus *Upeneus* (Mullidae) based on otolith shape analysis. *Journal of Ichthyology*, 56(1), 37-51.
- Pavlov, D., Emel'yanova, N., Ha, V. T. and Thuan, L.T.B., 2015.** Otolith morphology, age, and growth of freckled goatfish *Upeneus tragula* (Mullidae) in the coastal zone of Vietnam. *Journal of Ichthyology*, 55(3), 363-372.
- Pawson, M., 1990.** Using otolith weight to age fish. *Journal of Fish Biology*, 36(4), 521-531.
- Pinkerton, A., 2015.** The spy of the rebellion: being a true history of the spy system of the united states army during the Late Rebellion. Create Space Independent Publishing Platform, 667P.
- Ponton, D., 2006.** Is geometric morphometrics efficient for comparing otolith shape of different fish species? *Journal of Morphology*, 267(6), 750-757.
- Popper, A.N. and Lu, Z., 2000.** Structure–function relationships in fish otolith organs. *Fisheries Research*, 46(1), 15-25.
- Popper, A.N., Ramcharitar, J. and Campana, S.E., 2005.** Why otoliths? Insights from inner ear physiology and fisheries biology. *Marine and Freshwater Research*, 56(5), 497-504.
- Rooker, J.R., Secor, D.H., DeMetrio, G., Kaufman, A.J., R os, A.B. and Ti cina, V., 2008.** Evidence of trans-Atlantic movement and natal homing of bluefin tuna from stable isotopes in otoliths. *Marine Ecology-Progress Series*, 368, 231-239.
- Russ, J., 1990.** Computer assisted microscopy. The measurement and analysis of image, pp. 71–79. New York: Plenum Press Corp.
- Sabrah, M.M., 2015.** Fisheries biology of the Red Sea goatfish *Parupeneus forsskali* (Fourmanoir & Gu z , 1976) from the northern Red Sea, Hurghada, Egypt. *The Egyptian*

- Journal of Aquatic Research*, 41(1), 111-117.
- Sadighzadeh, Z., Tuset, V. M., Valinassab, T., Dadpour, M.R. and Lombarte, A., 2012.** Comparison of different otolith shape descriptors and morphometrics for the identification of closely related species of *Lutjanus* spp. from the Persian Gulf. *Marine Biology Research*, 8(9), 802-814.
- Sadighzadeh, Z., Valinassab, T., Vosugi, G., Motallebi, A., Fatemi, M.R., Lombarte, A. and Tuset, V.M., 2014.** Use of otolith shape for stock identification of John's snapper, *Lutjanus johnii* (Pisces: Lutjanidae), from the Persian Gulf and the Oman Sea. *Fisheries Research*, 155, 59-63.
- Short, J.A., Gburski, C.M. and Kimura, D.K., 2006.** Using otolith morphometrics to separate small walleye Pollock *Theragra chalcogramma* from Arctic Cod *Boreogadus saida* in mixed samples. *Alaska Fishery Research Bulletin*, 12(1), 147-152.
- Špiranec, S. and Banek Zorica, M., 2010.** Information Literacy 2.0: hype or discourse refinement? *Journal of Documentation*, 66(1), 140-153.
- Tuset, V., Lombarte, A., Gonzalez, J., Pertusa, J. and Lorente, M., 2003a.** Comparative morphology of the sagittal otolith in *Serranus* spp. *Journal of Fish Biology*, 63(6), 1491-1504.
- Tuset, V., Lozano, I., Gonzalez, J., Pertusa, J. and García-Díaz, M., 2003b.** Shape indices to identify regional differences in otolith morphology of comber, *Serranus cabrilla* (L., 1758). *Journal of Applied Ichthyology*, 19(2), 88-93.
- Tuset, V.M., Lombarte, A. and Assis, C.A., 2008.** Otolith atlas for the western Mediterranean, north and central eastern Atlantic. *Scientia Marina*, 72(S1), 1-198.
- Tuset, V.M., Rosin, P.L. and Lombarte, A., 2006.** Sagittal otolith shape used in the identification of fishes of the genus *Serranus*. *Fisheries Research*, 81(2), 316-325.
- Valinassab, T., Seifabadi, J., Homauni, H. and Afraie, A., 2012.** Relationships between fish size and otolith morphometric in some clupeids from the Persian Gulf and Oman Sea. *Cybiurn Journal*, 36(4), 505-509.
- Yilmaz, S., Yazicioglu, O., Saygin, S. A. and Polat, N., 2014.** Relationships of otolith dimensions with body length of European perch, *Perca fluviatilis* L., 1758 From Lake Ladik, Turkey. *Pakistan Journal of Zoology*, 46(5), 1231-1238.