

Zooplankton of Çat Dam Lake (Malatya-Turkey) with a new record for Turkish rotifers *Lecane intrasinuata* (Olofsson, 1917)

Saler S.¹; Bulut H.^{1*}; Karakaya G.²

Received: February 2015

Accepted: February 2017

1-Firat University, Fisheries Faculty, 23119 Elazığ-Turkey

2-Elazığ Fisheries Research Station, Elazığ-Turkey

*Corresponding author's Email: hilalhaykir@gmail.com

Keywords: Rotifera, *Lecane intrasinuata*, New record, Turkey, Çat Dam Lake

Introduction

There are many studies on the zooplankton of Turkish lakes, many of which are about seasonal fluctuations, distribution and hydrological changes. In recent years many scientists focused their attention on the zooplankton of Turkish inland waters (Dumont and De Ridder, 1987; Segers *et al.*, 1992).

The family Lecanidae consists of one genus, *Lecane* Nitzsch, 1827 with about 200 species. This genus is the most common rotifer species inhabiting in various aquatic environments (Segers 1994, 1995, 2008). They are from Monogonont rotifers and are diagnosed by the retractile head, the structure of the foot and toes and by the trophy in the female (Segers, 1995). Lecanids have got a loricate body and the body is compressed dorso-ventrally. The dorso-

ventral plates are connected by a flexible membrane. The feet have got two primitive segments of which only the posterior is movable. Toes varied remarkably within individuals of the same species (Arora, 1965). Their bodies resemble each other which causes difficulties in identifying species (Segers *et al.*, 1992).

Materials and methods

Çat Dam Lake was established on Abdulvahap Stream for irrigation and is located 68 km south of Malatya. The Dam lake volume is 240 hm³ and lake area is 14 km². Sampling stations are shown in Fig. 1, and listed with sampling coordinates in Table 1.




Figure 1: Çat Dam Lake and the sampling stations.

Table 1: Sampling coordinates.

Stations	Coordinates
I	38° 4'40.22"N, 38° 18'38.11"E
II	38° 4'12.73"N, 38° 18'45.48"E
III	38° 4'70.93"N, 38° 18'21.27"E
IV	38° 3'55.05"N, 38° 17'46.73"E
V	38° 3'30.98"N, 38° 16'38.55"E
VI	38° 2'40.49"N, 38° 14'17.52"E

Plankton samples were collected using a standard Hydro-bios plankton net (55 µm mesh size) from Çat Dam Lake on 02.20.2013 from six stations. Zooplankton samples were fixed in 4 % formalin solution. Specimens were analysed under Leitz inverted microscope and identified under Nikon Eclipse 80i microscope and drawings were made using camera lucida.

Results and discussion

From Çat Dam Lake, 19 zooplankton species (15 species from Rotifera, 2 species from Cladocera and 2 species from Copepoda) were identified. The species *L. intrasinuata* was not reported in any study in Turkey until now

(Ustaoğlu *et al.*, 2012; Ustaoğlu, 2015). Therefore, this species is a new record for the Turkish rotifer fauna.

List of zooplankton species recorded in Çat Dam Lake was given below.

Phylum Rotifera Cuvier, 1817
Class Euroatoria De Ridder, 1957
Subclass Monogononta Plate, 1889
Superorder Pseudotocha Kutikova, 1970

Order Ploimia Hudson & Gosse, 1886
Family Brachionidae Ehrenberg, 1838

Genus Keratella Bory de St Vincent, 1822

Keratella cochlearis (Gosse, 1851)

K. tecta (Lauterborn, 1900)

K. quadrata (Müller, 1786)

Genus *Notholca* Gosse, 1886*Notholca squamula* (Müller, 1786)**Genus *Kellicottia* Ahlstrom, 1938***Kellicottia longispina* (Kellicott, 1879)**Family Euchlanidae Ehrenberg, 1838****Genus *Euchlanis* Ehrenberg, 1832***Euchlanis dilatata* Ehrenberg, 1832**Family Lecanidae Remane, 1933****Genus *Lecane* Nitzsch, 1827***Lecane luna* (Müller, 1776)*L. lunaris* (Ehrenberg, 1832)*L. intrasinuata* (Olofsson, 1917)**Family Trichocercidae Haring, 1913****Genus *Trichocerca* Lamarck, 1801***Trichocerca capucina* Wierzejski & Zacharias, 1893*Trichocerca similis* (Wierzeski, 1893)**Family Synchaetidae Hudson & Gosse, 1886***Polyarthra dolichoptera* (Idelson, 1925)**Family Asplanchnidae Eckstein, 1883****Genus *Asplanchna* Gosse, 1850***Asplanchna priodonta* Gosse, 1850**Superorder Gnesiotrocha Kutikova, 1970****Order Flosculariacea Haring, 1913****Family Conochililidae Haring, 1913****Genus *Conochilus* Ehrenberg, 1834***Conochilus dossuarius* (Hudson, 1885)**Family Filinidae Haring & Myers, 1926****Genus *Filinia* Bory de St. Vincent, 1824***Filinia terminalis* (Plate, 1886)**Phylum Arthropoda Latreille, 1829****Subphylum: Crustacea Brünnich, 1772****Subclass Phyllopoda Preuss, 1951****Order Diplostraca Gerstaecker, 1866****Suborder Cladocera Latreille, 1829****Family Daphniidae Sars, 1865****Genus *Daphnia* Müller, 1785***Daphnia longispina* Müller, 1875**Family Bosminidae Baird, 1845****Genus *Bosmina* Baird, 1845***Bosmina longirostris* (Müller, 1785)**Class Maxillopoda Dahl, 1956****Subclass Copepoda H. Milne-Edwards, 1840****Infraclass Neocopepoda Huys & Boxshall, 1991****Superorder Gymnoplea Giebesbrecht, 1882****Order Calanoida Sars, 1930****Family Diaptomidae Sars, 1903****Genus *Acanthodiaptomus* Kiefer, 1932***Acanthodiaptomus denticornis* (Wierzejski, 1887)**Superorder Podoplea Giesbrecht, 1882****Order Cyclopoida Sars, 1918****Family Cyclopoidae Sars, 1913****Subfamily Cyclopinae Kiefer, 1927****Genus *Cyclops* Müller, 1785***Cyclops vicinus* Uljanin, 1875

Distribution of zooplankton species among 6 stations in Çat Dam Lake is given in Table 2.

Table 2: Distribution of zooplankton species among 6 stations in Çat Dam Lake.

Species	Stations					
	I	II	III	IV	V	VI
	Rotifera					
<i>A. priodonta</i>	+	+			+	+
<i>C. dossuarius</i>	+					
<i>E. dilatata</i>			+			
<i>F. terminalis</i>				+		

Table 2 continued:

<i>K. longispina</i>	+	+		+	+	+
<i>K. cochlearis</i>	+	+	+	+	+	+
<i>K. quadrata</i>	+	+	+	+	+	+
<i>K. tecta</i>			+			
<i>L. luna</i>			+			
<i>L. lunaris</i>	+		+			
* <i>L. intrasinuata</i>			+		+	
<i>N. squamula</i>				+		
<i>P. dolichoptera</i>	+		+	+		
<i>T. capucina</i>			+			
<i>T. similis</i>				+		
Cladocera						
<i>B. longirostris</i>	+	+	+	+	+	
<i>D. longispina</i>		+	+			
Copepoda						
<i>A. denticornis</i>		+				
<i>C. vicinus</i>	+	+	+	+	+	+

Family Brachionidae was found as the most dominant group (with 5 species, *K. cochlearis*, *K. quadrata*, *K. tecta*, *K. longispina* and *N. squamula*). In some dam lakes like Göksu (Bekleyen, 2003), Cip (Saler and Şen, 2000), Birecik (Bozkurt and Sagat, 2008), Beyhan (Bulut and Saler 2014), Kalecik (Bulut and Saler 2013) Dam Lakes species from Brachionidae represented most of the species.

In Çat Dam Lake *K. cochlearis* and *K. quadrata* were the most abundant species in this study followed by *Lecane* spp. Among zooplankton groups Rotifera was observed in high number of individuals and species diversity. This profile was in accordance with zooplankton distribution of dam lakes in Turkey (Bekleyen, 2003; Kaya and Altındağ, 2007; Özdemir Mis *et al.*, 2009; Saler and İpek Alish, 2014; Saler *et al.*, 2014).

The first Turkish inland water zooplankton list was made by Emir (1996) and she reported 167 species. Ustaoglu (2004), reported 229 rotifer species from Turkish inland waters. In

the Tigris River Bekleyen *et al.* (2011) identified 34 new Rotifer species. Ustaoglu *et al.* (2012) recorded 341 rotifer taxa from Turkey. Ustaoglu (2015), updated the rotifer species number as 417. Among these rotifer species *L. intrasinuata* has not been mentioned (Fig. 2).




Figure 2: General view of *Lecane intrasinuata* A. Ventral B. Dorsal.

Family Lecanidae Remane, 1933

Lecane intrasinuata (Olofsson, 1917)

Syn: *L. ephestra* Harring, 1921

Syn: *L. mylacris* Harring & Myers, 1926 (Segers, 2007).

The features of *L. intrasinuata* were given below.

Differential diagnosis

L. intrasinuata is confused with *L. stichaea*. *L. intrasinuata* has got a smooth lorica and the width of its lorica is greater than that of *L. stichae*. Its ventral plate width is about two thirds of its length and the plate is slightly swollen. The foot is shorter than in *L. flexilis*, and the presence of pseudoclaws distinguishes the species from *L. haliclysta*, *L. stichoclysta* and *L. verecunda*.

Description

Lorica is stiff, dorsal plate is anteriorly narrower, and medially wider than the ventral plate. The plate is smooth. Head aperture margins are dorsally and ventrally slightly convex or straight, coincident, with small antero-lateral spines. Ventral plate is longer than its width. Lateral margins curved, irregularly undulate, with anterior notches. Toes are parallel-sided, bearing pseudoclaws.

Measurements

DPI. 52-135, DPw. 68-101, VPI. 56-125, VPw. 36-76, toe 15-26, claw 4-6.

Distribution

According to Segers (1995), it was found in the northern temperate zone only. He also reported a record from a pond near Alexandrowsk from Norway Fig. 2.

References

Arora, H.C., 1965. Studies on Indian rotifera- part VI. On a collection of Rotifera from Nagpur, India, with

four new species and a new variety. *Hydrobiologia*, 26, 444- 456.

Bekleyen, A., 2003. A taxonomical study on the zooplankton fauna of Göksu Dam Lake (Diyarbakır). *Turkish Journal of Zoology*, 27, 95–100.

Bekleyen, A., Gökot, B. and Varol, M., 2011. Thirty-four new records and the diversity of the Rotifera in the Turkish part of the Tigris River watershed, with remarks on biogeographically interesting taxa. *Scientific Research and Essays*, 6, 6270–6284.

Bozkurt, A. and Y. Sagat, 2008. Vertical distribution of Birecik Dam Lake (Turkey), zooplankton, *Journal of Fisheries Sciences*, 2, 332-342

Bulut, H. and Saler, S., 2013. Kalecik Baraj Gölü (Elazığ-Türkiye) zooplanktonu. *Fırat Üniversitesi Fen Bilimleri Dergisi*, 25(2), 99-103

Bulut, H. and Saler, S., 2014. Zooplankton of Beyhan Dam Lake (Elazığ-Turkey). *Turkish Journal of Science and Technology*, 9(1), 23-28

Dumont, H.J. and De Ridder, M., 1987. Rotifers from Turkey. *Hydrobiologia*, 147, 65-74.

Emir, N., 1996. Türkiye zooplankton (Rotifera, Cladocera tür listesi ve yayılışları. 20.1: Rotifera, In: Genel ve Türkiye Zoocoğrafyası (Hayvan Coğrafyası) (ed. A. Demirsoy), Meteksan Yayınları, pp. 465-470.

Kaya, M. and Altındağ, A., 2007. Zooplankton fauna and seasonal changes of Gelingüllü Dam Lake (Yozgat, Turkey), *Turkish Journal of Zoology*, 31, 347-351.

- Özdemir Mis, D., Aygen, C., Ustaoglu, M.R. and Balık, S., 2009.** Tahtalı Baraj Gölü (İzmir)'nün zooplankton kompozisyonu. *E.Ü. Su Ürünleri Dergisi*, 26(2), 129-134.
- Saler (Emiroğlu), S. and Şen, D., 2000.** Cip Baraj Gölü (Elazığ) Rotifera faunasının taksonomik yönden incelenmesi. *Fırat Üniversitesi Fen ve Mühendislik Bilimleri Dergisi*, 12(1), 329-339.
- Saler, S. and İpek, Alış., 2014.** Zooplankton of Hancağız Dam Lake (Gaziantep - Turkey). *Journal of Survey in Fisheries Sciences*, 1(1), 45-54.
- Saler, S., Haykır, H. and Baysal, N., 2014.** Zooplankton of Uzunçayır Dam Lake (Tunceli-Turkey). *Journal of Fisheries Sciences*, 8(1), 1-7.
- Segers, H., Emir, N. and Mertens, J., 1992.** Rotifera from north and northeast Anatolia (Turkey). *Hydrobiologia*, 245, 179-189.
- Segers, H., 1994.** Redescription of *Lecane fadeevi* (Neiswestnowa-Shadina, 1935) (Rotifera, Lecanidae). *Mededelingen. Koninklijk Belgisch Instituut voor Natuurwetenschappen*, 64, 235-238.
- Segers, H., 1995.** Rotifera: the Lecanidae (Monogononta). In: guides to the identification of the microinvertebrates of the continental waters of the world Volume 2 (eds. H.J.F. Dumont and T. Nogrady). SPB Academic Publishing, 226 P.
- Segers, H., 2008.** Global diversity of rotifers (Phylum Rotifera) in freshwater. *Hydrobiologia*, 595, 49-59.
- Ustaoglu, M.R., 2004.** A check-list for zooplankton of Turkish inland waters, E.U. *Journal of Fisheries and Aquatic Sciences*, 21(3-4), 191-199.
- Ustaoglu M., R., Altındağ, A., Kaya M., Akbulut N., Bozkurt, A., Özdemir Mis D., Atasagun S., Erdoğan, S., Bekleyen, A., Saler, S. and Okgerman, H.C., 2012.** A check list of Turkish Rotifers. *Turkish Journal of Zoology*, 36(1), 607-622.
- Ustaoglu, M.R., 2015.** An updated zooplankton biodiversity of Turkish inland waters. *Journal of Limnology and Freshwater Fisheries Research*, 1(3), 151-159.