

Isolation and identification of antibacterial steroid compounds from *Ulva fasciata* in the Persian Gulf

Azizi R.¹; Motallebi Moghanjoughi A.A.^{1*}; Nazemi M.²; Sharif Ruhani M.³; Afshar Nasab M.⁴

Received: May 2019

Accepted: October 2019

Abstract

Due to numerous reasons, marine-obtained resources have attracted the interest of researchers. One of the subjects attracted the interest of researchers in recent years in the field of biologist marine algae due to their nutritional value, their benefits for health and their biological activities. In this study, the powder was extracted using acetone, then the extract purified by silica gel column chromatography with n-hexane and ethyl acetate combination to isolate the steroid compounds. Isolated compound was run through TLC and sprayed with vanillin-sulphuric acid reagent to detect the steroid compounds and profile of isolated compounds obtained by gas chromatography (GC). Antibacterial activity of steroids extracted of green algae *Ulva fasciata* from Qeshm Island in the Persian Gulf were assessed (in vitro). The antibacterial activity of the fractions was determined by Broth Dilution Methods against clinical Gram-negative bacteria: *Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonella typhi* and Gram-positive bacteria: *Staphylococcus aureus*, *Bacillus subtilis*, *Bacillus cereus*. The steroids; cholest-7-en-3ol and cholestan-3-one, cyclic 1, 2-ethanediyl acetal compounds in column fraction were identified by GC. The results showed the minimum bactericidal concentration (MBC) of the cholest-7-en-3ol and cholestan-3-one, cyclic 1,2-ethanediyl acetal for *S. aureus* and *Bacillus cereus* were 500 $\mu\text{g mL}^{-1}$, the MBC of the cholest-7-en-3ol for *B. subtilis* was 500 $\mu\text{g mL}^{-1}$ and 1000 $\mu\text{g mL}^{-1}$ detected for cholestan-3-one, cyclic 1,2-ethanediyl acetal. The compounds had neither antibacterial activity on gram negative bacteria. Based on our results, extracted steroids from *U. fasciata* can be considered as a source of novel antibiotics.

Keywords: Green algae, Antibacterial activities, Natural products, Persian Gulf

1-Department of Hygiene, Science and Research Branch, Islamic Azad University, Tehran

2-Persian Gulf and Oman Sea Ecological Research Center, Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research, Education and Extension Organization (AREEO), Bandar Abbas, Iran

3-Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research, Education and Extension (AREEO)

4- Kherad higher education center, Bushehr, Iran.

*Corresponding author's Email: motallebi@ifro.ir

Introduction

Marine organisms are potential sources of bioactive secondary metabolites for use in development of new pharmaceutical agents (Abedin and Taha, 2008; EL-Gamal, 2010) and many of these substances have been demonstrated to possess interesting biological activities (Faulkner, 2002; Abdel-Raouf *et al.*, 2008). Marine algae have been reported to produce a wide variety of bioactive secondary metabolites as antibacterial, and cytotoxic agents and the bioactive substances included alkaloids, polyketides, polysaccharide, phlorotannins, triterpenoids, sterols, Quinone's, lipids and glycerol (Cabrita *et al.*, 2010).

Secondary metabolites derivatives of algae offer as a potentially useful source of natural antibacterial agents for food and medical applications. Algae contain various inorganic and organic substances which have benefit for human. Antibacterial activity of algae extracts is generally assayed using various organic solvents which always provide a higher efficacy in extracting compounds for antimicrobial activity (Cordeiro *et al.*, 2006; Tüney *et al.*, 2006). Isolation of the natural compounds from algal could produce useful therapeutics, the harnessing and bioengineering of recently characterized allelochemicals represents a potential area of new marine antibacterial (Saritha *et al.*, 2013).

Ulva species are the most abundant algae in coastal benthic communities

around the world, Ulvacean are considered bio indicators species with increased importance in coastal ecosystem management, mainly related to green tides associated with eutrophication processes in shallow environments (Wichard, 2015). Exploring bacteria induced growth and morphogenesis in the green macro algae in order Ulvae (*Chlorophyta*) (Basham *et al.*, 2014). *Ulva fasciata* is one of the most common green algae and accrues almost every season (Makaremi *et al.*, 2011) also known as *sea lettuce* and is a common algae used in parts of the world for consumption. Due to the increased resistance to antibiotics, the study of new antibiotic is very important from the perspective of medical treatment (Daneshvar *et al.*, 2017). Many authors had found antimicrobial activities of macroalgae due to fatty acids contents (Daneshvar *et al.*, 2017). Antimicrobial activity depends on both algal species and solvents used for their extraction (Prakash *et al.*, 2011; Radika *et al.*, 2012). Various algae have shown to have antibacterial activity in vitro against gram positive and gram negative bacteria (Ostensvik *et al.*, 1998).

Due to the antimicrobial activity of steroids, this study investigates the isolation and identification of bioactive steroids and antibacterial activities of isolated compounds of *U. fasciata* from the Qeshm Island in the Persian Gulf.

Materials and methods

Sampling and identification

Ulva fasciata has been collected in February 2018 from the areas of

Shibderaz in Qeshm Island in the Persian Gulf (Fig. 1). After identification of species, the samples were dried in the shadow.




Figure 1: Geographical situation of sampling location in Qeshm Island.

Extraction

The powder of dried *Ulva fasciata* (100 g dry weight) was extracted with acetone solvent. After 72 hours of soaking in acetone, the solvent filtered and acetone evaporated to dryness, at low pressure at 30-35°C by using Rota vapor (Çitoğlu and Acıkara, 2012).

Isolation and identification of steroid compounds

Silica gel column chromatography with 70 cm height and 2 cm diameter used for purification of *Ulva fasciata* acetone extract (3/2 g). The packed column washed up by different combination (100:0, 90:10, 80:20, 70:30, 60:40, 50:50, 40:60, 30:70, 20:80, 10:90, 0:100) of hexane: ethyl acetate solvent and all fractions were collected in 10 ml

tube (108 fractions) (Çitoğlu and Acıkara, 2012).

Obtained fractions from the chromatography columns were performed by thin layer chromatography with a mobile phase including methanol-chloroform- n butanol solvents with a ratio of 70:20:10. To identify the steroid compounds, vanillin-sulphuric acid reagent was used as a 1% solution of vanillin in ethanol and 5% sulphuric acid solution in ethanol by spraying on a thin-layer chromatography plate. After spraying, thin-layer chromatography plates were placed in the oven at 110°C for 10 minutes; changes in visible light were observed; the portions of the steroids were changed to blue color (Attaway *et al.*, 1965). The gas chromatography-mass

spectrometry (GC-MS) spectrum was taken on GC-MS (Agilent7000 Series Triple Quad GC/MS MainFrame). The column part number 19091s-436 (60 m× 0.25 mm, film thickness 0.25 µm) was used for GC-MS. Helium was used as a carrier gas at a flow rate of 1.0 ml min⁻¹; the electron ionization mode with ionization energy of 70 eV was used for MS detection, with a mass range of 50–650 m z⁻¹. The compound was recognized using standard information in the Wiley 7.0 library using its retention time and mass fragmentation pattern.

Antibacterial assay

Antibacterial activity was determined against *Escherichia coli* ATCC 15224, *Pseudomonas aeruginosa* ATCC 27853, *Salmonella typhi* ATCC 1609, *Staphylococcus aureus* ATCC 1764, *Bacillus cereus* ATCC 11778 and *Bacillus subtilis* PTCC 1715 by performing the classic broth dilution susceptibility test. The amount of 1.5×10⁵ microorganisms in colony forming units [CFU] ml⁻¹, a 1:100 dilution of a suspension of turbidity equal to a McFarland standard 0.5, was added to an equal volume (1 ml) of each concentration (5, 10, 20, 30, 50, 100, 200, 300, 400, 500, 1000, 2000 µg ml⁻¹) of steroid components agent and to a tube of the growth control. After overnight, the tubes were examined for turbidity, indicating growth control of the microorganism. The lowest concentration of the agent that inhibits growth of the organism, as detected by lack of visual turbidity, was designated

as the minimum inhibitory concentration (MIC). Later, the MIC has been determined; a known quantity (0.1 ml) of inoculums from each tube of broth that showed on visible turbidity after 22 to 24 hours' incubation is subculture to solid agar plates. The number of colonies which grow up on the subculture after overnight incubation is then counted and compared to the number of CFU ml⁻¹ in the original inoculum. Since even bacterial extracts do not always totally sterilize a bacterial population, the lowest concentration of antimicrobial agent that allowed less than 0.1% of the original inoculums to survive is considered as the minimum bacterial concentrate (MBC) (Rosenblatt, 1991).

Kruskal-Wallis test was used to compare the mean concentration of each extracted material as well as tetracycline.

Results

Isolation of fractions containing steroids

In column chromatography, the acetone extract was packed and eluted with the distinct solvent system. Based on the TLC profile, 108 fractions have been gathered from column chromatography. The fractions C44 (Yield 121 mg) and B33 (Yield 92 mg) that gave a positive response for steroids by Vanillin-sulphuric acid reagent were selected and separated into eleven and ten sub fractions, by silica gel. The fraction of C44 E and C33F was subjected to TLC for isolation of active principle. After TLC run (Fig. 2), the band that gave a

positive response for steroids producing blue color with Aniline test, punched and dissolved in acetone, filtered and

brown crystals of compound were obtained (2.3 mg) and then comparing GC-MS with Wiley 7.0 library verified the compound's identity.




Figure 2: Thin layer chromatography of acetone extracts of *Ulva fasciata*.

Identification of steroids compounds

The fractions C44E and C33 F that gave a positive response for steroids by Vanillin-sulphuric acid reagent selected for GC-MS detection use. The cholest-7-en-3ol with the chemical formula of

$C_{27}H_{46}O$ (Fig. 3), molecular weight of $386.6535 \text{ g mol}^{-1}$ belonging to the steroids group identified with 92% purity in fraction C44 E (ethyl acetate-hexane 40: 60) at 32-34 min retention time of gas chromatography.

Hit 1 : Cholest-7-en-3-ol, (3 β ,5 α)-
C₂₇H₄₆O; MF: 824; RMF: 900; Prob 27.2%; CAS: 80-99-9; Lib: replib; ID: 26506.




Figure 3: GCMS analysis of cholest-7-en-3ol.

The cholestan-3-one, cyclic 1,2-ethanediyl acetal with the chemical formula of $C_{29}H_{50}O_2$ (Fig. 4), molecular weight of 430.7 g mol^{-1} belonging to

the steroids group identified with 94% purity in fraction C33 F (ethyl acetate-hexane 30: 70) at 34-38 min retention time of gas chromatography.

Hit 2 : Cholestan-3-one, cyclic 1,2-ethanediyl acetal, (5 α)-
C₂₉H₅₀O₂; MF: 651; RMF: 686; Prob 8.71%; CAS: 1858-14-6; Lib: mainlib; ID: 62030.




Figure 4: GCMS analysis of cholestan-3-one, cyclic 1, 2-ethanediyl acetal.

Antibacterial activities of fraction containing steroid compounds

As indicated in Table 1, the minimum inhibitory concentration of bacterial growth (MIC) of cholest-7-en-3ol and cholestan-3-one, cyclic 1, 2-ethanediyl acetal for *B. subtilis* was $500 \mu\text{g ml}^{-1}$ and for *E. coli* equal to $1000 \mu\text{g ml}^{-1}$. The MIC of cholest-7-en-3ol for *S.*

aureus and *B. cereus* were 300 and $500 \mu\text{g ml}^{-1}$, the MIC of cholestan-3-one, cyclic 1, 2-ethanediyl acetal for *S. aureus* and *B. cereus* were 200 and $300 \mu\text{g ml}^{-1}$. The cholest-7-en-3ol and cholestan-3-one, cyclic 1, 2-ethanediyl acetal compounds have not shown any inhibitory effects on growth of *S. typhi* and *P. aeruginosa*.

Table 1: Minimum inhibitory concentration of fraction containing steroid compounds from *U. fasciata*.

Tetracycline ($\mu\text{g ml}^{-1}$)	cholest-7-en-3ol concentration ($\mu\text{g ml}^{-1}$)	cholestan-3-one, cyclic 1,2-ethanediyl acetal concentration ($\mu\text{g ml}^{-1}$)	Bacteria
100	500	500	<i>Bacillus subtilis</i>
100	300	200	<i>Staphylococcus aureus</i>
100	500	300	<i>Bacillus cereus</i>
200	1000	1000	<i>Escherichia coli</i>

As shown in Table 2, the minimum bactericidal concentration (MBC) of the cholest-7-en-3ol and cholestan-3-one, cyclic 1,2-ethanediyl acetal for *S. aureus* and *Bacillus cereus* was $500 \mu\text{g}$

ml^{-1} , the MBC of the cholest-7-en-3ol for *B. subtilis* was $500 \mu\text{g ml}^{-1}$ and $1000 \mu\text{g ml}^{-1}$ detected for cholestan-3-one, cyclic 1,2-ethanediyl acetal.

Table 2: Minimum bactericidal concentration (MBC) of fraction containing steroid compounds extracted from *U. fasciata*.

Tetracycline ($\mu\text{g ml}^{-1}$)	cholest-7-en-3ol concentration ($\mu\text{g ml}^{-1}$)	cholestan-3-one, cyclic 1,2-ethanediyl acetal concentration ($\mu\text{g ml}^{-1}$)	Bacteria
200	500	1000	<i>Bacillus subtilis</i>
200	500	500	<i>Staphylococcus aureus</i>
200	500	500	<i>Bacillus cereus</i>
200	-	-	<i>Escherichia coli</i>

The results of Kruskal-Wallis test showed that there was no significant difference between the concentration of extracts and tetracycline in inhibiting the growth of each four strains ($p>0.05$). Similar results were observed ($p>0.05$) among these compounds in their lethality, indicating that the extracted compounds had a similar effect to tetracycline on bacterial growth and lethality.

Discussion

The marine secondary metabolites have received considerable attention over the last decades. There are large numbers of marine natural products which have been targeted by pharmaceutical industry (Saber *et al.*, 2016). Due to the increase of resistance to antibiotics, the study on new antibacterial activities of steroid derivatives is very important for human health (Doğan *et al.*, 2017; Cheng *et al.*, 2016). The aim of this study was to investigate the antibacterial activities of the green algae (*U. fasciata*) steroids; cholest-7-en-3ol and cholestan-3-one, cyclic 1,2-ethanediyl acetal.

In the present study, the cholest-7-en-3ol and cholestan-3-one, cyclic 1, 2-ethanediyl acetal, which are the steroid compounds, extracted from *U. fasciata*

using acetone solvent (Veeranan *et al.*, 2018; Fasya *et al.*, 2019). The sterol as same as steroid compounds which is found in our study has been identified from green algae *Ulva prolifera* named as cholest-5-en-3-ol (Wang *et al.*, 2016), 7-oxo-cholest-5(6)-en-3-ol from red algae *Jania rubens* (Ahmed *et al.*, 2011), 3 α , 6 α -dihydroxy-5 β -cholestan-12-one and 6 β -hydroxycholest-4-en-3-one from red alga *Laurencia papillosa* (Alarif *et al.*, 2011).

In this study, the antibacterial activities of cholest-7-en-3ol and cholestan-3-one, cyclic 1, 2-ethanediyl acetal has been examined. This research determined that fractions containing cholest-7-en-3ol and cholestan-3-one, cyclic 1, 2-ethanediyl acetal compounds extracted from *U. Fasciata* in a concentration of 500-1000 $\mu\text{g ml}^{-1}$ has antimicrobial activity on gram-positive bacteria of *B. subtilis*, *S. aureus* and *B. cereus* and they have not shown any inhibitory effects on growth of antimicrobial activity on gram-negative bacteria of *S. typhi* and *P. aeruginosa*. In this case, Chandrasekaran *et al.* (2014) reported that ethyl acetate extract of *U. fasciata* shows strong antibacterial activities; MIC ($125 \mu\text{g ml}^{-1}$) and MBC ($250 \mu\text{g ml}^{-1}$) against *B.*

subtilis. Chloroform and ethyl acetate fractions of *U. lactuca* have shown antibacterial activities against *Staphylococcus aureus* ($1.6 \mu\text{g ml}^{-1}$) and *Enterococcus faecalis* ($0.2 \mu\text{g ml}^{-1}$) (Habbu *et al.*, 2016). In another study, the fractions containing cholesterol derivative, 24-propylidene cholest-5-en-3 β -ol extracted from red seaweed *Laurencia papillosa*, showed the minimum inhibitory concentration ranged from 1.2 to 1.7 μg against *Staphylococcus aureus*, *Bacillus subtilis* and *Escherichia coli* (Kavita *et al.*, 2014).

The results of these studies are in accordance with the studies on the steroid compounds extracted from *U. fasciata* of Qeshm Island which has antimicrobial activities against gram-positive bacteria. The study of the new antibacterial of steroids; cholesterol derivatives have the potential of being used as antibacterial drugs. A steroid is a biologically active organic compound and they have two principal biological functions: cell membranes and signaling molecules. Since steroids resemble cholesterol, it is thought that they compete with this for the receptors and it can also disrupt cell integrity and permeability. It is thought that this is the reason why steroids and cholesterol derivatives can have antibacterial activities (Doğan *et al.*, 2017). Based on our results, the steroids extracted from green algae *U. fasciata* have shown a relatively strong antibacterial activity on gram positive bacteria and it could be considered as a source of novel antibiotic.

Acknowledgments

The authors take this opportunity to thank the Iranian Fisheries Science Research Institute. The project was conducted at the Persian Gulf and Oman Sea Ecological Research and acknowledges all personnel of this center.

References

- Abedin, R.M. and Taha, H.M., 2008.** Antibacterial and antifungal activity of cyanobacteria and green microalgae. Evaluation of medium components by Plackett-Burman design for antimicrobial activity of *Spirulina platensis*. *Global Journal of Biotechnology and Biochemistry*, 3(1), 22-31
- Ahmed, H.H., Hegazi, M.M., Abd-Allac, H.I., Eskander, E.F. and Ellithey, M.S., 2011.** Antitumour and antioxidant activity of some Red Sea seaweeds in Ehrlich ascites carcinoma in vivo. *Zeitschrift für Naturforschung C*, 66(7-8), 367-376.
- Alarif, W.M., Al-Lihaibi, S.S., Abdel-Lateff, A. and Ayyad, S.E.N., 2011.** New antifungal cholestane and aldehyde derivatives from the red alga *Laurencia papillosa*. *Natural Product Communications*, 6(12), 1934578X1100601208.
- Allwood, D.A., Xiong, G., Cooke, M.D., Faulkner, C.C., Atkinson, D., Vernier, N. and Cowburn, R.P., 2002.** Submicrometer ferromagnetic NOT gate and shift register. *Science*, 296(5575), 2003-2006.

- Al-Saif, S.S.A.L., Abdel-Raouf, N., El-Wazanani, H.A. and Aref, I.A., 2014.** Antibacterial substances from marine algae isolated from Jeddah coast of Red sea, Saudi Arabia. *Saudi Journal of Biological Sciences*, 21(1), 57-64.
- Attaway, J.A., Barabas, L.J. and Wolford, R.W., 1965.** Analysis of terpene hydrocarbons by thin layer chromatography. *Analytical Chemistry*, 37(10), 1289-1290.
- Bassham, D.C. and Crespo, J.L., 2014.** Autophagy in plants and algae. *Frontiers in Plant Science*, 5, 679.
- Cabrita, M.T., Vale, C. and Rauter, A.P., 2010.** Halogenated compounds from marine algae. *Marine drugs*, 8(8), 2301-2317.
- Chandrasekaran, M., Venkatesalu, V., Raj, G.A. and Krishnamoorthy, S., 2014.** Antibacterial activity of *Ulva fasciata* against multidrug resistant bacterial strains. *International Letters of Natural Sciences*, 14 P.
- Çitoğlu, G.S. and Acikara, Ö.B., 2012.** Column chromatography for terpenoids and flavonoids. In 'chromatography and its applications'. InTech, Rijeka. pp.13-49.
- Cordeiro, R.A., Gomes, V.M., Carvalho, A.F.U. and Melo, V.M.M., 2006.** Effect of proteins from the red seaweed *Hypnea musciformis* (Wulfen) Lamouroux on the growth of human pathogen yeasts. *Brazilian Archives of Biology and Technology*, 49(6), 915-921.
- Daneshvar, E., Vazirzadeh, A., Niazi, A., Sillanpää, M. and Bhatnagar, A., 2017.** A comparative study of methylene blue biosorption using different modified brown, red and green macroalgae—Effect of pretreatment. *Chemical Engineering Journal*, 307, 435-446.
- Doğan, A., Otlu, S., Çelebi, Ö. Aksu, P., Sağlam, A.G., Doğan, A.N.C. and Mutlu, N., 2017.** An investigation of antibacterial effects of steroids. *Turkish Journal of Veterinary and Animal Sciences*, 41(2), 302-305.
- El Gamal, A.A., 2010.** Biological importance of marine algae. *Saudi Pharmaceutical Journal*, 18(1), 1-25.
- Fasya, A.G., Baderos, A., Madjid, A.D.R., Amalia, S. and Megawati, D.S., 2019.** Isolation, identification and bioactivity of steroids compounds from red algae *Eucheuma cottonii* petroleum ether fraction. In AIP Conference Proceedings. 2120(1), 030025. AIP Publishing.
- Habbu, P., Warad, V., Shastri, R., Savant, C., Madagundi, S. and Kekare, P., 2016.** *In vitro* and *in vivo* antimicrobial activity of *Ulva lactuca* Linn. (Green algae) associated endophytic bacterial Strains. *Journal of Applied Pharmaceutical Science*, 6(10), 138-146.
- Kavita, K., Singh, V.K. and Jha, B., 2014.** 24-Branched Δ^5 sterols from *Laurencia papillosa* red seaweed with antibacterial activity against

- human pathogenic bacteria. *Microbiological Research*, 169(4), 301-306.
- Makaremi, M., Sabkara, J. and Mirzajani, A., 2011.** Investigation on *Nodularia* anomalous algal blooming (AAB) in south west Caspian Sea (2005-2006).
- Martínez-Banderas, A.I., Aires, A., Quintanilla, M., Holguin-Lerma, J.A., Lozano-Pedraza, C., Teran, F.J., Moreno, J.A., Perez, J.E., Ooi, B.S., Ravasi, T. and Merzaban, J.S., 2019.** Iron-based core-shell nanowires for combinatorial drug delivery and photothermal and magnetic therapy. *ACS Applied Materials and Interfaces*, 11(47), 43976- 988.
- Ostensvik, O., Skulberg, O.M., Underdal, B. and Hormazabal, V., 1998.** Antibacterial properties of extracts from selected planktonic freshwater cyanobacteria--a comparative study of bacterial bioassays. *Journal of Applied Microbiology*, 84(6), 1117-1124.
- Radhika, D., Veerabahu, C. and Priya, R., 2012.** Antibacterial activity of some selected seaweeds from the Gulf of Mannar Coast, South India. *Asian Journal of Pharmaceutical and Clinical Research*, 5(4), 89-90.
- Saberi, A., Zorriehzaha, M.J., Emadi, H., Kakoolaki, S. and Fatemi, S.M.R., 2017.** Effects of *Chlorella vulgaris* on blood and immunological parameters of Caspian Sea salmon (*Salmo trutta caspius*) fry exposed to Viral Nervous Necrosis (VNN) virus. *Iranian Journal of Fisheries Sciences*, 16(2), 494-510.
- Saritha, K., Mani, A.E., Priyalaxmi, M. and Patterson, J., 2013.** Antibacterial activity and biochemical constituents of seaweed *Ulva lactuca*. *Global Journal of Pharmacology*, 7(3), 276-282.
- Tüney, İ., Cadirci, B.H., Ünal, D. and Sukatar, A., 2006.** Antimicrobial activities of the extracts of marine algae from the coast of Urla (Izmir, Turkey). *Turkish Journal of Biology*, 30(3), 171-175.
- Veeranan, T., Kasirajaan, R., Gurunathan, B. and Sahadevan, R., 2018.** A novel approach for extraction of algal oil from marine macroalgae *Ulva fasciata*. *Renewable Energy*, 127, 64-73.
- Wang, Q., Wang, Z., Awasthi, M.K., Jiang, Y., Li, R., Ren, X., Zhao, J., Shen, F., Wang, M. and Zhang, Z., 2016.** Evaluation of medical stone amendment for the reduction of nitrogen loss and bioavailability of heavy metals during pig manure composting. *Bioresource Technology*, 220, 297-304.
- Wichard, T., Charrier, B., Mineur, F., Bothwell, J.H., Clerck, O.D. and Coates, J.C., 2015.** The green seaweed *Ulva*: a model system to study morphogenesis. *Frontiers in Plant Science*, 6, 72.
- Zhang, J., Ding, Y., Shen, S., Cheng, Y., Sun, M., Luan, H. and Liu, Y., 2017.** Thumt: An open source toolkit for neural machine translation. *arXiv preprint arXiv:1706.06415*.