

Isolation and characterization of microsatellite loci in the Persian sturgeon (*Acipenser persicus*, Borodine, 1897) and cross-species amplification in four commercial sturgeons from the Caspian Sea

Moghim M.^{1,3*}; Heist E. J.²; Tan S. G.³; Pourkazemi M.⁴; Siraj S. S.⁵
; Panandam J. M.⁶; Pourgholam R.¹; Kor D.¹;
Laloei F.¹; Taghavi M. J.¹

Received: January 2012 Accepted: March 2012

Abstract

In order to have a sustainable management on Persian sturgeon as a highly commercial species in the South Caspian Sea, we need to identify its population structure and the level as well as its conservation status in their natural habitat. To develop a conservation program for this all Caspian Sea' sturgeon species it requires knowledge of its genetic diversity using reliable molecular marker to study population genetic structure. For these purposes, an enriched library was prepared based on a modified biotin-capture method. Approximately 1800 positive clones were screened for microsatellites in an *Acipenser persicus* genomic library. Of these 350 positively hybridizing clones were sequenced, and 81 clones were identified as having microsatellites with adequate flanking regions. We developed and tested 68 microsatellite primer pairs for Persian sturgeon. Out of 68 primer pairs developed, 11 pairs resulted in poor or no amplification, 13 were ambiguous, 6 were monomorphic, 20 were tetrasomic and 18 were octosomic in Persian sturgeon. While none of the markers showed disomic inheritance in Persian sturgeon and Russian sturgeon (*A. gueldenstaedtii*). Several of the markers appeared useful for studies stellate sturgeon (*A. stellatus*), ship sturgeon (*A. nudiventris*) and beluga (*Huso huso*). Nearly all the polymorphic pattern for ship, stellate and beluga displayed the simple banding patterns characteristic of disomic loci, while those for Russian sturgeon displayed banding patterns characteristic of tetraploid or higher polyploid levels. These markers may prove useful in a variety of future sturgeon population genetic studies in the Caspian Sea.

Keywords: Persian sturgeon, *Acipenser persicus*, Caspian Sea, Microsatellite, Population genetic

1- Genetic Department of the Caspian Sea Ecology Research Center, P.O.Box: 961, Sari, Iran.

2- Fisheries and Illinois Aquaculture Center, Department of Zoology, Southern Illinois University, Carbondale, Illinois 62901-6501, USA.

3-Department of Cell and Molecular Biology, Faculty of Biotechnology and Biomolecular Sciences, University Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia.

4- International Sturgeon Research Institute, P.O. Box: 41635-3464 Rasht, Iran .

5-Department of Aquaculture, Faculty of Agriculture, University Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia.

6-Department of Animal Science, Faculty of Agriculture, University Putra Malaysia, 43400 UPM Serdang, Selangor, Malaysia.

*Corresponding author's email: mehdi.moghim@gmail.com

Introduction

The Persian sturgeon is an anadromous species living in the Caspian Sea, but it mainly inhabits in the southern Caspian region along the Iranian coast. Persian sturgeon enters the rivers for spawning, mainly the Sefid-Rud, Tajan and Gorganrud rivers in Iran and Kura river in Azerbaijan, less -the Volga, Ural, Samur, Terek, Lenkoranka and Astara rivers (Berg, 1948). Its population, after collapse in 1970's, has risen in the 1990 decade and comprises the largest proportion of the total Iranian sturgeon commercial catch in recent years (Pikitch et al., 2005; Moghim et al., 2006). While in 1980-s its catch did not exceed 5% of the total sturgeon catches at the Volga and the Ural rivers, the share of this species decreased to 0.03%, in the year of 2000 (Khodorevskaya et al., 2000). Persian sturgeon is listed as a critically endangered species by the International Union for Nature Conservation (IUCN 2011), due to continued overexploitation, illegal catch spawning habitat loss and pollution.

Persian sturgeon stocks are recovered mainly by artificial propagation and Iranian Fisheries, release millions of 3-5 g fingerlings to the adjacent rivers of Caspian Sea annually (Abdolhay and Baradaran Tahori, 2006; Moghim et al., 2006). The sustainable management and conservation plan of this unique species requires knowledge of its genetic structure and levels of each stock in its natural habitat. Several population genetic studies were conducted on five sturgeon species in the Caspian Sea using microsatellite markers (Pourkazemi, 2007; Safari et al., 2008; Noruzi et al., 2008; Khoshkholgh et al., 2008).

Cross-species amplification using microsatellite primers of *Scaphirhynchus* were applied in the Persian sturgeon by Moghim et al., (2009) but none of the loci exhibited disomic inheritance. While microsatellites are expensive to develop initially, because of the higher degree of

statistical power associated with codominant markers -microsatellite loci were developed for the Persian sturgeon to find disomic loci. The objective of the present research was to develop the Persian sturgeon specific microsatellite primers, and compare its application on other four sturgeon species in the Caspian Sea.

Materials and methods

An enriched library was prepared following a modification of the protocols of Hamilton et al., (1999) and Glenn et al., (2000) as described in Heist et al., (2003). Total genomic DNA from a single Persian sturgeon was digested with *RsaI*. Complementary linkers for use as polymerase chain reaction (PCR) primer sites were designed to contain an *RsaI* site when double-stranded (Linker-F: 5'-CTAAGGCCTTGATCGCAGAAGC-3'; phosphorylated Linker-R: 5'-pGCTTCTGCGATCAAGGCCTTAGAA AA-3') and ligated to genomic DNA fragments. Biotinylated (GT)₁₅, (GA)₁₅, (GATA)₅ and (GACA)₅ probes were hybridized to linker-ligated DNA fragments and microsatellite containing DNA was selectively retained by binding biotinylated DNA fragments to streptavidin coated MagneSphere™ paramagnetic particles (Promega, Madison, WI, USA). Microsatellite-containing fragments were then amplified using PCR reactions containing approximately 10 ng microsatellite-enriched genomic DNA and 1× PCR buffer (200 mM KCl, 100 mM Tris), 200 μm of each dNTP, 2 mM MgCl₂, 1 μm Linker-F as primer, and 2 units Taq DNA polymerase. PCR amplifications consisted of 94 °C for 5 min, followed by 40 cycles of 94 °C for 45 s, 62 °C for 1 min, and 72 °C for 1 min using an Quanta Biotec master cycler gradient thermocycler (Quanta Biotech Ltd, Surrey, United Kingdom). The PCR product was ligated

into a pUC19 cloning vector and used to transform DH5 α competent cells (Invitrogen, Carlsbad, Ca, USA). Colonies were transferred to a nylon membrane and probed with ^{32}P labeled (GT) $_{15}$, (GA) $_{15}$, (GATA) $_5$ and (GACA) $_5$. We isolated plasmid DNA from positive colonies using the Wizard miniprep kit (Promega). The positive clones were sequenced using M13 (F and R) universal sequence primers. Plasmid DNA was isolated from positive clones and sequenced with the ABI PRISM BigDye terminator cycle sequencing ready reaction kit using an ABI 377 automated sequencer (PE Applied Biosystems, Weiterstadt, Germany). Approximately 1800 positive clones were screened for microsatellites in an *Acipenser persicus* genomic library. Of these 350 positively hybridizing clones were sequenced, and 81 clones were identified as having microsatellites with adequate flanking regions. In total 68 microsatellite PCR primers were designed after omitting 13 clones with the same sequences. Microsatellite PCR primers were designed using the Primer3 (<http://www.genome.wi.mit.edu/cgi-bin/primer/primer3.cgi>) or the MacVector (Oxford Molecular) software package. These loci were tested in Persian sturgeon (n=12) to identify optimal annealing temperatures and to determine if disomic polymorphic products could be reliably amplified. Additional individuals (n=24) from different populations were used to confirm the ploidy status.

Amplification was performed using a gradient thermocycler at annealing temperatures ranging from 52 °C to 64 °C. The ten microlitre PCR reactions containing approximately 1-10 ng genomic DNA, 0.1 units Taq DNA polymerase, 0.5 mM of each primer, 200 mM of each dNTP, 2 mM MgCl $_2$, and 1 \times PCR buffer. Amplification consisted of a 5 min denaturing step at 95 °C, 40 cycles of 95 °C for 30 s, 56 - 64 °C for 30 s, and 72

°C for 30 s, followed by a single five-minute extension step at 72 °C. PCR products were suspended 1:1 in 98% formamide/loading dye, denatured at 95°C for 5 min, and separated in a 6% denaturing polyacrylamide gels on a BIO-RAD gel sequencer running at 70 W for 45 - 60 min and visualized via Silver staining (An et al., 2009). Allele sizes were estimated using a 50-bp ladder molecular size standard (Invitrogen).

Amplification results were characterized as monomorphic if a single band of the same size was observed in all individuals, disomic if one or two bands were seen in every individual, tetrasomic if some individuals exhibited three or four bands, octosomic if more than four bands were observed in some individuals, weak if products were too faint to resolve, and ambiguous if banding patterns were too complex for us to interpret.

All primer pairs (except Ape-01 to Ape-18) were tested for cross-species amplification efficiency with four sturgeon species of the Caspian Sea, under the same PCR conditions used for Persian sturgeon including; the Stellate sturgeon, Russian sturgeon, Ship sturgeon and Beluga. Six individuals from each species were screened for polymorphism at these loci.

Results

In total 68 microsatellites PCR primers were designed after omitting 13 clones with the same sequences. Out of 68 primer pairs developed, 10 resulted in poor or no amplification, 13 were ambiguous; six of loci that amplified successfully were monomorphic, 21 were tetrasomic and 18 were octosomic in Persian sturgeon. None of the loci exhibited disomic inheritance (Figure 1). Locus name, clone size, GenBank accession number, repeat motif, PCR annealing temperature, and primer sequences are listed for these loci in Table 1.

Table 1: Characterization of 68 microsatellite loci in Persian sturgeon (*Acipenser persicus*), including repeat motifs, primer sequences and GenBank accession numbers, and cross-amplification in Russian (*A. gueldenstaedtii*), stellatus (*A. stellatus*), ship (*A. nudiventris*) sturgeon and beluga (*Huso huso*).

Prime	Repeat type and length	primer sequences (5' to 3')	GenBank accession no.	Product size (bp)	<i>A. persicus</i>	<i>A. gueldenstaedtii</i>	<i>A. stellatus</i>	<i>A. nudiventris</i>	<i>H. huso</i>
Ape-01	(CAGA) ₁₄	F:CAATGTCACAAACACACACAGCG R:TTTCTCTCCAGTTCGTCAGATGC	JF773767	171	tetrasomic				
Ape_02	(GT) ₁₃	F:CAAACATACCGTTCTGTGGGAC R:CGTCTGCTGAAGAAGGTTAAATATC	JF773768	123	octosomic				
Ape_03	(CAGA) ₁₄	F:CAATGTCACAAACACACACAGCG R:GCAGAAAAACCAGCCACAGTC	JF773769	141	tetrasomic				
Ape_04	(CA) ₁₀	F:GATAAAGGCACGACGCTACAACCTAC R:CATCTCAACCTGACAAATACCGTG	JF773770	119	octosomic				
Ape_05	(CAGA) ₆	F:ACTGAACCATTGGAGTATTGAGGC R:ACAGTAAACGCACACCAACAAGG	JF773771	137	tetrasomic				
Ape_06	(CAGA) ₁₅	F:AAACCTTCAGAGAGAGAGGGAGCG R:GCAGAAAAACCAGCCACAGTC	JF773772	239	octosomic				
Ape_07	(CT) ₁₂	F:CACAATTCACAGTCAGGGCTGTGTC R:TGCCACAATTCACAGTCAGGG	JF773773	253	ambiguous				
Ape_08	(CT) ₄₁	F:AGCCCCTGTGTCTGTCTGTTTGG R:GGAAATTCCTTGGTGTGTGTGGG	JF773774	164	ambiguous				
Ape_09	(CT) ₃₅	F:GATCAGCTCCAGTTTGCAGTGC R:GGAGATAGATTCTGTTCTGCCAAGTC	JF773775	299	ambiguous				
Ape_10	(CAGA) ₁₃	F:AGGGAGCGACAACTTACTCCTG R:GCAGAAGCACAGCAATGTGAAATC	JF773776	275	octosomic				
Ape_11	(CAGA) ₇	F:AACCATTGGAGTATTGAGGCACTG R:ACAGTAAACGCACACCAACAAGG	JF773777	133	octosomic				
Ape_12	(CT) ₁₃	F:GCCTTCAACATTCTCCTTATTGAGG R:CGTTACGAAAAACAAGTGTCTTGCC	JF773778	112	octosomic				
Ape_13	(CTGT) ₁₃	F:TCGCAGAAAAACCAGCCAC R:AAACCTTCAGAGAGAGAGGGAGCG	JF773779	233	octosomic				
Ape_14	(GA) ₂₂	F:ATTTTCGTGTCTGTCTTAATTGGTG R:GTAAATCTCACAATGTCCGTGGC	JF773780	164	tetrasomic				
Ape_15	(CT) ₆₄	F:TTCCTGTTGCCAGACATTTTAAACAC R:TCCTTAATTGGTAAAATTCATACCG	JF773781	175	no amplify				
Ape_16	(GA) ₁₃	F:AATGGAGAGAGAGAGAGGGAGTG R:AAGTCTTACAAAACCCGTGGTGG	JF773782	230	tetrasomic				
Ape_17	(CTGT) ₁₅	F:TCGCAGAAAAACCAGCCAC R:GCATTTCCGAGAAAACCTTCAGAG	JF773783	248	octosomic				
Ape_18	(GA) ₁₄	F:CGCAGAAGCACTAAAAGTCAAAGTC	JF773784	202	tetrasomic				

Ape-19	(CA) ₁₄	R:GGAAGATTTTCAGAGAGCAGCACTC F:GGGGTTAGAAAGCACAGATGA R:CAAGGTGGCACAGTGGACTA	EU483155	172	octosomic	octosomic	ambiguo s s	disomic	disomic
Ape_20	(GACA) ₅	F:CACTGCCTGCTGCCTAAAAC R:ACTGTGGGGCTCTGTCTGTC	EU531732	176	tetrasomic	tetrasomic	disomic	mono	disomic
Ape_21	(GACA) ₅	F:GGAGACAGACGAGGGAGAGA R:ATTCGGGACGTGAGACACAT	EU531733	397	tetrasomic	weak	weak	weak	ambiguous
Ape_22	(GTCT) ₁₄	F:CAGAAAAACCAGCCCACAGT R:GAGAGAGAGGGAGCGACAAA	EU531734	245	octosomic	octosomic	disomic	mono	disomic
Ape_23	(CA) ₂₅	F:CCTGCCACACCTACACAGAC R:GCGCATGCCTACAACAATTT	JF781300	177	no amplify	ambiguous	disomic	ambiguous	ambiguous
Ape_24	(CA) ₁₄	F:TGAACACAAAACACGGGACA R:TAAGGCCTTGATCGCAGAAG	EU531735	237	no amplify	ambiguous	disomic	ambiguous	ambiguous
Ape_25	(GAGAG) ₅	F:CCCCTGTCTGTCTGTCTGTTT R:ATCTCAGCCAGGAAGAACGA	EU531736	159	tetrasomic	tetrasomic	disomic	disomic	mono
Ape_26	(GA) ₃₈	F:GAGAGAGAGGGAGCGACAAA R:CAGAAAAACCAGCCCACAGT	EU531737	225	tetrasomic	tetrasomic	mono	mono	disomic
Ape_27	(GA) ₃₈	F:AACGAGTCCATGCTGGAGAG R:CCCCGTGTCTGTTTGT	EU531738	171	tetrasomic	tetrasomic	disomic	disomic	mono
Ape_28	(CTGT) ₁₀	F:CTCAGACCCGTGAGACACAA R:GCATTTCCGAGAAACCTTCA	EU531739	192	tetrasomic	no amply	disomic	no amply	disomic
Ape_29	(GT) ₁₅	F:TGAACACAAAACACGGGACA R:CGCACACACACGCACATA	EU531740	215	mono	tetrasomic	mono	disomic	ambiguous
Ape_30	(GT) ₁₁	F:AGGGCTACCTCCAGCTGTGT R:TCGCTCCTCAGACTCTGGAC	EU531741	172	tetrasomic	tetrasomic	disomic	disomic	ambiguous
Ape_31	(CT) ₂₆	F:GCCCTGTGTCTGTCTGTTT R:CGTGTGTGAGCGAGATAGGA	EU531742	189	no amplify	_	mono	_	mono
Ape_32	(GACA) ₁₅	F:CAAAGAGAGAGGGAGCGACA R:CAGAAAAACCAGCCCACAGT	EU531743	227	octosomic	octosomic	disomic	mono	disomic
Ape_33	(CTAT) ₉	F:TGCTGATCTAACCATTTCTTTGC R:AAGGCACACCATCTTTGTCC	EU531744	190	tetrasomic	tetrasomic	disomic	disomic	disomic
Ape_34	(CA) ₁₀	F:CCACCACCCTCCCACAATA R:GGGCAAATTGACTGCTTGAT	EU531745	162	mono	_	mono	_	mono
Ape-35	(GACA) ₆	F:ACTGCCTGCTGCCTAAAACA R:CTAAGGCCTTGATCGCAGAA	JF740087	231	ambiguous	mono	disomic	mono	disomic
Ape-36	(CTGT) ₅	F:TAGCACTGGGAACAGAAGCA R:AAAGCTCCAACACATGGACA	JF740088	240	no amplify	ambiguous	disomic	ambiguous	no amplify
Ape-38	(GTCT) ₆	F:GTGCGTGTGTGTGTGTGTGT R:GTGTGACAGTGAAGCGGAGA	JF773785	352	no amplify	tetrasomic	mono	disomic	mono
Ape-39	(GA) ₃₆	F:GGAAGGGGAGAGAGAGAACG R:GCGCTGTATTGTGGTGA	JF773786	269	tetrasomic	ambiguous	mono	ambiguous	mono
Ape-40	(CA) ₁₈	F:CCGAAACACACATACGC R:GCGCTCTCGTAGACTGTGC	JF773787	250	ambiguous	ambiguous	disomic	ambiguous	disomic
Ape-42	(CT) ₁₈	F:CGTGCCCACTGTTTTACCTT R:TGGATTCTAGGACGGTTGG	JF773788	254	no amplify	no amplify	no amplify	no amplify	no amplify
Ape-43	(CT) ₂₅	F:GCCCTGTGTCTGTCTGTTT R:GCATGTCTTTTCCAAAGTGAA	JF773789	180	no amplify	ambiguous	no amplify	ambiguous	no amplify

Ape-46	(GA) ₂₇	F: TGTGCCACAATTCACAGTCA R: CAGAGAGAGTCAGCGGGTCT	JF773790	245	octosomic	octosomic	no amplify	disomic	mono
Ape-47	(GA) ₃₄	F: ATCTCAGCCAGGAAGAACGA R: GCCCTGTGTCTGTCTGTTT	JF773791	180	octosomic	tetrasomic	disomic	disomic	disomic
Ape-48	(GA) ₃₂	F: TGTGCCACAATTCACAGTCA R: CCACGTTTATTAACCCAAATCAA	JF773792	201	ambiguous	no amplify	no amplify	no amplify	no amplify
Ape-49	(GA) ₃₈	F: ATCTCAGCCAGGAAGAACGA R: GCCCTGTGTCTGTCTGTTT	JF773793	188	octosomic	tetrasomic	disomic	disomic	no amplify
Ape-50	(CA) ₂₄	F: CCTGCTGCTGTATAAACTATGGA R: CGGACTGTGTGTCTGTCTGTCTGTC	JF773794	249	mono	mono	disomic	mono	mono
Ape-51	(GA) ₁₈ G ₂ (GA) ₁₉	F: ATCTCAGCCAGGAAGAACGA R: CCCGTGTCTGTCTGTCTGTTT	JF773795	189	tetrasomic	tetrasomic	disomic	disomic	mono
Ape-52	(CAGA) ₆	F: CACTGCCTGCTGCCTAAAAC R: TATTAACCCATCGGCTCCAC	JF773796	151	no amplify	no amplify	mono	no amplify	mono
Ape-53	(CA) ₁₄	F: CGCACACACACGCACATA R: ACGGCACTATACGCCAAAAT	JF773797	196	ambiguous	weak	weak	weak	weak
Ape-55	(GA) ₂₅	F: ATCTCAGCCAGGAAGAACGA R: CCCGTGTCTGTCTGTCTGTTT	JF773798	165	tetrasomic	tetrasomic	disomic	disomic	mono
Ape-56	(CA) ₁₁	F: TCGTCCTGCTGAAGAAGGTAA R: CGTTCTGTGGACAGTGAGA	JF773799	146	tetrasomic	octosomic	ambiguous	tetrasomic	ambiguous
Ape-57	(CA) ₁₅	F: CCATGCACACGCACTAGTTT R: ATTGTCATGCCCGTTTCAGT	JF773800	218	mono	-	no amplify	-	no amplify
Ape-58	(CA) ₂₈	F: GGACTCCAGAGACAGTGCAA R: GGACACGCATAGGTGCTTCT	JF773801	155	ambiguous	ambiguous	disomic	ambiguous	disomic
Ape-59	(CA) ₁₁	F: CGTCCTGCTCAAGAAGGTAAA R: CGTCCTGCTCAAGAAGGTAAA	JF773802	110	no amplify	no amplify	no amplify	no amplify	no amplify
Ape-60	(CT) ₂₅	F: TTCAGGGATCCTGTCTCCAG R: GGGGAGCAGTCACAAAGAGT	JF773803	231	ambiguous	mono	no amplify	no amplify	no amplify
Ape-62	(CA) ₅ [(C ₂)(CA) ₂] ₄	F: GACTTCGCCTACAGCAGCTC R: TAGGAACCGGACACGCATAG	JF773804	385	octosomic	tetrasomic	disomic	disomic	disomic
Ape-63	(GGCA) ₆	F: GCACTTTGTTACGGCAGACA R: GACAGGAGGAAATGCTGGAA	JF773805	360	mono	tetrasomic	weak	mono	disomic
Ape-64	(CAGA) ₁₂	F: GAGAGAGGGAGCGACAAACTT R: TAGCTGAGTGGGTGTGGATG	JF773806	213	mono	mono	disomic	mono	weak
Ape-65	(GA) ₁₇ CA (CAGA) ₉ (GA) ₆	F: TTGAACCTTCCACATCCTGA R: CCCAAGGACCTACAGTCTGC	JF773807	154	ambiguous	ambiguous	disomic	ambiguous	weak
Ape-66	(GTCT) ₁₄	F: CAGAAAAACCAGCCCACAGT R: GAGAGAGAGGGAGCGACAAA	JF773808	225	ambiguous	octosomic	disomic	mono	disomic
Ape-68	(GACA) ₅	F: AGTTCGCACTGTAGGGATTCA R: TTCGCAATTAAGGTTAAAAAGACA	JF773809	300	ambiguous	mono	weak	disomic	disomic
Ape-70	(CA) ₁₁	F: AGTGACCCCTCTCTCCACT R: GTCAGGGTCAGGGTCTGTGT	JF773810	166	tetrasomic	mono	mono	mono	mono
Ape-71	(GACA) ₁₅	F: GAGAGAGAGGGAGCGACAAA R: CAGAAAAACCAGCCCACAGT	JF773811	296	octosomic	tetrasomic	mono	mono	-
Ape-73	(GACA) ₇ G 2(CAGA) ₆	F: GAGAGAGAGGGAGCGACAAA R: CAGAAAAACCAGCCCACAGT	JF773812	221	octosomic	octosomic	disomic	mono	no amplify

Ape-76	(GACA)15	F: GAGAGAGAGGGAGCGACAAA R: CAGAAAAACCAGCCCACAGT	JF773813	225	octosomic	tetrasomic	disomic	disomic	disomic
Ape-77	(GA)28	F: ATCTCAGCCAGGAAGAACGA R: CCCGTGTCTGTCTGTCTGTTT	JF773814	171	tetrasomic	disomic	disomic	disomic	ambiguous
Ape-78	(CAGA)6	F: CACTGCCTGCTGCCTAAAAC R: TATTAACCCATCGGCTCCAC	JF773815	151	tetrasomic	tetrasomic	disomic	disomic	disomic
Ape-80	(CTGT)14	F: GGGGTTCAGGAGGCTTTCTA R: GCACTTTGTTCAGGCAGACA	JF773816	228	ambiguous	disomic	_	mono	mono
Ape-81	(GA)28	F: GGTTCCAATGTATCAGGCAAA R: GCCGAGCAGCTCCATTAG	JF773817	152	tetrasomic	_	ambiguou s	_	ambiguous

Because no loci exhibited disomic inheritance in Persian sturgeon, standard tests for deviations from Hardy–Weinberg equilibrium and linkage disequilibrium could not be determined. Fifty two microsatellite primer pairs developed for Persian sturgeon were tested to generate polymorphic genetic markers for four Caspian Sea sturgeon species. In Russian sturgeon, forty six loci were screened in initial screening of samples. Only 32 (83%) of these primer pairs amplified successfully. Of these, 25 loci (54%) were found to be polymorphic in Russian sturgeon. Seven loci were monomorphic while eight loci failed to amplify. Of the 25 polymorphic loci identified, 18 loci were tetrasomic while seven loci were octosomic.

Of the 49 microsatellite loci that were tested in Stellate sturgeon, 39 loci (84%) amplified successfully of which 27 (69%) were polymorphic and seven loci (18%) were monomorphic. All polymorphic loci exhibited disomic banding patterns in stellate sturgeon. Ten loci failed to produce any bands. Forty six loci were tested for cross-species amplification in ship sturgeon. Thirty nine loci (85%) amplified successfully producing 18 polymorphic loci (39%), 13 loci were monomorphic and 8 loci failed to produce any bands. In addition, ambiguous bands were produced at eight loci. All polymorphic loci exhibited disomic banding patterns in Ship sturgeon.

Forty nine loci were screened in Beluga samples. Only 29 loci (83%) amplified successfully. 18 loci (37%) were polymorphic. 11 loci (24%) were monomorphic while 8 loci failed to amplify. All polymorphic loci showed disomic banding patterns. Thus all loci that amplified successfully and that were shown to be polymorphic in ship, stellate and beluga sturgeon species showed simple banding patterns characteristic of disomic loci, while those for Russian sturgeon(like Persian sturgeon) displayed

banding patterns characteristic of tetraploid or higher polyploid karyotypes. Examples of electrophoretic banding patterns at polymorphic loci in the four sturgeon species are presented in Figure 2. Detailed results of cross-species amplification efficiency of the SSR primer pairs developed for Persian sturgeon tested on four Caspian Sea sturgeon species are presented Table 1. Due to the polysomic nature of these loci and the small sample sizes screened in each species, it was considered not possible to test for conformation to hardy-Weinberg equilibrium or heterozygosity per locus. These data will require a more extensive study of larger populations per species.

Discussion

Traditionally, microsatellite marker are developed by extensive screening for microsatellite containing clones through repetitive hybridizations of a repeat motif probe to a large number of random clones (Rassmann et al., 1991). Such an isolation strategy resulted in low rate of the number of positive clones (containing microsatellites) detection. This traditional method usually that can be obtained by means of ranges from 12% to less than 0.04% (Zane et al., 2002).

Using modified protocols of Hamilton et al., (1999) and Glenn et al., (2000) to construct and clone genomic libraries increased proportions of inserts that contained tandem repeat arrays. Thus, a greater number of microsatellite repeat regions detected, sequenced and subsequently used to design species-specific flanking primers for microsatellite amplification. This technique reduced the time and effort as well as cost required for microsatellite isolation from Persian sturgeon. To date there has been no species specific microsatellite primers developed for the Caspian Sea sturgeon species and this is the first report for Persian sturgeon.

Developing microsatellite markers for sturgeon species can be challenging particularly in species that have experienced multiple polyploid events (i.e., 4n, 8n and 16n species) for example, Welsh and May (2006) found only nine reliable disomic microsatellites among 254 primer pairs tested in lake sturgeon (*A. fulvescens*), a species with the same ploidy level as Persian sturgeon.

Amplification results for Persian sturgeon and cross-species amplifications in four Caspian Sea sturgeon species were consistent with the reported ploidy levels of each species. Ship, Stellate and Beluga sturgeon are considered to be functional diploids ($2n= 120$), while Persian and Russian sturgeon are considered to be functional tetraploids ($2n= 240$) that are undergoing rediploidization (Ludwig et al., 2001; Fontana, 2002, Fontana et. al. 2008).

While none of the markers that amplified in Persian sturgeon were disomic, they may still prove to be useful as dominant markers (e.g. Israel et al., 2009) for this species. Several markers appear to show codominant inheritance

patterns in ship, stellate, and beluga sturgeon and may prove useful in a variety of future population genetic applications, ranging from stock assessment to mapping of quantitative trait loci in culture stocks. Testing more individuals and fine tuning optimization of PCR reactions, is likely to identify new alleles at polymorphic loci, as well as the possibility of detecting polymorphisms in loci that were recorded as being monomorphic in small test populations here. Results of these studies suggested that SSR DNA markers developed for Persian sturgeon were candidates for application in other sturgeon species in the Caspian Sea. This proved to be the case and suggests a high level of sequence homology among related species in the Caspian Sea, a result that is consistent with the results from studies on other sturgeon species (May et al. 1997; McQuown et al., 2000).

To eliminate the inherent difficulties associated with tetrasomic loci, future Persian sturgeon genetic marker development required identifying nuclear microsatellite loci that are disomic.





Figure 1: Electrophoretic banding pattern for locus Ape_19 in Persian sturgeon that exhibited octosomic inheritance. Relative allele's density would correspond to gene doses

Lane M: 50 bp DNA step ladder.



1-A: Persian sturgeon

1- B: Russian (lanes 1-5) and Ship (lanes 6-10) Sturgeon.

Figure 2: Electrophoretic banding pattern for locus Ape_20 in Persian (A), Russian (B: 1-5) and ship sturgeon (B: 6-10). This locus exhibited tetrasomy in Persian and Russian sturgeon but was monomorphic inheritance in ship sturgeon. Lane M: 50 bp DNA step ladder

Acknowledgements

The authors thank Mohamad reza Behroze Khoshghalb and Faramarez Bagherzadeh Afrozi for providing sturgeon samples. We thank Peter Mather for helpful comments on the manuscript. This research was funded by the Iranian Fisheries Research Organization (IFRO) and carried out in the Caspian Sea Ecology Research Center and Fish Genetic Laboratories of Southern Illinois University.

References

- Abdolhay, H. A. and Baradaran Tahori, H., 2006.** Fingerling production and Release for Stock Enhancement of Sturgeon in the Southern Caspian Sea: an overview. *Journal of Applied Ichthyology*, **22**, (suppl.1), 121-131.
- An, Z. W., Xie, L. L., Cheng, H., Zhou, Y., Zhang, Q., He, X. G. and Huang, H. S. 2009.** A silver staining procedure for nucleic acids in polyacrylamide gels without fixation and pretreatment. *Analytical Biochemistry*, **391**, 77-79.
- Berg, L. S., 1948.** The Freshwater Fishes of the USSR and Adjacent Countries, 4th edn, Part 1. Akademiya Nauk USSR, Moscow, Leningrad (in Russian, English translation published by Israel Program for Scientific Translations, Jerusalem), part 1. 504p.
- Fontana, F., 2002.** A cytogenetic approach to the study of taxonomy and evolution in sturgeons. *Journal of Applied Ichthyology*, **18**, 226-233.
- Fontana, F., Congiu, L., Mudrak, V. A., Quattro, J. M., Smith, T. I. J., Ware, K. and Doroshov, S. I. 2008.** Evidence of hexaploid karyotype in shortnose sturgeon. *Genome*, **51**, 113-119.
- Glenn, T. C., Cary, T. and Dust, M., 2000.** Microsatellite Isolation. www.uga.edu/srel/DNA_Lab/protocols.htm.
- Hamilton, M. B., Pincus, E. L., Fiore, A. D. and Fleischer, R. C. 1999.** Universal linker and ligation procedures for construction of genomic DNA

- libraries enriched for microsatellites. *BioTechniques*, **27**, 500–507.
- Heist, E. J., Jenkot, J. L. and Keeney, D. B. 2003.** Isolation and characterization of polymorphic microsatellite loci in nurse shark (*Ginglymostoma cirratum*). *Molecular Ecology Notes*, **3**, 59–61.
- Israel, J. A., Bando, K. J., Anderson, E. C. and May, B. 2009.** Polyploid microsatellite data reveal stock complexity among estuarine North American green sturgeon (*Acipenser medirostris*). *Canadian Journal of Fisheries and Aquatic Sciences*, **66**, 1491-1504.
- Khodorevskaya, R. P., Krasikov, E. V., Dovgopol, G. F. and Zhuravleva, O. L. 2000.** Formation of the stock of Caspian Acipenserids under present day conditions. *Journal of Ichthyology*, **40**, 602–609.
- Khosh kholgh, M. R., Pourkazemi, M., Kamali, A. and Rezvani Gilkolaie, S. 2008.** Investigation on genetic structure of Russian sturgeon (*A. gueldenstaedtii*) populations of the north (Volga River) and south Caspian Sea (coastal of Iran and Turkmenistan) using microsatellite techniques. *Iranian Scientific Fisheries Journal*, **16**(4), 68-80.
- Ludwig, A., Belfiore, N. M., Pitra, C., Svirsky, V. and Jenneckens, I. 2001.** Genome duplication events and functional reduction of ploidy levels in sturgeon (*Acipenser*, *Huso* and *Scaphirhynchus*). *Genetics*, **158**, 1203-1215.
- Moghim, M., Kor. D., Tavakolieshkalak, M. and Khoshghalb, M. B. 2006.** Stock status of Persian Sturgeon (*Acipenser persicus* Borodin, 1897) along the Iranian coast of the Caspian Sea. *Journal Applied Ichthyology*, **22**(suppl.1), 99-107.
- Moghim, M., Heist, E. J., Tan, S. G., Pourkazemi, M., Siraj, S. S. and Panadam, J. M. 2009.** Amplification of microsatellite in Persian sturgeon (*Acipenser persicus*). *Iranian Journal of Fisheries Sciences*, **8**(1), 97-102.
- Norouzi, M., Pourkazemi, M., Keyvan, A. and Kazemi, B. 2008.** Population genetic structure of stellate sturgeon (*Acipenser stellatus* Pallas, 1771) in the South Caspian Sea using Microsatellite markers. *Journal of Fisheries and Aquatic Science*, **3** (3), 158-166.
- Pikitch, E.K., Doukakis, P., Lauck, L., Chakrabarty, P. and Erickson, D. L., 2005.** Status, trend and management of sturgeon and paddlefish fisheries. *Fish and Fisheries*, **6**, 233–265.
- Pourkazemi, M., 2007.** Comprehensive Study on Assessment of Sturgeons Population Genetic Structure in the Caspian Sea. International Sturgeon Research Institute. 316p.
- Rassmann, K., Schlotterer, C. and Tautz, D., 1991.** Isolation of simple sequence loci for use in polymerase chain reaction-based DNA fingerprinting. *Electrophoresis*, **12**, 113-118.
- Safari, R., Pourkazemi, M., Rezvani, S. and Shabani, A., 2008.** Population structure of *Acipenser nudiventris* in the south coast of Caspian Sea and oral river using microsatellite method. *Iranian scientific fisheries journal*, **17** (1), 99-108.
- Welsh, A., May, B., 2006.** Development and standardization of disomic microsatellite markers for lake sturgeon genetic studies, *Journal of Applied Ichthyology*, **22**, 337-344.
- Zane, L., Bargelloni, L. and Patarnello, T., 2002.** Strategies for microsatellite isolation: a review. *Molecular Ecology*, **11**, 1-16.

شناسایی و جداسازی جایگاه های ریز ماهوار تاسماهی ایرانی (*Acipenser persicus*, Borodine, 1897) و بررسی امکان تکثیر آنها در ژنوم چهار گونه ماهیان خاویاری دریای خزر

مهدی مقیم^{1,3*}؛ ادوارد جی هیست²؛ سون گوان تن³؛ محمد پور کاظمی⁴؛ سیتی شاپور سراج⁵؛ جوتی مالار پانادام⁶؛ رضا پورغلام¹؛ داود کر¹؛ فرامرز لالوئی¹؛ محمدجواد تقوی¹

چکیده

به منظور مدیریت پایدار ذخایر تاس ماهی ایران که یک گونه با ارزش تجاری در جنوب دریای خزر است، ما نیاز به شناسایی جمعیت ها، ساختار جمعیتی و همچنین وضعیت حفاظتی آنها در زیستگاه طبیعی شان داریم. همچنین برای توسعه یک برنامه حفاظتی برای تاس ماهی ایران در دریای خزر نیاز به آگاهی از تنوع ژنتیکی آن داریم که داد های آن با استفاده از نشانگر مولکولی قابل اعتمادی جمع آوری شده باشد. نشانگر ملکولی ریزماهواره یا میکروستلایت، نشانگر مناسبی برای این منظور می باشد. برای این منظور، یک کتابخانه غنی شده از DNA تاس ماهی ایران بر اساس روش جذب بیوتین آماده شد. حدود ۱۸۰۰ کلونی سفید از کتابخانه ژنومی تاس ماهی ایران جدا سازی شد و برای کنترل وجود تکرار متوالی نوکلئید ها یا ریزماهواره غربال شدند. از این تعداد ۳۵۰ کلونی شناسایی و تعیین توالی شدند. از بین آن ها ۸۱ کلونی با داشتن ریزماهواره و مکان پهلوگیری مناسب (یا مناطق flanking) شناسایی شدند و ۶۸ جفت آغازگر ریزماهواره توسعه (develop) یافت. نتایج آزمایش PCR آغازگر ها با نمونه DNA تاس ماهی ایران نشان داد که از ۶۸ جفت آغازگر، ۶ آغازگر جایگاه مونومورف یا تک شکلی (monomorphic)، ۲۰ آغازگر جایگاه چند شکلی تتراسومیک (tetrasomic) و ۱۸ آغازگر جایگاه چند شکلی اکتاسومیک (octosomic) را تکثیر کردند. ۲۴ آغازگر هیچ جایگاهی را تکثیر نکردند یا الگوی باند ها ضعیف و مبهم بودند. اگرچه هیچ یک از آغازگرها جایگاه دیسومیک (disomic) در تاس ماهی ایران و تاس ماهی روس (A. *gueldenstaedtii*) نشان ندادند، تعدادی از نشانگرها برای مطالعات ماهیان خاویاری ازون برون (*A. stellatus*)، شیپ (*A. nudiventris*) و فیل ماهی (*Huso huso*) مناسب ظاهر شدند. باندهای حاصل از آغازگر ها در ازون برون، شیپ و فیل ماهی الگوی جایگاه های چند شکلی ساده که مشخصه جایگاه دیسومیک (disomic) است را نمایش دادند، در حالی که برای تاس ماهی ایران و تاس ماهی روس الگوی باندهای حاصل جایگاه های پلی سومیک چهار تایی یا بیشتر را نمایش دادند. این نشانگرها برای مطالعات ژنتیک جمعیت انواع مختلف ماهیان خاویاری دریای خزر مفید خواهند بود.

واژگان کلیدی: تاس ماهی ایران، *Acipenser persicus*، دریای خزر، ریزماهواره، ژنتیک جمعیت.

۱-بخش ژنتیک پژوهشکده اکولوژی دریای خزر، ساری، ایران. صندوق پستی: ۹۶۱، ساری، ایران.

۲-مرکز ماهی شناسی و آبی پروری ایلینویز، بخش جانورشناسی دانشگاه ایلینویز جنوبی، کاربوندل، ایلینویز، آمریکا، کدپستی: ۶۵۰۱-۶۲۹۰۱، آمریکا.

۳-بخش سلول و بیولوژی مولکولی دانشکده بیوتکنولوژی دانشگاه پوترا، کد ۴۳۴۰۰، سردانگ، سلانگور، مالزی.

۴-انستیتو تحقیقات بین المللی ماهیان خاویاری دکتر دادمان، کدپستی: ۳۴۶۴-۴۱۶۳۵، رشت، ایران.

بخش آبی پروری دانشکده کشاورزی دانشگاه پوترا، کد ۴۳۴۰۰، سردانگ، سلانگور، مالزی.

بخش دامپروری دانشکده کشاورزی دانشگاه پوترا، کد ۴۳۴۰۰، سردانگ، سلانگور، مالزی.

* پست الکترونیکی نویسنده مسئول Mehdi.moghim@gmail.com