Effects of different protein sources on growth performance and food consumption of goldfish, *Carassius auratus*

Bilen S.^{*}, Müge Bilen A.

Received: February 2012

Accepted: August 2013

Kastamonu University Inebolu Faculty of Fisheries Kastamonu/ Turkey

*Corresponding author's email: sbilen@kastamonu.edu.tr

Keywords: Aquarium, Goldfish, Carassius auratus, Food consumption, Growth performance

Protein is one of the most important components in a fish diet (De Silva and Andersoni, 1995; Kaushik, 1995). In aquaculture, the primary dietary animal protein source is fish meal, but its availability is limited and supply varies because of reductions in fish stocks related to factors such as climatic phenomena, and overexploitation and decline of ocean fisheries stocks. This variability can seriously affect aquaculture sustainability and provability, and therefore research in identifying alternative dietary protein sources has increased (Kissil et al., 2000; Maylor et al., 2000). Fish nutritionists have tried to use less expensive plant protein sources to partially or totally replace fish meal (Xie et al., 2000). In the context of research on substitutes for fish meal in diets of fish, numerous studies have been made in the past decades (Kaushik et al., 1995; Mambrini et al., 1999).

The market for ornamental fish in Turkey is increasing. To cater to the demand of quality feed for these species, a number of commercial feeds are available in the market. Feed formulation for goldfish is yet to be standardized. As a common practice in Turkey, commercial production of aquarium fish depends essentially on pelleted/granular form of fish feed.

Goldfish, Carassius auratus are the most popular choice of fish for a water garden. It can be found in a number of varieties and colors. This species reaches sexual maturity at a small size and the artificial insemination of goldfish eggs and rearing of larvae is comparatively easy (Wiegand et al., 1989). Therefore the goldfish provides a useful model for the study of nutrition in larval and juvenile Cyprinids. Some initial studies have been made on the nutritional requirements of warm water omnivore (Sales and Janssens, 2003).

The purpose of this study was to investigate food consumption and growth performance of goldfish fed diets containing only plant, only animal and plant-animal (50%-50%) derived protein source.

In the study, goldfish fingerlings were obtained from Kepez Research Institute of Aquatic Resource (Antalya/Turkey). The fish were acclimatized to experimental conditions for a period of two weeks before the start of the experiment. During this period, they were fed on a practical diet (Pinar Feed, 45% crude protein, 18% lipid) to satiation twice a day. In the trial, 81 fishes were divided into 9 aquariums (60 L for each aquarium) for the application of experimental diets. The aquariums were aerated and filtrated with GAP II filter. Water temperature was 25 °C and pH 7.3 was around. The fish were exposed to normal photoperiod – 9 Light: 15 Dark for 8 weeks. Each was weighted (average weight of 4.8 ± 0.05 g) at the beginning of the experiment. During feeding trial, fish were fed to satiation twice a day (at 08:30 and 15:30 h). The daily feed supplied was

recorded and unconsumed feed were collected 20 min after feeding, and dried. The trial lasted for 8 weeks. At the end of the trial, all fish were weighted individually. Five fish from each tank were taken for proximate analysis. Chemical composition and formulation of three experimental diets are given in Table 1.

Drotain course			
FIOLEIII SOUICE	Plant	Animal	Plant-Animal
Wheat Starch	0	35	17
Corn Gluten Meal	29.45	0	15.00
Fish Meal	0	52	26
Wheat Meal	14.97	0	7.50
Soybean Meal	29.45	0	15.00
Sunflower Meal	14.97	0	7.50
Soybean Oil	4.99	0	6.00
¹ Vit-Min Premix	2.99	3	3.00
Binder	2.99	3	3.00
Fish Oil	0	0	7
Lysin	0.17	0	0.00
Crude Moisture	4.04 <u>+</u> 0.50	4.04 ± 0.07	3.66 <u>+</u> 0.18
Crude Protein	36.32 <u>+</u> 0.24	37.08 <u>+</u> 0.34	36.05 <u>+</u> 0.13
Crude Lipid	10.67 <u>+</u> 0.40	11.63 <u>+</u> 0.50	10.69 <u>+</u> 0.60
Crude Ash	6.50 <u>+</u> 0.28	8.41 <u>+</u> 0.39	8.41 <u>+</u> 0.39
Gross Energy (MJ/kg)	20.82	20.75	20.48

Table 1: Diet formulation and chemical composition of experimental diet (% of dry wieght)

¹ Vit-Min Premix (mg kg ¹, NRC 1977): vitamin A, 5500IU; vitamin D₃, 1000 IU; vitamin E, 50 IU; vitamin K, 10 mg; choline, 550 mg; niacin, 100mg riboflavin 20mg; pyridoxine, 20 mg; thiamine 20mg; biotin, 0,1mg; folacin, 5mg; B₁₂, 20μg; inositol, 100 mg; choline chloride, 5000 mg. Mineral premix (mg kg ⁻¹ diets, H440): NaCl, 1,0; MgSO₄, 7; NaH₂PO₄ 25; KIO₃ 0,0003; ZnSO₄ 0,353; MnSO₄0,162.

Practical diets with three dietary animalplant protein source ratios 1:0, 0:1 and 1:1 (animal, A; plant, P and animal-plant, A-P) were formulated. Dry ingredients were blended to make a homogenous mixture and were mixed with boiled water. After moderate cooling, pellets (1 mm size) were prepared by over-dried at 40 °C and stored at 4 °C.

Moisture, crude protein, lipid and ash of the experimental feed and fish were determined by methods as described in A.O.A.C. (1995) methods. The moisture content was analyzed by drying in a hot air oven at 105 °C. Crude protein was estimated by the Kjeldahl method and crude lipid was estimated by soxhlet extraction with petroleum ether. Ash contents were determined from the residue remaining after incineration of the samples at 550 °C in a muffle furnace. Dietary gross energy was calculated using the conversion factors of 23.7 kJ g for protein,

719

39.5 kJ g for lipid and 17.2 kJ g 1 for carbohydrate (Brett and Groves, 1979). The growth parameters were calculated as follows: Specific Growth Rate (SGR) (% day⁻¹) = 100 x [(ln final fish weight) – (ln initial fish weight)] / days fed. Feed Intake (FI) (%) = daily feed intake (g) x 100 / biomass (g). Feed Conversion Ratio (FCR) = feed intake (g) / weight gain (g) x 100. Protein Efficiency Ratio (PER) = wet body mass gain/crude protein intake. Analyses were conducted using one-way ANOVA. For multiples comparisons of the means Duncan's multiple range was used (Zar, 1999). All the results were treated significant at the 5% level. The results of the study have been shown in Table 2.

1				
	А	Р	P-A	±SEM*
Survival (%)	100	100	100	-
Initial Mean Weight (g)	4.8	4.8	4.8	0.05
Final Mean Weight (g)	10.1 ^c	6.4 ^a	8.6^{b}	0.54
Feed Intake (%)	2.57 ^a	2.34 ^b	2.21 ^b	0.05
Weight Gain (%)	110.4 ^c	33.3 ^a	79.2 ^b	1.05
Feed Conversion Ratio (%)	2.9 ^a	$5.7^{\rm b}$	3.0 ^a	0.29
Specific Growth Ratio (%)	1.2^{c}	0.5^{b}	1.0^{a}	0.04
Protein Efficiency Ratio (%)	0.9^{a}	0.5^{b}	0.9^{a}	0.47

 Table 2: Growth performance, feed and nutrient utilization of goldfish fed the experimental diets

* Values in each row allocated common superscripts or without superscripts are not significantly different from each other (P > 0.05).

There were no statistically significant differences in survival rate (P>0.05) among the dietary treatments. Live weight SGR significantly gain and higher (P>0.05) for diet A than diets P and A-P. The mean feed intake of fish fed with diet A was significantly (P<0.05) higher than that of P and A-P fed fish. The FCR of P treatment was found highest (P>0.05), but there were not significant differences between A and A-P groups (P>0.05). The protein efficiency ratio of A and A-P groups approximately 80% higher than that

of fish fed with the P diet. The proximate compositions of the fish carcasses at the end of the trial are given in Table 3. There were no significant differences (P>0.05) between groups in crude lipid. However, dry matter of A and A-P group were significantly lower than P group. In terms of crude protein, P and A-P were similar (P>0.05), but A was higher than the other groups (P<0.05). Ash content of P and A were established similar and lower than A-P (P<0.05).

	•	-		
Grups	Dry Matter	Crude Protein	Crude Lipid	Ash
Р	69.74 <u>+</u> 3.77 ^a	40.92 ± 1.4^{a}	37.15 <u>+</u> 68 ^a	2.72 ± 52^{a}
А	65.18 <u>+</u> 1.31 ^b	43.16 <u>+</u> 31 ^b	34.79 <u>+</u> 1.21 ^a	2.65 ± 22^{a}
A -P	63.63 <u>+</u> 1.34 ^b	39.56 <u>+</u> 3.26 ^a	35.77 <u>+</u> 95 ^a	3.53 <u>+</u> 07 ^b

Table 3: Proximate analysis of goldfish fed different protein sources

Values are in percentage wet weight. A: Animal source; P: Plant source; A-V: Animal and Plant source (% 50-50). Means with different superscripts (a, b, c) are significantly different from each other (P<0.05).

In most studies, decreased growth was observed in fish when a large proportion of the fish meal was replaced by plant proteins (Fournier et al., 2004; Espe et al., 2006, Palti et al., 2006). Factors suggested to be the causes for decreased growth in fish fed plant protein include low palatability (Mohsen and Lovell, 1990), deficiency amino acid such as methionin and lysine (Dabrowski et al., 1989; Tantikitti and Chimsung, 2001), poor utilization of protein (Boonyaratpalin et al., 1998), and low energy (Opstvedt et al., 2003). Generally, it has been examined that a partial replacement of fish meal by plant protein source is well supported by fish, so currently more than 30% is already replaced in most commercial fish feeds (Forster et al., 1999; Burel et al., 2000; Vielma et al., 2000; Opstvedt et al., 2003). The goldfish in the experimental are omnivore species but adult goldfish prefer to eat plant; the young ones may require a greater amount of protein in the diet for optimum growth. The inclusion of fish meal in the diet of young goldfish, with a maximum digestibility of 93%, could promote their growth of young goldfish (Degani et al., 1997; Bandyopadhyay et al., 2005). Growth in terms of weight gain (WG) and specific growth rate (SGR) were highest in fish fed with animal protein and lowest in fish fed with plant protein diet (Table 2) during the study period (P<0.05). The lowenergy digestibility of plant feed ingredients can be attributed to their high-carbohydrate content and poor digestibility by carnivorous fish (Lupatsch et al., 1997). However, Abi-Avad and Kestemont (1994) reported that the growth performance were highest on Carassius auratus gibelio when the fish fed feed animal protein, that results also similarly to our results. In the present study, total feed intake was lower in goldfish fed

the plant and animal-plant diet than in the animal diet. Goldfish fed the animal diet and animal-plant diets had lower FCR than plant diet. The lower feed intake and highest FCR in plant diet fed fish were probably mainly caused by the existence of anti-nutritional factors and/or low palatability (Watanabe et al., 1992). Decreased growth of fish fed the diet with the inclusion of plant was reported to be due to low feed intake (Higgs et al., 1982), low digestibility of dry matter and/or protein (Hilton and Slinger, 1986), unbalanced amino acids and/or high fiber contents (Appler and Jauncey, 1983). In conclusion, the present study showed that inferior growth of fish fed animal protein was caused by different factors in goldfish fed diets with different protein contents.

References

- Abi-Ayad, A. and Kestemont, P., 1994. Comparison of the nutritional status of goldfish (*Carassius auratus*) larvae fed with live, mixed or dry diet. *Aquaculture*, 128: 163-176.
- A.O.A.C., 1995. Official methods of analysis of the association of the official analysis chemists. Association of Offical Analytical Chemists, Arlington, VA, USA.
- Appler, H. N. and Jauncey, K., 1983. The utilization of a filamentous green algae (*Cladophora glomerata* (L) Kutzin) as a protein source in pelleted feeds for *Sarotherodon niloticus* (Tilapia) fingerlings. *Aquaculture*, 30, 21-30.
- Bandyopadhyay, P., Swain, S. K. and Mishra, S., 2005. Growth and dietary utilizations in goldfish (*Carassius auratus* L.) fed diets formulated with various local agro-produces. *Bioresource Technology*, 96, 31-740.

- Boonyaratpalin, M., Suraneiranat, P. and Tunpibal, T., 1998. Replacement of fish meal with various types of soybean products in diets for the Asian sea bass. *Aquaculture*, 161, 67-78.
- Brett, J. R. and Groves, T. D. D., 1979. Physiological energetics, In: Fish Physiology. Bioenergetics and Growth. (Hoar, S., Randall, D. J. & Brett, J. R. eds), Academic PresS, New York: NY, pp. 279–352.
- Burel, C., Boujard, T., Tulli, F. and Kaushik, S. J., 2000. Digestibility of extruded peas, extruded lupin, and rapeseed meal in rainbow trout (*Oncorhynchus mykiss*) and turbot (*Psetta* maxima). Aquaculture, 188, 285-298.
- Dabrowski, K., Poczyczynski, P., Köck, G. and Berge, B., 1989. Effect of partially or totally replacing fish meal protein by soybean meal protein on growth, food utilization and proteolytic enzyme activities in rainbow trout (*Salmo gairdneri*). New in vivo test for exocrine pancreatic secretion. *Aquaculture*, 77, 29– 49.
- **De Silva, S. S. and Andersoni, T. A., 1995.** Fish Nutrition in Aquaculture. Chapman & Hall, 2-6 Boundary Row, London SE1 8HN, 319 p.
- Degani, G., Viola, S. and Yehuda, Y.,
 1997. Apparent digestibility coefficient of protein sources for carp *Cyprinus carpio* L. *Aquaculture Resource*, 28, 23–28.
- Espe, M., Lemme, A., Petri, A. and El-Mowafi, A., 2006. Can Atlantic salmon (*Salmo salar*) grow on diets devoid of fish meal? *Aquaculture*, 225, 255-262.
- Forster, I., Higgs, D. A., Dosanij, B. S., Rowshandeli, M. and Parr, J., 1999. Potential for dietary phytase to improve the nutritive value of canola protein

concentrate and decrease phosphorus output in rainbow trout (*Oncorhynchus mykiss*) held in 11 °C fresh water. *Aquaculture*, 179, 109-125.

- Fournier, V., Huelvan, C. and Desbruyeres, E., 2004. Incorporation of a mixture of plant feed stuffs as substitute for fish meal in diets of juvenile turbot (*Psetta maxima*). Aquaculture, 236, 451– 465.
- Higgs, D. A., McBride, J. R., Markert, J.
 R., Dosanjh, B. S., Plotnikoff, M. D.
 and Clarke, W. C., 1982. Evaluation of tower and candle rapeseed (canola) meal and bronowski rapeseed protein concentrate as protein supplements in practical dry diets for juvenile chinook salmon (*Oncorhynchus tshawytscha*). Aquaculture, 29, 1-31.
- Hilton, J. W. and Slinger, S. J., 1986. Digestibility and utilization of canola meal in practical-type diets for rainbow trout. *Canadian Journal of Fisheries and Aquatic Sciences*, 43, 1149-1155.
- Kaushik, S. J., 1995. Nutrient requirements, supply and utilization in the context of carp culture. *Aquaculture*, 129, 225-241.
- Kaushik, S. J., Cravedi, J. P., Lalles, J. P., Sumpter, J., Fauconneau, B. and Laroche, M., 1995. Partial or total replacement of fish meal by soybean protein on growth, protein utilization, potential estrogenic or antigenic effects, cholesterolemia and flesh quality in rainbow trout. *Aquaculture*, 133, 257– 274.
- Kissil, G. W. M., Lupatsch, I., Higgs, D. A. and Hardy, R. W., 2000. Dietary substitution of soy and rapeseed protein concentrates for fish meal, and their effects on growth and nutrient utilization

in gilthead seabream, *Sparus aurata* L. *Aquaculture Resource*, 31, 595-602.

- Lupatsch, I., Kissil, G. W., Sklan, D. and Pfeffer, E., 1997. Apparent digestibility coefficients of feed ingredients and their predictability in compound diets for gilthead seabream, *Sparus aurata* L. *Aquaculture Nutrition*, 3, 81– 89.
- Mambrini, M., Roem, A., Cravedi, J. P., Lalles, J. P. and Kaushik, S. J., 1999. Effects of replacing fish meal by soy protein concentrate and of DL-methionine supplementation in high energy extruded diets on growth and nutrient utilization in rainbow trout, Oncorhynchus mykiss. Journal of Animal Science, 77, 2990– 2999.
- Maylor, L. R., Goldburg, J., Primavera,
 H. J., Kautsky, N., Beveridge, M. C.
 M., Clay, J., Folke, C., Lubehenco, J.,
 Money, H. and Troell, M., 2000. Effect of aquaculture on world fish supplies. *Nature*, 405, 1017-1024.
- Mohsen, A. A. and Lovell, R. T., 1990. Partial substitution of soybean meal with animal protein sources in diets for channel catfish. *Aquaculture*, 90, 303-311.
- **Opstvedt, J., Aksnes, A., Hope, B. and Pike, I. H., 2003.** Efficiency of feed utilization in Atlantic salmon (*Salmo salar* L.) fed diets with increasing substitution of fish meal with vegetable proteins. *Aquaculture*, 221, 365–379.
- Palti, Y., Silverstein, T. J., Wieman, H.,
 Phillips, G. J., Barrows, F. T. and
 Parsons, J. E., 2006. Evaluation of
 family growth response to fishmeal and
 gluten-based diets in rainbow trout
 (Oncorhynchus mykiss). Aquaculture,
 255, 548-556.

- Sales, J. and Janssens, G. P. J., 2003. Nutrient requirements of ornamental fish. *Aquatic Living Resources*, 16, 533-540.
- Tantikitti, C. and Chimsung, N., 2001. Dietary lysine requirement of freshwater catfish (*Mystus nemurus* Cuv. & Val.). *Aquaculture Resource*, 32, 135–141.
- Vielma, J., Makinen, T., Ekholm, P. and Koskela, J., 2000. Influence of dietary soy and phytase levels on performance and body composition of large rainbow trout (*Oncorhynchus mykiss*) and algal availability of phosphorus load. *Aquaculture*, 183, 349-362.
- Watanabe, T., Viyakarn, V., Kimura, H., Ogawa, K., Okamoto, N. and Iso, N., 1992. Utilization of soybean meal as a protein source in a newly developed softdry pellet for yellowtail. *Nippon Suisan Gakkaishi*, 58, 1761–1773.
- Wiegand, M. D., Hataley, J. M., Kitchen, C. L. and Buchanan, L. G., 1989. Induction of developmental abnormalities in larval goldfish, *Carassius auratus* (L) under cool incubation conditions. *Journal* of Fish Biology, 35, 85-95.
- Xie, S., Zhu, X., Cui, Y. and Yang, Y., 2000. Utilization of several plant proteins by gibel carp (*Carassius auratus gibelio*). *Journal Applied Ichthyology*, 17, 70-76.
- Zar, H. J., 1999. Biostatistical Analysis. Prentice Hall (Upper Saddle River, N.J.), 929 p.