

Using RAPD markers potential to identify heritability for growth in *Fenneropenaeus indicus*

Rezvani Gilkolaei, S.^{1*}; Safari, R.²; Laloei, F.³; Taqavi, J.³; Matinfar, A.¹

Received: April 2009

Accepted: January 2010

Abstract

Sampling was done using 90 post larvae which were produced by reproduction of some broodstocks of *Fenneropenaeus indicus* in one day and reared in the same situation for 4 months. Samples were divided into 3 groups: high, medium and low growth (based on weight and length). Genomic DNA was extracted from muscle tissue using the phenol-chloroform method. The polymerase chain reaction (PCR) was carried out using 21 RAPD loci and PCR products were separated on 3% Agarose gel. From 21 loci studied, 12 produced polymorphic bands. The most polymorphic produced bands using OPAQ 9 and the least by OPAQ 7. Search for specific markers in *F. indicus* one specific band was observed in the low growth group using OPAQ4. The highest genetic distance (0.457) was between the low growth group and the medium and the lowest (0.091) between high growth and medium groups, therefore the highest genetic identity (0.912) was between high growth and medium groups and the lowest (0.633) between low growth group and the medium. Neighbor-joining resulted in two groups, the first including high and medium growth groups and the second low growth group. It appears that low growth group depended on separated population. Considering the mean weight of F₁ (mean weight of 90 specimens) (16.25±1.5 g), parental generation mean weight of 15 ±1.2 and mean weight of parent 31.6 g, response to selection (R) and heritability for growth in this species were estimated to be 1.2±0.2 and 0.07±0.01 respectively.

Keywords: *Fenneropenaeus indicus*, RAPD marker, Heritability, Growth, Iranian Fisheries

1-Iranian Fisheries Research Organization, P.O. Box: 14155-6116, Tehran, Iran.

2-Department of Fisheries, Golestan University, Golestan, Iran.

3-Genetic department, Ecology Research Institute of the Caspian Sea. Sari, Iran.

*Corresponding author's email: rezvani@ifro.ir

Introduction

The Indian white shrimp (*Fenneropenaeus indicus*) inhabits in the Indian Ocean coast from north of South Africa to India, South-East Asia, Indonesia, and Northern Australia (Benzie, 2009). In Iran this species has been distributed in the Persian Gulf and Oman Sea. *F. indicus* is non-burrowing, active in both day and night and prefers a sandy mud bottom (Afsharnasab et al., 2005). Its maximum total length is 184 mm (male) and 228 mm (female); its maximum carapace length is 56 mm (FAO, 2009). The shrimp mature and breed mostly in marine habitats and spend the juvenile and sub-adult stages in coastal estuaries from 2 to 90 m brackish waters or lagoons (FAO, 2009). Many natural shrimp stocks are in decline and farmed shrimps provided more than 50% of the world's production for several years (Benzie, 2009). Despite the high potential of aquaculture for increasing production, the sustainability of shrimp farming is treated by low production efficiency (Hetzl et al., 2000) and vulnerability to diseases (Afsharnasab et al., 2005; Gitterle et al., 2005; Lu and Sun, 2005; Olivier and Roel, 2009). The development of genetically improved stocks and domesticated breeds selected for commercial traits is one approach to overcoming these threats (Moss et al., 2007; Hoa, 2009; Pourkazemi et al., 2010), but the high coefficient of variation that comes from the strong environmental influence on the phenotypes make the selection of individuals with genetic advantages for these traits difficult (Falconer, 1998; Hoa, 2009) then the establishment of genetic marker to assist in selecting individuals for breeding would

be of great benefit in speeding the domestication of these species (Moore et al., 1999; Donato et al., 2005; Benzie, 2009). The benefits of domestication have been demonstrated in some aquatic species such as Rainbow Trout (Gjedrem, 1992), Atlantic Salmon (Gjedrem and Finland, 1995), Pacific Oyster (Taris et al., 2007) and Gaint Catfish (Kednapat et al., 2007). But despite the amenity of shrimps to genetic selections and because of their comparatively high fecundity, short generation time and larger genetic gain (Keys et al., 2004), the shrimp farming industry has been slow to adopt selective breeding programs (Goyaed et al., 2008; Cock et al., 2009). It is said by Pullin et al. (1998) that this reluctance has been due in part to the past perceptions of low genetic variability and difficulties in the domestication of shrimps (Benzie, 2009). Hence, it is necessary to challenge these perceptions and to quantify the responses to selection and heritability estimates for commercial traits. Low levels of protein variation and highly unlikely benefits of allozyme variation have been reported by Garcia and Benzie (1995) in shrimps. Although subsequent studies, using allozyme (Benzie et al., 1997; De la Rosa-Velez et al., 1999) have revealed considerable genetic variation within populations, but DNA analysis [RAPD (Garcia and Benzie, 1995); microsatellite (Wolfus et al., 1997); mtDNA and microsatellite (You et al. 2008)] would provide a better source of markers in penaeid prawns. Garcia and Benzie (1995) in a study of RAPD markers of potential use in penaeid prawn (*P. monodon*) breeding programs found that levels of

variation obtained by RAPD method is similar to those observed in other taxa, and are likely to be adequate for obtaining markers to assist selective breeding programs. RAPD producers were first developed by Welsh and McClelland; Williams et al.(1990) using PCR randomly amplifying anonymous segments of nuclear DNA with an identical pair of primers 8-10 bp in length that have been used as useful markers in breeding programs and gene mapping. The objective of this study was to assess RAPD markers in screening the high growth of wild and cultured brood stock before reaching the final maturation stage.

Material and methods

Sample collection

Sampling was done using muscles of 90 post larvae which were produced by reproduction of some brood stocks of *F. indicus* in one day and reared in the same conditions for 4 months in Bushehr province during Feb.-Aug. 2007. Samples were classified in 3 groups, high growth, medium and low growth. At first all samples were arranged in a row according to their size and divided into three groups, the first ten, low growth group; the second, medium and the third high growth group. Then the groups were weighed, the average weight for the selected low growth prawns was 13.74 ± 1.95 gr, medium growth 15.75 ± 1.5 gr and high growth 19.1 ± 1.4 gr and using SPSS there was a significant difference between them (Table 1). Selection response and heritability were measured for growth. Selection response was measured by the difference between mean body weight of the first and parental generations. Selection differential

was calculated by the difference between the mean body weight of the selected parents and the parental generations. Real heritability was calculated by the ratio of selection response to selection differential (Falconer, 1998).

DNA extraction

Genomic DNA was extracted from a 1cm^2 (50-60 mg) piece of muscle tissue using the phenol-chloroform procedure described by Hillis et al. (1996). The quality and concentration of DNA from samples were assessed by electrophoresis on a 1% agarose gel and Spectrophotometer, and then the samples were stored at -20°C for further analysis.

PCR amplification and electrophoresis

DNA from each individual was amplified through PCR. A single 10 nucleotides Oligomer of random sequence which existed in a general random amplified polymorphisms Kit (Table 2) were used for each reaction. A total of 12 different primers were tested. Each PCR reaction (final volume 25 μl) was composed of 5 μl of 10X reaction buffer, dNTPs 10 mM, MgCl_2 50 mM, primer 20 pmol, genomic DNA 100ng and 1.5-2 units of Taq polymerase. The temperature profile consisted of a 3-min initial denaturation at 94°C , followed by 30 cycles of 30 s at 94°C , 30 s at the respective annealing temperature, and 1 min at 72°C , ending with 5 mins at 72°C (Master cycler Eppendorf model 384). PCR products were separated on 3% agarose gel, and stained with ethidium bromide. RAPD PCR which was carried out using different PCR blocks and slightly varying conditions showed that the RAPD patterns were robust and reliably replicated.

Statistical analyses were carried out using Pop Gene Version (1.31) (Yeh et

al., 1999) and SPSS Version (10.5). Heterozygosity, genetic distance, identity between three groups and Dendrogram Based on (Nei, 1972) Genetic distance were measured using Pop Gene and selection response and heritability for weight were estimated using SPSS.

Results

From 21 studied loci, 12 produced polymorphic bands. The most produced band using OPAQ10 and the least by OPAQ11. The most polymorphic band produced using OPAQ9 and the least by OPAQ7 (Table 3). The estimated level of polymorphism in this study was 22%. Searching specific markers in *F. indicus* one specific band was observed just in low growth group using OPAQ4 (Fig. 1). The highest genetic distance (0.457) was

between low growth and medium groups and the lowest (0.091) between high growth and medium groups, therefore the highest genetic identity (0.912) was between high growth group and medium and the lowest (0.633) between low growth group and medium groups (Table 4). Neighbor-joining tree (Fig. 2) resulted in two groups, the first including high and medium growth groups and the second low growth group, it appears that low growth groups are depended on separated population of the two others. With considering the mean weight of F_1 (16.25 ± 1.5), mean weight of 15 ± 1.2 and mean weight of parent 31.6, response to selection (R) and heritability for growth in this species were estimated 1.2 ± 0.2 and 0.07 ± 0.01 , respectively.

Table1: Mean \pm SD weight for different groups

Factor	High growth	Medium growth	Low growth
Weight	19.1 ± 1.4 c	15.75 ± 1.53 b	13.74 ± 1.95 a

Different superscripts indicate significant differences ($p \leq 0.05$)




Figure 1: PCR product of *F.indicus* using OPAQ4, 13 samples of 3 groups (Small, Medium and Large), arrows show specific bands in small groups

Table2: List of the 12 primers used to amplify RDPD markers in *F. indicus*


Primer	Anneling (°C)	3'→5'
OPAQ1	35	GGTGGCGGGA
OPAQ2	39	GAGGTCCAGA
OPAQ3	37	GCTGCTGGAG
OPAQ4	37	GCTGTAGTGT
OPAQ5	35	GCGGTTGAGG
OPAQ6	34	CAAGGGAGGT
OPAQ7	35	GGGCACGCGA
OPAQ8	32	ACGGCCGACC
OPAQ9	51	CGGAGAGCGA
OPAQ10	50	TGGGCTCGCT
OPAQ11	53	ACTTGTGCGG
OPAQ12	53	GCGGGAGACC

Table3: Produced bands, Polymorphic bands and Monomorphic bands used primer

Primer's number	Polymorphic bands	Monomorphic bands	Produced bands
OPAQ1	4	3	7
OPAQ2	-	6	6
OPAQ3	3	3	6
OPAQ4	4	2	6
OPAQ5	2	4	6
OPAQ6	2	4	6
OPAQ7	1	5	6
OPAQ8	5	2	7
OPAQ9	8	-	8
OPAQ10	7	2	9
OPAQ11	5	-	5
OPAQ12	3	3	6
Total	43	34	78

Table 4: Distance matrix, upper rectangle is Nei's identity and lower rectangle is Nei's distance

	Medium growth	High growth	Low growth
Medium growth	***	0.912	0.633
High growth	0.0916	***	0.6469
Low growth	0.4571	0.4355	***

**Figure 2: Dendrogram Based Nei's Genetic distance: Method= UPGMA
Modified from Neighbor procedure of Phylip Version 3.5**

Discussion

Hetzel et al. (2000) did their experiments in different rearing environments of parent and offspring generations as they didn't find any reports in the literature on genetic correlations between growths of prawns in pond vs. tanks. Preston et al. (2004) in the study of growth rate in selected and non-selected *Metapanaeus japonicus* in a controlled environment and open pond found there is no significant differentiation between the two groups when the environment condition is in favorite range for this species. This study was done in open pond in environment condition in favorite range for this species during experiment time. Garcia and Benzie (1995) observed a specific band in one of the studied families, screened all families to find whether the band is truly family-specific marker, but results showed it was probably derived from bacterial or algal epicommsals contaminating the sample. Wolfus et al. (1997) in the study of specific markers in *Litopanaeus vannamei* found 23

specific markers in this species 2 of which were observed in single families from one population. In the current study in searching of specific markers in *F. indicus* one specific band was observed only in the low growth group, and we are continuing tests on sequencing specific bands to develop suite markers for this purpose. On the other hand according to Dendrogram Based Nei's Genetic distance, UPGMA method, it appears that low growth group is dependent on separated population of the two others. Nelson and Hedgecock, (1980) using Alozyme studies suggested that prawn have generally low levels of variation. Garcia and Benzie (1995) using RAPD reported 6-7% polymorphism in *Panaeus monodon* and suggested that randomly amplified polymorphic DNA (RAPD) approaches will be as useful in providing markers for prawn breeding programs as they have been for other species. Wolfus et al. (1997) saw high levels of heterozygosity (45-100%) in *L.*

vannamei and found the microsatellite technique to be a valuable tool for aquaculturists to use in analyzing the genetic diversity of breeding programs. Microsatellite genetic diversity and nucleotide divergence among haplotypes were demonstrated 0.63-0.74 and 0.2-16.3% respectively, by You et al. (2008). The estimated level of polymorphism in this study (22%) and previous data are promising and indicate RAPD markers are likely to access enough genetic variation for the establishment of a marker-assisted selective breeding program in prawn. The heritability of a metric character is one of its most important properties. It expresses, as we have seen the proportion of the total variance that is attributable to differences of breeding values, and this is what determines the degree of resemblance between relatives (Falconer, 1998) but the most important function of heritability in the genetic study of metric characters is its predictive role, expressing the reliability of the phenotypic value as a guide to the breeding value (Falconer, 1998; Moss et al., 2007). Recently, studies on genetics indicated a very good heritability on productive traits for penaeid shrimp and this could open a good outlook to detect related genes for marker assisted selections in future shrimp selective breeding programs (Hoa, 2009). Estimates of heritability for harvest weight have been reported 0.42 ± 0.15 in *L.vannamei* (Carr et al., 1997); 0.16 to 0.31 in *M. japonicus* (Hetzl et al., 2000); 1.32 ± 0.18 in *L.vannamei* (Perez-Rostro et al., 1999); 0.24 ± 0.05 (full-sib family), 0.17 ± 0.04 (half-sib family) in *L.vannamei* (Gitterle et al., 2005b) and 0.24-0.35 (univariate animal model), 0.37-0.45 (multivariate animal

model) in *L.vannamei* (Castillo-Juarez et al., 2007). Heritability of growth, TSV (Taura Syndrome Virus) resistance and tail percent were estimated 1 ± 0.12 , 0.28 ± 0.12 and 0.15 ± 0.12 respectively, in *L. vannamei* (Argue et al., 2002). Heritability value for body length was demonstrated 0.22 at 119 days of age (Perez-Rostro et al., 2003) and 0.43 at 25 days of age (Campos et al., 2006) in *L. vannamei*. The heritability for resistance to disease (Gitterle et al., 2005a; Ibarra et al., 2007) and reproductive traits (Arcos et al., 2004; Arcos et al., 2005; Macbeth et al., 2007) in *L. vannamei* have been evaluated. Data in different commercial hatcheries with different abilities confirm the role of genetical control on size and growth in penaeidae (Chow and Sandifer, 1991). The range of heritability (0-1) in the larval stage of *L.vannamei* and *L.stirostris* shows effects of environmental factors on growth (Lester and Lauser, 1990). Argue et al. (2002) studied heritability of sex ratio in *L.vannamei* and were not significantly different from zero. This differs from the results reported in turtles and fish, which exhibit significant heritability estimates for sex ratio (Lester et al., 1989), hence, instead of selective breeding, it may be possible to produce more females by manipulating the androgenic gland or exposing shrimp to exogenous hormones (Sagi and Cohen, 1990; Moss et al. 2002). In this study, heritability of weight for *F. indicus* was estimated 0.07 which is lower than those reported for other prawn that could be explained by lack of genetic care during the domestication period. Estimation of response to selection in *M. japonicus* (Hetzl et al., 2000), *L.stylirostris* (Goyard et al., 2002), and *L.*

vannamei (Argue et al., 2002) were averagely 4-18%, 10.7% and 25% after one generation. In this study, response to selection was estimated 1.2 ± 0.2 (8%) in one generation that is lower than those which have been observed in other marine species. Low rates of response observed in our study are presumably because of the relatively low selection intensity applied. Nevertheless, our study has demonstrated that growth will respond to selection in *P. indicus*. It is expected that the response to the selection is valid only in the first generation, but it has been shown in the experiments that response with little change has been maintained during several generations (up to 5, 10 or even more generations). Over the longer term, phenotypic variation as well as heritability may decrease, resulting in lower rates of genetic change. In addition, negative genetic correlation can arise and reduce long term genetic gain (Falconer, 1998). This preliminary investigation of RAPD's in panaeid has provided methods to obtain RAPD's from *F. indicus*, and is likely to be adequate to obtain markers to assist selective breeding programs. We are pursuing our study by sequencing the specific bands, designing primers and consequently examining them on the same age specimens.

Acknowledgements

This work was supported and funded by Agricultural and Natural Resources Affairs Bureau, Vice-President for Strategic Planning and Supervision and Iranian Fisheries Organization.

References

- Afsharnasab, M., Laloei, F. and Rezvani Gilkolaei, S., 2005.** Investigation of White Spot Syndrom Disease (WSSD) in *Penaeus indicus* by Polymerase Chain Reaction (PCR) in Iran. *Iranian Scientific Fisheries Journal*, 14(1), 1-12. (In Persian)
- Arcos, F. G., Racotta, I. S. and Ibarra, A. M., 2004.** Genetic parameter estimates for reproductive traits and egg composition in Pacific white shrimp *Penaeus (Litopenaeus) vannamei*. *Aquaculture*, 236, 151-165.
- Arcos, F. G., Racotta, I. S., Palacios, E. and Ibarra, A. M., 2005.** Ovary development at the onset of gametogenesis is genetically determined and correlated with reproductive traits at maturity in shrimp *Litopenaeus (Penaeus) vannamei*. *Marine Biology*, 148, 339-346.
- Argue, B. J., Arce, S. M., Lotz, J. M. and Moss, S. M., 2002.** Selective breeding of pacific white shrimp (*Litopenaeus vannamei*) for growth and resistance to Taura Syndrome Virus. *Aquaculture*, 204, 447-460.
- Benzie, J. A. H., Kenway, M. and Trott, L., 1997.** Estimates for the heritability of size in juvenile *Penaeus monodon* prawn from half-sib mating. *Aquaculture*, 152, 49-53.
- Benzie, J. A. H. 2009.** Use and exchange of genetic resources of penaeid shrimps for food and aquaculture. *Reviews in Aquaculture*, 1, 232-250.
- Caetano-Anolles, G., Bassam, B. Y. and Grasshoff, P. M., 1991.** High resolution DNA amplification .fingerprinting using very short arbitrary oligonucleotide primers. *Bio/technology*, 9, 553-557.

- Campos, G., Montaldo, H., Arechavaleta-Velasco, M. and Castillo-Juárez, H., 2006.** Heritability of body length at 25 days of age in the Pacific white shrimp (*Litopenaeus vannamei*). 8th World Congress on Genetics Applied to Livestock Production, August 13-18. Belo Horizonte, MG, Brazil.
- Carr, W. H., Fjalestad, K. T., Godin, D. M., Swingle, J., Sweeney, J. N. and Gjedrem, T., 1997.** Genetic variation in weight and survival in population of specific pathogen free shrimp. *Penaeus vannamei*. In: Book of Abstracts. World Aquaculture, Bangkok, Thailand. p. 63.
- Castillo-Juarez, H., Casares, J. C. Q., Campos-Montes, G., Villela, C. C., Ortega, A. M. and Montaldo, H. H., 2007.** Heritability for body weight at harvest size in the Pacific white shrimp, *Penaeus (Litopenaeus) vannamei*, from a multi-environment experiment using univariate and multivariate animal models. *Aquaculture*, 273, 42–49.
- Chow, S and Sandifer, P. A. 1991.** Differences in growth, morphometric traits and male sexual maturity among Pacific white shrimp. *Penaeus vannamei*, from different commercial hatcheries. *Aquaculture*, 92, 165-178.
- Cock, J., Gitterle, T., Salazar, M. and Rye, M., 2009.** Breeding for disease resistance of Penaeid shrimps. *Aquaculture*, 286, 1-11.
- De la Rosa-Ve'lez, J., Escobar, R., Correa, F. and Felix, E., 1999.** High allozyme variation and genetic similarity of two populations of commercial penaeids, *Penaeus brevisrostris* (Kingsley) and *P. vannamei* (Boone), from the Gulf of California. *Aquaculture Research*, 30, 459–463.
- Donato, M. D., Manrique, R., Ramirez, R., Mayer, L. and Howell, C., 2005.** Mass selection and inbreeding effect on a cultivated strain of *Litopenaeus vannamei* in Venezuela. *Aquaculture*, 247, 159-167.
- Falconer, D. S., 1998.** Introduction to Quantitative Genetics. 3Th Edition. Longman Scientific & Technical. 438p.
- FAO, 2009.** (Food and Agriculture Organization of the United Nations). The State of World Fisheries and Aquaculture. WWW. FAO.ORG.
- Garcia, D. K. and Benzie, J. A. H., 1995.** RAPD markers of potential use in penaeid prawn (*Penaeus monodon*) breeding programs. *Aquaculture*, 130, 137-144.
- Gjedrem, T., 1992.** Breeding plans for rainbow trout. *Aquaculture*, 100, 73-92
- Gjedrem, T. and Fimland, E., 1995.** Potential benefits from high health and genetically improved shrimp stocks. In Browdy, C. L., Hopkins, J.S., (Eds.). Proceedings of Special Session on Shrimp Farming Swimming Through Troubled Water. *Aquaculture*, 95, 235
- Goyard, E., Patrois, J., Peignon, J. M., Vanaa, V., Dufour, R., Viallon, J. and Bedier, E., 2002.** Selection for better growth of *Penaeus stylirostris* in Tahiti and Caledonia. *Aquaculture*, 204, 461-468.
- Goyard, E., Goarant, C., Ansquer, D., Brun, P., Decker, S. D., Dufour, R., Galinié, C. et al., 2008.** Cross breeding of different domesticated lines as a simple way for genetic improvement in small aquaculture industries: Heterosis and inbreeding effects on growth and

- survival rates of the Pacific blue shrimp *Penaeus (Litopenaeus stylirostris)*. *Aquaculture*, 278, 43-50
- Gitterle, T., Salte, R., Gjerde, B., Cock, F., Johansen, H., Salazar, M., Lozano, C. and Rye, M., 2005a.** Genetic (co) variation in resistance to White Spot Syndrom Virus (WSSV) and harvest weight in *Penaeus (Litopenaeus vannamei)*. *Aquaculture*, 246, 139-149
- Gitterle, T., Rye, M., Salte, R., Cock, J., Johansen, H. and Lozano, C., 2005b.** Genetic (co)variation in harvest body weight and survival in *Penaeus (Litopenaeus) vannamei* under standard commercial conditions. *Aquaculture*, 243, 83-92.
- Hetzel, D. J. S., Crocos, P. J., Davis, G. P., Moore, S. S. and Preston, N. C., 2000.** Response to selection and heritability for growth in Kuruma prawn, *Penaeus japonicus*. *Aquaculture*, 181, 215-223.
- Hillis, D. M., Moritz, C. and Mable, B., 1996.** Molecular systematics, Second edition. Sinauer Assoc., Inc. Sunderland, MA 655 pp.
- Hoa, N. D., 2009.** Domestication of black tiger shrimp (*Penaeus monodon*) in recirculation systems in Vietnam. PhD thesis, Ghent University, Belgium. 179P.
- Ibarra, A. M., Perez-Rostro, C. I., Ramirez, J. L. and Ortega-Estrada, E., 2007.** Genetics of the resistance to hypoxia in postlarvae and juveniles of the Pacific white shrimp *Penaeus (Litopenaeus) vannamei* (Boone 1931). *Aquaculture Research*. 38: 838- 846.
- Kednapat, S., Kamonrate, W. and Nakorn, V., 2007.** Genetic aspect in broodstock management of critically endangered Makong gaint catfish, *Pangasianodon gigas* in Tailand. *Aquaculture*, 264, 34-46
- Keys, S. J., Crocos, P. J., Burrridge, C. Y., Coman, G. J., David, G. P. and Preston, N. P., 2004.** Comparative growth and survival of inbred and outbred *Penaeus (marupenaeus) japonicas*, reared under controlled environment conditions: indications of inbreeding depression. *Aquaculture*, 241, 151-168.
- Lester, L. J., Lawson, K. S., Abella, T. A. and Palada, M. S., 1989.** Estimated heritability of sex ratio and sexual dimorphism in tilapia. *Aquatic Animal Health*, 10, 271-281
- Lester, L. J. and Lauser, K. S., 1990.** Inheritance of size as estimated by principal component analysis at two temperatures in *Penaeus vannamei*. *Aquaculture*, 83, 323-323.
- Lu, Y. and Sun, P. S. 2005.** Viral resistance in shrimp that express an antisense Taura syndrome virus coat protein gene. *Aquaculture research*, 67, 141-146
- Macbeth, M., Kenway, M., Salmon, M., Benzie, J., Knibb, W. and Wilson, K., 2007.** Heritability of reproductive traits and genetic correlations with growth in the black tiger prawn *Penaeus monodon* reared in tanks. *Aquaculture*, 270, 51-56.
- Moore, S. S., Whan, V., Davis, G. P., Byrne, K., Hetzel, D. J. S. and Preston, N., 1999.** The development and application of genetic markers for the Kuruma prawn *Penaeus japonicus*. *Aquaculture*, 173, 19-32
- Moss, D. R., Hennig, O. L. and Moss, S. M., 2002.** Sexual growth dimorphism in penaeid shrimp. Potential for all female

- culture. *Global Aquaculture Advocate*, 5, 60–61.
- Moss, D. R., Arce, S. M., Otoshi, C. A., Doyle, R. W. and Moss, S. M., 2007.** Effect of inbreeding on survival and growth of pacific white shrimp *Panaeus vannamei*. *Aquaculture*, 272S1, S30-S37
- Nei, M., 1972.** Genetic distance between populations. *The American Naturalist*, 106, 283-292.
- Nelson, K. and Hedgecock, D., 1980.** Enzyme polymorphism and adaptive strategy in the decapod Crustacea. *The American Naturalist*, 116, 238-280.
- Olivier, J. and Roel, B., 2009.** Typology of shrimp farming in Bac Lieu Province, Mekong Delta, using multivariate statistics. *Agriculture, ecosystems & environment*, 132, 153-159.
- Perez-Rostro, C. I., Ramirez, J. L. and Ibarra, A. M., 1999.** Maternal and cage effect on genetic parameter estimation for Pacific white shrimp *Penaeus (Litopenaeus) vannamei* Boone. *Aquaculture Research*, 30, 191-197.
- Perez-Rostro, C. I., Ibarra, A. M., 2003.** Quantitative genetic parameter estimates for size and growth rate traits in Pacific white shrimp *Penaeus vannamei* (Boone 1931) when reared indoors. *Aquaculture Research*, 34, 543–553.
- Pourkazemi M., Nazari S., Bakhshalizadeh S., 2010.** Karyotype analysis in white bream (*Blicca bjoerkna transcaucasica*) from north coast of Iran. *Iranian Journal of Fisheries Sciences*, 9(3), 454-463.
- Preston, N. P., Croscos, P. J. and Keys, S. J., Coman, G. J. and Koenig, R., 2004.** Comparative growth of selected and non-selected Kuruma shrimp *Penaeus (Marsupenaeus) japonicus* in commercial farm ponds; implications for broodstock production. *Aquaculture*, 231, 73-82.
- Pullin, R. S. V., Williams, M. J. and Preston, N., 1998.** Domestication of crustacean, *Asian Fisheries Sciences*, 11, 71-80.
- Sagi, A. and Cohen, D., 1990.** Growth, maturation and progeny of sex-reversed *Macrobrachium rosenbergii* males. *World Aquaculture*, 21(4), 87-90.
- Taris, N., Batista, F. M. and Boudry, P., 2007.** Evidence of response to unintentional selection for faster development and inbreeding depression in *Crassostrea gigas* Larvae. *Aquaculture*, 272, 69-79
- Welsh, J. and Meelelland, M., 1990.** Finger printing genomic using PCR with arbitrary primers. *Nucleic Acids Research*, 12, 7212-7213
- Williams, J. G. K., Kubelik, A., Livak, K., Rasfolshki, J. A., Tingey, S.V., 1999.** DNA polymorphism amplified by arbitrary primers are useful and genetic markers. *Nucleic Acid Research*, 18, 6531-6535
- Wolfus, G. M., Garcia, D. k and Alcivar-Warren, A. A., 1997.** Application of the microsatellite technique for analysing genetic diversity in shrimp breeding programs. *Aquaculture*, 152, 35-47
- Yeh, F. C., Yang, R. C. and Boyle, T., 1999.** Popgene Version 1.31. Microsoft Window-based Freeware for Population Genetic Analysis. Available: www.ualberta.ca/fyeh/. University of Alberta and the Center or International Forestry Research. Quick User Guide.
- You, E-M., Chiu, T-S., Liu, K-F., Tassanakajon, A., Klinbunga, S. and Triwitayakorn, K., 2008.**

Microsatellite and mitochondrial haplotype diversity reveals population differentiation in the tiger shrimp (*Penaeus monodon*) in the Indo-Pacific region. *Animal Genetics*, 39, 267–277.