# DOR: 20.1001.1.15622916.2011.10.1.16.5

# Effect of nanosilver particles on hatchability of rainbow trout (Oncorhynchus mykiss) egg and survival of the produced larvae

Soltani, M. 1\*; Esfandiary, M. 1; Sajadi, M. M. 2; Khazraeenia, S. 1; Bahonar, A. R.<sup>3</sup>; Ahari H.<sup>4</sup>

> Accepted: February 2010 Received: July 2009

### **Abstract**

Effect of nanosilver particles was studied on the hatchability of rainbow trout (Oncorhynchus mykiss) egg and survival of the produced larvae at about 12°C. In the first experiment the water-based nanosilver particles was used at concentrations of 0.5, 1, 2 and 4 mgL<sup>-1</sup> for 30 minutes per day starting 24 hour post egg incubation until the hatching time. The mean percentage of hatchability reached in 27.6±0.2, 38.2±0.1, 41.6±0.4 and 48.6±1.5 in troughs treated with 0.5, 1, 2 and 4 mgL<sup>-1</sup> nanocid, respectively compared with 64.7±0.2 % for trough treated with malachite green at 2 mgL<sup>-1</sup> as positive control (P<0.05) and 5.1± 0.2 % for trough without treatment (untreated group) (P<0.0.5). In the second experiment the effect of the nanosilver particles was evaluated in the form of nanosilver dressing i.e. coating nanocid on the surface of the incubator troughs. The mean percentage of hatchability was 69.4±0.1% compared with 52.5%±0.1 recorded for normal control (trough without nanosilver dressing) (P<0.05). When growing the produced larvae in the same troughs until one g in body weight, there was an insignificant increase in the weight of larvae kept in nanosilver trough ( mean weight 1.12±0.09 g) compared to control group (mean weight 1.03±0.02g) (P>0.05). These data suggest a possible application of nanosilver particles in aquaculture sector particularly using incubator troughs of trout containing nanosilver materials.

**Key words:** Silver nanoparticles, Rainbow trout egg, Hatchability, Survival

<sup>1-</sup>Department of aquatic animal health, faculty of veterinary medicine, university of Tehran, Tehran, Iran.

<sup>2-</sup>Department of marine biology, faculty of basic sciences, university of Hormozgan, Bandar Aabbas, Iran.

<sup>3-</sup>Department of food hygiene, faculty of veterinary medicine, university of Tehran, Tehran, Iran.

<sup>4-</sup>Veterinary section, Pars Nano Nasb, Co. Tehran, Iran.

<sup>\*</sup>Corresponding author's email: msoltani@ut.ac.ir

Downloaded from jifro.ir on 2025-11-15 ]

### Introduction

Nanotechnology, the science of producing and utilizing nano-sized particles is growing rapidly and in the last few years it begun to evolve in to a valuable science. In this view nanosilver particles have received more attention because of their antimicrobial properties and their applications to a wide variety of healthcare products including drug delivery systems, therapeutics, biosensors, environmental remediation and dressing materials (Kong et al.,2000; Aitken et al.,2006; Lansdown, 2006).

The small size of nanoparticles of metallic silver in solid or colloidal state allows for a high antimicrobial effect than silver salts (Lansdown, 2006; Atiyeh et al., 2007: Choi et al., 2008). The nanosilver particles inhibit growth can multiplication of the bacteria and fungi by binding to their DNA, suppress their respiration by poisoning the respiratory enzymes and component of electron transport system, and alter the microbial membrane function by binding to bacterial surface (Wright et al., 1994; Percival et al., 2005).

A relatively new field of nanoaquaculture can incorporate nano technology in fish farming for increasing diagnosing production, any disease, cleaning fish ponds, developing a system mass vaccination of fish ofusing nanocapsules containing DNA nanovaccination and fastening the growth of fish through iron nanoparticles consumption in fish feed (Wong et al., 1992; O'Hagan, 1998; Janes et al., 2001; Alan, 2006; Rajesh Kumar et al., 2008).

Despite the broad applications of nanoparticles in both human and terrestrial

animal veterinary medicines, minimum works have been done in aquatic medicine The occurrence of fungal and bacterial infections particularly during hatching period of trout egg and growing of the produced larvae is still one of the main obstacles encountered with the trout industry worldwide. Particularly such problem has been elaborated since the use effective of the most aquaculture antifungal, malachite green has been inhibited several years ago. Therefore, the investigation to find new environmentally friendly chemical substances is a present necessity.

The objective of this study was to assess the effect of nanosilver particles in the form of water soluble suspension and also as a dressing material coated on the egg incubator/troughs on the hatchability of rainbow trout (*Oncorhynchus mykiss*) egg and survival of the produced larvae.

## Matherials and mehthods Nanosilver

The nanocid P-series powder products of NanoNasb Pars Company, Tehran, Iran are comprised of titanium dioxide coated with 1-10% nanosilver particles and used as an additive in industrial products such as polymers, glue, ceramic and paints. In this study, the nanocid powder coated on the surface of the polyethylene trough (200 x50 x 40 cm) was used. This trough has a unique specification possessing silver ion on the surface.

Also, the water soluble colloidal nanosilver particles (4000 mgL<sup>-1</sup> stock solution) with average size of 5 nm containing Tween prepared in distilled water and brown in color were used.

Experimental design

Brood stock selection and spawning

Apparently healthy brood rainbow trout were used in this experiment. The ripen brood fish were spawned after anesthetizing them with clove oil essence at 100 mg L-1. The fertilization of eggs was performed according to the routine produced method described for trout eggs.

### **Trial 1**

Six troughs each containing 14700 fertilized eggs (each trough containing three trays and each tray containing 4900 eggs (350 g) were used. The dead eggs were removed carefully 24 hour postincubation and the remaining eggs were left untouched until the eyed-eggs stage. Four troughs were used as treatments and treated with nanosilver particles concentrations of 0.5, 1.0 and 2 and 4 mgL<sup>-1</sup> for 30 minutes per day until the eyed-egg stage. The dead eggs were removed carefully at eyed-egg stage, and the removal of dead eggs was continued daily until hatching time. One trough was treated with malachite green at 2 mgL<sup>-1</sup> for 20 minutes per day as the positive control and one trough without any treatment as the untreated group (Table 1). The experiment was repeated for the second time and the percentage of hatched eggs was then calculated at the end of hatching period as follow:

% hatched eggs= (number of hatched eggs/number of initial eggs) x 100

### **Trial 2**

One nanosilver trough containing four trays and each tray containing 6300 eggs (totally 25200 fertilized eggs) was used as the treatment. Two normal troughs similar to those in trial 1 were used as control groups. The dead eggs were removed

carefully 24 hours post-incubation and the remaining eggs were left untouched until the eyed egg stage. The dead eggs were then removed carefully daily until hatching time. The experiment was repeated for the second time and, the percentage of hatched eggs was then calculated at the end of hatching period as follow: Percent of hatched eggs/number of initial eggs) x 100

The hatched larvae were then grown in the same troughs until one g in body weight. The larvae were first fed at 4.8% of their body weight on a commercial diet of SFT00 up to 0.5 g body weight and then at 4.4% of their body weight on SFT0 and SFT1 diets up to 1 g weight. The of produced larvae percentage was calculated as follow: Percentage of produced larvae= (number of survived larvae at the end of the experiment/number of initial larvae) x 100. Also, the biometry of larvae was preformed every 10-15 interval days. Daily mortality was also recorded.

Water quality

Water quality parameters consisting of dissolved oxygen, temperature, pH, ammonia, nitrite, carbon dioxide and hardness were  $8\pm0.4~\text{mgL}^{-1}$ ,  $11.5\pm0.5^{\circ}\text{C}$ ,  $7.5\pm0.1$ ,  $<0.01\pm0.05~\text{mgL}^{-1}$ ,  $<0.1\pm0.1\text{mgL}^{-1}$ , ca  $9\text{mgL}^{-1}$  and  $170\pm1.2~\text{mgL}^{-1}$ , respectively.

Other hatchability parameters including inlet water flow (7.5-8 L.min<sup>-1</sup> for the eggs and 45-50 L.min<sup>-1</sup> for the larvae), protection of light during hatching period and no manipulation of eggs were performed during the experiment.

### **Statistics**

The statistical analysis of the obtained data was carried out using SPSS version 14

Downloaded from jifro.ir on 2025-11-15 ]

DOR: 20.1001.1.15622916.2011.10.1.16.5

software and T-test, Kruskal-Wallis and Mann-Whitnney test at P<0.05 as the confidence.

### Results

The results of hatchability of eggs treated with nanosilver particles are shown in Table 1. The mean survival of eggs up to eyed-stage and treated with nanosilver particles at concentrations of 0.5, 1, 2 and

4 mg/L were  $91.8\pm1.6$ ,  $91.1\pm0.3$ ,  $94.1\pm0.1$ , 94.5±0.1%, respectively, while those of positive (malachite green) and nontreatment group were 96±0.09% and 89.36±0.1 %, respectively. Statistical showed that no analysis significant difference was seen among the treated groups and also between untreated group and all nanosilver treated eggs up to eyedegg stage (P>0.05).

Table 1: Survival of fertilized rainbow trout eggs treated with different concentrations of nanosilver at about  $12^{\circ}$ C. The data are shown as Means  $\pm$  SE, based on two replications

Treatment (mgL <sup>-1</sup> )	Mean No. of	Mean No. of survived	Mean No. of survived
	initial fertilized	eggs up to eyed-egg	eggs from eyed-egg
	eggs $\pm$ SE	stage $\pm$ SE (%)	stage up to hatching
			time ± SE (%)
Nanosilver (0.5)	14029 ± 1.6	12881±2.2 (91.8±1.6)	3877±1.9 (27.6±0.2)
Nanosilver (1)	$14043\pm1.0$	12797±1.9 (91.12±0.3)	5372±2.1 (38.25±0.1)
Nanosilver (2)	$14001\pm0.9$	13175±1.6 (94.1±0.1)	5823±1.8 (41.6±0.4)
Nanosilver (4)	$13987\pm1.8$	13217±2.4 (94.49±0.1)	6810±1.1 (48.6±1.5)
Malachite green (2)	$14015 \pm 2.5$	13455±2.0 (96±0.09)	9080±2.2 (64.7±0.2)
Untreated group	$14001 \pm 2.3$	12515±2.4 (89.36±0.1)	719±2.1 (5.1±0.2)
Treatment (mgL <sup>-1</sup> )	Mean No. of	Mean No. of survived	Mean No. of survived
	initial fertilized	eggs up to eyed-egg	eggs from eyed-egg
	eggs $\pm$ SE	stage ± SE (%)	stage up to hatching
			time ± SE (%)
Nanosilver (0.5)	$14029 \pm 1.6$	12881±2.2 (91.8±1.6)	3877±1.9 (27.6±0.2)
Nanosilver (1)	$14043\pm1.0$	12797±1.9 (91.12±0.3)	5372±2.1 (38.25±0.1)
Nanosilver (2)	$14001 \pm 0.9$	13175±1.6 (94.1±0.1)	5823±1.8 (41.6±0.4)
Nanosilver (4)	$13987\pm1.8$	13217±2.4 (94.49±0.1)	6810±1.1 (48.6±1.5)
Malachite green (2)	$14015 \pm 2.5$	13455±2.0 (96±0.09)	9080±2.2 (64.7±0.2)
Untreated group	$14001 \pm 2.3$	12515±2.4 (89.36±0.1)	719±2.1 (5.1±0.2)

addition, eggs treated with malachite green (positive control) insignificantly resulted in a higher survival rate compared with both untreated and treated groups (P>0.05). Also, the mean survival of eggs from eyed-egg stage up to hatching period were  $27.6\pm0.2$ ,  $38.25\pm0.1$ , 41.6±0.4 and  $48.6 \pm 1.5$ percent at concentrations of 0.5, 1, 2 and 4 mgL<sup>-1</sup> nanosilver particles, respectively, while

those of malachite green (positive control) and untreated group were 64.7±0.2 and 5.1±0.2 percent, respectively (Table 1) (Fig. 1). Statistically, there was significant difference between untreated and treated groups (P<0.05). Also, there was a significant difference between positive control and all nanosilver treated groups (P<0.05).




Figure 1: Mean mortality pattern of fertilized eggs of rainbow trout treated with different concentrations of nanosilver particles for 30 minutes per day until eyed-egg stage at about 12°C

Table 2: Survival of the fertilized rainbow trout eggs incubated in nanosilver coated\_and normal troughs at about 12  $^{\circ}$ C. The data are shown as Means  $\pm$  SE, based on two replications

Treatment	Mean No.	Mean No of	Mean No of
	of initial	survived eggs	survived eggs
	fertilized	up to eyed-egg	up to hatching
	$eggs \pm SE$	$stage \pm SE$	time $\pm$ SE
	(g)	(%)	(%)
Nanosilver	24365±1.9	22503±1.7	16920±2.1
trough	$(1728\pm1.5)$	$(92.4\pm0.2)$	$(69.4\pm0.1)$
Normal trough	27606±2.0	22360±2.2	11627±1.9
	$(1958\pm1.6)$	$(81\pm0.3)$	$(52.5\pm0.1)$

mgl -1 Eggs treated with 2 4 and nanosilver resulted significantly in higher survival than treated groups at 0.5 mgl<sup>-1</sup> nanosilver particles (P<0.05), while no significant difference was seen between eggs treated at 0.5 and 1 mgl<sup>-1</sup> nanosilver particles (P>0.05). Also, no significant difference was seen between eggs treated with 1 and 2 mgl<sup>-1</sup> nanosilver particles (P>0.05). The results of survival of eggs incubated in nanosilver trough are shown in Table 2.

The mean survival rate of eggs up to eyedegg stage in nanosilver trough and normal trough was 92.4±0.2 and 81±0.3 percent, respectively which showed a significant difference (P<0.05). Also, the mean survival rate of eggs from the eyed-egg stage up to hatching time was 69.4±0.1 and 52.5±0.1 percent for nanosilver trough and normal trough, respectively which showed a significant difference (P<0.05). In addition the pattern of mortality of eggs from eyed-egg stage up to hatching time is shown in Figure 2. Based on this figure, the mean mortality of eggs incubated in nanosilver trough was significantly lower than the mean mortality of eggs incubated in normal trough during 7 days after eyedegg stage (P<0.05).




Figure 2: Mean mortality of rainbow trout egg incubated in nanosilver trough compared with normal trough at about 12  $^{\circ}\mathrm{C}$ 




Figure 3: Mortality progress of rainbow trout larvae grown in the nanosilver trough compared with normal troughs




Figure 4: Growth of produced rainbow trout larvae grown in nanosilver trough at about  $12^{\circ} C$ 

The pattern of mortality rate for the produced larvae is shown in Figure 3. According to this figure the mean mortality of larvae grown in nanosilver

trough was significantly lower than the normal trough during almost 30 days post-hatching period (P<0.05). At the end of growing period (up to one g weight), the

mean survival of grown larvae in nanosilver trough was  $91.8\pm1.5$  percent while that of normal trough was  $87.3\pm0.1$  percent (P<0.05). Also, the larval biometry (mean total weight) resulted in  $1.12\pm0.09$  g and  $1.03\pm0.02$  g for nanosilver trough and normal trough, respectively (P>0.05) (Figure 4).

### **Discussion**

The activity of colloid nanosilver particles against a wide range of gram positive/negative bacteria, viruses and fungi have been reported by a number of researchers including Panacek et al. (2006), Ju-Nam and Lead (2008), Zhang et al. (2008), Weir et al. (2008), Pallab Sanpui et al. (2008), Rai et al. (2009), Fayaz et al. (2009), Tai et al. (2009) and Barrena et al. (2009).

The mechanism behind the antimicrobial activity of nanosilver particles is basically due to interaction with sulfur containing proteins in bacterial cell wall as well as with phosphorus containing compounds like DNA and finally cell lysis and death occurs as a result of disrupting membrane permeability, deactivating cellular enzymes, impairment of energy cycle and DNA replication (Feng et al., 2000; Jeon et al., 2003; Sondi and Salopek-Sondi, 2004; Morones et al., 2005; Song et al., 2006).

Two major kinds of nanotechnology products are defined as nanomaterials fixed on a substrate i.e. coated devices and free nanoparticles (Piotrowska et al. 2009). In this study we used both forms of nanosilver particles to investigate their antimicrobial effects on the survival of rainbow trout eggs and the produced larvae. In the first experiment the nanosilver particles at used concentrations could inhibit growth and multiplication of

potential pathogenic micro organisms particularly Saprolegnia SO that percentage of egg survival and hatchability were significantly higher than untreated control (P<0.05). However such antimicrobial effect was less than malachite green after eyed-egg stage. Also, the use of nanosilver particles coated on the trough in the second experiment resulted in a higher rate of egg survival and hatchability indicative of its more efficient antimicrobial effects compared with the untreated control group.

Nanosilver particles coated devices are characterized with clear antimicrobial properties since they release silver ions gradually, thus provide a sustained supply of silver ions at the coating interface for prolonged prevention of bacterial growth (Furno et al., 2004; Roe et al., 2008; Chen and Schluesener ,2008). Accordingly the higher rate of trout egg survival and hatchability obtained in the present study may be due to the continuous release of silver ions from the polymeric silver providing coated trough suitable antimicrobial activity. Moreover it should nanosilver be noted that particles chemistry including their size distribution, morphology, surface area, charge, surface modifications, chemical composition (purity) have important role in their possible mechanisms of antimicrobial action (Singh et al., 2009). Nanosilver particles tend to aggregate in aqueous solutions because of their hydrophobic properties. Different stabilizers are being used to prevent their agglomeration and to obtain isolated nanosilver particles to produce more efficient antimicrobial effects (Hyung et al., 2007). The nanosilver particles used in this study were

in the form of silver ion with about 5 nm in size containing some Tween probably as according surfactant and to the manufacturer description more substance was added. Development of new nanoparticle surface engineering strategies should lead to more environmentally friendly nanoparticles. Biogenic synthesis of nanomaterials is an important of aspect current nanotechnology research, for instance use of fungus Trichoderma viride for the extracellular biosynthesis of silver nitrate particles from silver nitrate solution has been reported by Fayaz et al. (2009). It was observed that the aqueous silver (Ag+) ions, when exposed to filtrate of T. viride reduced in solution, thereby resulted in formation of extremely stable nanosilver particles.

In conclusion, this study suggests an application of nanosilver particles with their unique chemical and physical properties particularly in the form of nanosilver dressing troughs as a disinfectant in fish farming. Also, nanosilver particles may have a priority to current antibiotic/antifungal chemicals as according to Rai et al. (2009), the microorganisms are unlikely to develop resistance against silver nanoparticles, compared to antibiotics due to their large surface area in contact with and broad range of targets in the microbes. However, there are some questions, which need to be addressed, such as the exact mechanism of interaction of silver nanoparticles on the fungal and bacterial cells and how the surface area of nanoparticles influences its killing activity.

### Acknowledgements

This work was financially supported by Pars Nanonasb Comp. and a grant from the University of Tehran. Authors thank Mr Moghaddasi, Director of Mahyaran Aquaculture Company for his help on this work.

### References

- Aitken, R. J., Chaudhry, M. Q., Boxall, A. B. A. and Hull, M., 2006. Manufacture and use of nanomaterials: current status in the UK and global trends. *Occupational Medicine*, 56, 300–306.
- **Alan, S., 2006.** Opinion: Nanotech the way forward for clean water. *Filtration & Separation*, 43, 32-33.
- Atiyeh, B. S., Costagliola, M., Hayek, S. N. and Dibo, S. A., 2007. Effect of silver on burn wound infection control and healing: review of the literature. *Burns*, 33,139–148.
- Barrena, R., Casals, E., Colón, J., Font,X., Sánchez, A. and Puntes, V., 2009. Evaluation of the ecotoxicity of model nanoparticles .*Chemosphere*, 75, 850–857.
- Chen, X., Schluesener, H. J., 2008. Nanosilver: a nanoproduct in medical application.

*Toxicology Leterst*, 176, (1), 1–12.

- Choi, O. K., Deng, K., Kim, N. J., Ross, L. and Hu, Z. Q., 2008. The inhibitory effects of silver nanoparticles, silver ions, and silver chloride colloids on microbial growth. *Water Research*, 42, 3066–3074.
- Fayaz, M., Balaji, K., Girilal, M., Ruchi Yadav, P., Kalaichelvan, T. and Venketesan R. 2009. Biogenic synthesis of silver nanoparticles and its synergetic effect with antibiotics: A

- study against Gram positive and Gram negative bacteria. Nanomedicine: *Nanotechnology*, Reference: NANO 275 (article in press).
- Feng, Q. L., Wu, J., Chen, G. Q., Cui, F. Z., Kim, T. N. and Kim, J. O., 2000.

  A mechanistic study of the antibacterial effect of silver ions on Escherichia coli and Staphylococcus aureus. Journal of Biomedical Materials Research, 52, 662–668.
- Furno, F., Morley, K. S., Wong, B., Sharp, B. L., Arnold, P. L. and Howdle, S. M., 2004. Silver nanoparticles and polymeric medical devices: a new approach to prevention of infection. Journal of Antimicrobial Chemotherapy, 54(6), 1019–24.
- Hyung, H., Fortner, J. D., Hughes, J. B. and Kim, J. H., 2007. Natural organic matter stabilizes carbon nanotubes in the aqueous phase. *Environmental Science & Technology*, 41, 179–184.
- Janes, K. A, Calvo, P. and Alonso, M. J., 2001. Polysaccharide colloidal particles as delivery systems for macromolecules. Advanced Drug Delivery Reviews, 47, 83–97
- Jeon, H. J., Yi, S. C. and Oh, S. G., 2003. Preparation and antibacterial effects of Ag–SiO2 thin films by sol–gelmethod. *Biomaterials*, 24, 4921–4928.
- Ju-Nam, Y. and Lead., J. R. 2008.

  Manufactured nanoparticles: An overview of their chemistry, interactions and potential environmental implications. Science of the total environment, 400, 396-414

- Kong, J., Franklin, N. R., Zhou, C., Chapline, M. G., Peng, S., Cho, K. and Dai, H., 2000. Nanotube molecular wires as molecular sensors, *Science*, 287, 622–625.
- **Lansdown, A. B., 2006.** Silver in health care: antimicrobial effects and safety in use. *Current Problems in Dermatology*, 33, 17–34.
- Morones, J. R., Elechiguerra, J. L., Camacho, A., Holt, K., Kouri, J. B., Ramírez, J. T. and Yacaman, M. J., 2005. The bactericidal effect of silver nanoparticles. *Nanotechnology* 16, 2346–2353.
- **O'Hagan, D.T., 1998.** Microparticles and polymers for the mucosal delivery of vaccines. *Advanced Drug Delivery Reviews*, 34, 305–320.
- Pallab Sanpui, A., Murugadoss, P. V., Prasad,D., Sankar Ghosh,S. and Chattopadhyay,A., 2008. The antibacterial properties of a novel chitosan–Ag-nanoparticle composite. *International Journal of Food Microbiology*, 124, 142-146.
- Panacek, A., Kvitek, L., Prucek, R., Kolar, M., Vecerova, R. and Pizurova, N., 2006. Silver colloid nanoparticles: synthesis, characterization, and their antibacterial activity. The Journal of Physical Chemistry, 110(33), 16248–16253.
- Percival, S. L., Bowler, P. G., Russell, D., 2005. Bacterial resistance to silver in wound care. *Journal of Hospital Infection*, 60, 1–7.
- Piotrowska, G. B., Golimowski, J. and Urban, P. L., 2009. Nanoparticles: Their potential toxicity, waste and environmental management. Waste

- management (article in press).
- Rai, M., Yadav, A., Gade, A., 2009. Silver nanoparticles as a new generation of antimicrobials. Biotechnology Advances, 27, 76–83.
- Rajesh Kumar, S., Ishaq Ahmed, V. P., Parameswaran, V., Sudhakaran, R., Sarath Babu, V. and Sahul Hameed, A. S., 2008. Potential use of chitosan nanoparticles for oral delivery of DNA vaccine in Asian sea bass (*Lates calcarifer*) to protect from Vibrio (*Listonella anguillarum*. Fish & Shellfish Immunology, 25, 47-56.
- Roe, D., Karandikar, B., Bonn-Savage, N., Gibbins, B. and Roullet, J. B., 2008. Antimicrobial surface functionalization of plastic catheters by silver nanoparticles. *Journal of Antimicrobial Chemotherapy*, 61(4), 869–76.
- Singh, N., Manshian, B., Jenkins, G. J. S., Griffiths, S. M., Williams, P. M., Maffeis, T. G. G., Wright, C. J. and Doak, S. H., 2009. Nano Genotoxicology: The DNA damaging potential of engineered nanomaterials. Biomaterials 1–24 (article in press).
- Sondi, I. and Salopek-Sondi, B., 2004. Silver nanoparticles as antimicrobial agent: a case study on *E.coli* as a model for gram-negative bacteria. *Journal of Colloid and Interface*, 275,177–82.

- Song, H. Y., Ko, K. K., Oh, L. H. and Lee, B. T., 2006. Fabrication of silver nanoparticles and their antimicrobial mechanisms. *European Cells & Materials Journal*, 11, 58.
- Tai, C. Y., Wang, Y. H., Kuo, Y. W., Chang, M. H. and Liu, H.S., 2009. Synthesis of silver particles below 10 nm using spinning disk reactor. *Chemical Engineering Science*, 64, 3112 3119.
- Weir, E., Lawlor, A., Whelan, A. and Regan, F., 2008. The use of nanoparticles in antimicrobial materials and their characterization. *Analyst*, 133, 835–845.
- Wong, G., Kaattari, S. L., Christensen, J. M., 1992. Effectiveness of an oral enteric coated vibrio vaccine for use in salmonid fish. *Immunological Investigations*, 21, 353–364.
- Wright, J. B., Lam, K., Burrell, R. E., 1994. Wound management in an era of increasing bacterial antibiotic resistance: a role for topical silver treatment. *American Journal of Infection Control*, 26, 572–577.
- Zhang, Y., Peng, H., Huang, W., Zhou, Y. and Yan, D., 2008. Facile preparation and characterization of highly antimicrobial colloid Ag or Au nanoparticles. *Journal of Colloid and Interface Science*, 325,371–376.