

Effects of whey protein edible coating on bacterial, chemical and sensory characteristics of frozen common Kilka

Seifzadeh, M*

Received: July 2011

Accepted: July 2013

Abstract

The objective of current study was to investigate the effects of whey protein coating on the quality of common kilka during frozen storage period. For this study, common kilka was coated with 12% whey protein concentration. Also, non-coated kilka was used as a control sample. Coated and non-coated samples were then stored at -18 °C for six months. Bacterial, chemical and sensory properties of the samples were determined in each month. Results showed that total bacterial counts and *Staphylococcus* bacteria count were lower in the test samples compared to the control samples. *Coliform*, *Escherichia coli* and *Pseudomonas* bacterial contamination were negative till the end of the storage period in the covered samples. Chemical factors consisting of humidity, protein, lipid, ash and calorie were higher in the test samples compared to the control samples. Peroxide value, free fatty acids, thiobarbituric acid, TVB-N and pH were lower in the test samples. Sensory tests had a better quality in the test samples compared to the control samples. No statistically significant decreases were observed in chemical factors in the covered samples compared to the control samples ($p < .05$). In sensory specifications, there was a significant difference between the covered samples and the control samples ($p < .05$). According to the results of experiments and statistical analysis, the covered samples had a favorable quality till the end of the storage period but the control samples had lost their quality.

Keywords: Edible film, Whey protein, Chemical analysis, Sensory analysis, Kilka fish.

1-Iranian fisheries national fish processing center, P.O. Box 43145-1655, Anzali, Iran.

*Corresponding author's email: m_seifzadeh_1d@yahoo.com

Introduction

Kilka fish belong to the genus *Clupeonella*, family Clupeidae. These fish are composed of three species consisting of *Clupeonalla delicatula* • *C. engrauliformis* and *C. grimmi* (Naderi and Abdoli, 2004). They can be processed into salted, smoked, pickled conserved, dried and frozen fish. But in Iran, kilka products in the market are canned, packaged in frozen shape and/or fresh. The annual catch has increased from 19610 tons in 2004 to over 25483 tons in 2009. Around 10–12% of this catch is used for human consumption and the remaining 88 – 90 % is used for animal feeds. About 10260 to 20741 tons of the fish catch occurred in the Mazandaran Province, of which 5–12 % is used for human consumption and the remaining 88–95% for animal feeds (Iranian fisheries, 2008). Consumption of fresh kilka fish dropped during the years 2004-2009 (from 6.00 to 2.20%). Consumption of canned kilka also dropped for 5.2% to about 0.8% during the same period, whereas consumption of the frozen fish rose from 1.00% to 6.20% during these years (Seifzadeh, 2010). The frozen fish packs, due to their longer storage time as well as wider countrywide distribution, had much higher sales rate in comparison to the sales of fresh fish. Sales of frozen fish were also higher. The frozen fish packs were mostly frozen for less than three months because longer frozen storage time may lead to color changes, surface dryness and peroxide accumulation (Moeini, 2009). Despite this, the first sign of quality decline, even after only one month of frozen storage, was a decline in the weight of frozen packed fish. This will in turn

have a deteriorating effect on the texture and taste of the small sized fish. There was a 3.5% decline in fish weight after three months of frozen storage (Kochakian, 2002).

Kilka having a valuable protein and digestible fats, rich vitamins and minerals has attained an important position among animal products. In an overall view, the value of each food product such as kilka depends on nutritional specifications and its acceptability in the societies; therefore, accurate processing of this fish and preparation of favorable variety products have special importance for the market (Martin, 1994).

The final step of the food production chain is packaging. Packaging occupies the middle ground of production, distribution and consumption. Many small and large companies in all over the world are very active in packaging field. The competition of these companies with each other has led to great increase in the quality and quantity of packaging forms. In most fields, the producers as well as the consumers are not well aware of the high value of proper modern packaging techniques. Currently, numerous packaging and preservation methods such as non – biological decomposable synthetic chemicals are used for food preservation. Recently, new packaging materials such as edible films that are biologically decomposable have been presented to the market (Shariatifar, 2007).

It seems as though consumer market demand is fairly high for high quality seafood products especially those which can retain their superior quality of taste, texture and general fresh appearance following prolonged

period of cold or frozen storage(Ahmed,2006). Use of edible films for packaging of kilka fish seems to be an ideal method for proper long storage fish preservation. Edible coatings are completely water soluble, glossy, act just like a secondary skin and have the following favorable proportion such as rapid attachment to foodstuff, label attachment, anti-bacterial and anti-oxidant properties(Seifzadeh,2007). These coatings are protective of the aroma, taste and food color and help to maintain the nutritional components such as their vitamin, amino acid and fatty acid ingredients. Covering of food products with these films can lead to preservation of food moisture, and oxygen absorption lowering, which can substantially improve the appearance of food products. These coating are invisible to the naked eye (Ahvenian, 2003). Whey protein is derived from milk and is composed of protein, lactose and inorganic salts. It is anti bacterial, anti proteolysis and preservation of food moisture (Morrissey et al., 2009).

Edible films made of whey protein have been used for packaging of salmon, hot-dog, sausage, cracker and frozen fish-fillet (Stuchell and Krockta, 1995). Packaging of fish with edible film was originally used in Iran by Mirhashemi Rostami (2000). He used coconut oil for packaging of *Huso huso* fish in his project. But no research hasnot been carried out using whey protein films for packaging of fish in this country. In other countries, films consisting of whey proteins have been used for food packaging by Piyachonkwan (1995), Stuchell (1995), Krockta (1996), Shah (1999), Crapo (1999), Cagri (2002, 2003) and Min (2006 , 2007).The objective of current study

was to investigate the effects of whey protein coating on the quality of common kilka during frozen storage.

Materials and methods

180 kg of common kilka caught in spring were used for this study. Fresh fish landed from kilka fishing vessels were obtained early in the morning (5 a.m.), and the characteristics of fresh fish following the national standards of Iran (INS¹, 2002b) were recorded. The fish were chilled down to 0 °C under a cover of ice at a fish: ice ratio of 2:1 in insulated tubs and then transferred to the production line of the National Fish Processing Center under complete hygienic conditions. The method used in this study followed the protocol described by other researchers with a few modifications.

Before processing fish were washed with chlorinated water. The fish were first washed. Then the heads were cut, and viscera were taken out. The cleaned fish were washed again.

Experimental design (Treatments)

Two treatments were used for this research. The treatments included coated Kilka and non coated Kilka.

The cleaned fish immersed in whey protein at a concentration of 12% in no time. After this stage, the fish was packaged in disposable dishes and covered by cellophane.

Control samples processing

For preparation of the control sample, Kilka fish was washed at first. Then the fish heads were cut, and viscera were taken out. The fish were washed again after cleaning and

¹ Iranian National Standard

were packaged in disposable dishes with cellophane covers in 500 gr in weight.

The prepared samples (coating and control) were kept at -18 °C in cold-storage for a period of six months. Quality of these samples were studied similar to the test samples on the determined times by microbial, chemical and sensory tests. These samples were processed three times. Each step of the tests was repeated three times.

Their for experimental and control samples (48 packages) consisted of total bacterial counts (Andrews & Hammak, 2003 & Maturin, 2001) for *Staphylococcus* (Bennet and Lancette, 2001 & Holt et al ., 1994), *Coliform* (Feng et al, 2002), *Escherichia coli* (Feng et al., 2002) and *Pseudomonas* (Hasegawa, 1987). Microbial tests were carried out during nine steps. The first microbial tests on fish samples were conducted immediately after receiving the raw fish samples. The second test was conducted after cleaning. The third stage was conducted one day after processing and then the following tests were conducted on , the second step after cleaning the Kilka, the third step, one day after processing and the remaining steps, once a month after processing at determined times for a period of six months. In each step, these tests were repeated three times.

The chemical tests for the packaged samples with edible film and control (42

Results

The bacterial counts of the samples are shown in the Tables 1 and 2. As indicated no *Coliform*, *Escherichia coli* and *Pseudomonas* bacterial contamination was observed in the

packages) consist of measurement of calorie (Iranian National Standard No 8867, 2005, protein (Iranian National Standard No 924, 1994) humidity (Iranian National Standard No 5625, 2002), lipid (Iranian National Standard No 742, 2002), ash (Iranian National Standard No 744, 2001), peroxide value (Iranian National Standard No 493, 2003), TVN (Iranian National Standard No 5625, 2002), TBA (Iranian National Standard No 10494, 2006), FFA (Iranian National Standard No 493, 2003) and pH (Iranian National Standard No 1028, 2007). These tests were conducted in eight steps which included the raw fish samples, one day after processing and other steps once a month from the first month after processing up to the sixth month, on determined times. Each step of the tests was repeated three times.

Sensory tests for the test sample and control sample (14 packages) consisted of evaluation of tissue, odor, color and taste with the method of Ranking and through execution of Freedman test (Iso 85 – 87, 1988). The sensory tests on the packaged samples with edible film and the control samples were repeated once in each step.

The results of microbial and chemical tests were analyzed by SPSS Software and Two Way Variance Analysis, and results of the sensory tests were analyzed by Ranking and Freedman Test.

whey protein covered samples during storage time. The mean total bacterial counts and *Staphylococcus* bacteria count in processed samples by whey protein were 2.47 and 1.61

log CFU/g and in control samples 3.13 and 2.18 log CFU/g from 1 day until six months after processing, respectively.

There was a meaningful difference in the microbial results in the control samples and covered samples from the first day up to the sixth month ($p<.05$). Analysis of results obtained using Two – Way ANOVA at 95 % confidence level indicate significant differences in total bacterial counts and *Staphylococcus* bacteria count both in the covered samples as well as in the control samples ($p>.05$).

The chemical and proximate analysis results of the samples are shown in the Tables 3 and 4. The mean of moisture, peroxide value, TVB-N, pH, free fatty acids, thiobarbitouric acid , protein, fat, ash and calorie in the covered samples were 73.91%, 0/13meq/kgoil, 9.84mg/100g, 6.25, 1.15gr/100, 0/006 mg/kg, 19.00%, 4.25%, 2.10% and 120.73kcal/kg and in control samples 59.43%, 3.75 meq/kgoil, 16.22 mg/100g, 6.71, 9.21gr/100, 0/15mg/kg, 18.2%, 4.00%, 1.80% and 108.8 kcal/kg, respectively.

Variations of free fatty acids were meaningful from one day after keeping up to five months ($p<.05$) and at the sixth months. It was not meaningful in control samples.

Variations of pH and TBA showed significant differences between the first day with the sixth month after processing in control samples ($p<.05$).

Analysis of results obtained using Two – Way ANOVA at 95% confidence level indicate no significant differences in moisture, free fatty acids, thiobarbitouric acid, TVB-N and peroxide valuable in the covered samples ($p>.05$) compared with the control ones ($p<.05$).

According to Kolmogrof-Smirnof Test, distribution of microbial and chemical data was normal.

The sensory results of the samples are shown in the Table 5. Based on Freedman test in sensory properties, there was a meaningful difference between covered samples and control sample ($p<.05$).

According to the results of the tests and statistical analysis, better general quality was observed in the covered samples compared to the control sample.

Based on the statistical test, the covered samples had preserved their quality up to the end of storage period while the control samples had lost their quality.

Table 1: Microbial analysis results in fresh Kilka fish (log CFU/g)

Bacteria Samples	Total bacterial counts	<i>Staphylococcus</i> bacteria
Whole Kilka	4.49±0.12	2.39±0.23
Cleaned Kilka fish	3.81±0.11	2.51±0.15

Table 2: Microbial factors results in covered and control samples during storage period (log CFU/g)

Bacteria Samples		Total bacterial counts		<i>Staphylococcus</i> bacteria	
Time		control samples	Covered samples	control samples	Covered samples
First day		3.67±0.21 ^a	3.43±0.32 ^a	2.85±0.12 ^a	2.25±0.12 ^a
First month		3.46±0.11 ^b	3.34±0.22 ^b	2.57±0.42 ^b	2.1±0.22 ^b
Second month		3.32±0.25 ^c	3.25±0.42 ^c	2.32±0.32 ^c	1.84±0.31 ^c
Third month		3.20±0.32 ^d	3.07±0.18 ^d	2.17±0.17 ^d	1.59±0.18 ^d
Forth month		2.95±0.14 ^e	2.69±0.14 ^e	2.00±0.18 ^e	1.35±0.11 ^e
Fifth month		2.77±0.12 ^f	2.00±0.11 ^f	1.72±0.22 ^f	1.20±0.15 ^e
Sixth month		2.47±0.24 ^g	1.45±0.17 ^f	1.69±0.28 ^g	1.00±0.13 ^e

The different letters in the same column within the different storage time indicate significant differences ($p<.05$).

Values are mean + standard deviation.

Table 3: Chemical factors results in covered and control samples during storage period

Experiments	PV value		FFA		TBA		pH	TVN		
	meq/kg oil	gr/100	gr/100	mg/kg	mg/kg	mg/100gr		mg/100gr	mg/100gr	mg/100gr
Samples	control	Covered	control	Covered	Control samples	Covered	control	Covered	control	Covered
Time	samples	samples	samples	samples		samples	samples	samples	samples	samples
First day	0.20±0.01 ^a	0.08±0.10	4.10±0.25 ^a	1.1±0.15	0.03±0.024 ^a	0.006±0.001	6.2±0.10 ^a	6.2±0.20	9.8±0.36 ^a	9.8±0.35
First month	1.7±0.10 ^b	0.08±0.20	6.83±0.32 ^b	1.1±0.25	0.07±0.1 ^a	0.006±0.002	6.2±0.20 ^a	6.2±0.10	12±0.30 ^b	9.8±0.20
Second month	3.20±0.10 ^c	0.08±0.20	8.34±0.25 ^c	1.1±0.35	0.10±0.03 ^{ab}	0.006±0.002	6.4±0.10 ^a	6.2±0.20	14.6±0.30 ^c	9.8±0.25
Third month	4.50±0.10 ^d	0.08±0.15	9.52±0.28 ^d	1.1±0.02	0.14±0.01 ^{ab}	0.007±0.02	6.7±0.15 ^b	6.3±0.20	16.2±0.30 ^d	9.8±0.55
Forth month	6.00±0.25 ^e	0.09±0.26	10.96±0.46 ^e	1.2±0.05	0.17±0.01 ^{ab}	0.007±0.001	7±0.15 ^{bc}	6.3±0.20	18.5±0.30 ^e	9.9±0.41
Fifth month	5.60±0.6 ^f	0.09±0.15	12.37±1.31 ^f	1.2±0.15	0.25±0.03 ^{abc}	0.007±0.11	7.2±0.26 ^{bc}	6.3±0.10	20.8±0.50 ^f	9.9±0.30
Sixth month	5.10±0.30 ^g	0.1±0.25	12.38±0.10 ^f	1.3±0.35	0.32±0.02 ^{bc}	0.008±0.025	7.3±0.15 ^c	6.3±0.15	21.7±0.25 ^g	9.9±0.30

The different letters in the same column within the different storage time indicate significant differences ($p<.05$). Values are mean + standard deviation.

Table 4: Proximate analysis results in covered and control samples during storage period

Experiments Time	Moisture %		Protein %		Fat %		Ash %		Calorie kcal/kg	
Samples	Control	Coated	control	Coated	Control	Coated	Control	Coated	control	Coated
First day	72.2±0.35 ^a	73.93±0.04	18.2±0.65	19.00±0.07	4.00±0.36	4.25±0.03	1.80±0.15	2.10±0.25	108.8	120.73
First month	67.35±0.25 ^b	73.93±0.14	18.2±0.65	19.00±0.07	4.00±0.36	4.25±0.03	1.80±0.15	2.10±0.25	108.8	120.73
Second month	63.20±0.10 ^c	73.93±0.05	18.2±0.65	19.00±0.07	4.00±0.36	4.25±0.03	1.80±0.15	2.10±0.25	108.8	120.73
Third month	58.90±1.61 ^d	73.91±0.04	18.2±0.65	19.00±0.07	4.00±0.36	4.25±0.03	1.80±0.15	2.10±0.25	108.8	120.73
Forth month	54.15±0.16 ^e	73.91±0.04	18.2±0.65	19.00±0.07	4.00±0.36	4.25±0.03	1.80±0.15	2.10±0.25	108.8	120.73
Fifth month	50.43±0.14 ^f	73.90±0.04	18.2±0.65	19.00±0.07	4.00±0.36	4.25±0.03	1.80±0.15	2.10±0.25	108.8	120.73
Sixth month	46.16±0.17 ^g	73.90±0.06	18.2±0.65	19.00±0.07	4.00±0.36	4.25±0.03	1.80±0.15	2.10±0.25	108.8	120.73

The different letters in the same column within the different storage time indicate significant differences ($p<.05$). Values are mean + standard deviation.

Table 5: Sensory analysis and statistical analysis results in covered and control samples

Properties Samples	Color	Odor	Tissue	Taste	Total acceptance
Covered samples	75 ^a	70 ^a	79 ^a	79 ^a	79 ^a
Control samples	110 ^b	106 ^b	96 ^b	104 ^b	112 ^b
Score	110 – 75	106 – 76	99 – 69	104 – 64	112 – 68
LSD	35>19.6	36>19.6	23>19.6	40>19.6	44>19.6

The different letters in the same column within the different storage time indicate significant differences ($p<.05$).

Values are mean + standard deviation.

Discussion

The total bacterial counts and *Staphylococcus* bacteria in the covered samples have been decreased. This may be due to the effect of lactic acid, acetic acid, probiotic bacteria and production of ethanol and nisin. Pro-biotic micro-organisms of whey protein are able to produce organic acids such as lactic acid, acetic acid, nisin bacteriocin, hydrogen peroxide, ethanol, acetaldehyde, ammoniac, di-acetyl, Eh and pH reduction (Adams & Moss, 2002; Nasar *et al.*, 2008 & Seifzadeh, 2002). These compounds have antimicrobial specifications and are able to prevent the growth of *Staphylococcus*, *Pseudomonas*, *Coliform* and *Escherichia coli* (Zinoviadou *et al.*, 2009 & Cagri *et al.*, 2002 & 2003). Naturally, the Whey Proteins have a globule structure and high solubility and emulsification. This film has protein, lactose and minerals and is able to increase the ability of connection to water in Kilka (Stuchell & Krochta, 1995). Absorption of water by protein and connection and linking of protein chains to each other increase on the size of protein that

may cause to increase the viscosity, humidity and texture at the covered samples by this film, compared with the control sample (Sanker & Raghunath, 1995). In the control samples, because of presence of empty space between fish filets and also temperature rise and fall of the cold-storage, the Kilka in the packages has lost the humidity and will be dried and naturally it leads to decrease in weight about 3.5% after three months (Kochakian, 2002). This status may appear because of production of ice crystals at the products, too. Freezing is considered a basic action of dehydration that causes to exit the frozen humidity in vapor status from the food products. Air circulation at the cold- storage may intensify the humidity exit, too. This status may quicken the ruin of proteins and oxidation of lipids, and causes to decrease the quality of the taste and color change in Kilka without cover (Deman, 1999 & Safari, 1990 & Fatemi, 1990).

Amount of TVB-N in Kilka without cover during the storage period at the cold-storage showed an increase. This may be due

to the effect of decrease of humidity and production of free fatty acids on the denaturizing of the Protein. These factors decreased in covered samples compared with the control sample. We can relate its reason to the specification of whey protein that cause to prevention of the proteolyses, decrease of humidity and production of free fatty acids on denaturizing of protein and naturally increase in TVB-N(Morrissey et al., 2009; Piyachomkwan and Penner, 1995; Salmani et al., 2002).

Amount of ash showed an increase in the covered samples compared with control sample, That is because of presence of sodium, potassium, calcium, manganese and phosphate ions, and also presence of sulfur holder amino acids in the compound of whey protein and this point that these amino acids are considered as mineral elements of food products (Lyndon, 1999).

Amount of protein showed an increase in covered samples compared with the control sample. This may be due to the effect of some protein compounds such as α -lactalbomine and β -lactoglobine of this film (Shahidi and Botta, 1990).

Concentration of lipid showed an increase in the covered samples compared with control sample. That is because of presence of glycerides of milk lipid, phospho-lipids and lipoproteins in whey protein. Oxidation of the lipid arising from the reaction of lipid with oxygen and its hydrolyze are influenced by the action of lipolytic enzymes (Deis,2006 and Min et al., 2006). The enzyme of lipase of the tissue, the excreted lipolytic enzyme from the *Staphylococcus* bacteria and those enzymes

that are released from the dead and analyzed bacteria, have been able to be active at low water activity and may cause hydrolyze of lipids and production of non-saturated fatty acids through the lipolyze process(Cappuccino and Sherman, 1999; Min et al., 2005, 2007). Releasing of fatty acids with high numbers of carbons by lipase enzyme may not provide a specific bad taste but through passing the time, effects of accumulation of free fatty acids in muscles of fish because of their combination with protein of muscle may cause unfavorable taste and tissue damages (Crapo et al., 1999). In addition to fatty acids, because of protein denaturation, it provided some tissue changes and decrease of quality. In the control samples, concentration of these acids has been increased from the first month up to the fifth month. But the nearly constant concentration of these acids at the last processes of keeping has probably been because of decrease of raw materials and increase of oxidation in free fatty acids (Silva and Ammerman, 1993; Rezaei et al., 2002).

Based on the conducted experiments, the amount of the free fatty acids has been measured less in the samples covered by whey protein, compared with the control sample. This may be due to the prevention from losing the water from the surface and inside the body by the fine holes present at the body surface, prevention from the oxygen contact with the fish tissue and combination with non-saturated fatty acids and oxidation, and lack of absorption of light by the fish body surface (Lyndon, 1999; Ahvenainen, 2003).

Freezing caused the fish tissue denaturizing and moisture loses leads to the weight loss which may be due to the oxygen penetration

into the fish tissue and oxidation increasing of free fatty acids and peroxide value. But, peroxide, will start to induce decomposition, leading to the aldehydes, ketone and seton production, subsequently, in peroxide value reduction to over time (Krochta *et al.*, 1996). First products of lipid oxidation are hydroperoxide and they are broken to secondary products such as aldehydes that can react with proteins. This compound react with pigments and other molecules present at the fish, too. Therefore it leads to loss the color and producing bad smell (Rezaei *et al.*, 2003).

A suitable index for determination of progress in fat oxidation and producing of carbonyl compounds is measurement of TBA. Presence of such compounds in fish meat causes some changes in its sensory specifications such as taste and smell (Shah, 1999; Aubourg *et al.*, 1995).

Amount of peroxide and TBA was measured lower in the covered samples compared with control sample. We can say that it is because of prevention of decrease of water in fish tissue while freezing, decrease of water activity and free radicals production, covering specifications of edible film, prevention of oxygen absorption, prevention from humidity decrease, decrease of oxidation and production of secondary products of oxidation such as aldehydes and specifications of these compounds (Tall and Harris, 1995; Gigirey and Desousa, 1999).

In the present research, amounts of measured TBA had an increasing proceeding up to the end of cold - storage in the control Kilka. This is may be due to the effect of freezing on decrease of humidity, decrease of the water

activity factor, increase of oxidation at low water activity, production of free radicals, production of free fatty acids and being ready for oxidation (Hegenbart, 2006; Coles and McDowell, 2003).

The covered samples have better taste and smell compared with the control sample that is due to the effect of di-acetyl and lactose in the compound of this film. Proteins of α -lactalbomine and β -lactoglobine present at whey protein cause production and accumulation of big solution and masses. This process may lead to production of small sedimentations. These accumulations and connection of them with water, have increased the amount of water and consequently cause to increase the viscosity and development of the taste of the product (Bigelow and Lee, 2007; Chapman *et al.*, 1997).

Amount of calorie has shown an increase in the samples covered by edible film compared with the control samples. It is because of increase of the added value raised from the edible covers.

Based on the conducted experiments, the amount of protein, lipid and carbohydrates in the samples covered by this film is more than the control sample and with consideration to the conversion coefficient of each gram of protein, carbohydrate and lipid to energy. Amount of calorie of these samples has shown an increase (Marsh and Bugusu, 2007).

Through passing of time, lipid oxidation products such as hydro-peroxides have been analyzed and some compounds such as aldehyds and others have been produced in the control sample. These compounds have alkali specification and cause an increase in the pH

of the product. Therefore in the covered samples, changes of pH during the time storage were not significant (Anker and Hermansson, 2010).

Similar results were obtained by Rokwer, Bigelow, Fujki, Trout and Manish.

According to the results of the experiments, a meaningful difference in the index of total acceptance among the covered samples compared with the control sample were observed. No statistically significant differences were observed in the results of chemical experiments of the covered samples and presence of the meaningful difference at the results of the chemical experiments of the control sample. The covered samples up to the end of storage period at cold-room had a favorite quality but the control samples had lost their quality.

References

Adams, M. R. and Moss, M.O., 2002. Food microbiology. R. S.C. pp.37-44.

Ahmed, E. M., Cornell, J. A., Tomaszewski, F. B. and Deng, J. C., 2006. Effects of salt, Tripolyphosphate and sodium alginate on the texture and flavor of fish patties prepared from minced Sheep head. *Journal of food science*, 48(4), 1078–1080.

Ahvenainen, R., 2003. Novel food packaging techniques, Washington. CRC Pub. 590P.

Andrews, W. H. and Hammack, T. S., 2003. Food sampling and preparation of sample homogenate. FDA. 20P.

Anker, M. and Hermansson, A. M., 2010. Edible and Biodegradable Whey Protein Films as Barriers in Foods and Food Packaging. Nordic food pack. 30 P.

Aubourg, S. P., Manisilla, M. R. and Sotelo, C. G., 1995. Differential lipid damage in various muscle zones of frozen hake. *lebensm unters forsch. Chemistry and Materials Science Journal*, 208, 189–193.

Bennett, R. W. And Lancette, M., 2001. *Staphylococcus aureus*. FDA. 35P.

Gigirey, B. and Desousa, J., 1999. Chemical changes and visual appearance of albacore tuna as related to frozen storage. *Journal Food Science*, 64, 20–24.

Bigelow, W. and Lee, C. M., 2007. Evaluation of Various Infused Cryoprotective Ingredients for Their Freeze–Thaw Stabilizing and Texture Improving Properties in Frozen Red Hake Muscle. *Journal of Food Science*, 72, 56–64.

Cagri, A., Ustunol, Z. and Ryser, E. T., 2002. Inhibition of three pathogens on bologna and summer sausage using antimicrobial edible films. *Journal of Food Science*, 67, 2317-2324.

Cagri, A., Ustunol, Z., Osburn, W. and Ryser, E. T., 2003. Inhibition of *Listeria monocytogenes* on Hot dogs using antimicrobial whey protein based edible coating. *Journal of Food Science*, 68, 291-299.

Chapman, K. W., Xiaowen, L. U., Weilmeier, D. and Regenstein, J. M., 1997. Edible films on fish Seafood safety. Processing and Biotechnology. pp.139-150.

Coles, R., McDowell, D., 2003. Food packaging technology, Blackwell Publishing.

Crapo, C., Himelboom, B., Pfutzenreuter, R. And Lee, C., 1999. Texture modification processes for Giant Grenadier filets, *Journal of Aquatic Food Product Technology*, 8(4), 27-40.

Cappuccino, J. C. and Sherman, N., 1999. Microbiology, Benjamin/Cumming Science Publishing.

Deman, J. M., 1999. Food chemistry, (Translator: Akbarzadeh, B). Publishing Nemati. pp.35–39.

Deis, R. C., 2006. The complexity of shelf life stability, Virgo pub.

Fatemi, H., 1990. Food chemistry. Company publication, pp.255-256. (in Persian)

Fahim, H. R., 1994. Frozen prepared fish cutlet dough Kilka. Kilka Company. pp (in Persian)

Feng, P., Weagant, S. D. and Grant, M. A., 2002. Enumeration of *Escherichia coli* and the *Coliform* bacteria. FDA, 30P.

First consulting engineers.,1998. Kilka products complex. Fisheries and Aquaculture Study Inbox. 146P.

Hasegawa, H.,1987. Laboratory manual on analytical methods and procedures for fish and fish products. Southeast Asian Fisheries Development Center. pp.130–180.

Hegenbart, S., 2006. The changing face of shelf life, Virgo Publishing.

Holt, J. G., Krieg, R. N., Sneath, P. H. A., Staley, J. T. And Williams, S. T., 1994. Bergeys manual of determinative bacteriology ninth edition, Williams & Wilkins. 895P.

Iranian National Standards No. 493., 2003. Edible fats and oils – sampling. Institute of Standards and Industrial Research of Iran. (in Persian)

Iranian National Standards No. 742., 2002. Meat and meat products – determination of total fat - test method. Institute of Standards and Industrial Research of Iran. (in Persian)

Iranian National Standards No. 744., 2001. Meat and meat products - Determination of total ash - test method. Institute of Standards and Industrial Research of Iran. (in Persian)

Iranian National Standards No. 924., 1994. Measuring total protein in meat and its products. Institute of Standards and Industrial Research of Iran. (in Persian)

Iranian National Standard No. 1028., 2007. Meat and meat products – measurement of pH. Institute of Standards and Industrial Research of Iran. .pp?? (in Persian)

ISO85_ 87.,1988. Sensory analysis _ methodology first edition. ISO.

Iranian National Standard No. 5625., 2002. Cleaned and frozen Kilka - specification and test methods. Institute of Standards and Industrial Research of Iran. (in Persian)

Iranian National Standard No 8867., 2005. Animal products, Animal feeding stuffs and feaces or urine – Determination of gross calorific value–Bomb calorimeter method. Institute of Standards and Industrial Research of Iran. (in Persian)

Iranian National Standard No.10494., 2006. Animal and vegetabla fats and oils determination of 2 thiobarbituric acid value direct method. (in Persian)

Kochakian, A., 1992. Kilka fish canned. Iranian Fisheries Company. (in Persian)

Kochakian, A., 1995. Preparation of fish products and Kylka Fish ball and its storage. MSc thesis, Islamic Azad University, North Tehran. (in Persian)

Kochakian, A., 2002.a Preparation of fish meat without bones from Kilka and its packing and distribution. Organization for Research and Education Center. (in Persian)

Kochakian, A., 2002.b Preparation of fish sauce Kilka. Iranian Fisheries Research Organization. (in Persian)

Kochakian, A., 2002.c Physico-chemical variations in the fish Kilka after fishing and protective

materials effects on it. Organization for Research and Education Center. (in Persian)

Krochta, J. M., German, J. B. and Carthy, M. J., 1996. Edible films for preventing loss of quality in frozen fish, California sea grant. Biennial Report of Completed Projects pp.1992-1994. 73-78.

Lyndon, A., 1999. Biochemistry of food, (Translator: Abromand, E). Publishing Ramand and Agricultural Sciences. 260P.

Marsh, K. And Bugusu, B., 2007. Food packaging roles materials and environmental issues. *Journal Food Packaging*, 72(3), 39-56.

Martin, A. M., 1994. Fisheries processing. Chapman and Hall.

Maturin, L. J. and Peeler, J. T., 2001. Aerobic plate counts. FDA. 10P.

Min, S., Harris, J. and Krochta, J., 2005. *Salmonella enteritica* and *Escherichia coli* O157:H7 inhibition by lactoferrin, lysozyme and lactoperoxidase systems and by edible Whey protein films incorporating lactoperoxidase systems. IFT Annual Meeting. Pp??

Min, S., Rumsey, T. R. and Krochta, J., 2006. Lysozyme diffusion in smoked salmon coated with whey protein films incorporation lysozyme, *China Development Research Foundation*, 1, 5-29.

Min, S., Harris, J. and Krochta, J., 2007. *Listeria monocytogenese* inhibition by whey protein films and coatings incorporatin the lactoperoxidase system, IFIS Publishing.

Min, S., Harris, J. and Krochta, J., 2007. Time to talk turkey inhibition of *Salmonella enteritica* and *Escherichia coli* O157:H7on rooted turkey by

Safari, M., 1990. Physicochemical principals of food storage. *Tehran University Publications*. 209(221), 224-226. .(in Persian)

edible whey protein coating incorporating the lactoperoxidase system. *Journal of Food Protection*, 69(4), 784-793.

Morrissey, M. T., Chung, Y. C. and An, H., 2009. Whey protein concentrate as a proteinase inhibitor in pacific whiting surimi. *Journal of Food Science*, 61(2), 367-371.

Nasar, A., Kasra Kermanshah, R. and Nahava, A., 2008. The effect of organic acids and Nisin inhibitor concentrations on the growth of *Bacillus cereus* isolated from cheese. *Journal of Food Sciences and Nutrition*, pp.21-30.

Piyachomkwan, K. and Penner, M. H., 1995. Inhibition of pacific whiting surimi – associated protease by Whey protein concentrate. *Journal Food Biochem*. 18(1), 345-358.

Rezaei, M., Sahari, M., Moeene, S., Safari, M., Rezaeian, M. and Ghafouri, F., 2003. Evaluation of qualitative properties in Kilka in freezing Storage . *Journal of Marine Science*. 55-64. (in Persian)

Rezaei, M., Sahari, M. and Moeeni, Q., 2006. Quality assessment of Kilka fish fat during storage in different freezing temperatures. *Journal Of Science and Technology Of Agricultural and Natural Resources*, pp.35-45. (in Persian)

Salmani, H. and Nikopur, E., 2003. Review of Sturgeon storage time in the cold room. *Journal of pejouhesh and sazandegi*, pp.98-102. .(in Persian)

Salmani, H., Gholamipoor, S. and Yosefian, M., 2002. Nitrogen and histamine variations of Kilka fish in storage period. *Iranian Journal of of Fisheries*, 2, 31-43. (in Persian)

Sedaghat, N., 1996. Technology, food packaging. Barsava Publkishing. 260P. .(in Persian)

Seifzadeh, M., 1381. Role of lactic acid bacteria as protective food. Thirteenth Congress of the food industry. 178P. .(in Persian)

Seifzadeh, M., 2007. Application of edible Films in fish products packaging. *Journal of World Aquaculture* seventh. pp.34-40. .(in Persian)

Shah, A. J., Hansen, B. and Larsen, R. B., 1999. Fish crackers produced by extrusion with addition of whey protein concentrate. *Food Australia*, 51(3), 104–106.

Shahidi, F. and Botta, J. R., 1994. Seafood's, chemistry, processing technology and quality. Chapman & Hall.

Shariatifar, M., 2007. Use of edible films for increasing shelf life of food products. 16th National Congress Of Food Industry. 110P. .(in Persian)

Shbanpor, B. and Shabani, A., 2006. Effect of different washing conditions on chemical composition and properties of Kilka Surimi gel. 16th National Congress Of Food Industry. 65P. .(in Persian)

Shojaei, A. M.,1996. Preparation of Kilka crackers. Institute of Fisheries Research and Education. pp.?? .(in Persian)

Shojaei, A. M., 2002. Evaluation of Kilka oil for food uses. Iranian Fisheries Research Organization. (in Persian)

Silva, J. and Ammerman, G. R., 1993. composition, lipid change and sensory evaluation of two sizes of channel catfish during frozen storage. *Journal Applied Aquaculture*. 2(2), 39–49.

Stringer, M. and Dennis, C., 2000. Chilled foods. CRC press.

Stuchell, Y. M. and Krochta, J. M., 1995. Edible coating on frozen King salmon: Effect of Whey protein isolate and acetylated monoglycerides on moisture loss and lipid oxidation, *Journal of Food Science*, 60(1), 28-31.

Tall, J. and Harris, P., 1995. Rancidity in frozen fish. *Journal Fish Oil: Technology, Nutrition and Marketing*. pp.35–48.

Zare Gashti, Gh., 1995. Evaluation of sturgeon and Kilka products in vacuum packaging. Institute of Fisheries Research and Education.

Zinoviadou, K. G., Koutsoumanis, K. P. And Biliaderis, C. G., 2009. Physico – chemical properties of Whey protein isolate films containing oregano oil and their antimicrobial action against spoilage flora of fresh beef. *Meat Sciences*, 82, 338–345.