# Identification and probiotic properties of lactic acid bacterial isolated from freshwater fish

Hanol Bektas Z.1\*; Ucar F.B.2; Giray B.2

Received: February 2016 Accepted: December 2018

## **Abstract**

Since chemotherapeutic agents used to control microbial fish diseases have led to resistance in bacteria, alternative methods have emerged in recent years. In this regard, identification of lactic acid bacteria with good probiotic properties and determination of their probiotic properties are extremely important. In this study, 25 lactic acid bacteria were isolated from freshwater fish and identified at species level using phenotypic, biochemical and molecular tests. The pH tolerance, antagonistic activity and antibiotic sensitivity of these 25 strains were examined, and they were considered as bacteria displaying the best activity in the potential probiotic treatment of fish diseases. *Lactococcus lactis* species numbered F2, F4, F9 and F10 were determined to have a potential probiotic capacity.

Keywords: Probiotic, Molecular identification, Lactic acid bacteria, ARDRA technique

<sup>1-</sup>Aquaculture Research Station Directorate, Egirdir-ISPARTA/TURKEY

<sup>2-</sup>Department of Biology, Basic and Industrial Microbiology Section, Faculty of Science, Ege University, İzmir/TURKEY

<sup>\*</sup>Corresponding author's Email: zubeydehanol17@gmail.com

### Introduction

Nowadays, fish diseases emerging due to the increase in commercial-scale aquaculture operations have become a major limiting factor in aquaculture (Bondad-Reantaso al.. 2005: et Kesarcodi-Watson 2008: al., et2009). Subasinghe *et al.*, Major bacterial pathogens leading to fish diseases are Aeromonas salmonicida (Furunculosis), Yersinia ruckeri (enteric red mouth disease), Vibrio anguillarum (vibriosis), Aeromonas hydrophila (minor surface lesions, septicemia, internal bleeding) and Lactococcus garvieae' (hemorrhagic septicemia and meningoencephalitis) (Balcazar et al., 2007 ab; Öztürk and Altınok, 2014). These diseases are quite common all over the world and cause serious economic losses in salmon and trout aquaculture (Austin and Austin, 2012). Vaccination antibiotics and extensively used combat fish to diseases. When administered to immunologically immature fish. vaccination is ineffective. Antibiotics cause an increase in resistance among pathogenic bacteria and when these resistant bacteria enter the intestinal tract of humans, they pose a threat to the treatment of diseases (Denev et al., 2009). Therefore, during the last 10 years, there has been an increased interest in using probiotics to prevent the growth of pathogenic microorganisms and to decrease the rate of fish diseases (Ringo and Gatesoupe, 1998; Ringo, 2004; Perez-Sanchez et al., 2011). Good probiotic stomach acids should generate resistance to damage caused by bile salts and

proteases; reduce the intestinal pH by producing lactic acid and thereby prevent the growth of pathogenic bacteria. reduce the production of several toxic and carcinogenic metabolites, help the absorption of minerals such as calcium due to the increased intestinal acidity, and be able produce compounds such bacteriocins. organic acids. and hydrogen peroxide which inhibit the growth of pathogenic microorganisms, and vitamin B and K. Lactic acid bacteria (LAB) have the abovementioned properties and they are abundantly found in nature (Perez-Sanchez et al., 2011; Corcionivoschi et al., 2010). In studies conducted so far, lactic acid bacteria have demonstrated to greatly differ from one species of fish to another and from one geographic region to another (Buntin et al., 2008; Bushell and Burns, 2012). Therefore, isolation and identification of bacteria in the gastrointestinal system of fish have gained importance. Since phenotypic and biochemical tests used for the identification of LABs are inadequate, molecular biological methods have been used recently. Through molecular biological methods, it has been possible to understand the characterization of the microbiota of the stomach and gut, and interactions of bacteria with bacteria and bacteria with host in sickness and in health (Brunvold et al., 2007; Liu et al., 2008).

In this study, the aim was to determine phenotypic and molecular characterization, antagonistic effects, pH tolerance and antimicrobial activities of LAB isolated from freshwater fish due to the need for new probiotics to fight diseases.

## Materials and methods

Isolation of bacteria

Gut samples obtained from healthy rainbow trout (Oncorhynchus mykiss), carp (Cyprinus carpio), pool fish (Carassius gibelio) and zander (Sander lucioperca) living in Egirdir Lake were diluted for the isolation of LAB. The LAB isolates were inoculated in the selective media and in TSA to enhance the chances of isolation. All the fish were treated in 1 liter of water containing 150 triacine mg methanesulphonate (MS-222) for 15 min and killed (the ethic committee with 68385072-604/0224 number in 01.03.2014 date obtained from Mediterranean Aquaculture Research Directorate-TURKEY) before the start of the study. Then one gram of the sample obtained from the fish gut was placed in 10 ml PBS (phosphatebuffered saline) and diluted 10<sup>-7</sup> times. 0.1 ml of dilutions were seeded on TSA (Tryptic soy agar: Merck KGaA Darmstadt Germany) (this medium was used to increase the chances of isolation), MRS (de Man Rogosa and Sharpe agar (Merck KGaA Darmstadt Germany) and M17 (Conda Pronadisa) agar and incubated at 22 °C under aerobic and anaerobic conditions (The MRS petri was incubated in anaerobic jar-Anaerocult A to produce anaerobic conditions) for 24-48 hours (Balcazar et al., 2007a). Colonies isolated from TSA, MRS and M17 agars were stored at -80°C in TSB (Tryptic Soy Broth: Merck KGaA Darmstadt Germany) containing 15% glycerol (Balcazar *et al.*, 2007 b; Perez-Sanchez *et al.*, 2011).

Phenotypic biochemical and of characteristics the 25 isolates obtained from the colonies grown on TSA and M17 MRS, agar were identified with phenotypic and biochemical tests using the Bergeys Manual of Systematic Bacteriology (Hammes and Hertel, 2009; Teuber, 2009).

DNAextraction and molecular identification of the genus Lactobacillus DNA isolation was carried out with the rapid phylogenetic analysis (Liu et al., 2000). For the molecular biological identification, Lacto 16S-F (GGA ATC TTC CAC AAT GGA CG) and Lacto 16S-R (CGC TTT ACG CCC AAT AAA TCC GG) primers specific to the genus Lactobacillus were used and a partial region (216 bp) of 16 S rDNA was amplified using PCR (Polymerase Chain Reaction) (Abdulamir et al., 2010).

Amplified ribosomal DNA restriction analysis (ARDRA) and sequencing Then, after the 16S rDNA about 1.5 kb regions of the 25 isolates were (5'amplified with 27F AGAGTTTGATCCTGGCTCAG-3') 1492R (5'and GGYTACCTTGTTACGACTT-3') primers using PCR through ARDRA (Amplified Ribosomal DNA Restriction Analysis). They were cut with the Msp I (Promega), Hae III (Promega), Hinf I (Promega) enzymes and the fragments were formed through electrophoresis

(Soto et al., 2010). The images were grouped as follows. After the 16srDNA regions isolated from the organisms were cut with 3 different restriction enzymes, electrophoresis was carried out. As clearly described in the electrophoresis images, samples showing the same band patterns were collected in the same group and one sample from each group was sent to the sequence. Since there were 2 groups in our study, obtaining the accession numbers of the sequence results of the two samples representing these 2 groups was sufficient. However, as noted in the text, to avoid errors due to slight differences and to be sure, not one sample but 6 samples from one of the groups (F2, F4, F40, F53, F66, F69), and two samples from the other group (F70, F36) were sent the sequence. Three strains selected to represent each group were sequenced with the same primers used in PCR. The sequence data obtained were compared with the sequences in the GenBank database using the BLAST algorithm and then sent to NCBI (National Center for Biotechnology Information) to receive an access number.

# Detection of antagonistic activity

Antagonistic activities of these 25 strains against the test pathogen organisms of Salmonella typhimirium (ATCC 14028), Escherichia coli O157:H7, Pseudomonas aeruginosa ATCC 27853, Yersinia ruckerii, L. garviae ATCC 49156 were determined with the disc diffusion technique and was performed as triplicates. After the

pathogenic bacteria were incubated on the TSA agar at 22 °C for 24 hours, their inoculum density was adjusted to 0.5 to Mac Farland (50-100 µl), and they were seeded on the general medium through swab cultivation. At the same time, fresh cultures of lactic acid bacteria prepared by incubating LAB on the MRS broth for 24 hours were centrifuged at 8000 rpm, at 4 °C for 5 min, and cell-free solutions were obtained. After the sterile discs placed in petri dishes containing pathogenic bacteria were soaked in the obtained supernatant, the petri dishes were incubated at 22 °C for 24 hours. The inhibition zones which formed after incubation were measured and the results were recorded (Allameh et al., 2012; Chemlal-Kherraz et al., 2012).

## Antimicrobial susceptibility testing

Antimicrobial susceptibility testing of isolates was performed with the disc diffusion method in accordance with the Clinical guidelines for and Laboratory Standards Institute (CLSI), and the following antibiotics were tested with the disk diffusion method: Doxcycline (30 μg), Enoxacine (10 μg), Erythromycin (15µg), Florfenicol (30 Trimethoprim/sulfamethoxazole μg),  $(1.25/23.75 \mu g)$ , Enrofloxacin  $(5\mu g)$ , Oxytetracycline  $(30 \mu g)$ , Chloramphenicol (30 µg) (CLSI, 2008).

# pH tolerance

The isolates' tolerance towards different pH conditions (pH 1 and pH 3) was determined using the modified method implemented by Perez-Sanchez *et al.* (2011).

# Experimental infection

For experimental infection the method used by N. Haines et al. (2013) was used with a minor modification. To determine if F4 isolate which has the most proper probiotic potential from fish would produce infection in rainbow trout, a group of 20 trout (9-10g) were inoculated intraperitoneally with 1 x10<sup>5</sup> bacteria by optical density in a final volume of 10 ul sterile PBS. Animals were held in 2 (PVC  $\pi$  150cm=880L) at 16°C and monitored daily for clinical of streptococcosis, including evidence of external hemorrhage, exophthalmia, lethargy, and loss of appetite. Two weeks after inoculation all fish were euthanized and the spleens were aseptically removed, macerated and suspended in liquid media for reisolation as described above.

# Results

Upon the completion of the phenotypic tests, it was found that 22 of the 25 isolates belonged to the genus Lactobacillus whereas 3 of them belonged to the genus Lactococcus. After being amplified with primers specific to the genus Lactobacillus, all of these 25 isolates yielded 216 bp band specific to the genus Lactobacillus (Fig. 1). Definitive identification was made **ARDRA** after the method implemented and sequence analysis of the selected strains exhibiting identic restriction pattern were performed (Figs. 2, 3, 4). However, after the sequence analysis, while 23 of these 25 strains were identified, 2 were not (F17 and F48). The 23 identified strains were divided into two groups. One of the groups included 19 L. lactis strains, and the other included 4 Carnobacterium maltoramaticum strains. Due to the small visually detectable differences within these two groups, accession numbers of these 8 strains were obtained in order to avoid possible errors. Thev are as follows: Lactocococcus lactis strain F2. KM017400; L. lactis strain F4. KM017401; L. lactis strain F40. F17402; L. lactis strain F53, F17403; L. lactis strain F66, 17404; L. lactis strain 17405; F69. Carnobacterium maltoramaticum strain F36. KM017406; C. maltoramaticum strain F70, KM017407.

Results of antagonistic effects of the 25 isolates are given in Table 1. Of the strains, those numbered F2, F4, F9 and F38 showed the highest antagonistic effects against four of the five test bacteria. Strains numbered F7, F37 and F38 showed the highest antagonistic activity against L. garviae ATCC 49156; F4, F10, F30 and F38 against E. coli 0157: H7; F2, F4, F7, F8, F9, F33, F38. F54 F53 and against typhimirium; F2, F4, F6, F8, F9, F10, F30, F38, F39, F40, F48, F53, F54, F70 and F75 against P. aeriginosa ATCC 27853; and F2, F6, F9, F38, F39, F68, F69 and F75 against Yersinia ruckerii. The results of the antibiotic susceptibility test are given in Table 2. Of the samples studied, only those numbered F8 and F48 were sensitive to antibiotics tested. The highest antibiotic resistance was to Enoxacin, Oxytetracycline and Chloramphenicol. All the strains were sensitive to Vancomycin. Except for those

numbered F70 and F72, all the strains were sensitive to penicillin.

pH tolerance results are given in Table 3. While no growth was observed in the majority of the samples at pH 1, most of the samples proliferated at pH 3

after a three-hour incubation. The strains which displayed the highest survival rate at pH 3 were those numbered F2, F4, F8, F9, F10, F48, F53 and F69.

Table 1. Diameter of inhibition zone (mm).

Numbers of isolates	E. coli	S.typhimirium	P.aeruginosa ATCC 27853	Y. ruckerii	L.garviae ATCC 49156
F2	9	12	14	16	10
F4	12	10	14	2	14
F6	11	9	10	12	16
F7	11	11	4	6	18
F8	8	11	10	8	2
F9	10	10	11	10	8
F10	13	9	11	4	6
F17	8	7	-	2	6
F30	14	11	10	2	2
F33	11	10	-	-	-
F36	8	4	-	-	2
F37	9	5	-	6	16
F38	10	10	12	10	18
F39	-	8	11	10	10
F40	-	6	10	-	2
F48	2	7	12	-	-
F53	4	13	12	4	2
F54	3	12	16	4	4
F66	-	8	8	6	6
F68	4	6	8	12	6
F69	5	7	8	12	6
F70	-	-	10	4	6
F71	4	9	-	2	2
F72	6	4	-	2	2
F75	8	9	10	10	6

Table 2: Results of antibiotic susceptibility testing.

İsolates	Doxycyline	Enoxacin	Erithromycin	Florfenicol	Trimethoprim	Oxytetracycline	Enrofloxacine	Chloramphenicol
F2	S	R	S	S	S	R	R	R
F4	S	R	R	S	S	R	S	R
F6	S	R	S	S	S	R	S	S
F7	S	R	S	S	S	R	S	S
F8	S	S	S	S	S	S	S	S
F9	S	S	S	S	S	R	S	S
F10	S	S	S	S	S	R	S	R
F17	S	I	S	S	S	S	S	R
F30	S	I	S	S	S	I	S	R
F33	S	S	S	S	S	R	S	R
F36	S	S	S	S	S	S	S	R
F37	S	S	S	S	S	S	S	R
F38	S	R	S	S	S	R	I	S
F39	S	R	S	S	S	R	S	S
F40	S	I	S	S	S	R	S	R
F48	S	S	S	S	S	S	S	S
F53	S	R	S	S	S	R	S	R
F54	S	S	S	S	R	R	S	I
F66	S	R	S	S	S	R	S	R
F68	S	R	S	S	S	R	I	S
F69	S	R	S	S	S	R	S	S
F70	S	S	S	S	S	S	S	R
F71	S	S	S	S	S	S	S	R
F72	S	S	S	S	S	S	S	R
F75	S	S	S	S	S	R	S	R

\*S: sensitive, \*R: resistant, \*I:Intermediate sensitive

Table 3: Results of pH resistance.

Number of	рН 1			рН 3		
isolates	0. h	3.h	%alive number	0. h	3. h	%alive number
F2	1.3x 10 <sup>9</sup>	0	0	$1.3x10^9$	$1.0x10^9$	82
F4	$1.0x\ 10^7$	$4.0x10^5$	4.1	$1.0 \times 10^{7}$	$5.0 \times 10^6$	50
F6	$2.5 \times 10^9$	0	0	$2.5 \times 10^9$	$7.4 \times 10^8$	29
F7	$1.0 \times 10^8$	0	0	$1.0x10^{8}$	$1.4x10^{7}$	15
F8	$1.3x10^9$	0	0	$1.3x10^9$	$5.3x10^8$	40
F9	$8.1 \times 10^7$	0	0	$8.1 \times 10^7$	$6.1 \times 10^7$	74
F10	$2.4x10^8$	0	0	$2.4 \times 10^8$	$1.5 \times 10^8$	61
F17	$2.1 \times 10^9$	0	0	$2.1 \times 10^9$	0	0
F30	$2.7x\ 10^7$	0	0	$3.0x10^{7}$	$5.2x10^5$	1.2
F33	$3.6 \times 10^8$	0	0	$3.6 \times 10^9$	0	0
F36	$8.1 \times 10^{8}$	$3.4 \times 10^6$	1.7	$9.1x10^{8}$	$6.7x10^8$	29
F37	$9.0x10^{8}$	$4.4 \times 10^6$	1.9	$9.0x10^{8}$	$6.7x10^7$	7.4
F38	$1.8 \times 10^7$	$3.4x10^4$	0.2	$1.8 \times 10^{7}$	$8.2 \times 10^{5}$	4.4
F39	$2.1x10^{8}$	$1.2 \times 10^6$	0.5	$2.1x10^{8}$	$1.8 \text{x} 10^7$	7.3
F40	$2.5 \times 10^9$	$3.6 \times 10^7$	1.4	$2.5 \times 10^9$	$5.8 \times 10^8$	22
F48	$8.4 \times 10^{8}$	$2.6 \times 10^8$	28	$8.4 \times 10^{8}$	$8.4 \times 10^{8}$	48
F53	$2.6 \times 10^9$	0	0	$2.6 \times 10^9$	$1.5 \times 10^9$	51
F54	$3.2x10^9$	0	0	$3.2x10^9$	$2.1x10^9$	46
F66	$3.9x10^9$	$5.2x10^3$	0	$3.8 \times 10^9$	$8.5 \times 10^{8}$	21
F68	$3.8 \times 10^9$	$3.8 \times 10^9$	0.42	$3.8 \times 10^9$	$3.0x10^9$	39
F69	$3.8x10^9$	$4.0x10^{7}$	0.4	$3.8 \times 10^9$	$2.7x10^9$	65
F70	$1.5 \times 10^8$	$2.8 \times 10^3$	0	$1.5 \times 10^{8}$	$5.8x10^{7}$	37
F71	$3.4x10^9$	$1.1 \times 10^6$	0	$3.4x10^9$	$1.3x10^9$	40
F72	$2.1x10^8$	0	0	$2.1x10^{8}$	0	0
F75	$7.6 \times 10^9$	$1.0 \times 10^5$	0	$7.6 \times 10^9$	$2.0x10^9$	28




Figure 1: T; Reference strain, L. acidaphilus ATCC 4356, F 2-F75: LAB, NK: negative control




Figure 2: Restriction fragment profiles of the 16 S rDNA regions of the strains cut by *Hae*III enzyme. *Lactococcus lactis* group; F2, F4, F6, F7, F8, F9, F10, F30, F33, F37, F38, F39, F40, F53, F54, F66, F68, F69, F75 *Carnobacterium maltoramaticum* group; F36, F70, F71, F72, Uncharacterized bacterium: F17, F48.




Figure 3: Restriction fragment profiles of the 16 S rDNA regions of the strains cut by *MspI* enzyme. *Lactococcus lactis* group; F2, F4, F6, F7, F8, F9, F10, F30, F33, F37, F38, F39, F40, F53, F54, F66, F68, F69, F75 *Carnobacterium maltoramaticum* group; F36, F70, F71, F72, Uncharacterized bacterium: F17, F48.




Figure 4: Restriction fragment profiles of the 16 S rDNA regions of the strains cut by *Hinf*I enzyme. *Lactococcus lactis* group; F2, F4, F6, F7, F8, F9, F10, F30, F33, F37, F38, F39, F40, F53, F54, F66, F68, F69, F75 *Carnobacterium maltoramaticum* group; F36, F70, F71, F72, Uncharacterized bacterium: F17, F48.

### Discussion

The results of PCR performed using primers specific to the Lactobacillus confirmed the phenotypic test results. Three isolates identified as Lactocococcus through phenotypic identified methods were Lactobacillus with these primers. However, the sequence results of the selected samples obtained through ARDRA did not support the results of phenotypic tests and the PCR. This result indicated that the primers were not specific to the genus Lactobacillus. Indeed, determined in the NCBI showed that 16S rDNA regions of the genera Lactobacillus and Lactococcus were similar at 84%, and that the selected primers could amplify the same size region in the genus Lactococcus. Antibiotic resistance and sensitivity in lactic acid bacteria vary depending on the strains and the source of isolation (Salminen et al., 1998). In some studies conducted so far (Coppola et al., 2005; Kim and Austin, 2008; Chemlal-Kherraz et al., 2012), it was reported that most of the antibiotic-resistant probiotics might be helpful when fish are administered antibiotics and that beneficial microorganisms could stay in the gut of the fish for a long time; therefore, their resistivity could be an advantage. Since LAB which have the potential for being used as probiotics should generally be sensitive antibiotics (Karapetkov et al., 2011), their survival rate at pH 1 and pH 3 should be high (Chemlal-Kherraz et al., 2012), and they should display a good antagonistic activity over pathogenic bacteria (Balcazar et al., 2006; Jini et al., 2011; Zapata and Lara-Flores, 2013; Chemlal-Kherraz et al., 2012),

Bacterial isolates were obtained from healthy fish and experimentally inoculated rainbow trout didn't show any signs of disease. L. lactis species (F2, F4, F9 and F10) identified in this study were thought to have a potential probiotic capacity. In further studies, with experiments these potential probiotic strains will be continued. The species with probiotic features were determined to be sensitive to antibiotics used in the treatment of fish diseases in the present study because Lake Egirdir is a natural freshwater lake. There is a need for low-cost, broad-spectrum potential probiotics which have no adverse effects on humans and the environment, which are effective against various fish diseases and which can be used all kinds environmental conditions aquaculture. The study is expected to provide information on the reduction of of antibiotics the use and chemotherapeutic agents and raise the awareness of the manufacturers. It is also expected to help manufacturers to better understand eco-friendly, production sustainable. organic systems. The potential of LABs to be used as probiotics will be investigated in the further studies.

## Acknowledgements

This study was supported by the Ege University Scientific Research Project coded as 2013 FEN 050. The author would like to thank the Fisheries Research Station Directorate, Isparta/TURKEY.

#### References

Abdulamir, A.S., Yoke, T.S., Nordin, N., Bakar, F.A., 2010. Detection and quantification of probiotic bacteria using optimized DNA extraction, traditional and real-time PCR methods in complex microbial communities. *African Journal of Biotechnology*, 9, 1481-1492.

Allameh, S.K., 2012. Isolation, identification and characterization of *Leuconostoc mesenteroides* a new probiotic from intestine of sneakhead fish (*Channa striatus*). *African Journal of Biotechnology*, 11, 3810-3816. DOI: 10. 5897/AJB11.1871.

Austin, B. and Austin, D.A., 2012.

Bacterial fish pathogens: Disease of farmed and wild fish. Springer Dordrecht Heidelberg, New York, London. 651 P.

Balcazar, J.L., De Blas, I., Ruiz-Zarzuela, I., Vendrell, D., Dolores-Evora, M., Muzquiz, J.L., 2006.
Growth inhibition of *Aeromonas* species by lactic acid bacteria isolated from salmonids. *Microbial Ecology in Health Disease*, 18, 61-63.

DOI:10.1080/08910600600761331.

Balcazar, J.L., de Blas, I., Ruiz-Zarzuela, I., Vendrell, D., Girones, O. and Muzquiz, J.L., 2007a. Sequencing of variable regions of 16S r **RNA** gene identification of lactic acid bacteria isolated from the intestinal microbiota of healthy salmonids. **Comparative** Immunology, Microbiology and Infectious 111-118. Diseases, 30. DOI:10.1016/j.cimid.2006.12.001.

Balcazar, J.L., Vendrell, D., de Blas, I., Ruiz-Zarzuela, I., Girones, O. and Muzquiz J.L., 2007b. In vitro competitive adhesion and production of antagonistic compounds by lactic acid bacteria against fish pathogens. *Veterinary Microbiology*, 122, 373-380.

DOI:10.1016/j.vetmic.2007.01.023

- Bondad-Reantaso, M.G., Subasinghe, R.P., Arthur, J.R., Ogawa, K., Chinabut, S., Adlard, R, Tan. Z. and Shariff, M., 2005. Disease and health management in Asian aquaculture. *Veterinary Parasitology*, 132, 249-272.
- L., Brunvold, Sandaa, **R.A.**, Mikkelsen, H., Welde, E., Bleie, Qivind, В., 2007. Н., Characterization of bacterial communities associated with early stages of intensively reared cod (Godus morhua) using denaturing gradient gel electrophoresis (DGGE). Aquaculture, 272, 319-327. Doi: 10.1016/j.aquaculture.2007.08.053.
- Buntin, N., Canthachum, S., Hongpattarakere, T., 2008. Screening of lactic acid bacteria from gastrointestinal tracts of marine fish for their potential use as probiotics. Songklanakarin Journal of Science and Technology, 30, 141-148.
- Bushell, C. and Burns, M., 2012. Feasibility study into the use of DNA sequencing for the identification of probiotic bacteria. *Journal of Association of Public Analysts*, 40, 28-38.

- Chemlal-Kherraz, D., 2012. The probiotic potential of lactobacilli isolated from Nile tilapia (*Oerochromis niloticus*)'s intestine. *African Journal of Biotechnology*, 11, 13220-13227. DOI: 10.5897/AJB12.748
- Clinical and Laboratory Standards
  Institute., 2008. Performance
  standarts for antimicrobial
  susceptibility testing; 18<sup>th</sup>
  Infromational Supplement CLSI
  Document. M100-S18.Wayne (PA).
- Coppola, R., Succi, Mariantonietta., Tremonte, P., Reale, A., Salzano, G. and Sorrentino, E., 2005. Antibiotic susceptibility of *L. rhamnosus* strains isolated from Parmigiano Reggiano cheese. Lait, 85(3), 193-204. DOI: 10.1051/lait:2005007.
- Corcionivoschi, N., 2010. Probioticsidentification and ways of action.

  Innovative Romanian Food
  Biotechnology, 6, 1-11.Deney,
  S., Staykov,Y., Moutafchieva, R.
  and Beev, G., 2009. Microbial
  ecology of the gastrointestinal tract
  of fish and the potential application
  of probiotics and prebiotics in finfish
  aquaculture. International Aquatic
  Research, 1, 1-29.
- Hammes, W.P. and Hertel, C., 2009.

  Lactobacillus Beijerinck 1901,
  212<sup>AL</sup>. In: W. B. Whitman, A.C,
  Parte (Eds.), Bergey's Manual of
  Systematic Bacteriology, 2nd
  edition, Springer Dordrecht
  Heidelberg London, New York, pp.
  711-722.
- Jini, R., Swapna, H.C., Kumar-Rai, A., Vrinda, R., Halami, P.M.,

- Sachindra, N.M., Bhaskar, N.,2011. **Isolation** and characterization of potential lactic acid bacteria (LAB) from freshwater fish processing wastes for application in fermentative utilisation of fish processing waste. Brazilian Journal of Microbiology, 42. 1516–1525. DOI: 10.1590/S1517-838220110004000039.
- Karapetkov, N., Georgieva, R.,
  Rumyan, N., Karaivanova, E.,
  2011. Antibiotic susceptibility of different lactic acid bacteria strains.
  Benefical Microbes. 2, 335-339.
  DOI: 10.3920/BM2011.0016.
- Kesarcodi-Watson, A., Kaspar, H., Lategan, M.L.J. and Gibson, L., 2008. Probiotics in aquaculture: The need, principles and mechanism of action and screening processes, *Aquaculture*, 274, 1-14.
- Kim, D.H. and Austin, B. 2008.

  Characterization of probiotic carnobacteria isolated from rainbow trout (*Oncorhynchus mukiss*) intestine. *Letters in Applied Microbiology*, 47, 141-147. DOI: 10.1111/j.1472-765X.2008.02401.x.
- **Lara-Flores, M., 2011.** The Use of Probiotic in Aquaculture: an overview. *International Research Journal of Microbiology*, 2, 471-478.
- Liu, D., Coloe, S., Baird, R., Pedersen, J., 2000. Rapid minipreparation of fungal DNA for PCR. *Journal of Clinical Microbiology*, 38, 471.
- Liu, Y., Zhou, Z., Yao, B., Shi, P., He, S., Benjaminsen Holvold, L., Ringo, E., 2008. Effect of

- intraperitoneal injection of immunostimilatory substances on allochtonus gut microbiota of Atlantic solmon (Salmo salar L.) determined using denaturing gel electrophoresis gradient (DGGE). Aquaculture Research, 39, 635-646. DOI: 10.1111/j.1365-2109.2008.01934.x.
- Haines, A.N., Gauthier, **D.T.**, Nebergall, E.E., Cole, S.D., Nguyen, K.M., Rhodes. M.W., Vogelbein, W.K., 2013. First report of Streptococcus parauberis in wild finfish from North America, *Veterinary Microbiology*, 166(1-2), 270-275.,

DOI:10.1016/j.vetmic.2013.05.002

- Öztürk, R.Ç. and Altınok, I., 2014. Bacterial and viral fish diseases in Turkey, *Turkish Journal of Fisheries and Aquatic Sciences*, 14, 275-297 DOI: 10.4194/1303-2712-v14\_1\_30.
- Perez-Sanchez, T., Balcázar, J.L., García, Y., Halaihel, N., Vendrell, D., de Blas, I., Merrifield, D.L. and Ruiz-Zarzuela. I. 2011. Identification and characterization of lactic acid bacteria isolated from rainbow trout, Oncorhynchus mykiss (Walbaum), with inhibitory activity Lactococcus against garviae. Journal of Fish Diseases, 34, 499-507. Doi: 10.1111/j.1365-2761.2011.01260.x.
- **Ringo, E. and Gatesoupe, F.J., 1998.**Lactic acid bacteria in fish: a review. *Aquaculture*, 160, 177-203. DOI: 10.1016/S0044-8486(97)00299-8
- **Ringo, E., 2004.** Lactic acid bacteria in fish and fish farming. In. : S. Salminen, A. Von Weight, A.

- Ouwehand (Eds.), Lactic Acid Bacteria:microbiology and functional aspects. Marcel Dekker Inc., New York, USA. pp. 581-610.
- Salminen, S., Von Wright, A., Morelli, L., Marteau, P., Brassart, D., de Vos W.M., Fonden, R., Saxelin, M., Collins, K., Mogensen, G., Birkeland, S.E., Mattila-Sandholm, T., 1998. Demonstration of safety of probiotics. *International Journal of Food Microbiology*, 44, 93-106.
- Soto, L.P., Frizzo, L.S., Bertozzi, E., Avataneo, E., Sequeira, G.J., Rosmini, M.R., 2010. Molecular microbial analysis of *Lactobacillus strains* isolated from the gut of calves for potential probiotic use.

- *Veterinary Medicine International*, 2010, 1-7. DOI:10.4061/2010/274987.
- **Subasinghe, R., Soto, D. and Jia, J., 2009.** Global aquaculture and its role in sustainable development. *Reviews in Aquaculture*, 1, 2-9.
- Teuber, M., 2009. Genus II. schleifer, Lactococcus Kraus, Dvorak, Klipper-Baelz, Collins and Fischer 1986, 354. In: W. В. Whitman, A.C. Parte (Eds.), Bergey's Manual of Systematic Bacteriology, 2nd edition, Springer Dordrecht Heidelberg London, New York. pp.711-722.