# Acclimation responses of gill ionocytes of red tilapia (Oreochromis mossambicus × O. niloticus) to water salinity and alkalinity

Jeerawat Thammaratsuntorn J.; Jin-Liang Zhao J.L.\*; Zhao L.H.; Zhuang Q.Q.; Guo J.T.; Larbi Ayisi Ch.

Received: May 2014 Accepted: September 2014

#### **Abstract**

To understand the acclimation strategies of red tilapia to different environments, this study aimed to evaluate different responses of red tilapia (O. mossambicus  $\times$  O. niloticus) to salinity (10-30%), alkalinity (1-3 gL<sup>-1</sup> NaHCO<sub>3</sub>) and salinity and alkalinity (10/1-30/3 %/gL<sup>-1</sup>NaHCO<sub>3</sub>) environments. Localization, type, size, and numeration of gill ionocytes were investigated on the same specimens by scanning electron immunohistochemistry microscopy (SEM) and (IHC) with antibodies Na<sup>+</sup>/K<sup>+</sup>-ATPase Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>-</sup>contransporter (NKCC). (NKA). cvstic transmembrane conductance regulator (CFTR) and carbonic anhydrase (CA). Ionocytes were only located on filaments conducted by SEM. Four types of ionocytes namely pit, convex, concave and transitory types were determined morphologically by their apical openings of which concave and transitory type were not present in freshwater (FW) and saltwater (SW) fish (10). Both ionocytes size and number increased with elevated stress levels. In comparison to FW, density of ionotypes increased to about 4.75, 3.00 and 3.44 fold in SW (30), AW (3) and S&AW (30/3) respectively. Immunoreactive cells on gill filaments confirmed branchial distribution of ionocytes. Immunoreaction of NKA, NKCC and CA appeared in FW except for CFTR while they all appeared in SW (30), AW (3) and S&AW (30/3).

**Keywords**: Acclimation response, Ionocytes, Salinity, Alkalinity, Red tilapia.

Key Laboratory of Freshwater Fisheries Germplasm Resources, Ministry of Agriculture, Shanghai Ocean University, Shanghai 201306, China

<sup>\*</sup>Corresponding author's email: jlzhao@shou.edu.cn

### Introduction

Osmoregulation is one of the most vital functions to all euryhaline fish (Evans, which 2008), provide hyperosmo-regulatory and hypoosmo-regulatory abilities to keep osmolality of body fluids within a narrow physiological range in freshwater (FW) and seawater (SW) fish, respectively (Kang et al., 2013). Osmoregulation is mainly achieved by the gills, kidney and intestine. Among these osmoregulatory organs, the gills are the unique to aquatic animals (Evans et al., 2005; Gilmour and Perry, 2009).

The transporting epithelia contains a specialized subset of cells called ionocytes (formally called mitochondria-rich cells or chloride cells; Keys and Willmer, 1932), the major cells in fish gills that transport ions actively (Perry et al., 2003; Evans et al., 2005; Hwang, 2009), these ionocytes could secrete and absorb ions in SW and FW environments, respectively (Hirose et al., 2003; Hwang and Lee, 2007), in addition to carrying out acid-base regulation and other functions (Perry and Gilmour, 2006; Gilmour and Perry, 2009).

basolateral The membranes of ionocytes form extensive tubular systems (Karnaky et al., 1976; Sardet et al., 1978), the primary site of ion transportation between ionocytes and the external environment is the apical membrane (Christensen et al., 2012). On the apical and basolateral membranes, ionocytes possess a specific complement of transporters or channels that allow for directional movement of ions (Dymowska et al., 2012). The apical

surface morphology of the gill ionocytes are usually found to vary across salinity environments (Uliano et al., 2010). The biochemical cellular, and molecular mechanism of fish ionocytes acclimation to salinity have been well developed (Hwang et al., 2011). Meanwhile. in some inland waters. alkalinity is another form environmental factors that fish always encounter, which also has adverse effects on fish survival and growth (Fielder et al., However, the physiological mechanism of fish in adaption to water alkalinity or salinity & alkalinity is less understood.

Tilapia is a group of euryhaline fish and has been often served as model for ionoregualtion studies. Work on tilapia has been mostly performed Mozambique tilapia O. mossambicus, and different types of ionocytes gills have been revealed in tilapia based on their apical openings (Lee et al., 1996; Chang et al., 2001; Inokuchi et al., 2009, Choi et al., 2011). Also, there is an increase in size and number of ionocytes in response to increasing salinity (Ayson et al., 1994; Kültz et al., 1995; Van Der Heijden et al., 1997; Hiroi et al., 2005). Due to their specific abilities to grow and reproduce in hypersaline environments, black-chin tilapia Sarotherodon melanothern was used as a model for hypersaline adaption studies (Lorin-Nebel, 2012). The small cichlid fish, O. alcalicus graham are found in hot, alkaline lagoons surrounding Lake Magadi in Kenya (Walsh et al., 2001). An early work on morphology of ultra-structure of the chloride cells and their modification on extreme alkalinity revealed that the ionocytes decreased in number and size in lower alkalinity environments (Maina, 1990).

Red tilapia was a cross hybrid of a reddish-orange 0. mutant female mossambicus and a normal male O. niloticus (Galman, 1983). It has been developed as a popular commercial cultured species in Southeast Asia, as well as the Middle East, the Caribbean, Central and South America (Watanabe et al., 2002). Since their fast growth and high tolerance, they are of great potential to be developed and cultured in broad environments. The purpose of this study to investigate was therefore the localization, morphological and numeration change of gill ionocytes of Red tilapia in response to different environmental treatments, including alkalinity, salinity salinity, and & alkalinity scanning electron by microscopy (SEM). In the meantime, immunohistochemistry (IHC) of some important ion transporters were also carried out to verify the external observation. Taken together, these data would contribute to better understanding of osmoregulation strategies of Red tilapia gill ionocytes to these different environments.

#### Materials and methods

Experimental fish

Red tilapia (O.  $mossambicus \times O$ . niloticus) fish, averaged  $5.5\pm0.3$  cm in total length, were obtained from Fish Germplasm Station, Shanghai Ocean University, taken to laboratory and maintained in recirculated aquariums

freshwater weeks with for two adjustment. Salt waters (SW) were prepared using local fresh water with sodium chloride (NaCl). Alkaline waters (AW) were prepared using local fresh with sodium water bicarbonate (NaHCO<sub>3</sub>). Both NaCl and NaHCO<sub>3</sub> were used in preparing salt & alkaline waters (S&AW). Three treatment levels were also designed for each kind of environment: (1) SW: 10‰, 20‰, and 30 % salinity, (2) AW: 1 gL<sup>-1</sup>, 2 gL<sup>-1</sup>, and 3 gL<sup>-1</sup> NaHCO<sub>3</sub>, (3): S&AW: 10/1, 20/2, and 30/3 ‰/gL<sup>-1</sup> NaHCO<sub>3</sub>, using Mettler (SG7-ELK, USA) Toledo concentration of salinity and Hanna instruments (HI755, USA) concentration of alkalinity.

All fish were gradually acclimated from freshwater to 5, 10, 15, 20, 25, 30 (‰) salt waters, 1, 2, 3 (gL<sup>-1</sup> NaHCO<sub>3</sub>) alkaline waters, and 5/1, 10/1, 15/2, 20/2, 25/3, 30/3 (% /gL<sup>-1</sup> NaHCO<sub>3</sub>) salt & alkaline waters, and stayed at each treatment for 3 days before next transfer. During the experiment, fish were fed two times per day and water temperature was maintained at 25-27 °C, half of the water was changed every three days to preserve optimal water quality. After having acclimated to different treatments for one week, 10 fish were randomly sampled in each treatment level, the outmost pair of gills were removed for scanning electron microscopy analysis. In the meantime, 5 individuals at FW, SW (30), AW (3), and S&AW (30/3)treatments were also sampled for immunohistochemistry analysis.

Scanning electron microscopy (SEM)

The gills were fixed in a fixative, which consisted of 5% (v/v) glutaraldehyde and 4% (w/v) paraformaldehyde (PFA) in 0.1 M phosphate buffer (PB, pH 7.2), at 4°C for 12 hours. Fixed gills were rinsed with 0.1 M PB for 15 min three times, and then dehydrated in a series of ascending concentrations of ethanol from 30%. The samples were dried at room temperature for 1 hour, and then mounted on specimen stubs, and coated with platinum-palladium in an ion sputter (Hitachi E-1010, Tokyo, Japan). examined Ionocytes were and photographed with a Hitachi S-3400N Scanning Electron Microscopy (Tokyo, Japan). For each individual, ionocytes were counted and measured within 10 continuous filaments by image analysis software (NIS-Elements F program).

## Antibody

Primary antibodies include: (1) a mouse monoclonal antibody (α5; Developmental Studies Hybridoma Bank), raised against  $\alpha$ -subunit the of the avian Na<sup>+</sup>/K<sup>+</sup>-ATPase (NKA); (2) a mouse (T4; monoclonal antiserum Developmental Studies Hybridoma Bank) raised against the C-terminus of human Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>-</sup> contransporter (NKCC); (3) a mouse monoclonal antibody (R&D Systems, Boston, MA, USA) directed against 104 amino acids C-terminus of the human cystic fibrosis transmembrane conductance regulator (CFTR); (4) a rabbit polyclonal antibody against a synthetic peptide corresponding to the N-terminal end of human Carbonic anhydrase 1 (CA; Boster company,

China). Samples were stained with Cell and Tissue Staining Kit (Mouse kit; HRP-DAB system, R&D Systems®, Boston, MA, USA).

# Immunohistochemistry (IHC)

The gills were fixed in 4% paraformaldehyde for 12-16 hours, and washed 0.01 in M phosphate-buffered saline (PBS) twice for 40 min each. The samples were dehydrated through a graded ethanol series, infiltrated with xylene, and embedded in paraffin. Cross sections of the gills were provided at 8 µm thickness and mounted on Poly-D-lysine and 3-Aminopropyl-Triethoxysilane (APES) coated glass slides. The deparaffinized sections were incubated with xylene and rehydrated sections were incubated with a graded ethanol series and dH<sub>2</sub>O, respectively. The antigen retrieval (AR) technique, which is predominantly based on high-temperature heating of tissues, is used as a non-enzymatic pretreatment for immunohistochemical staining formalin-fixed, paraffin-embedded tissue sections for 5 min at 100°C and then the sides were cooled at room temperature for 30 min and washed in PBS for 5 min twice. They were immunohistochemically stained with primary antibody, a monoclonal antibody ( $\alpha$ 5) to Na<sup>+</sup>/K<sup>+</sup>-ATPase  $\alpha$ -subunit and Cell & Tissue Staining Kit (Mouse kit; HRP-DAB system, R&D Systems®, Boston, MA, USA).

## Statistical analysis

Data were presented as means  $\pm$  standard error of the mean. The significance of

differences at *p*<0.05 was examined using one-way analysis of variance (ANOVA), followed by Tukey–Kramer *post-hoc* test.

#### Results

Scanning electron microscopy

Red tilapia has four gill arches at each side. Each gill arch supports one set of paired gill filaments. Each gill filament in turn supports numerous secondary lamellae, which extend out from both sides of the filament body (Fig 1a.). Under observation in scanning electron microscopy, each ionocytes were characterized with an apparent apical opening, and they were only located in the gill filaments of FW, SW, AW and S&AW acclimated fishes (Fig 1b.)

Four types of ionocytes with different apical openings were observed during the acclimation experiments: pit, concave, convex and a transitory apical surface (Fig 2.), these apical structures also showed some variation within the same types. The apical opening of pit type was ellipse-shaped with a narrow and deep hole; the internal structure could not be observed (Fig 2a.). The concave type was somewhat ellipse, its surfaces was slightly dented or flat with a mesh-like structure (Fig 2b.). The convex type was a rough surface, curved or bowed outward like the outside of a bowl or sphere or circle (Fig 2c.). The transitory apical surface was somewhat similar to both concave surface and enlarged pit and its opening was more deeply dented and appeared as a large crater (Fig 2d.). The convex and transitory types were not observed in FW and SW (10) treatments, however, these four types were presented in SW (20), SW (30) treatments and all AW and S&AW treatments (Fig 3.).

The mean size (area) of different ionocytes in acclimation to SW, AW and S&AW are shown in Table 1 and Fig 3. The size of pit type averaged  $1.14\pm1 \mu m^2$ in FW and increased about 11.70 fold in SW, whereas it was relatively constant in AW and S&AW. The mean size of concave type increased significantly about 5.74, 8.39, and 6.11 fold in SW, AW and S&AW, respectively, with an increasing trend. The convex transitory types were absent in FW and SW (10), convex type increased about 1.84 fold from SW (20) to SW(30), 3.29 fold from AW (1) to AW (3), and 3.62 fold from S&AW (10/1) to S&AW (30/3), the transitory type increased about 2.85 fold from SW (20) to (30), 3.94 fold from AW (1) to AW (3), and 4.53 fold from S&AW (10/1) to S&AW (30/3). In comparison to FW, the mean size of ionotypes increased to about 13.66, 14.19, and 14.07 fold in SW (30), AW (3) and S&AW (30/3), respectively.

The mean densities (number) of different ionocytes in acclimation to SW, AW and S&AW treatments are shown in Table 2 and Fig 4. In FW fish, the pit type was dominant with mean density of  $5.54\times10^3$  cells/mm<sup>2</sup>, and the concave type accounted for  $0.7\times10^3$  cell/mm<sup>2</sup>. In acclimation to SW, the mean density of ionocytes increased significantly and became about 4.75-fold in SW (30) (pit type 37%, concave type 23%, convex type 19%, and transitory type 21%).




Figure 1: Structure of the gill of red tilapia (A) and location of ionocytes (B) identified by scanning electron microscopy. Gill arch (GR), gill filaments (GF), lamella (LM), ionocytes were showed by arrow.




Figure 2: Four types of ionocytes in gill filaments of red tilapia identified by scanning electron microscopy (SEM). (A) pit, (B) concave, (C) convex, (D) transitory.




Figure 3: Change of the mean size of four types of gill ionocytes of red tilapia in acclimation to freshwater (FW), salinity water (SW), alkalinity water (AW), salinity & alkalinity water (S&AW).




Figure 4: Change of the mean density of four types of gill ionocytes of red tilapia in acclimation to freshwater (FW), salinity water (SW), alkalinity water (AW), salinity & alkalinity water (S&AW).

Table 1: The mean size of four types of gill ionocytes of red tilapia in different environmental treatments  $(\mu m^2)$ .

Туре	FW	SW(‰)			AW(gL <sup>-1</sup> )			S&AW(%₀/gL <sup>-1</sup> )		
	0	10	20	30	1	2	3	10/1	20/2	30/3
pit	1.14±1	6.34±0.3	11.88±1.2	13.34±0.3	1.6±1.3	1.72±0.4	2.07±0.33	1.64±0.44	1.97±1	2.06±0.4
concave	1.07±0.8	2.28±2	3.51±1	6.15±0.55	2.55±0.78	5.67±1	8.98±0.3	2.3±0.5	5.02±0.3	6.54±0.5
convex			2.75±2.3	5.08±0.75	2.87±2.1	6.3±2	9.46±0.5	2.99±1	7.58±0.8	10.85±1
transitory			1.97±0.8	5.62±23	2.75±0.3	7.543±0.8	10.85±1	2.57±1	8.59±1	11.65±2

Table 2: The mean density of four types of ionocytes of Red tilapia in o different environmental treatments (10<sup>3</sup> cell /mm<sup>2</sup>)

Type	FW	SW(‰)			AW(gL <sup>-1</sup> )			S&AW(%₀/gL <sup>-1</sup> )		
	0	10	20	30	1	2	3	10/1	20/2	30/3
pit	5.54±0.23	6.59±0.7	8.83±0.12	9.8±0.04	5.34±0.15	5.75±0.77	5.85±0.76	5.34±0.09	5.44±0.93	5.45±0.04
concave	0.07±0.02	2.28±0.39	3.51±0.44	6.15±0.08	0.89±0.59	2.93±0.89	4.78±0.33	2.3±0.94	5.02±0.5	6.54±0.35
convex			2.75±0.05	5.08±0.3	0.54±09	2.74±0.45	3.9±0.44	1.04±0.16	4.7±0.2	4.04±0.78
transitory			1.97±0.32	5.62±0.49	0.18±28	1.8±0.22	2.3±0.02	0.76±0.65	2.99±0.66	3.3±0.03

The mean density of ionocytes increased about 3.00-fold in AW (3) (pit type 35%, concave type 28%, convex type 23%, and transitory type 14%), and about 3.44-fold in S&AW (30/3) (pit type 28%, concave type 34%, convex type 21%, and transitory type 17%). In comparison to FW, the densities of ionotypes increased to about 4.75, 3.00, and 3.44 fold in SW (30), AW (3) and S&AW (30/3), respectively.

# Immunohistochemistry (IHC)

Whole-mount immunohistochemistry with anti-NKA, anti-NKCC, anti-CFTR, and anti-CA of gill filaments were investigated in FW, SW (30), AW (3), and S&AW (30/3) treatments. In comparison with the control (without the first antibody), immunoreactions of NKA, NKCC, CFTR, and CA were found only

at the inter-lamellar region of the gill filaments (Fig 5.). In FW treatment, immunoreactions of NKA, NKCC and CA appeared, while CFTR disappeared. In SW (30), AW (3) and S&AW (30/3) treatments, immunoreactions of NKA, NKCC, CA and CFTR were spotted.

#### Discussion

Distribution of ionocytes in FW, SW, AW and S&AW

The present study explored the chronic responses of gill ionocytes of Red tilapia in acclimation to salinity, alkalinity and salinity & alkalinity in waters, the ionocytes were only located on the filaments in FW, SW, AW and S&AW adapted fishes under SEM observation, which was also confirmed by immuoreaction detection in IHC.




Figure 5: Whole-mount immunohistochemistry with anti-Na<sup>+</sup>/K<sup>+</sup>-ATPase (NKA), anti-Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>-</sup> cotransporter (NKCC), anti- cystic fibrosis transmembrane conductance regulator (CFTR), and anti- carbonic anhydrase (CA) of gill filaments of Red tilapia in freshwater (FW), salt water (SW 30), alkaline water (AW 3) and salinity & alkalinity in water (S&AW 30/3) Scale bar, 10μm.

This observation was similar to that described for Mozambique tilapia (O. mossambicus), where ionocytes in SW had always been located in the filaments (Kültz et al., 1995; Lin et al., 2004a; Sardella et al., 2004; Inokuchi et al., 2008; Choi et al., 2011). In contrast, the ionocytes of the black-chin tilapia *melanotheron*) (Sarotherodon were localized on the filaments and expanded extensively to the lamellae acclimatizing to seawater or hyper saline waters (Ouattaraa et al., 2009).

Different patterns of ionocytes in the gills have been revealed in fish: (1) ionocytes were found on filaments and

lamellae in freshwater and seawater (Zydlewski and McCormick, 2001; Lin et al., 2006; Christensen et al., 2012), (2) ionocytes were distributed on both filaments and lamellae in freshwater, where lamellae ionocytes disappeared following seawater transfer (Varsamos et al., 2002; Nebel et al., 2005; Hiroi and McCormick, 2007), (3) ionocytes were found only in the filament and rarely in the lamellae in both freshwater and seawater (Uchida et al., 2000; Lin et al., 2004b; Yang *et al.*, 2009). The function of gill lamellae is to improve the gill function. Thin plates of tissue on gill filaments that contain the capillary across

which gases are exchanged, increase the total area of the gills (Ito *et al.*, 2005; Jeffery *et al.*, 2007). The different distributional patterns of ionocytes might reflect a more suitable mechanism for different euryhaline teleosts to cope with environmental fluctuations, or an adapted mechanism acquired during the teleosts evolution.

Response of types, size and densities of ionocytes to SW, AW and S&AW

Different types of ionotypes have been described in tilapia fish encountering various environments, and involved in different ionic transportation processes. Pisam et al. (1995) firstly reported that there were two types of MR cell ( $\alpha$  and  $\beta$ the cells) in gill epithelium freshwater-adapted tilapia, the apical surface of  $\alpha$  cells were slightly depressed to form shallow apical cavities while the apical surface of  $\beta$  cells were flattened and hardly depressed. During seawater acclimation,  $\alpha$  cells were transformed into seawater-type MR cells, whereas  $\beta$ cells degenerated and disappeared. Later, three types of MR cells, wavy convex, shallow basin and deep holes, were identified scanning electron by micrographs, wavy-convex and shallow-basined MR cells were evident in FW adapted fish, whereas the deep hole type was dominant in SW (5) -adapted fish (Lee et al., 2000), and SW (30)-adapted fish (Lee et al., 1996). Recently, Choi et al., (2011) proposed that the ionocytes could be classified into four types: an apical pit, a convex apical surface, a concave apical surface and a transitory apical surface. fish acclimated to freshwater, three types of apical opening were observed, the pit, the convex and the concave. Following transfer to 70% seawater, the convex was absent completely; and the transitory appeared (Choi et al., 2011). Based on the newly classification method, our SEM observations also revealed there were four types of ionocytes with different apical openings in red tilapia in acclimation to SW, AW, and S&AW, pit, a convex, concave, and transitory type. Only the pit type and concave type were observed in FW and low SW (10) treatments. Since SW (10) is about the iso-osmotic for many tilapias, these two type ionocytes might be enough to maintain homeostasis at hypo- and iso-osmotic environments, where there was no or little load for ionic excretion. In response to higher salinities (20 and 30), alkalinities, salinity and alkalinities treatments, all four kinds of ionocytes emerged. Therefore, the convex type and transitory type were of great necessity to Red tilapia in adaption to (higher) salinity. alkalinity and salinity-alkalinity in waters to cope effectively with higher load of ionic excretion.

The size change different of ionocytes types in acclimation to SW. AW and S&AW showed different responses. In SW, the mean size of pit type increased greatly than that of the concave, convex, and transitory type with increasing salinity. On the contrary, the mean size of the concave, convex, and transitory type increased highly than that of the pit type in AW and S&AW, respectively. Moreover, the size of convex and transitory types increased greatly than that of concave type in S&AW (Fig 3). In acclimation to different environments, enlarged ionocytes could be a fast and effective way to improve their transportation capacities and responsible for more ionic transportation.

The ionocytes number of red tilapia also changed highly in acclimation to SW, AW, and S&AW, the density of ionocytes in SW (30), AW (3), and S&AW (30/3) treatments was about 4.75, 3.00, and 3.44 fold in comparison to that in FW, respectively, which was consistent with some previous work, supporting their significance of ionocytes number in hyperosmotic conditions in the cichlid (Mozambique tilapia mossambicus, Pratapa and Wendelaar, 1993; Hiroi et al., 2005; Inokuchi et al., 2008; Choi et al., 2011; Sarotherodon melanotheron, Ouattara et al., 2009). The elevated number of ionocytes was probably a higher and long-termed response, which is a result of the proliferation and differentiation of newly recruited ionocytes. Increase of ionocytes number also improved transportation capacities. For comparison of different environments, the density in AW (3) was smaller than that in SW (30), which probably due to less stress strength. An interesting finding was the density in salinity and alkalinity treatment (30/3) was not, but less than the sum of the single salinity (30) and single alkalinity (3). There might be some interaction among different ions in salt and alkaline water, that decrease the total stress strength, or ionocytes might use the same transports or exchangers simultaneously to regulate both salinity and alkalinity.

The different functions of inocytes

The ionocytes were first identified as possessing chloride secreting function (Keys and Willmer, 1932), currently, ionocytes have been known to be involved in multiple functions (Marshall, 2003; Evans *et al.*, 2005; Wang *et al.*, 2009; Hwang *et al.*, 2011). Thus, they were also functionally classified by immunoreactive of various transport proteins (Hiroi *et al.*, 2005; Ouattara *et al.*, 2009).

IHC revealed that immunoreaction of NKA, NKCC, CFTR, and CA were found only at the inter-lamellar region of the gill filaments of red tilapia in FW, SW, AW, S&AW, except for CFTR which was not detected in FW. NKA was sodium pump and responsible for the active Na<sup>+</sup> transportation, it also provided a driving force for other transporting systems (Tang et al., 2012). NKCC was a membrane transport protein that contransported Na<sup>+</sup>, K<sup>+</sup>, and Cl<sup>-</sup> into or out of cells (Haas, 1994). CFTR functioned as cAMP-activated a ATP-gated anion channel (e.g. Cl<sup>-</sup>) (David et al., 2006). CA catalyzed the rapid dehydration of HCO<sub>3</sub> or hydration of  $CO_2$  ( $HCO_3^- + H^+ \leftrightarrow CO_2 + H_2O$ ) in acid-base regulation (Badger and Price, 1994). In contrast to SW, AW, and S&AW, the absence of CFTR in FW tilapia suggested there was no or little anions transportation in FW for Red tilapia. In different ionic transportation conditions in SW, AW, and S&AW, various transporters and exchangers were located in different kinds of ionocytes needed for effective transportation, as a result, NKA, NKCC, CFTR, and CA all present in SW,

AW. S&AW and without obvious difference. Since the functions different types of ionocytes are still uncertain, we could not clearly explain which specific transporters in each and what ionocytes types specific transportation functions involved osmoregulation. Therefore, more ionic transporters/channels and integrated work should be implemented to better understand the molecular mechanism of fish in acclimation different to environments.

# Acknowledgements

This work was supported by the National Key Technology Support Program, China (2012BAD26B03), China Agriculture Research System (CARS-49-04B) and Shanghai University Knowledge Service Platform (ZF1206).

#### References

- Ayson, F.G., Kaneko, T., Hasegawa, S. and Hirano, T., 1994. Development of mitochondria-rich cells in the yolk-sac membrane of embryos and larvae of tilapia, *Oreochromis mossambicus*, in fresh water and seawater. *Journal of Experimental Zoology*, 270(2), 129-135.
- Badger, M.R. and Price, G.D., 1994. The role of carbonic anhydrase in photosynthesis. *Annual Review of Plant Physiology and Plant Molecular Biology*, 45(1), 369-392.
- Chang, I.C., Lee, T.H., Yang, C.H., Wei,
  Y.W., Chou, F.I. and Hwang, P.P.,
  2001. Morphology and function of gill mitochondria-rich cells in fish acclimated to different environments.

- Physiological and Biochemical Zoology, 74(1), 111-119.
- Choi, J.H., Lee, K.M., Inokuchi, M. and Kaneko, T., 2011. Morphofunctional modifications in mitochondria-rich cells of Mozambique tilapia transferred from freshwater to 70% seawater, detected by dual observations of whole-mount immunocytochemistry and scanning microscopy. Comparative electron Biochemistry and Physiology. Part A, 158(1), 132-142.
- Christensen, A.K., Hiroi, J., Schultz, E.T. and McCormick, S.D., 2012. Branchial ionocyte organization and ion-transport protein expression in juvenile alewives acclimated to freshwater or seawater. *The Journal of Experimental Biology*, 215(4), 642-652
- **David, C.G., Paola, V. and László, C., 2006.** The ABC protein turned chloride channel whose failure causes cystic fibrosis. *Nature*, 440(**7083**), 477-483.
- **Dymowska, A.K., Hwang, P.P. and Goss, G.G., 2012.** Structure and function of ionocytes in the freshwater fish gill. *Respiratory Physiology & Neurobiology*, 184(3), 282-292.
- Evans, D.H., Piermarini, P.M. and Choe, K.P., 2005. The multifunctional fish gill: dominant site of gas exchange, osmoregulation, acid-base regulation, and excretion of nitrogenous waste. *Physiological Reviews*, 85(1), 97-177.
- **Evans D.H, 2008.** Teleost fish osmoregulation: what have we learned since August Krogh, Homer Smith,

- and Ancel Keys. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 295(2), 704-713.
- **Fielder, D.S., Bardsley, W.J. and Allan, G.L., 2001.** Survival and growth of Australian snapper, *Pagrus auratus*, in saline groundwater from inland New South Wales, Australia. *Aquaculture*, 201(1-2), 73-90.
- Galman, O.R. and Avtalion, R.R., 1983.

  A preliminary investigation of the characteristics of red tilapias from the Philippines and Taiwan. Proceeding of the International Symposium on Tilapia in Aquaculture, Nazareth, Israel, May. 1983.
- Gilmour, K.M. and Perry, S.F., 2009. Carbonic anhydrase and acid-base regulation in fish. *The Journal of Experimental Biology*, 212(11), 1647-1661.
- Haas, M., 1994. The Na-K-Cl cotransporters. *The American Journal of Physiology*, 267(4), 869-885.
- Hiroi, J. and McCormick, S.D., 2007.

  Variation in salinity tolerance, gill Na<sup>+</sup>/K<sup>+</sup>-ATPase, Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>-</sup> cotransporter and mitochondria-rich cell distribution in three salmonids *Salvelinus namaycush*, *Salvelinus fontinalis* and *Salmo salar*. The *Journal of Experimental Biology*, 210(6), 1015-1024.
- Hiroi, J., McCormick, S.D.,
  Ohtani-Kaneko, R. and Kaneko, T.,
  2005. Functional classification of mitochondrion-rich ells in euryhaline Mozambique tilapia (*Oreochromis mossambicus*) embryos, by means of

- triple immunofluorescence staining for Na<sup>+</sup>/K<sup>+</sup>-ATPase, Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>-</sup>cotransporter and CFTR anion channel. *The Journal of Experimental Biology*, 208(1), 2023-2036.
- Hirose, S., Kaneko, T., Naito, N. and Takei, Y., 2003. Molecular biology of major components of chloride cells. *Comparative Biochemistry and Physiology. Part B*, 136(4), 593-620.
- Hwang, P.P. and Lee, T.H., 2007. New insights into fish ion regulation and mitochondrion-rich cells. *Comparative Biochemistry and Physiology. Part A*, 148(3), 479-497.
- **Hwang, P.P., 2009.** Ion uptake and acid secretion in zebrafish (*Danio rerio*). *The Journal of Experimental Biology*, 212(11), 1745-1752.
- Hwang, P.P., Lee, T.H. and Lin, L.Y., 2011. Ion regulation in fish gills: recent progresses in the cellular and molecular mechanisms. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 301(1), 28-47.
- Inokuchi, M., Hiroi, J., Watanabe, S., Lee, K.M. and Kaneko, T., 2008. Gene expression and morphological localization of NHE3 NCC and NKCC1a in branchial mitochondria-rich cells of Mozambique tilapia (Oreochromis mossambicus) acclimated to a wide salinities. Comparative range of Biochemistry and Physiology. Part A, 151(**2**), 151-158.
- Inokuchi, M., Hiroi, J., Watanabe, S., Hwang, P.P. and Kaneko, T., 2009.

  Morphological and functional

- classification of ion-absorbing mitochondria-rich cells in the gills of Mozambique tilapia. *The Journal of Experimental Biology*, 212(7), 1003-1010.
- Ito, N., Watanabe, H. and Hashimoto, Y., 2005. Angioarchitecture of the Branchial Arterial System of Carp (Cyprinus carpio L.). Anatomia, Histologia, Embryologia, 34(1), 23.
- Jeffery, K.R., Bateman, K., Bayley, A., Feist, S.W., Hulland, J., Longshaw, C., Stone, D., Woolford, G. and Way, K., 2007. Isolation of a cyprinid herpesvirus 2 from goldfish, Carassius auratus (L.), in the UK. Journal of Fish Diseases, 30(11), 649-656.
- Kang, C.K., Yang, W.K., Lin, S.T., Liu, C.C., Lin, H.M., Chen, H.H., Cheng, C.W., Lee, T.H. and Hwang, P.P., **2013.** The acute and regulatory phases time-course changes of in gill mitochondrion-rich cells of seawater-acclimated medaka (Oryzias dancena) when exposed hypoosmotic environments. Comparative **Biochemistry** and Physiology. Part A, 164(1), 181-191.
- Karnaky, K.J., J<sub>R.</sub>, Kinter, L.B., Kinter, W.B. and Stirling, C.E., 1976.

  Teleost chloride cell. II.

  Autoradiographic localization of gill

  Na, K-ATPase in killifish *Fundulus*heteroclitus adapted to low and high salinity environments. The Journal of Cell Biology, 70(1), 157-177.
- Keys, A. and Willmer, E.N., 1932. "Chloride secreting cells" in the gills of fishes, with special reference to the common eel. *The Journal of*

- Physiology, 76(3), 368-378.
- Kültz, D., Jürss, K. and Jonas, L., 1995. Cellular and epithelial adjustments to altered salinity in the gill and opercular epithelium of a cichlid fish (*Oreochromis mossambicus*). Cell and Tissue Research, 279(1), 65-73.
- Lee, T.H., Hwang, P.P., Lin, H.C. and Huang, F.L., 1996. Mitochondria-rich cells in the branchial epithelium of the teleost, *Oreochromis mossambicus*, acclimated to various hypotonic environment. *Fish Physiology and Biochemistry*, 15(6), 513-523.
- Lee, T.H., Hwang, P.P., Shieh, Y.E. and Hwang, P.P., 2000. The relationship between "deep-hole" mitochondria-rich cells and salinity adaptation in the euryhaline teleost, *Oreochromis mossambicus. Fish Physiology and Biochemistry*, 23(2), 133-140.
- Lin, C.H., Huang, C.L., Yang, C.H., Lee, T.H. and Hwang, P.P., 2004a. Time-course changes in the expression of Na. K-ATPase and the morphometry of mitochondrion-rich cells in gills of euryhaline tilapia (Oreochromis mossambicus) during freshwater acclimation. Journal of *Experimental* Zoology. **Part** Comparative Experimental Biology, 301(1), 85-96.
- Lin, C.H., Tsai, R.S. and Lee, T.H., 2004b. Expression and distribution of Na, K-ATPase in gill and kidney of the spotted green pufferfish, *Tetraodon nigroviridis*, in response to salinity challenge. *Comparative Biochemistry and Physiology. Part A*, 138(3), 287-295.

- Lin, Y.M., Chen, C.N., Yoshinaga, T., Tsai, S.C., Shen, I.D. and Lee, T.H., 2006. Short-term effects of hyposmotic shock on Na<sup>+</sup> /K<sup>+</sup>-ATPase expression in gills of the euryhaline milkfish, *Chanos chanos. Comparative Biochemistry and Physiology. Part A*, 143(3), 406-415.
- Lorin-Nebel, C., Avarre, J.C., Faivre, N., Wallon, S., Charmantier, G. and Durand, J.D., 2012. Osmoregulatory strategies in natural populations of the black-chinned tilapia *Sarotherodon melanotheron* exposed to extreme salinities in West African estuaries. *Comparative Biochemistry and Physiology. Part B*, 182(6), 771-780.
- Maina, J.N., 1990. A study of the morphology of the gills of an extreme alkalinity and hyperosmotic adapted teleost *Oreochromis alcalicus grahami* (Boulenger) with particular emphasis on the ultrastructure of the chloride cells and their modifications with water dilution. A SEM and TEM study. *Anatomy and Embryology*, 181(1), 83-98.
- Marshall, W.S., 2003. Rapid regulation of NaCl secretion by estuarine teleost fish: Coping strategies for short-duration freshwater exposures. *Biochimica et Biophysica Acta*, 1618(2), 95-105.
- Nebel, C., Romestand, B., Nègre-Sadargues, G., Grousset, E., Aujoulat, F., Bacal, J., Bonhomme, F. and Charmantier, G., 2005. Differential freshwater adaptation in juvenile sea-bass *Dicentrarchus labrax*: involvement of gills and urinary system. *The Journal of*

- *Experimental Biology*, 208(**20**), 3859-3871.
- Ouattaraa, N., Bodinierb, C., Nègre-Sadarguesb, G., D'Cottaa, H., Messada, S., Charmantierb, G., Panfilic, J. and Baroiller, J., 2009. Changes in gill ionocyte morphology and function following transfer from fresh to hypersaline waters in the tilapia Sarotherodon melanotheron. Aquaculture, 290(1-2), 155-164.
- Perry, S. F. and Gilmour, K. M., 2006.
  Acid-base balance and CO<sub>2</sub> excretion in fish: unanswered questions and emerging models. *Respiratory Physiology & Neurobiology*, 154(1-2), 199 -215.
- Perry, S.F., Shahsavarani, A., Georgalis, T., Bayaa, M., Furimsky, M. and Thomas, S.L., 2003. Channels, pumps, and exchangers in the gill and kidney of freshwater fishes: their role in ionic and acid-base regulation. *Journal of Experimental Zoology*, 300(1), 53-62.
- Pisam, M., Moal, L.E.C., Auperin, B., Prunet, P. and Rambourg, A., 1995. Apical structures of "mitochondria-rich" α and β cells in euryhaline fish gill: their behavior in various living conditions. *The Anatomical record*, 241(1), 13-24.
- **Pratapa, H.B. and Wendelaar, B.S.E., 1993.** Effect of ambient and dietary cadmium on pavement cells, chloride cells, and Na<sup>+</sup>/K<sup>+</sup>-ATPase activity in the gills of the freshwater teleost *Oreochromis mossambicus* at normal and high calcium levels in the ambient water. *Aquatic Toxicology*, 26(**1-2**), 133-149.

- Sardella, B.A., Matey, V., Cooper, J., Gonzalez, R.J. and Brauner, C.J., 2004. Physiological, biochemical and morphological indicators of osmoregulatory stress in California Mozambique tilapia (*Oreochromis mossambicus x O. urolepis hornorum*) exposed to hypersaline water. *The Journal of Experimental Biology*, 207(8), 1399-1413.
- Sardet, C., Pisam, M. and Maetz, J., 1978. The surface epithelium of teleostean fish gills: cellular and junctional adaptations of the chloride cell in relation to salt adaptation. *The Journal of cell biology*, 80(1), 96-117.
- Tang, C.H., Lai, D.Y. and Lee, T.H., 2012. Effects of salinity acclimation on Na (+)/K (+)-ATPase responses and FXYD11 expression in the gills and kidneys of the Japanese eel (Anguilla japonica). *Comparative Biochemistry and Physiology. Part A*, 163(3-4), 302-310.
- Uchida, K., Kaneko, T., Miyazaki, H., Hasegawa, S. and Hirano, T., 2000. Excellent salinity tolerance of Mozambique tilapia (*O. mossambicus*): elevated chloride cell activity in the branchial and opercular epithelia of the fish adapted to concentrated seawater. *Zoological Science*, 17(2), 149-160.
- Uliano, E., Cataldi, M., Carella, F., Migliaccio, O., Iaccarino, D. and Agnisola, C., 2010. Effects of acute changes in salinity and temperature on routine metabolism and nitrogen excretion in gambusia (Gambusia affinis) and zebrafish (Danio rerio). Comparative Biochemistry and

- Physiology. Part A, 157(3), 283-290.
- Van Der Heijden, A.J.H., Verbost, P.M., Eygensteyn, J., Li, J. and Wendelaar Bonga, S.E. and Flik, G., 1997. Mitochondria-rich cells in gills of tilapia (*Oreochromis mossambicus*) adapted to fresh water or sea water: quantification by confocal scanning laser microscopy. The Journal of Experimental Biology, 200(1), 55-64.
- Varsamos, S., Diaz, J.P., Charmantier, G., Flik, G., Blasco, C. and Connes, R., 2002. Branchial chloride cells in sea bass (*Dicentrarchus labrax*) adapted to fresh water, seawater, and doubly concentrated seawater. *Journal of Experimental Zoology*, 293(1), 12-26.
- Walsh, P.J., Grosell, M., Goss, G.G., Bergman, H.L., Bergman, A.N., Wilson, P., Laurent, P., Alper, S.L., Smith, C.P., Kamunde, C. and Wood, C.M., 2001. Physiological and molecular characterization of urea transport by the gills of the Lake Magadi tilapia (*Alcolapia grahami*). The Journal of Experimental Biology, 204(3), 509-520.
- Wang, P.J., Lin, C.H., Hwang, L.Y., Huang, C.L., Lee, T.H. and Hwang, P.P., 2009. Differential responses in gills of euryhaline tilapia, *Oreochromis mossambicus*, to various hyperosmotic shocks. *Comparative Biochemistry and Physiology. Part A*, 152(4), 544-551.
- Watanabe, W.O., Losordo, T.M.,
  Fitzsimmons, K. and Hanley, F.,
  2002. Tilapia production systems in the Americas: Technological advances, trends, and challenges. Reviews in

Fisheries Science, 10(3-4), 465-498.

Yang, W.K., Hseu, J.R., Tang, C.H., Chung, M.J., Wu, S.M. and Lee, T.H., 2009. Na<sup>+</sup> /K<sup>+</sup>-ATPase expression in gills of the euryhaline sailfin molly, *Poecilia latipinna*, is altered in response to salinity challenge. *Journal of Experimental Marine Biology and Ecology*, 375(1-2), 41-50.

**Zydlewski, J. and McCormick, S.D., 2001.** Developmental and environmental regulation of chloride cells in young American shad, *Alosa sapidissima*. The Journal of Experimental Zoology, 290(2), 73-87.