## **Research Article**

# Analyses of macro benthic fauna in rocky habitats along the coast of Bushehr, Persian Gulf, Iran

Azizi N.1\*; Sari A.2; Fatemi S.M.1; Farshchi P.1; Mousavi-Nadushan R.1

Received: October 2017 Accepted: December 2017

#### **Abstract**

The present study considers community variability and structure at taxa-groups of macro-fauna. Sampling was carried out in three seasons at three zones in intertidal rocky shores of Bushehr, Persian Gulf. In this study, 1936 individuals belonging to 66 species were collected. These are classified into 12 groups. The mollusks were abundant in all three sites in winter and also low zone in all seasons. In contrast, Portunid crabs and Polychaets were the lowest abundant groups. All zones in three stations were occupied by mollusks and were dominated by *Planaxis sulcatus* and hence the community is named "*P. sulcatus*". Due to habitat diversity, the most abundant decapod groups were xanthoid and porcelanid crabs.

Keywords: Biodiversity, Rocky shore, Persian Gulf, Iran

<sup>1-</sup>Department of Marine Biology, College of Marine Science and Technology, Tehran Science and Research Branch, Islamic Azad University, Tehran, Iran

<sup>2-</sup>School of Biology and Center of Excellence in Phylogeny of Living Organisms, College of Science, University of Tehran, Tehran, Iran

<sup>\*</sup>Corresponding author's Email: Azizi.nn@gmail.com

#### Introduction

The marine areas around the world are threatened by human activities (Halpem et al., 2008). The coastal areas of Bushehr Province, Persian Gulf, Iran, are not exempt and affected by activities including construction of oil platforms such as south and north Parses and Emam-Hassan port in vicinity of the Khark Island, the main oil port of Iran. In the eastern part of province, the Nayband Marine Coastal Natural Park is located which was studied recently by Badri (2007) and had been identified 8 species of echinodermata in Khark and Karkho islands and Nayband Bay, Ashja-Ardalan et al. (2011)

Ashja-Ardalan et al.(2011) had been identified 97 species of mollusks, nine species of crabs and 6 species of Echinodermata, and Kohan et al. (2012) studied gastropods and reported community of Planaxis sulcatus. The other studies on Bushehr coastal waters include Sharifipour et al. (2005) on evaluation of ecological susceptibility, Vazirizadeh et al. (1997) on benthos from Bushehr intertidal zone, Azizi (2007) on biodiversity and identified 57 species of bivalvia in Khark and Karkho islands and Nayband Bay, and Savari et al. (2010) on Decapods, Vazirizadeh et al. (2011) on faunal community as an indicator of sewage pollution, Aarebi et al. (2012), Boulder type to be influenced by pollutions and very Bandar-Rostami polluted, Vazirizadeh et al. (2012) studied ecological assessments of mollusks and found out community of P. sulcatus and reported high diversity in Nayband Bay.

Nourinezhad et al. (2013) emphasized on macro-fauna, Naderloo et al. (2013), on hermit crabs, Dabbagh et al. (2012), on first record of Holothuria scabra, Farsi (2013) on biodiversity of Interand subtidal macrobenthic assemblages, Shahdadi et al. (2014) considered current status of intertidal barnacle in Persian Gulf and Gulf of Oman. Aghajanpour al.(2015)et on combining biological and geomorphological data in order to identify the biotopes and Nateghi-Shahrokni et al. (2016) on echinoderm fauna. Overall, Intertidal rocky shores are considered as biodiversity hotspots (Naderloo et al., 2013) some of which are threatened by major developments and anthropogenic born pollutants interacting with these vulnerable ecosystems. Macro-faunal community composition is a valuable index for biotope comparison. The previous mostly studies were devoted taxonomy and ecology, and therefore there was a need for current knowledge about population variability and habitat diversity. The shoreline of Bushehr province comprised of several typical intertidal habitats including rocky, muddy, sandy, and rocky/sand. These habitats are home to variety species. Therefore, the aims of this study were to demonstrate distribution of benthic macro-fauna in different intertidal levels from high tide to low tide zone and habitat diversity using community structure.

## Sampling methods

The present study was carried out in three intertidal locations at Bushehr Province, Persian Gulf, Iran in three seasons including winter and summer 2014, and spring 2015. These were selected based on substrate types. From east to west, these are Shirinu; N 27° 37′ 10.5″, E 52° 28′ 48.5″, Bandar-Amerie; N 28° 30′ 11.6″, E 51° 05′ 41.9″ and Bonak; N 29° 43′ 53.4″, E 50° 19′ 47.7″ (Fig. 1).

## Sampling locations




Figure 1: Sampling locations in Bushehr province, Iran.

Samplings were carried out at the lowest low waters levels of the spring tides. The tide tables were obtained from official site of Easy Tide (www.lukho.gov.uk/easytide). The Physical properties of the localities were similar, comprising bedrock, boulder and cobble with little or no sediment on substrate.

## Sampling

In each location, samples were collected from upper, mid and lower intertidal zones using three random quadrates 0.25 m<sup>2</sup> at each zone perpendicular to the coastline. In quadrates, all taxa including mobile, encrusting and hidden in rock crevices were collected. Samples were first separated and then transferred to the

laboratory. Finally, specimens were identified to the lowest possible taxonomic level and their abundances were recorded. The use key for identification of: Molluska (Bosch et al., 1995) and the others species identified by Naderloo and Nateghi-Shahrokni. Geographical positions were recorded hand-held **GPS** by  $(GPSMAP^{\otimes}78_{S}).$ 

## Statistical analyses

The univariate and multivariate analyses were carried out using SPSS (v.18) and PRIMER 5, respectively. The multivariate analyses include Multidimensional Scaling (MDS), Kdominance curves, cluster **SIMPER** analysis. Normality was tested using the Shapiro-Wilk's test. Comparisons of observed frequency were considered none significantly differentces at a level of p>0.05 and therefore analysis was carried out by nonparametric test. The discriminant analysis was applied to consider species compositions between three locations. The SIMPER and ANOSIM routines were applied within the PRIMER v5.0 software package (Clarke Gorley, 2001) and investigate community variability and structure in taxa-groups and order Macro-faunal communities levels. assessment based on dis-similarities were evaluated using multidimensional (MDS) and analyses scaling similarity. In both analyses, Bray-Curtis similarity indices were calculated.

Due to scarcity and lesser abundances of some species, using variable analysis, associated taxa were recognized and pulled together (Table 1). This resulted in revealing to 12 groups, which is used for community structure evaluations.

Based on frequency distribution of taxa group on each zones per locations, the partial frequency of groups are illustrated.

#### Result

This study revealed that the intertidal zone at Bushehr province considered a rich fauna. In total, 1936 individuals from 216 quadrates (0.25 m<sup>2</sup>) covering 54 m<sup>2</sup> of benthic macro-fauna have been recorded. The results have been classified into 12 groups consist of 66 species (Table 1).

Table 1: 12 taxa groups recorded in each location during three different seasons in 2014-2015.

Season	Winter	Summer	Spring																								
Station *	1	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	2	3	3	3	3	3	3	3	3	3
Zone	High	High	High	Mid	Mid	Mid	Low	Low	Low	High	High	High	Mid	Mid	Mid	Low	Low	Low	High	High	High	Mid	Mid	Mid	Low	Low	Low
Anaemons	Spore						*	*										*					*	*		*	*
Mollusks	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Polychaeta Mollusks											*															*	
Barnacles								*					*						*	*			*	*		*	*
Alpheid	duming			*			*										*										
Xanthoid	500	*	*		*	*	*	*			*			*	*		*	*		*			*	*		*	*
Porcellanid			*	*		*	*						*		*												*
Portunid	5011					*									*												

Tal	ole 1	coı	ntin	ued	l <b>:</b>																						
Season	Winter	Summer	Spring	Winter	Summer	Spring	Winter	Summer	Spring	Winter	Summer	Spring	Winter	Summer	Spring	Winter	Summer	Spring	Winter	Summer	Spring	Winter	Summer	Spring	Winter	Summer	Spring
Station *	1	1	1	1	1	1	1	1	1	2	2	2	2	2	2	2	2	2	3	3	3	3	3	3	3	3	3
zone	High	High	High	Mid	Mid	Mid	Low	Low	Low	High	High	High	Mid	Mid	Mid	Low	Low	Low	High	High	High	Mid	Mid	Mid	Low	Low	Low
Grapsid crabs										*					*						*						
Other crabs		*			*		*											*	*				*	*		*	*
Other Crustaceans			*		*		*				*				*									*			
Echinoderms				*	*		*	*										*						*			*

\* 1. Shirinu, 2. Bandar-Amerie, 3. Bonak

We observed, based on presence and absence of taxa at 12 groups, that the mollusks are abundant in three sites in all seasons and Portunid crabs and Polychaets are the least abundant groups. The dominancy of taxa groups were as follows:

Mollusks >Xanthoid crabs > other crabs >Barnacle>Porcellanid crabs, Anemones/Sponges, Echinoderms >Peracarid>Grapsid crabs, Alpheid shrimps>Portunid crabs>Polychaeta.

In total, 66 species are recorded in all seasons and localities. The species composition for the groups is shown as the presence and absence of species. This includes 44 species of mollusks (bivalves, gastropods and chitons). Others presented as echinoderms and

major crustaceans. Anemones and sponges were considered as one group because of low density and their habitat similarity.

Due to habitat diversity, crabs were divided into five groups including: Xanthoid, Portunid, Grapsid and others including less abundant ones and Porcellanids (Anomurans). The most abundant groups were Xanthoid and Porcelanid crabs.

In the winter, mollusks were dominant at the high Zone and Echinoderms at the Low and Mid Zones. In the summer, mollusks, Cirripedia (Barnacle) and other Crustacean (Xanthoid crabs) were dominant at the all zones. In the spring, mollusks and

Xanthoid crabs were dominant at the each zone (Table 1).

the Shirinu, mollusks dominant at three zones. In the Bandar-Amerie, mollusks and Xanthoid crabs were dominant at all zones. In Bonak, mollusks were dominant at the high zone. At the mid zone and low zone, from abundance point of view, anthoid mollusks. barnacles crabs. and anemones were dominant and are arranged in the following taxa-groups dominancy order using table 1:

mollusks>xanthoid crabs>barnacles >anemones -sponges

In the winter, at Shirinu, more taxa groups were found with relativly higher abundances. At Bandar-Amerie, the most abundant group was mollusks while others groups were less abundant. In Bonak, the most abundant taxa groups were mollusks (13%) while others were less abundant (alpheid shrimp, 2.3% and echinoderm 3%) (Fig. 2).

## Winter




Figure 2: Abundances of taxa groups in winter 2014 for three localities.

In the summer, in Shirinu, mollusks and Xanthoid crabs were abundant. Bandar-Amerie was similar to Shirinu based on taxa group abundance. In Bonak, more taxa groups were found with relatively higher abundances. The aAlpheid shrimp abundance was different and more than other stations (e. g. mollusks, 7.4%, barnacle,4.5%) (Fig. 3).

#### Summer




Figure 3: Abundances of taxa groups in summer 2014 for three localities.

Results of frequency distribution (Fig. 4) show significant abundances of taxa

groups in Bonak.

# **Spring**




Figure 4: Abundances of taxa groups in spring 2015 for three localities.

In Bonak, six groups were found with relatively higher abundances. These are mollusks, Xanthoid, other crabs, aAlpheid shrimp, echinoderms and Anemone-Sponge group. The Xanthoid crabs were the most abundant group (Fig. 4) in spring (8.4%) and mollusks placed in close frequency (7%).

Result of MDS, based on 12 taxa group's abundance data, showed that samples collected in summer were partially separated but in the other seasons presented overlapping and stress level indicating close similarities (Fig. 5).




Figure 5: MDS ordination of seasons using abundance of taxa groups. ▲, Winter; ■, Summer; ▼, Spring

Table 2: One-way ANOSIM results (global and pair-wise comparison between levels of each factor) for each of factor of the factors considered (999 random possible permutations from a large number possible).

Tests (factors)	Global R	R Statistic	Significance level%
Substrate type boulder- cobble boulder- bedrock cobble- bedrock	0.123	0.1 0.003 0.157 0.213	33.5 0.1 0.1
Intertidal_zone High- Mid High –Low Mid -Low	0.03	7.6 0.02 0.088 -0.021	17.7 0.7 77.9
Season Winter- Summer Winter - Spring Summer - Spring	0.106	0.1 0.107 0.103 0.116	0.2 0.2 0.2

In Table 2, results of ANOSIM tests performed the square-root transformed data matrix considering separately three summarized factors. The results are showing differences in substrate type have been observed in bedrock. While boulder and cobble showed this differences for community composition. For some seasons, factor plotted in Fig. 1, the level winter is equivalently different from summer and spring (R-values of the same magnitude). These results show there were significant differences between two levels, high and low zones.

Table 3 shows the two-way crossed ANOSIM results for the two factors "substrate type" and "season" at each levels. These results show that bedrock is an important substrate type due to high diversity and also there have been significant differences between this habitat and others. Assuming substrate type as the main discriminating factor of samples here, separate analyses were performed for each level.

Table 3: Two-way crossed ANOSIM results (global and pair-wise comparison between levels of each factor) for factors considered substrate type and season of the zone intertidal factor (999 random possible permutations from a large number possible).

Tests (factors)	Global R	R Statistic	Significance level%
substrate type		0.1	
boulder- cobble	0.269	0.087	2.7
boulder- bedrock	0.268	0.288	0.1
cobble- bedrock		0.441	0.1
Season		0.1	
Winter- Summer	0.239	0.298	0.1
Winter – Spring	0.239	0.224	0.1
Summer -Spring		0.235	0.2

The SIMPER procedure for taxa abundance contributions showed that

there are no main differences in three seasons (Table 4).

Table 4: SIMPER analysis of taxa-groups contribution (%) for each group at each season according to abundances during the study period.

Taxa groups	C	ontribution	%	Taxa groups	Contribution%				
Taxa groups	winter	summer	Spring	Taxa groups	winter	summer	Spring		
Mollusks	97	86.6	78	Xanthoid crabs			11		
Barnacle				Porcellanid crabs		4.7	4.7		
Other Crustaceans				Portunid crabs					
Polychaeta				Grapsid crabs					
Alpheid shrimp				Echinoderm					
Other crabs				Anemones/Sponges					

Also the SIMPER procedure showed differences related to substrate type in three sites. In the cobble dominated shore (Bandar-Ameri), similarity was 44 but the other substrate types

presented some overlapping indicating similarities most probably due to close abundances of mollusks. While, in the substrate dominated with boulder (Shirinu) this value was 90.6 (Table 5).

Table 5: SIMPER analysis of taxa- groups contribution (%) for each group at each substrate type according to abundance during the study period.

	C	ontribution	%		Contribution%	/ <sub>0</sub>
Taxa groups	Boulder Shirinu	Cobble Amerie	Bedrock Bonak	High zone	Mid zone	Low zone
Mollusks	90.6	92	75	96.7	86	84.8
Barnacle			14.8			
Other Crustaceans						
Polychaeta						
Alpheid shrimp						
Other crabs						
Xanthoid crabs						4
Porcellanid crabs				4.7		3
Portunid crabs						
Grapsid crabs						
Echinoderm						
Anemones/Sponges						

There are dissimilarities in abundances among localities (Shirinu and Bonak) of the present study, which was %69. This value was 67.78 for three zones mostly between low and mid levels.

The K-dominance curves on taxa group abundances for two studied levels showed that the dominance was higher in the winter level (Fig. 6) and also it was higher in the high-shore

level (Fig. 7). The differences in cumulative dominance among regions (mid and low zones) were not so strong (Fig. 7). The differences in cumulative dominance among stations showed that this value dominated in Bandar-Amerie (Fig. 8).




Figure 6: Cumulative dominance for season levels. Season: ▲, Winter; ■, Summer; ▼, Spring.




Figure 7: Cumulative dominance for zone levels. Zone: ▲, High; ■, Mid; ▼, Low.




Figure 8: Cumulative dominance for stations: ▲; Shirinu, ■; Bandar-Amerie; ▼;Bonak.

The studied community was included 12 macro-faunal groups. The total recorded number was 1936 individual in three locations. The maximum

abundance were in winter=538, summer=602 and spring=796.

Based on data (Table 8) the frequency percentages were categorized into 12 groups (Table 6).

Table 6: Taxa group seasonal dominancy (%) (based on Bouderesque 1971, quoted by Albano *et al.*, 2009). Categories: abundant (x > 15 %), common ( $5\% < x \le 15\%$ ), occasional ( $1\% < x \le 5\%$ ), scarce ( $x \le 1\%$ ).

070); setti ee (A_ 170)	•		
Taxa groups	Winter (%)	Summer (%)	Spring (%)
Anemones-Sponges	scarce (0.05)	scarce (0.3)	occasional (3.5)
Mollusks	abundant (20)	abundant (20)	abundant (18)
Polychaeta	scarce (0)	scarce (0.5)	scarce (0)
Barnacle	scarce (0.2)	occasional (5)	occasional (3)
Alpheid shrimp	occasional (2.3)	scarce (0.05)	scarce (0)
Xanthoid crabs	scarce (0.05)	occasional (1.4)	common (10)
Porcellanid crabs	scarce (0.6)	occasional (3)	occasional (2)
Portunid crabs	scarce (0)	scarce (0)	scarce (0.1)
Grapsid crabs	scarce (0.05)	scarce (0)	scarce (0.2)
Other crabs	scarce (1)	scarce (1)	occasional (1.3)
Other Crustaceans	scarce (0.05)	scarce (0.4)	scarce (0.25)
Echinoderm	occasional (3)	scarce (0.3)	occasional (3.5)

Among taxa groups, mollusks have been found in all sites in different seasons with relatively higher frequencies. The Xanthoid crabs have been common in spring and scarce in winter. The others taxa were scarce or occasionally found in seasons.

Table 7: Species locational dominancy (%) (based on Bouderesque 1971, quoted by Albano *et al.*, 2009). Categories: abundant (x>15 %), common (5% <x  $\leq$ 15%), occasional (1% <x  $\leq$  5%), scarce (x  $\leq$  1%).

Mollusks	Shirinu (%)	Amerie (%)	Bonak (%)
Planaxis sulcatus	common (13)	abundant (19)	occasional (5)
Nerita albicilla	occasional (4)	scarce (1)	occasional (1.5)
Lunella coronate	scarce (1)	occasional (5)	scarce (1)
Xanthoida			
Epixaltnus frontalis	scarce (0.5)	occasional (4)	abundant (70)

Among species, *P. sulcatus* (mollusks) in Bandar-Amerie has been found with relatively higher frequency than Shirinu and considered as common species. *Epixanthus frontalis* (Xanthoida) found with relatively higher frequency in Bonak (%).

Based on analysis of data presented in Table 8, three locations and three zones were compared. Among 12 taxa groups in three stations, significant differences observed in were echinoderm anemones/sponges and group  $(0.007 \le p \le 0.009)$ . The high between difference observed was Bandar-Amerie Shirinu and  $(0.04 \le p \le 0.07)$ .

The evident significant differences at three zones were in other crabs and anemones/sponges group  $(0.01 \le p \le 0.04)$ . In Mid-Low zones the

most significant differences between groups were observed  $(0.002 \le p \le 0.02)$ .

Kruskal-Wallis analysis among 12 taxa groups in three seasons showed significant differences in polychaetes, barnacle, Xanthoid crabs and Porcellanid crabs  $(0.03 \le p \le 0.028)$ .

The most significant differences were observed in Winter- Summer for four groups including polychaetes, barnacles, Xanthoid and Porcellanid crabs  $(0.04 \le p \le 0.008)$ , while only polychaetes were found to be different among spring-summer (p=0.04).

Table 8: Frequency of taxa groups per zone.

station	Zone	Anemones-Sponges	Mollusks	Polychaeta	Barnacle	Alpheid shrimp	Xanthoid crabs	Porcellani dcrabs	Portunid crabs	Grapsid crabs	Other crabs	Other Crustaceans	Echinoderm
	High	0	117	0	0	0	1	7	0	0	3	1	0
Shirinu	Mid	0	143	0	0	15	1	12	1	0	2	5	16
	Low	4	96	0	2	29	18	20	0	0	14	1	49
	High	0	99	2	0	0	3	15	0	1	0	3	0
Amerie	Mid	0	130	0	2	0	8	25	2	2	2	3	0
	Low	2	213	0	0	1	16	7	0	0	1	0	2
	High	0	136	0	20	0	1	0	0	2	2	0	0
Bonak	Mid	26	48	0	42	0	54	0	0	0	20	1	25
	Low	43	124	7	85	0	112	18	0	0	14	0	40

The schematic vertical distribution of taxa grouping patterns is presented (Table 8), based on proportional

dominance on the three zones of the different locations, in Bushehr province.

Table 9: Diagram of macro-fauna in three zones at three stations. Combined proportional representative of taxa group. Each sketch represents about five individual based on data in table 8. The frequency of mollusks ( ) multiplied by numbers shows number of comprising five individuals.

	Shirinu	Bandar-Amerie	Bonak		
	Boulder	Cobble	Bedrock		Legend
	800	28°	<b>3</b>		Degena
				Mollosks	٨
	♦× 23	Ô× 19	◇ × 27	Polychaeta	V
	€3€	<u>&amp;                                    </u>	6)t 6)t 6)t 6)t	Barnacie	ege
				Others.crabs	\$1 kg
eu u				Xanthoidae.crabs	& <sup>2</sup> &
High Zone				Porcellanidae.crafts	£ 2
_				Anemone-Sponge	*
				Echinoderm	*
				Alpheid shrimp	- alg
					A
				Perecarid	

	Shirinu	Bandar-Amerie	Bonak	
	Boulder	Cobble	Bedrock	Legend
	820	<i>?</i>	83	
	○×28	○ × 26	0×9 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	
Midzone			95 95 95 95 95 95 96 97 98 98 98 98	
Mio	1×××	\$ 2 £ 3	多多米米米米	
	******	£ 2	考考考考	
	Ô× 19	♦ × 42	×24 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
		\$ 2 \$ 2 \$ 2		
	\$ 2 \$ 2 \$ 2 \$ 2	\$ 2 P		
2	\$ \$ \$ \$ \$ \$ \$	\$ 3 P	8282 V	
Lo	& A		£32222	
			એક એ	
			<b>考考考考考考考 メンシン</b>	
	****			

We selected the dominant groups and defined dominant species based on species' abundance to evaluate the community. Mollusks are the most common forms per locality while polychaetes have been found with low abundance. All zones in three stations occupied by mollusks and were dominated by *P. sulcatus*. Many Xanthoid and Porcelanid crabs are

present and *Leptodius exeratus* (13%) and *Petrolisthes rufescens* (69%) were the abundant species, respectively.

Among these groups, some species were dominant. The most abundant mollusks were belonging to *P. sulcatus* (38%), *Lunella coronate* (7.5%), *Nerita albicilla* (6%), *Osilinus kotschyi* (6%) and others (42.5%).




Figure 9: The mean (+/-SE) dominant species group number in high, mid and low zones for three localities. Mean based on three replicate quadrates.

In Fig. 9, the mean dominant species are shown for three zones. Among mollusks, *P. sulcatus*, *Lunella coronate* 

and Nerita albicilla were dominant species.




Figure 10: The mean (+/-SE) dominant species in three seasons and three localities. Mean based on three replicate quadrates.

In echinoderms, the sea cucumber, *Holothuria lecouspilota* was dominant species.

According to data presented in Table 9, total mean numbers of mollusk in the winter were slightly higher than the other seasons. The total individuals of P. sulcatus in the winter were higher than the summer, and spring, showed the highest abundance whiles this value Nerita albicilla and Lunella coronate in the winter were lower than the summer. Also in the echinoderms and Alpheid shrimp taxa groups, total individuals in the winter were higher than the summer. The individuals of H. leucospilota were abundant in winter. The Porcellanid crabs group was found mostly in winter. The others groups in the summer were abundant than the winter. Among the Porcellanid crabs, P. rufescens were almost found in the summer, only one individual was found in the winter.

In the spring, Xanthoid, Porcellanid, Portunid and Grapsid crabs. echinoderms, Anemones-Spognes groups, and other crabs show total individuals higher than the two others seasons. The total number individuals of *P. sulcatus* represents the first major dominant species in all localities. The individuals of $\boldsymbol{E}$ frontalis and H. leucospilota were found mostly in the spring.

#### **Discussion**

In this section we will discuss first, taxa groups are compared per stations, second total numbers of individuals in community and also species are demonstrated for each season and third,

dominant species and community structure is evaluated per seasons and station.

Our previous knowledge and data, as stated in introduction, are mostly limited to faunistic studies. The present study is a survey on grouping of taxa based on abundances in intertidal zones of rocky shores.

In these localities, absence/presence data per seasons was considered to evaluate richness and total abundance in the communities. Here, due to impractical comparison of individual taxa at species level, variations among taxa group and distribution of benthic macro-fauna in different intertidal levels and seasons on rocky shores at Bushehr province intertidal zones have considered. These types habitats and biotope pattern has been studied less and need to be assessed to comparable substrate in the other shores from the Persian Gulf and the world (see introduction). Also, the type of shore and season seems to affect the abundance. The species diversity pattern is common in many rocky shores (Raffaelli and Hawkins, 1996; Underwood, 1997).

The variations in diversities and abundances are affected by local features of the studied shores (Cardoso et al., 2011). Results of several studies in different shores indicate that some morphodynamics have affects on the coastal macrofaunal patterns (Reichert et al., 2008; Van Colen et al., 2009; Cardoso et al., 2011). In Brazil, Cardoso et al. (2011) noted effects of morphodynamics such as grain size, wave height on abundance and

biodiversity of mollusks, which were more than the values in the sheltered shores. They noticed crustaceans were dominant on exposed shores.

In current study, common taxa groups in intertidal shores of Bushehr province such as Porifera, Cnidaria, Polychaeta, Mollusca, Arthropoda and Echinodermata have been considered. The analysis of assemblages of three localities can be explained by relationship between morphodynamics (substrate types or grain size) and population of macrofauna (Cardoso *et al.*, 2011).

The substrate type in Shirinu was boulder dominated and 49 species were recorded (in nine replicated 50×50 centimeters quadrate), while Bandar-Amerie was mainly cobble by 39 Bonak differentiable species. bedrock were recorded with 39 species (Personal observation N.A). The Rocky stations bearing crevices and rock pool provide better living opportunity for taxa. In similar study by Azizi (2007), 57 bivalve species were identified in the Nayband Bay. Later, in Khark and Kharku Islands, Badri (2007) identified eight species of Echinoderms. Ashja-Ardalan et al. (2011) identified 97 species of mollusks, nine species of crabs and six species of Echinoderms at Shirinu, one of the localities at present study.

In this study, in total 1936 individuals of benthic macro-fauna were recorded and, based on their common characteristics and functions, have been classified into 12 groups. Mollusks were more abundant taxa group. It seems that high values in

terms of abundance of certain groups belong to gastropods, which here comprised of 48% of total community in different biotic substrate. The results of other studies in the region in different years (Emam, 2005; Azizi, 2007; Badri, 2007; Ashja-Ardalan *et al.*, 2011; Vazirizadeh *et al.*, 2011; Kohan *et al.*, 2012; Vazirzadeh *et al.*, 2012; Fahimi, 2013; Aghajanpour *et al.*, 2015; Farsi, 2013) shows fluctuations in species composition.

Also, in a case study by Vazirizadeh et al. (2011), 41 species of mollusks were recorded and gastropods were found to be dominant group. In their study, continuous and intensive competition was observed in rocky stations. In another study by Kohan et al. (2012), 87 species were identified in seven rocky-sand stations, 37 species found in one station and rich diversity different habitats of Bushehr province was recorded. Nourinezhad et al. (2013) collected 29 species in Bushehr province and accordingly Asalouyeh was found as one of the habitats with a rich diversity.

There differences are between communities in high, mid and low zones. In this study, zonal pattern of community structure is considered. In Bonak lower zone, many crevices in bedrock were occupied by anemones mostly Diadumene sp. Whereas, intertidal flat in Bonak mid shore, were at tidal flow regularly and this provide suitable habitat in mid and low zones for filter feeders such common barnacle species, Amphibalanus amphitrite.

Based on all taxa evaluations, mollusks are the most contributed in three

seasons (N=1124). We observed high number of species and individual richness within community in three locations when data pulled together from three seasons.

In similar study, Ashja-Ardalan *et al*. (2011) showed consistency of mollusks in four stations rocky-sand shores comprised of 65% mollusks in winterspring.

In this survey, the most occurrences of mollusks were 13% in winter. Mollusks can survive at low temperature even if the body water gets frozen up to 70% (Raffaelli and Hawkins, 1996). The rate of water in low slope face and algae in surface area helps to increase tolerance on flow and ebb tides. Also seaweed area on rock and sand retain water for food and mollusk attachment.

In winter, mollusks in Shirinu were with similar value compare to Bonak (N<100) while in Bandar-Amerie these were more abundant (N>200).

Mollusks in summer and spring were observed with nearly similar frequency in three stations. Xanthoid crabs were more abundant in spring and showed the highest value in Bonak. In Bonak, similar to spring, more individuals of Barnacle, *A. amphitrite* was recorded in summer. While in winter there was no record of this species possibly due to cold winter and its effects of intertidal community at low tide. Porcellanid contributed with high frequency in summer and spring but appeared in lower abundance in summer.

When the three seasons in various localities have been compared, the Shirinu showed more significant

differences in summer and winter. While in Bandar-Amerie and Bonak significant differences have observed in spring. In a study by Ghiasnejad et al. (2008), autumn was recorded with high abundance of mollusks; substrate type and suitable attachment area on rocks which were important factors to support higher diversity. While Savari et al. (2010) observed decreased diversity and abundance of decapods in the Bushehr city coast. They demonstrated that even if the substrate was similar, high temperature was important on diversity status. The high biodiversity on the rocky shores seems to be due to the substrate stability and this is more stable in tropical regions (Savari et al., 2010).

As stated by Raffaelli and Hawkins association (1996)the between temperature and respiration gastropods are complex and depends on food availability. It seems movement in tidal flow is an important factor in accessibility to food in gastropod communities. In the study by Amini-Yekta et al. (2013), winter was recorded with the high abundance of mollusks species, results of the current survey also showed that winter was of high value for mollusks probably due to decrease in tidal regime in March and September.

While gastropods formed large part of mollusks (84%), some factors supported their distinct abundance in winter. These are aggregated on rock surfaces during winter for foraging, sunbathing and breeding. The wave action and the time in which these are out of water and also behavioral

variations have influences on the spatial patterns in gastropods.

In the winter, mollusks show the high value in Bandar Amerie. Possibly, the cobble coverage of the substrate in this locality provides enough suitable habitats by providing the stone surfaces and spaces among stones temperature tolerable microhabitats. While in Bonak, substrate is less divers and hence showing lesser diversity. The other two localities provide more habitat diversity to accommodate rich macro-fauna.

The results of one-way Analysis of Similarity (ANOSIM) test showed significant differences between seasons that are shown by Multidimensional Scaling (MDS) plot. Overall using pulled data, winter had low diversity and a trend level in low temperature while in spring higher diversity was observed (Fig. 4). In Figs. (2-4) the same results has been observed and probably higher diversity in spring depend temperature, food on availability and the breeding season in which more individuals are looking for appropriate mate.

In three locations, 66 species of macro-fauna (Table 1) have been identified, from which *P. sulcatus* was the most abundant of mollusk taxa groups (Tables 2, 5). Vazirizadeh *et al.* (2012) reported the high contribution of mollusks. It seems, the high diversity of gastropods is due to effective biological adaption and foraging behavior. The most important appears to be temperature because, too many *P. sulcatus* individuals have been observed resting on stone and rocks. According

to Rao and Sundaram (1972), these active in the were afternoon. Comparatively, Bonak was different from individuals species abundance point of view, as species diversity and richness are possibly linked to substrate type which was bedrock comprised of tidal pool and many crevices (Fig. 4). In mobile animals such as gastropods, the pedal attachment and mucus secretion allows firm attachment while water is at flow and ebb at stationary tidal phase (Raffaelli and Hawkins, 1996).

In the present study, crabs were categorized into five groups. The Grapsid crabs were grazer and the other predators. The Xanthoid crabs were dominant groups regarding common factors such as zone and slope face. The comparison of three zones revealed differences at high and low intertidal echinoderms. zones for crustacean (Xanthoid crabs) and anemones/sponges groups 6). Biotic and abiotic factors such as water substrate type, and human economic activities in some localities on rocky shores, can attribute to observation of these differences. Effects human activities. which were observed Bandar-Amerie and adiacent breakwater. have influences on cobble substrates community.

In Fig. 8, dominancy of *P. sulcatus* at high zone is shown while among mid and low zones there are no differences. The results of SIMPER analysis agree with those of Kruskal-Wallis test, considerable dissimilarity value between three zones were in mid-low zones (67.78). Also, as shown in Fig. 8,

the greatest dissimilarity in values between stations has observed in Shirinu and Bonak (69).

We observed aggregation of *P. sulcatus* on open surface of boulders in Shirinu between cobbles in Bandar-Amerie. Intact open bedrock in Bonak and strong temperature affect the abundance of *P. sulcatus* and the individuals were stayed in rock pool to avoid heat. Total abundance of 12 taxa groups was the highest in the spring and the lowest in winter (Fig. 4).

Recruitment for large number of P. sulcatus requires the water movement for feeding in the summer- winter transition (Denley and Underwood, 1979). This species has also been found at more sheltered region (Denny, 1988; Denley and Underwood, 1979; Leung, 2012) and its world distribution is in the northwest Pacific and Indian Ocean (Hong Kong, Thailand, Australia, the Red sea, India, the Suez Canal and the Philippines). P. sulcatus is common species and mostly was found in different seasons and zones at the Bandar-Amerie. These are more abundant in winter than the summer 2012). In winter. (Leung, dominancy was reported by high value of mollusks abundance in the Hong Kong. While in India and Japan this dominancy is reported in spring and summer (Leung, 2012). Results of the present study show that P. sulcatus are the most abundant species and observed significant difference in winter, while Leung (2012) found no variations between seasons, tidal condition and times of day (p>0.05).

Ghiasnejad *et al.* (2008), reported 29 species of gastropods from intertidal zone at in Qeshm Island and *P. sulcatus* was one of the abundant species.

P. sulcatus with frequency of more than 800 individuals per m<sup>2</sup> was reported in autumn (Ashja-Ardalan et al., 2011). Aarebi et al. (2012) showed there was significant difference in winter, summer and spring of macrofauna in polluted sediments. Also, Kohan et al. (2012) recorded the highest gastropod species in the eastern Bushehr and demonstrated that P. sulcatus is the dominant species. In Shirinu, Farsi (2013) observed 17 species of mollusks. According to Kohan et al. (2012) gastropods could be attributing as a variable to show the similarity of substrate and as these are of high value because of adaption efficiently to fluctuating environmental factors. Differences between zones such as the time spent under seawater or being out of water were real element for variability in species richness. The average density of Xanthoid crabs shows these were more abundant group after mollusks. Badri (2007) and Ashja-Ardalan et al. (2011) reported same echinoderms species in Shirinu.

Fahimi (2013) studied intertidal xanthoid crabs in Hormoz Island, *L. exetratus* and *E. frontalis* were found to be the dominant species with higher abundance in summer. Present study reports Xanthoid crabs as dominant group abundant in spring. Two species namely, *P. rufescens* (Anomura) and *L. exetratus* (Xantoid) were recorded as dominant species.

The coastal crustaceans include both littoral and supralittoral species (Defeo and McLachlan, 2013). These results confirm Defoe's opinion, as echinoderms exist in mid and low zones. The echinoderms were abundant in winter in Shirinu and spring in Bonak.

Barnacle individuals were collected at high more than the mid and low zones in the summer in Bonak and *Amphibalanus amphitrite* was dominant species. Shahdadi *et al.* (2014) reported nearly similar results. High mortality observed in barnacles could be due to flood run off, 2014 in the region and also cold winter affecting the out of water barnacles at low tide.

Since echinoderms, Xanthoid crabs and anemones need the areas with water movement for their feeding, this is probably the reason for their aggregation in the low zone and significant consequently there are differences between abundances in high and low zones.

Also, Aghajanpour *et al.* (2015), using Costal and Marine Classification Standard (CMECS), have found dominant species and defined bedrock substrate including bedrocks, crevices, cobbles and tide pools in the eastern coast of Bushehr province. In their study, seven substrate subgroups, five microhabitats in rock substrate and nine biotic groups were identified.

Future studies needed in these coastal areas to examine temporal changes, environmental parameters and industrial activities and their effects on macro-fauna.

Macro-fauna absence/presence and abundance of

species at intertidal sandy beach is a observed due to physical pattern features (McLachlan et al., 2007). According to different aforementioned report from shores at different longitude and latitude, no similar patterns for species abundance and composition are expected. Ecological features including tide time /range, beach exposure, slope, sand size, breakwater and wave power, gas or oil sources and tar balls, and season changes have marked effects on the organisms and level zones (see McLachlan et al., 2007; Defeo et al., 2009).

In conclusion, present study identified macro-fauna diversity and dominant species in the three zones, seasons at three localities. In total, 66 species were categorized in 12 taxa groups. Mollusks were the contributed in three seasons. The spring was observed to serve the highest entire diversity possibly due to higher productivity and breeding season. Among 66 species, only one gastropod was dominant species and accordingly the assemblage named "P. sulcatus community". These results are important in finding any possible patterns taxa group zonal distribution. Human and economical activities in this area have influences on the community structure and the species composition alteration, which deserve further survey.

## Acknowledgements

The authors express their gratitude to Professor M. Malek and Dr. R Naderloo, University of Tehran for their invaluable help in this study. We would like to acknowledge the following organizations for their financial, logistic and instrumental supports namely, Islamic Azad University, Science and Technology Branch, Tehran; Zoological Museum, University of Tehran; The port and Maritime Organization and Department of Environment of Iran.

#### References

- Aghajanpour, F., Savari, A., Danehkar, A. and Chegini, V., 2015. Combining biological and geomorphological data to introduce biotopes of Bushehr province, the Persian Gulf. Environmental **Monitoring** Assessment, and 187(12), 740. DOI: 10.1007/s10661-015-4956-x.
- Albano, M. and Obenat, S.A., 2009.
  Assemblage of benthic macrofauna in the aggregates of the tubiculous worm *Phyllochaetopterus socialis* in the Mar del Plata harbour, Argentina. *Journal of Marine Biological Association of the United Kingdom*, 89, 1099-1108.
- Amini Yekta, F., Kiabi, B., Ashja-Ardalan, A. and Shokri, M., 2013. Temporal variation in rocky intertidal gastropods of the Qeshm island in Persian Gulf. *Journal of the Persian Gulf*, 4(13), 9-18.
- **Arebi, I., Savari, A., and Vazirizadeh, A. 2012.** Ecological investigation of macrofauna community along Delvar sediments intertidal. *Journal of Oceanography*, 12, 27-36.
- Ashja-Ardalan, A., 1993.

  Identification and distribution of bivalvia in the Chabahar Bay. PhD thesis Tehran Azad University,

- Science and Technology branch. 345 P.
- Ashja-Ardalan, A., Zibaseresht, R. and Badri, S., 2011. Studying of distribution, identification and biodiversity of Molluscs, Crabs and Echinodermata in intertidal zone of the Gulf of Nayband. Research project, 262 P.
- Azizi, N., 2007. Identification and Distribution of Bivalvia in the Khark and Kharku Islands and Nayband Bay, Persian Gulf. MSc thesis Tehran Azad University, Marine Sciences branch. 98 P.
- Badri, S., 2007. Density and transmittal of echinoderms in Nayband Bay, Khark and Kharko Islands. MSc thesis. Khorramshahr University of Marine Science and Technology. 120 P.
- Boaventura, D., Pedro, R., Foneca, L. and Hawkins, S., 2002. Intertidal rocky shore communities of the continental Portuguese Coast: Analysis of distribution patterns. *Marine Ecology*, 23, 69-90.
- **Bouderesque, Ch., 1971.** Méthodes d'étude qualitative et quantitative du benthos (en particulier du phytobenthos). *Tethys*, 3, 79-104.
- Bolam, S.G and Eggleton, J., 2008.

  Spatial distribution of macrofaunal assemblages along the English Channel. *Journal of Marine Biological Association of the United Kingdom*, 88, 675-687.
  - **Bosch, D. Dance, P. Moolenbeek, R. and Oliver. G. 1995**. Seashells of Easter Arabia. Motivate., Dubai, UAE.

- Brown, C.J., Hewer, A.J., Meadows, W.J., Limpenny, D.S., Cooper, K.M., Rees, H.L. and Vivian, C.M.G., 2001. Mapping of gravel biotopes and an examination of the factors controlling the distribution, type and diversity of their biological communities; DEFRA Research project AE0908, Final report, Centre for Environment, Fisheries and Aquaculture Science. Lowestoft. NO 114. 300 P.
- Cardoso, R.S., Mattos, G., Caetano, C.H.S., Cabrini, T.M.B., Galhardo, L.B. and Meireis, F., 2011. Effects of environmental gradients on sandy beach macrofauna of a semi- enclosed bay. *Marine ecology*, 33, 106-116.
- Clarke, K.R. and Gorley, R.N., 2001.
  PRIMER (Plymouth Routines In Multivariate Ecological Research)
  v5: Manual/Tutorial.
- A.R., Dabbagh, Keshavarz, M., Mohammadikia, D., Afkhami, M. Nateghi Shahrokni, and S.A., 2012. Holothuria scabra (Holothuroidea: Aspidochirotida): first record of a highly valued sea cucumber, in the Persian Gulf, Iran. Marine Biodiversity Records, 5, 1-6. DOI:
  - org/10.1017/S1755267212000620
- Defeo, O., McLachlan, A., Schoeman, D.S., Schlacher, T.A., Dugan, J., Jones, A., Lastera, M. and Scapini, F., 2009. Threats to study beach ecosystems: Areview. ELSEVIER. *Estuarine, Costal and Shelf Science*, 81, 1-12.
- **Defeo, O. and McLachlan, A., 2013**. Global patterns in sandy beach

- macrofauna: Species richness, abundance, biomass and body size. *Geomorphology*, 191(1), 106-114.
- **Denley, E.J. and Underwood, A.J., 1979.** Experiments on factors influencing settlement, survival and growth of two species of barnacles in New South Wales. *Journal of Experimental. Marine Biology and Ecology*, 36, 269-293.
- **Denny, M.W., 1988.** Biology and the mechanics of the wave-swept environment. Princeton University Press, Princeton, NJ. 367 P.
- Emam, R., 2005. Identification and distribution of Bivalvia in the Khark Island, Persian Gulf. Ghasemi, S., Zakaria, M. and Mola Hoveizeh, N., 2011. Abundance of molluscs (Gastropods) at mangrove forests of Iran. *Journal of American Science*, 7(1), 660-669.
- **Fahimi, N., 2013.** Identification and distribution of xanthid crabs and reproduction of the dominant species in the intertidal zones of Hormuz Island. M.Sc. Thesis, for M. Sc. in Marine biology. Tarbiat Modares University, Noor branch. E69. https://doi.org/10.1017/S17552672120 00620.
- Farsi, A., 2013. Distribution of macrobenthic community along sub intertidal intertidal and of Bushehr. Quarterly **Journal** of Fisheries, 67, 1.
- Ghiasnejad, G., Fatemi, M.R., Jvanshir Khohi, A. and Sari, A.R., 2008. Identification and Abundance and diversity of rocky intertidal Gastropod in southern Qeshm Island.

- Journal of Environmental Science and Technology, 10(1), 205-215.
- Halpem, B., Walbridge, S., Selkoe, K., Kappel, C., Micheli, F., D'Agrosa, C., Bruno, J., Casey, K., Ebert, C., Fox, H., Fujita, R., Heinemann, D., Lenihan, H., Madin, E., Perry, M., Selig, E., Spalding, M., Steneck, R. and Watson, R., 2008. A global map of human impact on marine ecosystems. *Science*, 319, 948-952.
- **Kohan, A., Badparast, Z. and Shokri, M., 2012.** The Gastropod fauna along the Bushehr province intertidal zone of the Persian Gulf. *Journal of the Persian Gulf,* 3, 33-42.
- Leung, N.H., 2012. Spatial dispersion patterns of *Planaxis sulcatus*: Patterns and consequences. University of Hong Kong. Pokfulam, Hong Kong SAR. p5.
- McLachlan, A. and Dorvlo, A., 2007.

  Global patterns in sandy beach macrobenthic communities:

  Biological Factors. *Journal of Coastal Research*, 23, 5.
- Naderloo R., Türkay, M. and Sari, A., 2013. Intertidal habitats and decapods (Crustacea) diversity of Qeshm Island, a biodiversity hotspot within the Persian Gulf. *Marine Biodiversity*, 43(4), 445-462.
- Nateghi-Shahrokni, S.A., Fatemi, S.M.R., Nabavi, S.M.B. and Vosoughi, G.H., 2016. Contribution to the knowledge of echinoid fauna from Persian Gulf (Echinodermata: Echinoidea). *Iranian Journal of Animal Biosystematics*, 11, 1-14.
- Nourinezhad, M., Nabavi S. M. B., Vosughi Gh., Fatemi M. R.,

- **Sohrabi M., 2013.** Identification and estimation of Macrofauna in low tides of Bushehr province. *Journal of Fisheries Sciences*, 12(2), 411-429.
- Raffaelli, D. and Hawkins, S., 1996.

  Intertidal ecology. First edition.

  Chapman and Hall, London. pp. 300-350.
- Rao, K.S. and Sundaram, K.S., 1972. Ecology of intertidal mollusks of Gulf of Manner and Palk Bay. Indian National Science Academy, Part B, 38(5,6), 462-474.
- Reichert, K., Buchholz, F., Bartsch, I., Kersten, T. and Gimenez. L., 2008. Scale-dependent patterns of variability in species assemblages of the rocky intertidal at Helgoland (German Bight, North Sea). *Journal of Marine Biological Association of the United Kingdom*, 88(7), 1319-1329.
- Savari, A., Jahanpanah, M. and Vazirizadeh, A., 2010. The investigation of species diversity and dominance of Decapoda of the intertidal zone of Bushehr rocky shores, the Persian Gulf. *Journal of Oceanography*, 1(3), 7-16.
- **Shahdadi, A., Sari, A. and Naderloo, R., 2014.** A checklist of the barnacles of Persian Gulf and Gulf of Oman with nine new records. *Zootaxa*, 3784(3), 201-223.
- Sharifipour, R., Danekar, A. and Nouri, J., 2005. Determination of ecological susceptibility of the Bushehr province shoreline in the northern Persian Gulf. *Environmental Science*, 7, 45-52.
- **Underwood, A.J., 1997.** Experiments in ecology: their ecological design

and interpretation using analysis of variances. Cambridge: Cambridge university press. 482 P.

- Van Colen, C., Snoeck, F., Struyf, K., Vincx, M. and Degraer, S., 2009. Macrobenthic community structure and distribution in the Zwin nature reserve (Belgium and The Netherlands). *Journal of Marine Biological Association of the United Kingdom*, 89(3), 431-438.
- Vazirizadeh, A., 1997. Study on benthos collected from Bushehr intertidal zone. M.Sc. thesis. Chamran University. 146 p.
- Vazirizadeh, A. and Arebi, E., 2011. Study of Macrofaunal communities as indicators of sewage pollution in intertidal ecosystem: A case study in Bushehr (Iran). World Journal of Fish and Marine Sciences, 3(2), 174-182.
- Vazirizadeh, A. Mohamadi, M. and Fahkri, A., 2012. Ecological assessment community of mollusks in rocky shores, Bushehr province. . *Journal of Oceanography*, 9, 55-61.7.