Population genetic structure studies of *Liza aurata* based on mtDNA control region sequences analyses in the southern coasts of the Caspian Sea

Saeidi Z.^{1*}; Rezvani Gilkolaei S.²; Soltani M.³

Received: December 2015

Accepted: April 2017

1- Faculty of Agriculture and Natural Resource, Tehran University, Karaj, Iran.

- 2-Iranian Fisheries Research Organization, Agricultural Research Education and Extension Organization (AREEO), Tehran, Iran.
- 3-Department of Aquatic Animal Health, Faculty of veterinary medicine, University of Tehran, Tehran, Iran.

*Corresponding author's email: zohreh.saeidiii@gmail.com

Keywords: Liza aurata, Control region, Polymorphic site, Haplotype diversity

Introduction

Nowadays many species are endangered as a result of habitat loss. Decreases in population lead to reduced genetic diversity, which can cause survival crisis in a population (Cecconi et al.. 1995). Nowadays optimal management of fish stocks needs information on population structure of species that is provided to researchers through genetic science. Bereavement of science about stock composition will lead to the fracture of fisheries management and unsuitable harvest of stocks (Papasotiropoulos et al., 2007). One of the beneficial methods to diversity demonstrate genetic is haplotype analysis of the D-loop region, an index which is very important and determinant for the preservation of species. Significant genetic variation is found in the D-loop region, even among individuals within a given species. Grey mullets are not endemic species of the Caspian Sea. Juveniles of L. aurata, L. saliens and Mugil cephallus were introduced from the Black Sea into the Caspian Sea during the years 1930-1934. But only the introduction of L. aurata and L. saliens was successful and they adapted well to the ecological conditions of the Caspian Sea (Fazli et al., 2008). Papasotiropoulos et al. (2007) used mtDNA sequences analysis to explore phylogenetic relationship among five species of the Mugilidae family. Furthermore Erguden et al., 2010 examined eight Mugilid species from the Mediterranean Sea and the Black Sea on the basis of the 16S rDNA gene of mitochondrial DNA. To date there is still not enough genetic analysis about L. aurata stock except that Ghodsi et al. (2011) studied genetic diversity of L. aurata in the coasts of the Golestan province by microsatellite method and Nematzadeh et al. (2012) determined genetic differences and phylogenic relationships among six Mugilidae species (Mugil cephalus, M. capito, Liza subviridis, L. saliens, L, Valmugil buchanani) using aurata, PCR-sequencing. In this study, we used partial sequences of the mtDNA control region to estimate the genetic diversity and differentiation of L. aurata. Accordingly, the main objectives of this study were to examine the population genetic structure and evolutionary history of golden grey mullet stocks in Iranian waters of the Caspian Sea.

Materials and methods

In total, 33 fin samples were taken from L. aurata collected by beach seine from Gilan (Anzali), Mazandaran (Sari) , and Golestan (Gomishan) Provinces in January and April of 2012. The specimens were taken to Molecular Genetics Laboratory located at the Caspian Sea Ecology Research Center and stored under -20°C. DNA was extracted from 50 mg of fin sample using the method of ammonium acetate (McQuown et al., 2000). PCR was carried out using the primers D-loop F1 Forward-GGCATTTGGTTCCTACTTCAGG and 12S1-H Reverse-TGCGGAGACTTGCATGTGTAAGT for the D-loop fragment, which were designed based on Atabeyoglu (2007). PCR reactions were prepared in 50 µL total volume as follows: PCR Buffer (1X), MgCl₂ (1.8 mM), dNTP_S (0.1 mM), Primer F (1.5 Pmol), Primer R (1.5 Pmol), Tag DNA polymerase CinnaGen company (unit 1) and DNA template (100-200 ng). Reactions of PCR amplification were conducted in a thermal cycler (Auto-Q, **O**uanta biotech, England) using the following conditions: denaturation step at 95°C for 30 sec, annealing at 49°C for 30 sec and extension at 72°C for 30 sec for 30 cycles. PCR products were purified by electrophoresis in a 1.5 % agarose gel. Lastly, the positive PCR products were used as templates for **mtDNA** sequencing by dNTP method (Pherson et al., 2000). The purified DNAs of each sample with primer were transferred to the BIONEER Company in South Korea for sequencing. The mtDNA control region sequences of all samples were aligned using Clustal X (Thomson et al., 1997) in BioEdit. Genetic diversity indices, number of haplotypes (N), haplotype diversity (h), nucleotide diversity (P), polymorphic sites (s) and fixation index (Fs_T) were estimated using the software DnaSP (Rozas *et al.*, 2003). The mean difference of paired nucleotide (Tamura et al., 2007) within and among samples of regions and the Neighbor-Joining (NJ) tree was constructed using MEGA (ver.5.05). Genetic distance within samples was estimated using (Nei, 1978) model. Estimation of gene flow (Nm) was derived using the equation: $Nm=[(1/Fs_T)-1]/2$ (Weir and Cockerham, 1984).

Results and discussion

After alignment, DNA sequences of the control region (D-loop) in golden grey mullet were 900 bp. One sequence data from each region were submitted to GenBank (accession numbers. KF418242 . KF465679 . KJ769204). Individual nucleotides were compared to identify conserved and mutated nucleotides. From a total of 923 study areas, a total of 489 protected sites and 420 polymorphic sites were observed in the gene sequence. The haplotype diversity (h) values of golden grey mullet were 0.989±0.002. 1.000±0.044. and 1.000 ± 0.052 in the Gilan. Mazandaran, and Golestan regions, respectively. The nucleotide diversity (P) values were 0.085, 0.023, and 0.024 in the Gilan, Mazandaran, and Golestan regions. respectively. The highest nucleotide diversity was observed in the Gilan region (0.085) and lowest in Mazandaran (0.023)(Table 1). The genetic diversity was calculated respectively(0.989±0.002, 1.000±0.044, 1.000±0.052) in the Gilan, Mazandaran and Golestan Provinces, according to Nie (1987) model. Based on the AMOVA (Analysis of Molecular Variance), the highest genetic variations (0.71)were observed within populations and lowest (0.09) was among populations within regions (Table 2).

The high rates of Fs_T value were observed between Gilan and (0.695),Mazandaran also between Gilan and Golestan (0.692), and the lowest was among Golestan and Mazandaran (0.067) Provinces. Based on the Nei (1987) model, the highest rate of gene flow (3.470) was observed between Mazandaran and Golestan regions, and in addition the lowest rate (0.958) was between Mazandaran and Gilan (Table 3).

 Table 1: Haplotype diversity (h), nucleotide diversity (P), observed heterozygosity (H_o), expected heterozygosity (H_e) and Hardy-Weinberg equilibrium (HWE) of golden grey mullet samples in the Caspian Sea.

Location	N	h	P	H	H _e	HWE	Tajima'D
Gilan	14	0.989	0.085	0.374	0.353	0.024	-1.389
Mazandaran	10	1.000	0.023	0.301	0.298	0.021	-0.869
Golestan	9	1.000	0.024	0.281	0.203	0.033	-0.443

Table 2: Analysis of molecular variance (AMOVA) of golden grey mullet in the Caspian Sea. Sum square (SS), p value (Pro).

Source of variation	SS	Est.var.	%	Pro
Among regions	23.47	0.024	0.20	0.030
Among populations within regions	13.41	0.092	0.09	0.021
Within populations	96.58	0.132	0.71	0.026

Downloaded from jifro.ir on 2025-08-02

Table 3: Gene flow (Nm) of golden mullet samples in the Caspian Sea.						
Nm	Gilan	Mazandaran	Golestan			
Gilan	-	0.958	1.020			
Mazandaran	0.958	-	3.470			
Golestan	1.020	3.470	-			

The Nei (1978) model was used to estimate the genetic distance. Hence the genetic distance estimated between Gilan and Mazandaran, Gilan and Golestan, as well as between Golestan and Mazandaran Provinces were 0.458, 0.452, and 0.081, respectively. Phylogenic relationships among golden grey mullet were calculated by MEGA software and divergence time was estimated using Tahjima's test (Tajima, 1993). The results obtained from genetic differences showed that there were significant differences among Gilan with Mazandaran and Golestan Provinces (Fig. 1).



Figure 1: Neighbor-Joining (NJ) tree drawn for golden grey mullet in the Gilan, Golestan and Mazandaran Provinces.

Grey mullets are distributed worldwide, inhabiting coastal and brackish waters of all temperate and tropical regions of the globe (Papasotirpoulos et al., 2007), so they can be compared with marine species. Zhongduo et al.. 2014 investigated high sequence variation and low population differentiation of mitochondrial control regions of wild large yellow croaker as a marine species in South China Sea. they found that the haplotype diversity values of different populations were all higher than 0.980. Furthermore Jean et al. (2014) surveyed population genetic structure in the endemic cyprinid fish Microphysogobio alticorpus in Taiwan by the cyt b and d-loop region analyses and found the average haplotype diversity value was 0.896. Our study indicated high levels of haplotype and nucleotide diversities of L. aurata in the southern coasts of the Caspian Sea. Our results obtained from this study on halplotype diversity are in agreement with the results on mullet fish of the Mediterranean Sea (Erguden et al., 2010) and golden grey mullet (Rossi et al.. 1998; Caldara et al., 2002; Papasotirpoulos et al., 2007). A similar conclusion was reached by Naderi (2011) who observed high levels of halplotype diversity and more than one population of L. saliens in the southern coasts of the Caspian Sea using microsatellite method. Two different populations of L.aurata in the southern coasts of the Caspian Sea is a sign of our data analysis in our study. The genetic distance between Gilan

Province and Mazandaran and Golestan Provinces indicates the high difference between the Gilan population and other populations in Mazandaran and Golestan regions. In this study, the gene flow rates between Gilan and Mazandaran and that between Gilan and Golestan Provinces were low which indicated the infrequent rates of immigration between Gilan region and the two other regions. Hereupon the reproductive isolation occurred in the Gilan population which is a factor for the constitution of different populations. Inbreeding of populations within a species produced a unique gene pool of population. and reproductive that characteristics are the main elements for population differentiation (Turan et al., 2005). High levels of gene flow were observed among grey mullets in Ghaneh (2011) (5.05) and Ghodsi et al. (2011) (5.153-39.264) investigations. In this study the observed genetic diversity for golden grey mullet was 1.000 in the Golestan and Mazandaran Province, which is in agreement with marine species such as Sciaenops ocellatus (Gold and Richardson, 1991), Clupea harengus and Brevoorita tyrannus (Kornfield and Boydanowicz, 1987; Avise et al, 1989) but in contrast with Anguilla rostrata, Arius felis and Cynoscion nebulosus (Avise et al., 1989). The Fs_T was calculated at 0.692 and 0.695 between Gilan province and Golestan and between Gilan and Provinces Mazandaran (p < 0.05). respectively. The overall standardized Fs_T value among all samples in Jean et al. (2014) was 0.876. In contrast to the present study, Ghodsi *et al.* (2011) calculated the Fs_T value at 0.016. As a conclusion two populations of *L.aurata* exist in the Iranian waters of the Caspian Sea based on our study as a result of different ecological status in these regions.

References

- Atabeyoglu, K., 2007. Determination genetic differences between of mtDNA D-Loop F1 and 12S1-H region of native salmons (Salmo trutta) caught in the Rivers of Aras, Karasu and Coruh in our district using PCR-RFLP and microsatellite methods. MS Thesis, Department of Fisheries, institution of Natural and Applied Sciences. Ataturk University, 62P.
- Avise, J.C., 1989. A role for molecular geneticists in the recognition and conservation of endangered species. *Trends in Ecology and Evolution Journal*, 4, 279–281.
- Caldara, F., Bargelloni, L., Ostellari, L., Penzo, E., Colomb, L. and Patarnello, T., 2002. Molecular phylogeny of grey mullets based on mitochondrial DNA sequence analysis: Evidence of a differential rate of evolution at the intra family level. *Molecular Phylogenetics and Evolution*, 6, 416-424.
- Cecconi,F.,Giorgi,M.andMariottini,P.,1995.Uniquefeatures in the mitochondrial D-loopregionoftheEuropeanseabass

Dicentrarchus labrax. Gene, 160(2), 149-155.

- Erguden, D., Gurlek, M., Yaglioglu, D. and Turan, C., 2010. Genetic identification and taxonomic relationship of Mediterranean Mugilid species based on Mitochondrial 16S rDNA sequence data. Journal of Animal and *Veterinary Advances*, 9(2), 336-341.
- Excoffier, L., Guillaume, L. and Schneider, S., 2005. Arlequin (version 3.0): An integrated software package for population genetics data analysis. *Evolutionary Bioinformatics*, 1, 47-50.
- Fazli, H., Ghaninejad, D., Janbaz, A.
 and Daryanabard, R., 2008.
 Population ecology parameters and biomass of golden grey mullet (*Liza aurata*) in Iranian waters of the Caspian Sea. *Fisheries Research*, 93, 222-228.
- Ghaneh, R., 2011. Study of genetic diversity of *Liza saliens* in the southern coasts of the Caspian Sea by mtDNA method. Thesis report submitted to the Office of Graduate Studies of Islamic Azad University, Science and Research Branch, in partial fulfillment of the requirements for the Master degree.
- Ghodsi, Z., Shabani, A. and Shabani Pour, B., 2011. Study of genetic diversity of *Liza aurata* in the coasts of the Golestan Province by microsatellite method. *Taxonomic and Biosystematics Journal*, 3(6), 35-49.

- Gold, J.R. and Richardson, L.R., 1991. Genetic studies in marine fishes. Ananalysis of population structure in the red drum (*Sciaenops ocellatus*) using mitochondrial DNA. *Fisheries Research*, 12, 213-241.
- Jean, C.T., Wu, C.Y., Tsai, K.C., Wang, W.K., Hsu, Y.Y., Chang, Y.M., Lin, H.D., 2014. Population *Microphysogobio alticorpus* in Taiwan: Evidence for a new phylogeographical area. *Biochemical Systematic and Ecology*, 57, 108-116.
- Kornfield, I. and Boydanowicz, S.M., 1987. Differentiation of mithochondrial DNA in Atlantic herring, *Clupea harengus. Fishery Bulletin*, 85, 561-568.
- McQuown, E.C., Sloss, B.I., Sheehan, R.J., Rodzen, J., Tranah, G. and May, B., 2000. Microsatellite and analysis of genetic variation in sturgeon: new sturgeon primer sequences for Scaphirhynchus and Acipenser. Transactions of the American Fisheries Society, 139, 1380-1388.
- Naderi, L., 2011. Study of genetic diversity of *Liza saliens* in the southern coasts of the Caspian Sea by microsatellite method. Master Degree Thesis. Gorgan University of Agricultural Sciences and Natural Resources.
- Nei, M., 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics*, 89, 583-590.

- Nei, M., 1987. Molecular evolutionary genetics. Columbia University Press, New York .512 P.
- Nematzadeh, M., Rezvani, S., Khalesi, M.K., Laloei, F. and Fahim, A., 2012. A phylogeny analysis on six mullet species (Teleosti: Mugillidae) using PCRsequencing method. *Iranian Journal* of Fisheries Sciences, 12(3), 669-679.
- Papasotiropoulos, V., Klossa-Kilia, E., Kilias, G., Alahiotis, S. and G., 2007. Molecular Kilias, phylogeny of grey mullets (Teleostei: Mugilidae) in Greece: Evidence from sequence analysis of mtDNA segments. **Biochemical** Genetics Journal, 45, 623-636.
- Pherson, M.M., Moller, S. and Moller, S.G. 2000. PCR (Basic: From background to bench). BIOS Scientific Publishers Ltd. population structure. Evolution, 38, 1358-1370. Raymond, M. and Russet, E., 1995. GENEPOP (version 1.2): Population genetics software for tests and ecumenicism. exact Journal of Heredity, 86, 248-249.
- Rossi, A.R., Capula, M., Crosetti, D., Campton, D.E. and Sola, L., 1998. Genetic divergence and phylogenetic inferences in five species of Mugilidae (Pisces: Perciformes). *Marine Biology*, 131, 213-8.
- Rozas, J., Sanchez, J.C., Delbarrio, X. and Rozas, R., 2003. DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics*, 19, 2496-2497.

DOR: 20.1001.1.15622916.2017.16.4.19.6

- Tajima, F., 1993. Simple methods for testing molecular clock hypothesis. *Genetics*, 135, 599-607.
- Tamura, K., Dudley, J., Nei, M. and S., Kumar, 2007. Mega 4: Molecular evolution genetics analysis (MEGA) software version Molecular 4.0. **Biology** and Evolution, 24, 1596-1599. PMID: 17488738. http://www.ncbi. nlm.nih. gov/ pubmed/17488738.
- Thomson , J.M., 1997. The Mugilidae of the world. Meme Queens Us. 41, 457-62.
- Turan, C., Caliskan, M. and Kucuktas, H., 2005. Phylogenetic relationships of nine mullet species (Mugilida) in the Mediterranean Sea. *Hydrobiologia*, 532, 45-51.
- Weir, B.S. and Cockerham, C.C., 1984. Estimating F-statistics for the analysis of population structure. *Evolution*, 38, 1358–1370.
- Zhongduo, W., Chong, C., Yusong, G. and Chuwu, L., 2014. High sequence variation and low population differentiation of mitochondrial control regions of wild large yellow croaker in South China Sea. *Biochemical Systematic and Ecology*, 56, 151-157.