# **Research Article**

# Comparative effect of different vaccines on immune-related gene expressions of rainbow trout (*Oncorhynchus mykiss*) and experimentally infected with *Streptococcus iniae* and *Lactococus garvie*

Khaj H.1\*; Mesbah M.2; Tabandeh M.R.3; Mohammadian T.2; Dadar M.4

Received: March 2019 Accepted: November 2019

#### **Abstract**

To understand how different vaccines, i.e., bivalent formalin-inactivated vaccine (Lactococus garvie and Streptococcus iniae), could affect the immune-related gene expressions, the TNF-alpha, IL-1β and IL-8 genes of head kidney were measured in Oncorhynchus mykiss. To address this, 630 juvenile fish weighing 26±3 g were initially (day 0) injected by three different treatments, including control group (normal saline only), AquaVac<sup>TM</sup> Garvetil (Merck) and Iranian vaccine (Jehad-e-Daneshagahi) and followed by bath immersion into the same vaccine on day 30. The head kidney was withdrawn to measure immune-related gene expressions at the beginning of experiment (day 0), and following 14, 30, 45 and 60 days of post-vaccination. In the next step, control and both vaccinated groups were also subjected to either L. garvie or S. iniae challenge. Similar samples were taken immediately following bacteria injection (time= 0) and 12, 48, 72 h, and following 7 and 10 days of challenged test. The real-time PCR indicated up-regulation of all cytokine genes following vaccination at day 14 as compared to initial day (p<0.05). In pre-challenged experiment, the higher induction in the level of those genes in the head kidney was related to AquaVac vaccine (p<0.05). Injection by S. iniae and L. garviae induced the level of immune-related gene expression in the head kidney within the first few days with higher intensity in case of unvaccinated control group. Although the AquaVac produces higher up-regulation of inflammatory gene expression rather than Iranian vaccine, the immunostimulatory effects of both vaccines is a time-restricted.

**Keywords**: Cytokine genes, Pro-inflammatory genes, Bacterial infection, Passive vaccination, *Oncorhynchus mykiss* 

<sup>1-</sup>Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Karaj, Iran

<sup>2-</sup>Department of Clinical Science, Faculty of Veterinary, Shahid Chamran University of Ahvaz, Ahvaz, Iran

<sup>3-</sup>Department of Basic Science, Faculty of Veterinary, Shahid Chamran University of Ahvaz, Ahvaz, Iran

<sup>4-</sup>Department of Brucellosis, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Karaj, Iran

<sup>\*</sup>Corresponding author's Email: Hosseinkhaj@yahoo.com

## Introduction

Oncorhynchus mykiss is considered as one of the most important commercial fish species for aquaculture in many countries, including Iran. Bacterial infections induced heavy economic losses on this species all over the world (Vendrell et al., 2008; Raissy et al., 2018). Although the antibiotic treatment was futile to some extent. aquaculturists are still eager to apply different types of antibiotic even for prevention. Moreover, the emergence of antibiotic-resistance for these pathogen strains (Park et al., 2009; Marti et al., 2018) negated the success of this method for outbreak control (Sakai et al., 1995), and consequently, suggesting the necessity of an alternative way such as vaccination to protect fish against the infections (Dimitroglou et al., 2011; Faber et al., 2019).

Amongst trout infectious diseases. Lactococcus garvieae and Streptococcus iniae, the causative agent for lactococcosis and streptococcosis, respectively accounted massive outbreaks in the aquaculture industry especially during warm seasons (Diler et al., 2002; Didinen et al., 2014; Raissy et al., 2016). It is also considered to be responsible for more than 50% of losses in trout farms (Vendrell et al., 2006; Shiry et al., 2019). These gram-positive non-motile bacteria affect different freshwater and marine species (Nakai et al., 1999; Chen et al., 2001; Evans et al., 2006; Bastardo et al., 2012). The lactococcal septicemia not only produces clinical signs but also some lesions (such as fibroplasia) could be observed at the serous membrane of different organs (Chang *et al.*, 2002). The infection caused by *S. iniae* is responsible for several pathological changes, including meningitis, lesions in internal organs as well as sepsis among different species (Eldar *et al.*, 1999; Eldar and Ghittino, 1999). It is believed that they can threat the public health by producing some chronic diseases like skin lesions in human (Elliott *et al.*, 1991; see the review written by Novotny *et al.*, 2004; Wang *et al.*, 2007).

Besides the costs forced by drug treatments in large aquaculture systems, bacterial resistance and lack efficiency could constrain antibiotics application to aquaculture industry (Harikrishnan et al., 2010). Protection of aquatic species against lactococcosis streptococcosis were already examined both in experimental and field studies (Eldar et al., 1997; Ooyama et al., 1999; Soltani et al., 2019). Previously, it has been shown that the i.p. vaccination with formalinkilled Streptococcus sp. in the presence of adjuvant could protect O. mykiss against this pathogen while the other kind of immunization, i.e., bath immersion did not act the same (Akhlaghi et al., 1996). The combined vaccines are also developed to prevent fish from bacterial and viral pathogens (Sakai et al., 1995). However the efficacy of combined vaccines for lactococcosis and streptococcosis did not investigate well on O. mykiss.

Currently, the importance and success of vaccinations against bacterial

diseases in fish are obvious; however, little attempts are fulfilled to understand the mechanisms of vaccine-induced disease resistance (Harun et al., 2011). One of the possible mechanisms for pursuing how immune system was altered after vaccination is, however, to examine the immune-related expression. Activation of macrophages by fish vaccination results in releasing various cytokines such as IL-1, IL-6 and TNF-alpha and enhancing T-cell mediated cytotoxicity and proliferation (Dieraba and Ouere, 2000; Evans et al., 2004; Mohammadian et al., 2019). It typically determined following vaccination against infectious haematopoietic necrosis virus yersiniosis in O. mykiss by Purcell et al. (2004) and Harun et al., (2011), respectively, but similar studies on lactococcal or streptococcal vaccinations are quite rare and limited to other fish species (Sun et al., 2012). In this paper, we aimed to examine how O. mykiss can be immunized following injection with repetitive bath immersion of two different bivalent formalin-inactivated vaccines (L. garvie and S. iniae) in a time-course manner. To do this, some immune-related gene expression was evaluated to obtain an insight into the possible underlying mechanisms that immunized fish might respond to vaccination. The ability of i.p and bath vaccinations was also followed up by experimental infections with either L. garvie or S. iniae.

## Materials and methods

#### Fish maintenance

Juvenile rainbow trout, O. mykiss (n=630)weighing  $26 \pm 3$ g were purchased from a local supplier in Loresatn province (Iran) and transferred to the Aquatic Animal Health Lab at Shahid Chamran University of Ahvaz. All fish were acclimated for at least 2 weeks before experiments commenced in 300 l tanks. Acclimation tanks were consisted air supplier to remain dissolved oxygen higher than 10 mg/l and located in a temperaturecontrolled room (16±2°C). The fish initially examined bacterial pathogens and parasites. All fish were fed three times a day ad libitum with diet containing 36.81% crude protein, 11.33% crude 11.58% crude fiber and 3.50% (GFT1- Fardaneh Company).

## **Treatments**

O. mykiss juveniles were randomly divided into three groups (treatments). Each group containing 210 fish (each group included three tanks and each tank contained 70 fish). The first group only received normal saline was (similar to vaccine volume) and considered as control. The second group was injected by AquaVac<sup>TM</sup> Garvetil (Merck, Germany) and third group was injected by Iranian vaccine (Jehad-e-Daneshagahi). The vaccines contained 1×109 S. iniae cells /mL and  $1\times10^9$  L. garviae cells /mL. The injected dose was similar for both vaccines and included 0.1 mL per fish (according manufacture manual). All abovementioned treatments were followed by bath immersion into the same vaccine on day 30. To do this, 1000 g fish were immersed into 9 l water containing 1 l of vaccine for 5 min (all procedures were done according to manufactures' protocol). All fish were deprived of food for two days before either injection and/or immersion bath.

## Sample collection

Three fish from each tank were randomly selected and anesthetized by 2-phenoxyethanol (0.3 mL/l, Merck, Germany). The head kidney was -70°C withdrawn and stored at (Matsuyama et al., 2007) for further measuring of immune-related gene expression, including tumor necrosis factor alpha (TNF-alpha), interleukin 1 beta (IL-1β) and interleukin 8 (IL-8) at the beginning of experiment (day 0), and following 14, 30, 45 and 60 days. In the next step, the control group, Iranian AguaVac and vaccine treatments were also subjected to either L. garvie or S. iniae challenge test. Similar samples were taken immediately following bacteria injection (time= 0) and 12, 48, 72 h, and following 7 and 10 days of challenged test.

# Gene expression

RNA isolation and cDNA synthesis

At each exact time, the head kidneys' RNA was isolated (see the sampling section), using the Tri Pure isolation reagent according to the manufacturer's

procedure (Roche, Canada). Fifty mg of each sample was taken under sterile condition. The RNA concentration was determined by using nano-drop (Eppendorf, Germany) and the purity of the RNA was determined by the optical density (OD) absorption ratio at 260/280 nm. Reverse transcription was carried out with the Rocket Script RT PreMix Kit using 1 µg of RNA and oligo dT based on the manufacturer's protocol (Bioneer Corporation, South Korea).

# Real-time quantitative PCR

To evaluate the expression levels of IL-1β TNF-alpha, and IL-8 (Inflammatory genes) (Secombeset al., 1998; Secombeset al., 2011) mRNA in the head kidneys, real-time PCR was performed using qPCRTM Master Kit for SYBR Green I® (Jena Biosciense, Germany) on a Light cycler® Detection System (Roche, USA). Relative expression levels of the all transcripts were compared to  $\beta$ -actin. Specific sets of primers (Bioneer Co., South Korea) were designed based on O. mykiss (Table 1). Reactions were performed in a 12.5 µL mixture containing 6.25 µL qPCR TM Green Master Mix (2X), 0.25 µL of each primer (10 μM), 3 μL (100 ng) cDNA, and 2.75 µL nuclease-free water. The PCR protocol consisted of a 5 min denaturation at 94 °C followed by 45 cycles at 94 °C and 60° C for 30 sec (in triplicate). Two separate reactions without cDNA or with RNA were performed as control groups in parallel with experimental groups. According to the comparative  $2^{-\Delta\Delta Ct}$  method, the relative quantification was performed using Light cycler 96® software (Mohammadian *et al.*, 2018).

All qPCR analysis was performed according to the Minimum Information for Publication of Quantitative Real-Time PCR Experiments (MIQE) guideline (Bustin *et al.*, 2009).

Table 1: Primers used for detection of target genes.

Genes	Product size	Primer (5'-3')	Access Number
EF1α	327bp	Forward CAAGGATATCCGTCGTGGCA	AF498320
		ReverseACAGCGAAACGACCAAGAGG	
TNF-	102 bp	Forward TGTGTGGGGTCCTCTTAATAGCAGGTC	AJ277604
alpha		ReverseCCTCAATTTCATCCTGCATCGTTGA	
IL-8	226 bp	Forward GAATGTCAGCCAGCCTTGTC	HG917307.1
		ReverseTCCAGACAAATCTCCTGACCG	
IL1-β	239 bp	Forward CCTGGAGCATCATGGCGTG	AJ278242
		ReverseGCTGGAGAGTGCTGTGGAAGAACATATAG	

# Bacterial challenge

The *L. garvieae* and *S. iniae* strains were originally isolated from naturally infected trout and cultured routinely on tryptone soy agar (Difco) plates, at 27°C for 48 h. The isolates were identified as *L. garvieae* and *S. iniae* by standard methods like colony and cell morphology, gram staining and biochemical characteristics.

Prior to challenge test of O. mykiss with either L. garviae or S. iniae, the lethal dose (LD) of each bacterium was determined. First, each bacterium was grown on TSA at 27°C for 48 h. The culture medium was then centrifuged at 3500 rpm for 10 min. The bacteria were washed twice with normal saline and concentration of bacteria was adjusted to 1×10<sup>9</sup> CFU/mL with McFarland tubes. Serial dilutions  $(1\times10^5)$  up to  $1\times10^9$  CFU/mL) were selected for i.p injection. The mortality was recorded daily for 4 continuous days and the obtained data was subjected to probit regression analysis calculate LD<sub>50</sub>(Sun *et al.*, 2011). The amounts of LD<sub>50</sub> for each bacterium, i.e., *L. garviae* and *S. iniae* were injected ( $10^7$  CFU/mL and  $10^7$  CFU/mL, respectively) as i.p. doses for further experiments.

## Statistical procedure

If data approved for normality test (Kolmogorov–Smirnov) and homogeneity of variance (Levene), further statistical test have been Two-way performed. Analysis Variance was applied to determine the effects of treatments and time on different parameters in the first experiment (before challenge test). Multi-way of Variance Analysis (MANOVA) performed was determine the combined effect treatments, time and bacterial strains on gene expressions after challenge test. The multiple comparisons (Duncan) were followed if the p value was statistically significant (SPSS, 18). Logarithmic transformations of bacterium doses were used to calculate the LD<sub>50</sub> at exact time durations using the probit regression analysis (SPSS, 18). All experimental data were presented as the mean $\pm$ SD, and the level of significance for all tests was set at p<0.05.

## **Results**

In order to evaluate whether different vaccines and time interval may composed of any changes in immunerelated gene expression of *O. mykiss* 

prior to the challenge test, the data were subjected to Two-way ANOVA. In spite of any possible statistical difference originated from abovementioned single factor, the significant combined effects were observed in case of TNF-alpha, IL-1β and IL-8 genes of head kidney. Moreover, in case of challenged test, the MANOVA revealed significant combined effects for all measured parameters (Tables 2 and 3).

Table 2: Two-Way ANOVA performed for each parameter with P value in pre-challenge test.

	Time	Treatment	Time× Treatment
TNF-alpha	< 0.001	< 0.001	< 0.001
IL-1β	< 0.001	< 0.001	< 0.001
IL-8	< 0.001	< 0.001	< 0.001

Table 3: MANOVA performed for each parameter with P value in post-challenge test

Table 5: MANOVA performed for each parameter with P value in post-challenge test.										
	Time	Treatment	Bacterial Strain	Time× Treatment	Time× Bacterial Strain	Treatment× Bacterial Strain	Time× Treatment ×			
							Bacterial Strain			
TNF- alpha	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001			
IL-1β	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001			
IL-8	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001	< 0.001			

Pre-challenge test The TNF-alphagene expression in the head kidney of fish, vaccinated with AquaVac showed a significant increase at different time interval as compared to day 0 (p<0.05). The higher TNF-alpha was also observed following 14 days vaccination as compared to other sampling times (p < 0.05). This gene was restored to its initial level on day 45 and remained unchanged even on day 60. This pattern was also observed in TNFalpha gene expression in the head kidney of fish vaccinated by Iranian treatments. Regarding that, the only significant increase was observed on days 14 and 30 as compared to other sampling times (p<0.05), indicating a peak at those time intervals. Although there was a significant difference in TNF-alphaamong different treatments at the beginning of experiment, this difference was much obvious on day 14 with higher level for AquaVac (p<0.05). The higher TNF-alphagene expression was also observed in both vaccine treatments on day 30 as compared with control but this change did not last following 45 and 60 days of experiment (Fig. 1).




Figure 1: The effects of different vaccines on head kidney TNF- alpha gene expression at different time intervals. All values were obtained from 9 individual fish (3/replicate) and expressed as mean $\pm$ SD. Different lowercase alphabetic letters on each bar indicate significant difference among different treatments and different capital letters express significant difference among different sampling times (p<0.05).

IL-1 $\beta$  gene expression in the head kidney showed a significant increase in all vaccinated and unvaccinated groups following day 14 (p<0.05). As it is shown in Figure 2, the only significant increase in IL-1 $\beta$  gene expression was related to day 14 for Iranian and AquaVac vaccines in comparison with control group (p<0.05).

The results obtained from IL-8 gene expression exhibited a similar trend for both sampling time and treatment as we observed for IL-1 $\beta$  (Fig. 3). In this regard, an up-regulation of IL-8 was

observed in all groups following 14 days of experiment. The higher level of this gene was also observed at that time in treated fish with AquaVac (p<0.05).

## Post-challenge test

The real-time PCR shown that TNF-alphagene expression was elevated following 12 h after i.p. injection of *S. iniae* in control untreated group and this parameter was declined significantly following 48 and 72 h as compared to 12 h (p<0.05). This decreasing trend was last for even 7 and 10 days after *S.* 

*iniae* injection in control group and reached the value similar to injection

time (day 0).




Figure 2: The effects of different vaccines on head kidney IL-1B gene expression at different time intervals. All values were obtained from 9 individual fish (3/replicate) and expressed as mean $\pm$ SD. Different lowercase alphabetic letters on each bar indicate significant difference among different treatments and different capital letters express significant difference among different sampling times (p<0.05).




Figure 3: The effects of different vaccines on head kidney IL-8 gene expression at different time intervals. All values were obtained from 9 individual fish (3/replicate) and expressed as mean $\pm$ SD. Different lowercase alphabetic letters on each bar indicate significant difference among different treatments and different capital letters express significant difference among different sampling times (p<0.05).

A significant increase (p<0.05) in TNF-alpha gene was also observed in both vaccinated groups (especially Iranian vaccine) in which they reached a peak at 48 and 72 h after injection. The upregulation of this gene in the head kidney did not last for 10 days and came back to their initial level.

Interestingly, the higher (p<0.05) level of TNF- alpha gene expression was observed in control challenged group (S. iniae) as compared with two vaccinated groups in all sampling times, except for time 0. This difference was also attenuate on day 10 in which a slight decrease (p<0.05) was only for Iranian vaccine (Fig. 4).




Figure 4: The effects of different vaccines on head kidney TNF- alpha gene expression following challenge test with S. iniae at different time intervals. All values were obtained from 9 individual fish (3/replicate) and expressed as mean $\pm$ SD. Different lowercase alphabetic letters on each bar indicate significant difference among different treatments and different capital letters express significant difference among different sampling times (p<0.05).

Similar to *S. iniae* challenge test, the control group in *L. garviae* was also showed a significant up-regulation in all sampling points except for day 10 as compared with injection time (p<0.05). Similar trend was also observed for

other treatments but with lesser intensity. Although the up-regulation of this gene was mostly higher in control group (i.e., at 12 and 48 h), significant increase was also found in AquaVac treatment in which showed similar or

even higher (p<0.05) level of TNF-alpha as compared to control (Fig. 5). The pattern observed for IL-1 $\beta$  was similar to what we observed for TNF-alphafollowing *S. iniae* or *L. garviae* i.p. injections in which the higher level

was obtained following 48 h of injection (p<0.05). In addition, the vaccine treatments resulted in significant down-regulation (p<0.05) of this gene approximately at all sampling times with much lower level for Iranian vaccine (Figs. 6 and 7).




Figure 5: The effects of different vaccines on head kidney TNF- alpha gene expression following challenge test with L. garviae at different time intervals. All values were obtained from 9 individual fish (3/replicate) and expressed as mean $\pm$ SD. Different lowercase alphabetic letters on each bar indicate significant difference among different treatments and different capital letters express significant difference among different sampling times (p<0.05).

Similar to other measured genes, the IL-8 was also shown the same trend following *S. iniae* injection (Fig. 8). The control group indicated the higher (p<0.05) level of this gene at all sampling points, except for day 10.

The *L. garviae* injection resulted in significant rise in IL-8 gene expression in the head kidney in all treatments following 48 and 72h of injection as compared with time 0 (p<0.05) and this

increase did not continue until the end of sampling time. The down-regulation of IL-8 was much more obvious in the case of Iranian vaccine in all sampling points as compared to other groups (p<0.05). Although the higher level of this gene was mostly observed in control group, there was no significant change at 48 h between AquaVac and control (Fig. 9).




Figure6: The effects of different vaccines on head kidney IL-1B gene expression following challenge test with S. iniae at different time intervals. All values were obtained from 9 individual fish (3/replicate) and expressed as mean $\pm$ SD. Different lowercase alphabetic letters on each bar indicate significant difference among different treatments and different capital letters express significant difference among different sampling times (p<0.05).




Figure 7: The effects of different vaccines on head kidney IL-1B gene expression following challenge test with L. garviae at different time intervals. All values were obtained from 9 individual fish (3/replicate) and expressed as mean $\pm$ SD. Different lowercase alphabetic letters on each bar indicate significant difference among different treatments and different capital letters express significant difference among different sampling times (p<0.05).




Figure 8: The effects of different vaccines on head kidney IL-8 gene expression following challenge test with S. iniae at different time intervals. All values were obtained from 9 individual fish (3/replicate) and expressed as mean $\pm$ SD. Different lowercase alphabetic letters on each bar indicate significant difference among different treatments and different capital letters express significant difference among different sampling times (p<0.05).




Figure 9: The effects of different vaccines on head kidney IL-8 gene expression following challenge test with L. garviae at different time intervals. All values were obtained from 9 individual fish (3/replicate) and expressed as mean $\pm$ SD. Different lowercase alphabetic letters on each bar indicate significant difference among different treatments and different capital letters express significant difference among different sampling times (p<0.05).

## **Discussion**

In aquaculture, passive vaccines are routinely used to immunize fish against bacterial infections owing to several beneficial effects such as their costbenefit advantages (Vinitnantharat et al., 1999; Munang'andu and Evensen, 2019). Protective efficacy of vaccines is strongly related to their ability to stimulate the immune system with wide-ranging effects (Bercovier et al., 1997). The immunostimulatory effects of vaccines are thought to be related to their high antigenlevels (Leong et al., 1997). It. however. reflects the important status of vaccination against bacterial diseases in fish though little attempt was performed to understand the molecular mechanisms of vaccineinduced disease resistance or immunization (Harun et al., 2011). Therefore. identifying genes related to immune functions basically perceive the level of immune response following vaccination (Huang et al., 2014b).

In the current study, although the expression of TNF-alphain control group was slightly elevated on day 14, the increases were much higher in AquaVac and Iranian vaccines at the same sampling point, supporting our hypothesis that bivalent formalininactivated vaccinecan stimulate O. mykiss immune system after two weeks of vaccination. Similar trends were also observed for IL-1B and IL-8 gene expressions. These increases probably due to increased phagocytosis and chemotaxis activity and production of effective molecules, free radicals of oxygen and nitrogen (Ma, 2001). which, in turn, can help degradation of extracellular pathogens and therefore, increasing the non-specific defense mechanisms. In addition, an increase in pro-inflammatory response might be the results of up-regulation of IL-1B, antimicrobial activity of leukocytes, and induction of nitric oxide production by monocytes/ macrophages (Yang et al., 2013). TNF-alphaand IL-1ß genes are of induce inflammatory capable response by regulating the expression of other cytokines (Whyte, 2007). Similar increases in the expression of other cytokines, including IL-8 expression were reported in different studies in Paralichthys olivaceus with a peptide subunit vaccine (Sun and Hu, 2015), Cyprinus carpio in the early postvaccination period (Jiang et al., 2016) and in the liver of commercially vaccinated trout fed with ergosan immune stimulant after the introduction of antigen into the body (Gioacchini et al., 2008). IL-8 gene expression was found to increase significantly in Gadus during vaccination with morhua anguillarum Listonella (Caipang, 2013).

Nonetheless, the up-regulation immune gene no longer continued, i.e., following day 30 the immunization has been weakened, suggesting immunization is restricted to the history of vaccination. This might, however, manifest the ineffectiveness of the booster applied here since antigen absorption is a vital process activation of the immune system (Joosten et al., 1997). Furthermore, the lower level of immune-related gene on day 30 and afterward demonstrated the lack of effectiveness or even negligible effects for bath immersion, which has been done at the same day, i.e., day 30. Repeated injection of vaccine in O. mykiss could raise a secondary immune response whereas similar immersion vaccination against yersiniosis did act the same (Jaafar et al., 2018). Similarly, Akhlaghi et al. (1996) examined the passive immunization with mammalian antistreptococcal antibodies mykiss. They found only one month protection in this species by measuring the relative survival rate, confirming our findings that exhibited a peak in immune-related gene expression of head kidney after immunization. In support of the present finding, Soltani et al. (2007) reported higher antibody production in i.p immunized O. mykiss than fish immunized by immersion and oral routes. In another study, the effect of i.p. injection of formalin-inactivated vaccine on S. iniae infection in Epinephelus coioides was performed. It was observed that expression of IL-1β and TNF- alpha genes reached to the peak following 14 days of postimmunization (Huang et al., 2014a). Bastardo et al. (2012) reported high serum antibody level in O. mykiss immunized by two bivalent vaccines against L. garvieae and A. hydrophila 15 day after immunization. Moreover, the only initial up-regulation of IL-1β gene expression was reported in different organs of Salmo salar and Danio rerio(Haugland et al., 2005,

Zhang et al., 2012). This might however be the results of low antigen uptake through bath immersion that diminished the efficiency of vaccination procedure as compared to only one-step injection (Jaafar et al., 2018). Klesius et al. (2000) hypothesized that efficiency of vaccines following to different administration routes could be related to the antigenic composition of pathogen, especially in *S. iniae*.

In the next part of the study, we focused on the immune-related gene expression of post-challenged vaccinated fish. Our findings indicated that TNF- alpha, IL-1β and IL-8 expression in head kidney was up-regulated in all challenged vaccinated and unvaccinated control treatments in which the control was shown higher amount of those with more steady level over the measured times. Similarly but with lesser extent, increased levels could observed up to 72 h after L. garvieae and S. iniae challenge test in most treatments. Yang et al. (2016) reported that when fish were subjected to with Edwardsiella challenge test ictaluri, lower level of IL-1ß gene was observed as compared with control infected fish. Similarly, O. mykiss may also increase their immune-related gene expression, especially TNF- alpha and IL-1 $\beta$  in the spleen and gill vaccinated group after 6 challenging with Yersinia ruckeri. This increasing trend reached the maximum level following 24 and 48 h postinfection (Harun et al., 2011). However, this increase was not as strong as our observation in control infected fish. Reduction in the expression of TNFalpha and IL-1β genes were reported in the intestine of O. mykiss after with challenging Aeromonas salmonicida in comparison to noninfected fish (Mulder et al., 2007). Down-regulation of immune-related gene expression after E. tarda challenge was observed in vaccinated P. olivaceus when compared with control (Matsuyama et al., 2007). Similar to our findings, it is also known that infected catfish with E. ictaluri can increase the IL-8 gene expression of different organs while it was lower than that of the control group (Chen et al., 2005). In addition, IL-1\beta gene expression was upregulated following 6 and 12 h postchallenging with Edwarssiella tarda in Labeo rohita, but after 7 days it returned to its normal value, although its value was lower than that of the control group (Mohanty and Sahoo, 2010), confirming our findings. It seems that the lower expression of TNF- alpha, IL-1β, and IL-8 genes in the vaccine treatment in comparison with the control treatment is only subjected to experimental challenge with S. iniae and L. garvieae bacteria, which might probably manifest the effective role of the used vaccine in the reduction of the microbial load of kidney cells. This consequently induces a reduction in the anti-inflammatory responses in vaccine treatment, which ultimately provides better protection for vaccinated treatments (Ballesteros et al., 2015).

Other reports on the effects of viral and bacterial challenges in other infectious agents as well as species were seldom revealed no significant changes or even a slight increase when compared the vaccinated fish to control. The results of these studies are not consistent with the present study, which is probably because of the positive effects of the vaccine due to the presence of antigens to activate macrophages and white blood cells, resulting in the initiation of pro-inflammatory reactions, increased TNF- alpha, IL-1\beta and IL-8 cytokines in the body of fish. Ultimately, it increases the host's protection against S. iniae and L. garvieae bacteria. Thus, results in this kind of immunological responses are indeed different among species well experimental as procedure.

In summary, the selected immune genes in the present study revealed an initial up-regulation in both vaccinated groups with higher intensity in AquaVac treated fish. However, this suggests that the inflammatory gene response might be changed in line with the immunity response, nominating them as marker to prove the efficiency of vaccination in O. mykiss instead of conventional parameters. Although different vaccination methods, applied here. indicated significant gene upregulation, the i.p. injection is much efficient rather than bath immersion in case of inactivated S. iniae and L. garvieaevaccines. According to the obtained results, the higher amount of TNF-alpha, IL-1β and IL-8 gene

expressions in unvaccinated control fish was discerned with more stabilized level over the measured times. However, the conclusion cannot be generalized for either lower and/or higher ranges of vaccines in other species. More investigation needs to be addressed to elucidate the possible effects of other pathogenic bacteria in fish.

## Acknowledgement

We would like to thank the faculty and staff members of Department of Clinical Sciences of Shahid Chamran University of Ahvaz for their assistance.

## References

- Akhlaghi, M., Munday, B. and Whittington, R., 1996. Comparison of passive and active immunization of fish against streptococcosis (enterococcosis). *Journal of Fish Diseases*, 19, 251-258. DOI:10.1111/j.1365-2761.1996.tb00132.x
- Ballesteros, N.A., Saint-Jean, S.R. and Perez-Prieto, S.I., 2015.

  Immune responses to oral pcDNA-VP2 vaccine in relation to infectious pancreatic necrosis virus carrier state in rainbow trout *Oncorhynchus mykiss. Veterinary Immunology and Immunopathology*, 165, 127-137.DOI:10.1016/j.vetimm.2015.04.001
- Bastardo, A., Ravelo, C., Castro, N., Calheiros, J. and Romalde, J.L., 2012. Effectiveness of bivalent

vaccines against Aeromonas hydrophila and Lactococcus garvieae infections in rainbow trout *Oncorhynchus mykiss* (Walbaum). *Fish and Shellfish Immunology*, 32, 756-761.

DOI:10.1016/j.fsi.2012.01.028

- **Bercovier, H., Ghittino, C. and Eldar, A., 1997**. Immunization with bacterial antigens: infections with streptococci and related organisms. *Developments in Biological Standardization,* 90, 153-160.
- Bustin, S.A., Benes, V., Garson, J.A., Hellemans, Huggett, J., Kubista, M., Mueller, R., Nolan, T., Pfaffl, M.W. and Shipley, G.L., 2009. The **MIOE** guidelines: minimum information for publication of quantitative real-time **PCR** experiments. Clinical 55, 611-622. Chemistry, DOI:10.1373/clinchem.2008.112797
- Caipang, C.M.A., 2013. Expression of genes involved in the early immune response at the distal segment of the gut in Atlantic cod, *Gadus morhua* L. after vaccination with a bacterial antigen. *Aquaculture International*, 21, 591-603.DOI:10.1007/s10499-012-9591-8
- Chang, P., Lin, C. and Lee, Y., 2002. Lactococcus garvieae infection of cultured rainbow trout, *Oncorhynchus mykiss* in Taiwan and associated biophysical characteristics and histopathology. *Bulletin-European Association of Fish Pathologists*, 22, 319-327.

- Chen, S.C., Lin, Y.D., Liaw, L.L. andWang, P.C., 2001. Lactococcus garvieae infection in the giant freshwater prawn Macrobranchium rosenbergii confirmed bv polymerase chain reaction and 16S rDNA sequencing. Diseases Aquatic Organisms, 45, 45-52. DOI:10.3354/dao045045
- Chen, L., He, C., Baoprasertkul, P., Xu, P., Li, P., Serapion, J., Waldbieser, G., Wolters, W. andLiu, Z., 2005. Analysis of a catfish gene resembling interleukin-8: cDNA cloning, gene structure, and expression after infection with Edwardsiella ictaluri. *Developmental and Comparative Immunology*, 29, 135-142.

DOI:10.1016/j.dci.2004.06.011

- Didinen, B., Yardimci, B., Onuk, E., Metin, S. and Yildirim, P., 2014. **Naturally** Lactococcus garvieae infection in rainbow trout (Oncorhyncus mykiss Walbaum, 1792): new histopathological observations, phenotypic and molecular identification. Revue de Medecine Veterinaire, 165, 12-19.
- Diler, O., Altun, S., Adiloglu, A., Kubilay, A. and Istklt, B., 2002. First occurrence of streptococcosis affecting farmed rainbow trout (Oncorhynchus mykiss) in Turkey. Bulletin-European Association of Fish Pathologists, 22, 21-26.
- Dimitroglou, A., Merrifield, D.L., Carnevali, O., Picchietti, S., Avella, M., Daniels, C., Güroy, D. andDavies, S.J., 2011. Microbial

- manipulations to improve fish health and production—a Mediterranean perspective. *Fish and Shellfish Immunology*, 30, 1-16. DOI:10.1016/j.fsi.2010.08.009
- Eldar, A., Horovitcz, A. and Bercovier, H., 1997. Development and efficacy of a vaccine against Streptococcus iniae infection in farmed rainbow trout. *Veterinary Immunology and Immunopathology*, 56, 175-183. DOI:10.1016/S0165-2427(96)05738-8
- Eldar, A.A. and Ghittino, C., 1999. Lactococcus garvieae and Streptococcus iniae infections in rainbow trout Oncorhynchus mykiss: similar, but different diseases. Diseases of Aquatic Organisms, 36, 227-231. DOI:10.3354/dao036227
- Eldar, A., Perl, S., Frelier, P. and Bercovier, H., 1999. Red drum Sciaenops ocellatus mortalities associated with Streptococcus iniae infection. Diseases of Aquatic Organisms, 36, 121-127. DOI::10.3354/dao036121
- Elliott, J., Collins, M., Pigott, N. and Facklam, R., 1991. Differentiation Lactococcus lactis of and Lactococcus garvieae from humans by comparison of whole-cell protein ofpatterns. **Journal** Clinical Microbiology, 29, 2731-2734. DOI:10.1128/JCM.29.12.2731-2734.1991
- Evans, J.J., Pasnik, D.J., Klesius, P.H. and Al-Ablani, S., 2006. First report of *Streptococcus agalactiae* and Lactococcus garvieae from a

- wild bottlenose dolphin (*Tursiops truncatus*). *Journal of Wildlife Diseases*, 42, 561-569. DOI:10.7589/0090-3558-42.3.561
- Faber, M.N., Holland, J.W. and Secombes, C.J., 2019. Vaccination strategies and IgM responses against PKD in rainbow trout. Fish and Shellfish Immunology, 91, 423. DOI:10.1016/j.fsi.2019.04.159
- Gioacchini. G., Smith. P. and Carnevali, O., 2008. Effects of Ergosan on the expression cytokine genes in the liver of juvenile rainbow trout (Oncorhynchus mykiss) exposed to enteric red mouth vaccine. Veterinary *Immunology* and Immunopathology, 123, 215-222. DOI:10.1016/j.vetimm.2008.01.037
- Harikrishnan, R., Balasundaram, C. and Heo, M.S., 2010. Effect of probiotics enriched diet on Paralichthys olivaceus infected with lymphocystis disease virus (LCDV). Fish and Shellfish Immunology, 29, 868-874.

DOI:10.1016/j.fsi.2010.07.031

- N.O., Wang, T. Harun, and **C.J.**, 2011. Secombes, Gene expression profiling in naïve and rainbow vaccinated trout after Yersinia ruckeri infection: insights into the mechanisms of protection seen in vaccinated fish. Vaccine, 29, 4388-4399.
  - https://doi.org/10.1016/j.vaccine.201 1.04.003
- Haugland, Ø., Torgersen, J., Syed, M. and Evensen, Ø., 2005. Expression

profiles of inflammatory and immune-related genes in Atlantic salmon (*Salmo salar* L.) at early time post vaccination. *Vaccine*, 23, 5488-5499.

DOI:10.1016/j.vaccine.2005.07.034

- Huang, H.Y., Chen, Y.C., Wang, P.C., Tsai, M.A., Yeh, S.C., Liang, H.J. and Chen, S.C., 2014a. Efficacy of a formalin-inactivated vaccine against Streptococcus iniae infection in the farmed grouper **Epinephelus** coioides bv intraperitoneal immunization. 32. 7014-7020. Vaccine, DOI:10.1016/j.vaccine.2014.08.039
- Huang, L., Wang, K., Xiao, D., Chen,
  D., Geng, Y., Wang, J., He, Y.,
  Wang, E., Huang, J. and Xiao, G.,
  2014b. Safety and immunogenicity of an oral DNA vaccine encoding Sip of Streptococcus agalactiae from Nile tilapia Oreochromis niloticus delivered by live attenuated Salmonella typhimurium. Fish and Shellfish Immunology, 38, 34-41.
  DOI:10.1016/j.fsi.2014.02.017
- Jaafar, R., Al-Jubury, A., Chettri, J.K., Dalsgaard, I., Kania, P. and Buchmann, K., 2018. Secondary immune response of rainbow trout following repeated immersion vaccination. *Journal of Fish Diseases*, 41, 117-123 DOI:10.1111/jfd.12682
- Jiang, X., Zhang, C., Zhao, Y., Kong,X., Pei, C., Li, L., Nie, G. and Li,X., 2016. Immune effects of the vaccine of live attenuatedAeromonas hydrophila screened by

.108

rifampicin on common carp (*Cyprinus carpio* L). *Vaccine*, 34, 3087-3092.

DOI:10.1016/j.vaccine.2016.04.075

- Joosten, P., Tiemersma, E., Threels, A., Caumartin-Dhieux, C. and Rombout, J., 1997. Oral vaccination of fish againstVibrio anguillarumusing alginate microparticles. Fish & Shellfish Immunology, 7, 471-485. DOI:10.1006/fsim.1997.0100
- Klesius, P.H., Shoemaker, C.A. and Evans, J.J., 2000. Efficacy of single and combined Streptococcusiniae vaccine administered intraperitoneal and intramuscular tilapia routes in (Oreochromis niloticus). Aquaculture, 188(3-4). 237-246. DOI:10.1016/S0044-8486(00)00345-8
- Leong, J., Anderson, E., Bootland, L., Chiou, P., Johnson, M., Kim, C., Mourich, D. and Trobridge, G., **1997**. Fish vaccine antigens produced or delivered by recombinant **DNA** technologies. **Developments** in Biological Standardization, 90, 267-277.
- Ma, X., 2001. TNF-α and IL-12: a balancing act in macrophage functioning. *Microbes and Infection*,
  3, 121-129. DOI:10.1016/S1286-4579(00)01359-9
- Marti, E., Huerta, B., Rodríguez-Mozaz, S., Barceló, D., Marcé, R. and Balcázar, J.L., 2018. Abundance of antibiotic resistance genes and bacterial community composition in wild freshwater fish

- species. *Chemosphere*, 196, 115-119.
  DOI:10.1016/j.chemosphere.2017.12
- T.. Fujiwara, Matsuyama, A., Kamaishi, Nakayasu, **C...** T., Oseko, N., Hirono, I. and Aoki, T., 2007. Gene expression of leucocytes in vaccinated Japanese flounder (Paralichthys olivaceus) during the course of experimental infection with Edwardsiella tarda. Fish and Shellfish Immunology, 22, 598-607. DOI:10.1016/j.fsi.2006.08.006
- Mohammadian, T., Alishahi, Tabandeh, M.R., Ghorbanpoor, M. and Gharibi, D., 2018. Changes in Immunity, Expression of some Immune-Related Genes of Shabot Fish. Tor grypus, **Following Experimental** Infection with Aeromonas hydrophila: Effects of Autochthonous Probiotics. Antimicrobial **Probiotics** and Proteins, 10, 616-628. DOI: 10.1007/s12602-017-9373-8
- Mohammadian, T., Nasirpour, M., Tabandeh, M.R. and Mesbah, M., **2019.** Symbiotic effects of  $\beta$ -glucan, oligosaccharide mannan and Lactobacillus casei growth on performance, enzymes intestine activities, immune-hematological parameters and immune-related gene expression in common carp, Cyprinus carpio: An experimental infection with Aeromonashydrophila. Aquaculture. х, X-X.

- DOI:10.1016/j.aquaculture.2019.06.
- Mohammadian, T., Nasirpour, M., Tabandeh, M.R., Heidary, A.A., Ghanei-Motlagh, R. and Hosseini, 2019. Administrations **S.S.**. probiotics autochthonous altered rainbow iuvenile trout Oncorhynchus mykiss health status, growth performance and resistance Lactococcus garvieae. experimental infection. Fish and Shellfish Immunology, 86, 269-279. DOI:10.1016/j.fsi.2018.11.052
- Mohanty, B. and Sahoo, P., 2010. Immune responses and expression profiles of some immune-related genes in Indian major carp, Labeo rohita to Edwardsiella tarda Fish infection. and Shellfish Immunology, 28, 613-621. DOI:10.1016/j.fsi.2009.12.025
- Mulder, I., Wadsworth, S. C., **2007**. Cytokine Secombes, intestine of expression in the rainbow trout (Oncorhynchus mvkiss) during infection Aeromonas salmonicida, Fish and Shellfish Immunology, 23, 747-759. DOI:10.1016/j.fsi.2007.02.002
- Nakai, T., Sugimoto, R., Park, K.H., Matsuoka, S., Mori, K.I., Nishioka, T. and Maruyama, K., 1999.

  Protective effects of bacteriophage on experimental *Lactococcus garvieae* infection in yellowtail. *Diseases of Aquatic Organisms*, 37, 33-41. DOI:10.3354/dao037033
- Novotny, L., Dvorska, L., Lorencova, A., Beran, V. and Pavlik, I., 2004.

- Fish: a potential source of bacterial pathogens for human beings. A review. Veterinarni Medicina-UZPI (Czech Republic). DOI:10.17221/5715-VETMED
- Ooyama, T., Kera, A., Okada, T., Inglis, V. and Yoshida, T., 1999. The protective immune response of yellowtail *Seriola quinqueradiata* to the bacterial fish pathogen *Lactococcus garvieae*. *Diseases of Aquatic Organisms*, 37, 121-126. DOI:10.3354/dao037121
- Park, Y.K., Nho, S.W., Shin, G.W., Park, S.B., Jang, H.B., Cha, I.S., Ha, M.A., Kim, Y.R., Dalvi, R.S. and Kang, B.J., 2009. Antibiotic susceptibility and resistance of *Streptococcus iniae* and *Streptococcus parauberis* isolated from olive flounder (Paralichthys olivaceus). *Veterinary Microbiology*, 136, 76-81.
  - DOI:10.1016/j.vetmic.2008.10.002
- Purcell, M.K., Kurath, G., Garver, K. A., Herwig, R.P. and Winton, J.R., 2004. **Ouantitative** expression profiling of immune response genes in rainbow trout following infectious haematopoietic necrosis virus infection **DNA** (IHNV) or Fish vaccination. and Shellfish Immunology, 17. 447-462. DOI:10.1016/j.fsi.2004.04.017
- Raissy, M., Hashemi, S., Roushan, M., Jafarian, M., Momtaz, H., Soltani, M. and Pirali Kheirabad, E., 2018. Effects of essential oils of Satureja bachtiarica and Nigella sativa on the efficacy of

- lactococcosis vaccine in rainbow trout (*Oncorhynchu smykiss*). *Iranian Journal of Fisheries Sciences*, 17(1), 95-106. DOI:10.22092/IJFS.2018.115587
- Sakai, M., Atsuta, S. and Kobayashi, M., 1995. Efficacies of combined vaccine for *Vibrio anguillarum* and *Streptococcus* sp. *Fisheries Science*, 61, 359-360. DOI:10.2331/fishsci.61.359
- Secombes, C., Zou, J., Laing, K., Daniels, G. and Cunningham, C., 1998. Cytokine genes in fish. *Aquaculture*, 171, 93-102. DOI:10.1016/S0044-8486(98)00441-4
- Secombes, C., Wang, T., and Bird, S., 2011. The interleukins of fish. Developmental and Comparative Immunology, 35, 1336–1345. https://doi.org/10.1016/j.dci.2011.05.001
- Shiry, N., Soltanian, S., Shomali, T., Paknejad, H.and Hoseinifar, S.H., **2019.** Immunomodulatory effects of orally administrated florfenicol in rainbow trout (Oncorhynchus mykiss) following experimental challenge with streptococcosis/lactococcosis. International Immunopharmacology, 73, 236-245. DOI:10.1016/j.intimp.2019.05.007
- Shoemaker, C.A., Klesius, P.H. and Evans, J.J., 2001. Prevalence of Streptococcus iniae in tilapia, hybrid striped bass, and channel catfish on commercial fish farms in the United States. *American Journal of*

- *Veterinary Research*, 62, 174-177. DOI:10.2460/ajvr.2001.62.174
- Soltani, M., Alishahi, M., Mirzargar, and Nikbakht, G., 2007. Vaccination of rainbow trout against infection: Streptococcus iniae comparison of different routes of administration and different vaccines. Iranian **Journal** of Fisheries Sciences, 7(1), 129-140. http://hdl.handle.net/1834/11186
- M., Soltani, Kane, A., Taheri-Mirghaed, A., Pakzad, K. and Hosseini-Shekarabi, P., Effect of the probiotic, Lactobacillus plantarum on growth performance haematological indices rainbow trout (Oncorhynchus mykiss) immunized with bivalent streptococcosis/lactococcosis vaccine. Iranian Journal of Fisheries *Sciences*, 18(2), 283-295. 10.22092/ijfs.2018.117757
- Sun, Y., Yang, H., Ma, R., Song, K. and Li, J., 2011. Effect of Lactococcuslactis and Enterococcus faecium on growth performance, digestive enzymes and immune response of grouper *Epinephelus coioides*. Aquaculture Nutrition, 165, 1-9. DOI:10.1111/j.1365-2095.2011.00894.x
- Sun, Y., Zhang, M., Liu, C.-S., Qiu, R. and Sun, L., 2012. A divalent DNA vaccine based on Sia10 and OmpU induces cross protection against *Streptococcus iniae* and Vibrio anguillarum in Japanese flounder. *Fish and Shellfish*

- *Immunology,* 32, 1216-1222. DOI:10.1016/j.fsi.2012.03.024
- Sun, Y. and Hu, Y.H., 2015. Cell-penetrating peptide-mediated subunit vaccine generates a potent immune response and protection against Streptococcus iniae in Japanese flounder (*Paralichthys olivaceus*). Veterinary Immunology and Immunopathology, 167, 96-103. DOI:10.1016/j.vetimm.2015.07.008
- Vendrell, D., Balcázar, J.L., Ruiz-Zarzuela, I., De Blas, I., Gironés, O. and Múzquiz, J., L. 2006.

  Lactococcus garvieae in fish: a review. Comparative Immunology, Microbiology and Infectious Diseases, 29, 177-198.

  DOI:10.1016/j.cimid.2006.06.003
- Vendrell, D., Balcázar, J.L., De Blas, I., Ruiz-Zarzuela, I., Gironés, O. and Múzquiz, J.L., 2008. Protection of rainbow trout (Oncorhynchus mykiss) from lactococcosis by probiotic bacteria. Comparative Immunology, Microbiology and Infectious Diseases, 31, 337-345. DOI:10.1016/j.cimid.2007.04.002
- Vinitnantharat, S., Gravningen, K. And Greger, E., 1999. Fish vaccines. Advances in Veterinary Medicine, 41, 539-550. DOI: 10.1016/s0065-3519(99)80040-8
- Wang, C.Y.C., Shie, H. S., Chen, S.C., Huang, J.P., Hsieh, I.C., Wen, M.S., Lin, F.C. and Wu, D., 2007. Lactococcus garvieae

- infections in humans: possible association with aquaculture outbreaks. *International Journal of Clinical Practice*, 61, 68-73. DOI:10.1111/j.1742-1241.2006.00855.x
- Whyte, S.K., 2007. The innate immune response of finfish—a review of current knowledge. *Fish and Shellfish Immunology*, 23, 1127-1151. DOI:10.1111/j.1742-1241.2006.00855.x
- Yang, K., Zhang, S., Chen, D., Zhang, A., Wang, X. and Zhou, H., 2013. IFN-γ-activated lymphocytes boost nitric oxide production in grass carp monocytes/macrophages. *Fish and Shellfish Immunology*, 35, 1635-1641. DOI:10.1016/j.fsi.2013.09.017
- Yang, Q., Pan, Y.L., Wang, K.Y., Wang, J., He, Y., Wang, E.L., Liu, T., Geng, Y., Chen, D.F. and Huang, X.L., 2016. OmpN, outer membrane proteins of Edwardsiella ictaluri are potential vaccine candidates for channel catfish (Ictalurus punctatus). Molecular Immunology, 78. 1-8. DOI:10.1016/j.molimm.2016.08.011
- Zhang, Z., Wu, H., Xiao, J., Wang, Q., Liu, Q. and Zhang, Y., 2012. Immune responses of zebrafish (Danio rerio) induced by bathvaccination with a live attenuated Vibrio anguillarum vaccine Fish candidate. and Shellfish 33, Immunology, 36-41. DOI:10.1016/j.fsi.2012.03.031