Effect of the probiotic, Lactobacillus plantarum on growth performance and haematological indices of rainbow trout (Oncorhynchus mykiss) immunized with bivalent streptococcosis / lactococcosis vaccine

Soltani M.^{1,2}*; Kane A.¹; Taheri-Mirghaed A.¹; Pakzad K.¹; Hosseini-Shekarabi P.³

Accepted: May 2017 Received: March 2017

Abstract

The present study evaluated the effect of Lactobacillus plantarum as a probiotic on the growth performance and haematological parameters of juvenile rainbow trout (Oncorhynchus weighing 29.6±1.8 immunized with mykiss) streptococcus/lactococcus bivalent vaccine for 60 days at 16±1.5°C. Fish were randomly allocated to 4 equal groups: probiotic (g kg⁻¹ feed equal to 10⁸ cells g⁻¹) supplemented diet group (group P), vaccinated fish fed with a normal diet (group V), vaccinated fish fed with probiotic (group P+V) and unvaccinated fish fed normal diet (group C). Results showed that weight gain, final weight, condition factor and thermalunit growth coefficient were improved in P+V group compared with other groups but was significantly different from groups V and C (p<0.05). Also, feed conversion ratio in the P+V group was lower than other trails (p<0.05). In addition, white blood cell count (WBC) in groups V and P+V were significantly higher than in both P and C groups (p<0.05). Also, PCV was significantly increased (p<0.05) in both P and P+V groups compared to group V and the control group. No significant difference (p>0.05)was found in RBC, Hb, MCH, MCHC and MCV levels among the different experimental groups. These data show that application of L. plantarum could be a benefit of vaccinated trout, enhancing fish growth and haematological parameters.

Keywords: Lactobacillus plantarum, probiotic, trout, haematology, growth, vaccine, streptococcosis/lactococcosis

¹⁻Department of Aquatic Animal Health, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

²⁻School of Veterinary and Life Sciences, Murdoch unviersity, WA, Australia

³⁻Departmet of Fisheries Science, Faculty of Agriculture and Natural Resources, Science and Research Branch, Islamic Azad University, Tehran, Iran.

^{*}Corresponding author's Email: msoltani@ut.ac.ir

Introduction

Aquaculture is an important food producing sector, but is faced with significant constraints due to several infectious diseases including streptococcosis and lactococcosis (Austin and Austin, 2007; Bostock et al., 2010). These bacterial infections are widely spread in many regions, including Iranian trout (O. mykiss) farms causing huge economic losses (Soltani et al., 2005; Soltani et al., 2008; Erfanmanesh et al., 2012). To date, control of these diseases has been focused upon the use of antibiotics, and more recently on vaccinations (Evans et al., 2004; Soltani et al., 2007; Sun et al., 2010). Application of antibiotics is the common way to treat bacterial infections in aquaculture however, this leads to overuse and gives rise to resistant strains (Karunasagar et al., 1994; Akinbowale et al., 2006). A local vaccine is currently available to protect Iranian rainbow trout farms with encouraging results so far.

Besides vaccination, the use of probiotics, in preference to antibiotics, is increasingly gaining acceptance for the control of pathogens in aquaculture (Sheehan *et al.*, 2009; Raissy *et al.*, 2018). Probiotics have been associated with many useful effects on growth performance (Giannenas *et al.*, 2015; Sivani *et al.*, 2016), disease resistance by enhancing innate immunity (Irianto and Austin, 2002; Cha *et al.*, 2013; Safari *et al.*, 2016), reduction in pathogenic microbiotia of fish gut (Abid *et al.*, 2013), and with a consequence in a less chemotherapy

(Irianto and Austin, 2002; Azad and Al-Marzouk, 2008).

Several studies have shown lactic (LAB) acid bacteria such especially Lactobacillus plantarum, from the gastrointestinal tract of fish, to beneficial for their probiotic properties in aquaculture (Ringo and Gatesoupe, 1998; Ali and Sahu, 2002; Balcázar et al., 2008; Harikrishnan et al., 2010; Giri et al., 2013).

Application of L. plantarum probiotic illustrated to increase the disease resistance, non-specific immune defence and growth performance in fish (Son et al., 2009; Giri et al., 2014; Dawood et al., 2015). However, only a few data available on the effect of probiotics on growth and immunephysilogy of vaccinated fish. Tilapia vaccinated with an A. hydrophila bacterin and fed Lactobacillus at various concentrations significantly activated neutrophils, and lymphocytes compared to vaccine only (Venkatalakshmi & Ebanasar 2015). Oral application of the mixture of Lactobacillus acidophilus, Bacillus subtilis, Saccharomyces cervisiae and Aspergillus oryzae to tilapia vaccinated against A. hydrophila significantly increased survival from the challenge infection (Alv et al. 2016). Oral use of L. plantarum in trout bath-vaccinated against Yersinia ruckeri elevated some immunological responses and growth of vaccinated fish (Soltani et al. 2017a, b). The present study aimed to explore the effects of L. plantarum as probiotic on haematological parameters and growth performance in trout immunized with streptococcus/lactococcosis bivalent vaccine.

Materials and methods

Probioite

L. plantarum (kc426951) previously isolated from the intestinal tract of rainbow trout was used for this study. The probiotic cell suspension was added to commercial fish diet (Faradaneh, Shahrekord, Iran) at a concentration of 1g (10⁸cells g⁻¹) kg⁻¹ feed.

Fish culture and experimental design Juvenile rainbow trout (*O. mykiss*) weighing 29.6±1.8 g obtained from a local hatchery in Arak province, Iran were allowed to acclimatize for 14 days in a 5000-L tank with a flow-through system (7 L min⁻¹), and were fed a

commercial diet (Faradaneh, Shahrekord, Iran). Fish were then randomly distributed into twelve 1000-L tanks with 80 fish per tank, four treatments each in three replicates.

Four groups were included: fish fed with the probiotic at g kg⁻¹ feed equal to 10⁸ cells g⁻¹ (group P), vaccinated fish fed with the probiotic (group P+V), vaccinated fish fed with a normal diet (group V) and unvaccinated fish fed normal diet (group C). Bivalent vaccine against streptococcosis/lactococcosis (formalin inactivated cells at 1x10⁹ cells ml⁻¹ of each Streptococcus iniae and Lactococcus garviae strains) was used. Vaccination was preformed using the bath method for 1 min under good aeration. The protocol of experimental groups is given in Table 1.

Table 1: Protocol of experimental groups studied in the present study.

Immunization regime	Feeding regime	Feeding rate
Vaccinated fish	Probiotic (1g kg ⁻¹ food)	2%
Vaccinated fish	Normal diet without probiotic	2%
Unvaccinated fish	Probiotic (1g kg ⁻¹ food)	2%
Unvaccinated fish	Normal diet without probiotic	2%

During the feeding trial, the fish were fed three times a day (09:00, 12:00 and 15:00 hours) at the rate of 2% of body weight. The feeding rate was corrected every 12 days. The water quality parameters including pH, dissolved oxygen, water temperature, NO₂ and NH₃ were monitored daily and maintained at 7.5±0.5, 8±0.4 mg L⁻¹, 16±1.5 °C, <0.1 mg L⁻¹ and 0.01 mg L⁻¹

¹, respectively during the 60 days of the experimental period.

Growth performance

At the end of the experiment the weight and size of individual fish (n=9 per replicate) in the tanks of each treatment group (n=27 per group) were measured, and feed utilization and growth

parameters were calculated using the following formulae:

Condition factor=weight/(length)³×100 Feed conversion ratio (FCR)=food consumption (g)/weight gain (g)

Weight gain (g)=final weight-initial weight

Thermal-unit growth coefficient (TGC) = $(W_2(^{1/3})-W_1(^{1/3}))/(T\times\Delta t)\times 1000$

Where W_1 is the group average initial weight (g), W_2 is the group average weight at time t (g), T is the water temperature (${}^{\circ}C$) and Δt is the duration of the experiment (days).

Haematological assay

On days 12, 24, 36, 48 and 60 of the experiment, nine fish from each group were anesthetized with clove oil (200 mg L⁻¹), and blood was collected from caudal vein. Haematological parameters including white blood cells erythrocyte count (RBC), (WBCs), cell volume packed (PCV), haemoglobin (Hb), mean erythrocyte haemoglobin (MCH), mean erythrocyte haemoglobin concentration (MCHC) and mean erythrocyte cell volumes (MCV) were measured (Svobodova et al., 1991).

Statistical analysis

The mean and SEM were determined, and the data were analysed using one-way ANOVA using SPSS package (SPSS 1998) at the 0.05 significance level.

Results

Growth performance

The results of feed utilization and growth performance parameters are given in Table 2. All groups showed above 97 % survival rates (p<0.05). The results revealed that growth performance parameters including weight gain, final weight, TGC and significantly FCR were (p < 0.05)improved in all treatment groups compared to the control, and the highest values of final weight (156.24±2.29 g), weight gain (126.74±2.29 g) and TGC (0.201 ± 0.003) plus the lowest level of FCR (0.96 ± 0.02) were obtained in Group P+V. In addition, Group P+V significantly enhanced condition factor compared to the other groups (1.1 ± 0.05) (p<0.05).

Haematological parameters

Results of WBC are given in Fig. 1. On days 12, 24, 36, 48 and 60, WBC levels in groups V and P+V, were significantly higher than in the other two groups. However, no significant change was found in group P (Fig. 1).

Also, results of RBC are shown in Fig. 2. On days 12 and 48, RBC level in group P+V was significantly higher than C group (Control) (p<0.05), (Fig. 2). No significant differences were seen among other trails during the experiment (p>0.05).

Table 2: Mean ± SD of growth performance and feed utilization parameters in trout immunized
with streptococcosis/lactococosis vaccine and fed Lactobacillus plantarum for 2 months at
16+1.5 °C.

Growth factor	C (control)	P	V	V+ P
Initial weight (g)	29.5 ± 1.98	29.54±1.86	29.88±1.81	29.50±1.76
Final weight (g)	140.22±2.91 ^a	152.83±2.59 ^{bc}	148.47 ± 2.67^{b}	156.24±2.29°
Weight gain (g)	110.72±2.91 ^a	123.29±2.59 ^{bc}	118.59±2.67 ^b	126.74±2.29 °
Feed (g fish ⁻¹)	127.67±7.02	120±5.29	123.17±3.88	121.33±5.13
FCR	1.15 ± 0.03^{a}	$0.97\pm0.03^{\text{ by}}$	1.04 ± 0.03^{b}	0.96 ± 0.02^{c}
Initial size (cm)	14.50 ± 0.22	14.40 ± 0.2	14.55±0.28	14.40 ± 0.23
Final size (cm)	23.99±0.18 a	24.13 ± 0.34^{a}	24.05 ± 0.45^{a}	24.23±0.48 a
Condition factor	1.02±0.02 ^a	1.09 ± 0.03^{ab}	$1.07{\pm}0.05^{\ ab}$	1.1±0.05 b
TGC	0.185 ± 0.003^a	0.198 ± 0.003^{bc}	0.192 ± 0.003^{b}	0.201 ± 0.003^{c}

FCR: Feed conversion ratio, TGC: Thermal-unit growth coefficient, P: fish fed with probiotic (L. plantarum), V: fish immunized with streptococcosis/lactococosis vaccine, P+V: fish fed with probiotic (L. plantarum) and immunized with streptococcosis/lactococosis vaccine, Control: fish fed with normal diet. Values (n=27) within the same row with different superscripts denotes significant differences (p<0.05).




Figure 1: White blood cells (WBC) in vaccinated rainbow trout with streptococcosis / lactococcosis vaccine and fed with probiotic (*Lactobacillus plantarum*) for 2 months at 16±1.5°C. P: unvaccinated fish fed with the probiotic (*L. plantarum*), V: fish immunized with streptococcosis / lactococcosis vaccine and fed with a normal diet, P+V: fish immunized with streptococcosis / lactococcosis vaccine and fed with the probiotic supplemented in diet, Control: unvaccinated fish fed with normal diet. Values (n=27) within the same row with different superscripts denotes significant differences (p<0.05).




Figure 2: Red blood cells (RBC) in vaccinated rainbow trout with streptococcosis / lactococcosis vaccine and fed with probiotic (*Lactobacillus plantarum*) for 2 months at $16\pm1.5^{\circ}$ C. P: unvaccinated fish fed with the probiotic (*L. plantarum*), V: fish immunized with streptococcosis / lactococcosis vaccine and fed with a normal diet, P+V: fish immunized with streptococcosis / lactococcosis vaccine and fed with the probiotic supplemented in diet, Control: unvaccinated fish fed with normal diet. Values (n=27) within the same row with different superscripts denotes significant differences (p<0.05).

In addition, group P+V showed a significant increase in Hb level at 36 days (p<0.05). Results also showed that fish in groups P and P+V had significantly higher levels of PCV on

days 24 and 48 (p<0.05). No significant differences were found (p>0.05) in MCH, MCHC and MCV levels among the trails during 2 month experiment (Table 2).

Table 3: The mean±SD of derived hematological parameters in trout immunized with streptococcus / lactococosis vaccine and fed a probiotic (*Lactobacillus plantarum*) for 2 months at16±1.5 °C.

months atto±1.5°C.							
Treatments	Hb (g dL ⁻¹)	PCV (%)	MCHC (g dL ⁻¹)	MCH (pg)	MCV (fl)		
12 days							
Control	5.69 ± 0.3^{a}	32±1 ^a	17.78 ± 0.4^{a}	54.75±2.46 ^a	307.79 ± 8.16^{a}		
P	5.86 ± 0.45^{a}	33.33±1.53 ^a	17.56 ± 056^{a}	54.58±2.92 ^a	310.67 ± 9.29^{a}		
V	5.38 ± 0.6^{a}	32.67 ± 3.06^{a}	16.46 ± 0.38^{a}	56.48 ± 5.22^{a}	343.15±31.74 ^a		
V+P	6.45 ± 0.84^{a}	36±3°	17.87 ± 0.87^{a}	56.31±4.38 ^a	314.79±9.97 ^a		
24 days							
Control	5.19±0.15 ^a	$31.33{\pm}0.58^a$	16.57±0.39 ^a	50.66±2.66 ^a	305.61±11.43 ^a		
P	5.75 ± 0.4^{a}	35.33 ± 2.08^{b}	16.28±0.66 ^a	52±5.46 ^a	318.81±21.9 ^a		
V	5.84 ± 0.32^{a}	33.33±0.58 ^a	17.51±0.66 ^a	54.31±4.21 ^a	309.79±12.46 ^a		
V + P	6.2±0.32 ^a	36 ± 1^b	17.21±0.75 ^a	52.82±1.75 ^a	306.94±5.09 ^a		
36 days							
Control	6.99 ± 0.17^{ab}	33 ± 2^{a}	21.24 ± 1.8^{a}	60.57 ± 6.36^{a}	285.66 ± 6.82^{a}		
P	7.09 ± 0.38^{ab}	33.67 ± 2.52^{a}	21.16 ± 2.08^{a}	68.34 ± 5.88^{a}	321.95 ± 16.02^{a}		
V	6.5 ± 0.21^{a}	34.67 ± 1.53^{a}	19.36±1.51 ^a	64.37±5.42 ^a	323.47±10.25 ^a		
V+P	7.31 ± 0.35^{b}	32.33 ± 2.08^{a}	22.64±0.71 ^a	70.21 ± 6.46^{a}	318.6±19.7 ^a		
48 days							
Control	7.25 ± 0.46^{a}	36.33 ± 1.53^{a}	19.95±0.46 ^a	62.14±1.44 ^a	311.48±5.58 ^a		
P	8.06 ± 0.34^{a}	42.67 ± 1.53^{b}	18.92 ± 1.13^{a}	65.66±2.11 ^a	348.25±30.75 ^a		
V	7.1 ± 0.64^{a}	34.67 ± 2.08^{a}	20.45 ± 0.64^{a}	69.45±1.57 ^a	339.91 ± 15.87^{a}		
V + P	8.42±0.95 ^a	43 ± 1^b	18.73±2.24 ^a	62.91±5.12 ^a	337.01±17 ^a		
60 days							
Control	7.62 ± 0.25^{a}	41.33 ± 1.53^{a}	18.47 ± 1.19^{a}	60.87±5.29 ^a	329.23±7.72 ^a		
P	7.78 ± 0.24^{a}	43.33 ± 1.53^{a}	17.96±1.19 ^a	60.9 ± 4.24^{a}	339.14±23.06 ^a		
V	7.34 ± 0.37^{a}	43.33 ± 1.53^{a}	17.07 ± 0.49^{a}	55.28±3.16 ^a	324.21 ± 25.35^{a}		
V + P	8.02 ± 0.42^{a}	43.57 ± 1.53^a	18.35 ± 0.36^{a}	60.74 ± 2.62^{a}	331.18±19.66 ^a		

P: unvaccinated fish fed with the probiotic (*L. plantarum*), V: fish immunized with streptococcosis /lactococcosis vaccine and fed a normal diet, P+V: fish immunized with streptococcosis/lactococcosis vaccine and fed with the probiotic supplemented in diet, Control: unvaccinated fish fed a normal diet. Values (n=27) within the same row with different superscripts denotes significant differences (p<0.05). Values (n=27) within the same row with different superscripts denotes significant differences (p<0.05).

Discussion

In vertebrates, it has been shown that maintaining and improving an active immune system can be energetically too costly as it is necessary to adjustment physiological activities (Fair *et al.*, 1999; Pilorz *et al.*, 2005). It has been

shown that increases in immune response cause adverse effects on biological functions, possibly through an increase in metabolic rate (Pilorz *et al.*, 2005). Moreover, Ackerman *et al.* (2000) reported that administration of a monovalent, adjuvanted *Aeromonas*

salmonicida vaccine increases the metabolic rate of iuvenile trout. Considering the negative correlation between growth performance metabolic rate, probiotics may help to compensate this deficiency vaccinated fish. While there have been some individual works investigating the effects of vaccination and probiotic diets on the immune response and growth performance of fish, our study evaluated the dietary administration of probiotic L. plantarum alone concurrently with streptococcosis/lactococcosis vaccine haematological growth performance of rainbow trout. results of the present study showed that the streptococcus/lactococcosis vaccine and dietary supplementation with L. plantarum $(10^8 \text{ CFU g}^{-1})$ improved the feed utilization and growth performance parameters in trout. This might be in part due to an attribution of the increased digestibility coefficient of the diet (Dawood et al., 2015). Also, modulating the digestive physiology of fish by enhancing intestinal enzymes activity can improve the growth of fish fed a probiotic as mentioned by several workers (Yanbo and Zirong, 2006; Lazado et al., 2012; Sun et al., 2012). For example, use of diets supplemented with L. plantarum improved feed efficiency and growth in different fish species (Son et al., 2009; Jatobá et al., 2011; Parthasarathy and Ravi, 2011; Giri et al., 2013). In the present study the growth performance including weight gain, final weight, and TGC significantly (p < 0.05)were improved in all treated groups

compared with the control. However, the highest values of final weight (156.24 ± 2.29) weight g), gain (126.74 ± 2.29) and **TGC** g) (0.201 ± 0.003) were obtained in vaccinated fish fed with the probiotic (group V+P). Also, the lowest level of FCR (0.96 ± 0.02) were seen in this group. The reason why the application of the probiotic can provide such better growth conditions in vaccinated fish is not clear and warrants further studies. However, it may in part be due to the enchantment of some non-specific factors, e.g. lysozyme, complement component, phagocytosis vaccinated fish that can provide a better condition for fish physiology challenge with the stress condition during the rearing period. This is supported by a higher level immunocompetent cell population (WBC) in V+P group through the experiment (Fig. 1). Neutrophils and lymphocytes were increased in the vaccinated tilapia fed with Lactobacillus (Venkatalakshmi & Ebanasar 2015). Also, our results showed that the weight gain, final weight TGC in P group were higher than V group and the control (Table 1). This is because when a fish population is immunized with an antigen/vaccine, the growth condition can be negatively influenced at least for a short time (Ward et al., 1985; Midtlying and Lillehaug, 1998; Koskela et al., 2004) because of a force on fish immunephysiological activities to develop the immune responses. Therefore. application of probiotic in the diet of immunized fish can prevent such negative effects on the growth condition as clearly seen in this study (a better growth condition in V+P group compared to others) as similar findings were seen by Soltani *et al.* (2017a, b) when vaccinated trout against *Yersinia ruckeri* were fed the same probiotic, an enhancement was seen in the growth factors and immunological responses compared to vaccinated fish fed normal feed.

Haematological parameters are important indicators of the general of health condition fish (Bandyopadhyay and Mohapatra, 2009; Kader et al., 2010). In this study, the WBC level in vaccinated trout, either with or without L. plantarum dietary supplement was significantly higher than other groups. Similar increases were also reported in Nile tilapia (Oreochromis niloticus) and sturgeon (Huso huso) vaccinated against A. hydrophila, and trout vaccinated with polyvalent vaccine against three pathogenic species (Khosh-bavar-Rostami et al., 2007, Nikoskelainen et al., 2007; Silva et al., 2009). Also, a slight enhancement was seen in RBC and Hb and PCV in vaccinated fish fed with the probiotic. Such enchantment can be attributed to an increase in oxygen requirement due to the increase in metabolite rate in response to immunization (Fair et al., 1999; Pilorz et al., 2005). However, such differences were not significant compared to other trails. Yazici et al. (2015) reported that plantarum did not influence leukocytes counts of trout because their rearing condition was run in sub-lethal water temperatures (25-26C°) that can negatively affect the immune status of the fish.

In conclusion, the present study explores the effects of L. plantarum on growth performance haematological parameters of trout immunized with streptococcosis/lactococcosis vaccine. These results show that growth performance and some haematological indices i.e. WBC were improved in the vaccinated trout fed with the probiotic showing a synergistic effect by the probiotic and the vaccine. To our knowledge, this is the first study evaluating the beneficial efficacy of this probiotic as a dietary supplementation haematology growth and vaccinated trout against streptococcosis and lactoocosis.

Acknowledgement

This work was financially supported by research council of University of Tehran and centre of excellence of aquatic animal health, University of Tehran, Iran.

References

Abid, A., Davies, S.J., Waines, P., Emery, M., Castex. M.. Gioacchini, G., Carnevali, O., Bickerdike, R., Romero, J. and Merrifield, D.L., **2013.** Dietary synbiotic application modulates Atlantic salmon (Salmo salar) intestinal microbial communities and intestinal immunity. Fish and Shellfish Immunology, 35, 1948-1956.

Ackerman, P.A., Iwama, G.K. and Thornton, J.C., 2000. Physiological

- and immunological effects of adjuvanted *Aeromonas salmonicida* vaccines on juvenile rainbow trout. *Journal of Aquatic Animal Health*, 12, 157-164.
- Akinbowale, O.L., Peng, H. and Barton, M.D., 2006. Antimicrobial resistance in bacteria isolated from aquaculture sources in Australia. *Journal of Applied Microbiology*, 100, 1103–1113.
- Ali, S. and Sahu, N.P., 2002. Response of *Macrobrachium rosenbergii* (de Man) juveniles to fish silages as substitutes for fish meal in dry diets. *Asian Fisheries Science*, 15, 59-69.
- Aly S., Mohamed A.A.Z., Rahmani A.H., Nashwa M.A.A. 2016. Trials to improve the response of *Orechromis niloticus* to *Aeromonas hydrophila* vaccine using immunostimulants (garlic, Echinacea) and probiotics (Organic Green TM and Vet-Yeast TM). *African Journal of Biotechnology* 15: 989–994.
- Austin, B. and Austin, D.A., 2007. Bacterial fish pathogens: disease of farmed and wild fish, 4th Edn. Springer Praxis, Godalming, UK, pp 411.
- Azad, I.S. and Al-Marzouk, A., 2008.

 Autochthonous aquaculture probiotics-A critical analysis.

 Proceeding of the 1st International Society of Biotechnology Conference. pp. 171–177.
- Balcázar, J.L., Vendrell, D., de Blas, I., Ruiz-Zarzuela, I., Muzquiz, J.L. and Girones, O., 2008.

 Characterization of probiotic properties of lactic acid bacteria

- isolated from intestinal microbiota of fish. *Aquaculture*, 278, 188–191.
- Bandyopadhyay, P. and Das Mohapatra, P.K., 2009. Effect of a probiotic bacterium *Bacillus circulans* PB7 in the formulated diets: on growth, nutritional quality and immunity of *Catla catla* (Ham.). *Fish Physiology and Biochemistry*, 35(3), 467–478.
- Bostock, J., McAndrew, B., Richards, R., Jauncey, K., Telfer, T., Lorenzen, K., Little, D., Ross, L., Handisyde, N., Gatward, I. and Corner, R., 2010. Aquaculture: global status and trends. *Philosophical Transactions of the Royal Society, B.*, 365, 2897–2912.
- Cha, J.H., Rahimnejad, S., Yang, S.Y., Kim, K.W. and Lee, K.J., **2013.** Evaluation of *Bacillus* spp. as dietary additives on growth performance, innate immunity and disease resistance of Olive Flounder (Paralichtys olivaceus) against Streptococcus iniae water additive. Aquaculture, 402-403, 50-57.
- Dawood, M.A.O., Koshio, Sh., Ishikawa, M. and Yokoyama, S., 2015. Effects of heat killed Lactobacillus plantarum (LP20) supplemental diets growth performance, stress resistance and immune response of red sea bream, Pagrus major. Aquaculture, 442, 29-36.
- Erfanmanesh, A., Soltani, M., Pirali, E., Mohammadian, S. and Taherimirghaed, A., 2012. Genetic characterization of *Streptococcus iniae* in diseased farmed rainbow

- trout (*Onchorhynchus mykiss*) in Iran. *Scientific World Journal*,2012, DOI: 10.1100/2012/594073.
- Evans, J.J., Klesius, P.H. and Shoemaker, C.A., 2004. Efficacy of Streptococcus agalactiae (Group B) vaccine in tilapia (Oreochromis niloticus) by intraperitoneal and bath immersion administration. Vaccine, 22, 3769–3773.
- **Fair, J.M., Hansen, E.S. and Ricklefs, R.E., 1999.** Growth, developmental stability and immune response in juvenile Japanese quails (*Coturnix coturnix japonica*). *Proceedings of the Royal Society of London B*, 266, 1735-1742.
- Giannenas, I., Karamaligas, I., Margaroni, M., Pappas, I., Mayer, E., Encarnacao, P. and Karagouni, E., 2015. Effect of dietary of incorporation a multi-strain probiotic on growth performance and health status in rainbow trout (Oncorhynchus mykiss). Fish Physiology and Biochemistry, 41, 119-128.
- Giri, S.S., Sukumaran, V. and Oviya, M., 2013. Potential probiotic Lactobacillus plantarum VSG3 improves the growth, immunity, and disease resistance of tropical freshwater fish, *Labeo rohita*. Fish and Shellfish Immunology, 34, 660–666.
- Giri, S.S., Sukumaran, V., Sen, S.S. and Jena, P.K., 2014. Effects of dietary supplementation of potential probiotic *Bacillus subtilis* VSG1 singularly or in combination with *Lactobacillus plantarum* VSG3 or/and *Pseudomonas aeruginosa*

- VSG2 on the growth, immunity and disease resistance of *Labeo rohita*. *Aquaculture Nutrition*, 20, 163-171.
- Harikrishnan, R., Balasundaram, C. and Heo, M.S., 2010. Effect of probiotics enriched diet on *Paralichthys olivaceus* infected with lymphocystis disease virus (LCDV). *Fish and Shellfish Immunology*, 29, 868–874.
- Irianto, A. and Austin, B., 2002. Probiotics in aquaculture. *Journal of Fish Diseases*, 25, 1–10.
- Jatobá, A., Vieira, F.N., Buglione-Neto, C.C., Mouriño, J.L.P., Silva, B.C., Seiffert, W.Q. and Andreatta, E.R., 2011. Diet supplemented with probiotic for Nile tilapia in polyculture system with marine shrimp. *Fish Physiology and Biochemistry*, 37, 725-732.
- Kader, M.A., Koshio, S., Ishikawa,
 M., Yokoyama, S. and Bulbul, M.,
 2010. Supplemental effects of some crude ingredients in improving nutritive values of low fishmeal diets for red sea bream, *Pagrus major*.
 Aquaculture, 308. 136–144.
- Karunasagar, I., Pai, R., Malathi, G.R. and Karunasagar, I., 1994.

 Mass mortality of *Penaeus monodon* larvae due to antibiotic-resistant *Vibrio harveyi* infection. *Aquaculture*, 128, 203–209.
- Khoshbavar-Rostami, H.A., Soltani, M. and Hassan, H.M.D., 2007. Immune responses of great sturgeon *Huso huso* to *Aeromonas hydrophila* bacterin. *Journal of Fish Biology*, 70, 1931-1938.
- Koskela, J., Rahkonen, R., Pasternack, M. and Knuutinen,

- H., 2004. Effect of immunization with two commercial vaccines on feed intake, growth, and lysozyme activity in European whitefish (*Coregonus lavaretus* L.). *Aquaculture*, 234, 41–50.
- **Lazado, C.C., Caipang, C.M.A. and Kiron, V., 2012.** Enzymes from the gut bacteria of Atlantic cod, *Gadus morhua* and their influence on intestinal enzyme activity. *Aquaculture Nutrition*, 18, 423-431.
- Midtlying, P.J. and Lillehaug, A., 1998. Growth of Atlantic salmon Salmo salar after intraperitoneal administration of vaccines containing adjuvants. Diseases of Aquatic Organisms, 32, 91–97.
- Nikoskelainen. S., Verho, S., Jarvinen, S., Madetoja, J., Wiklund, T. and Lilius, E., 2007. Multiple whole bacterial antigens in polyvalent vaccine may result in inhibition of specific responses in rainbow trout (Oncorhynchus mykiss). Fish and Shellfish Immunology, 22, 206-217.
- Parthasarathy, R. and Ravi, D., 2011. Probiotic bacteria as growth promoter and biocontrol agent against *Aeromonas hydrophila* in *Catla catla* (Hamilton, 1822). *Indian Journal of Fisheries*, 58, 87-93.
- Pilorz, V., Jäckel, M., Knudsen, K. and Trillmich, F., 2005. The cost of a specific immune response in young guinea pigs. *Physiology and Behavior*, 85, 205-211.
- Raissy, M, Hashemi, S., Roushan, M., Jaafarian, M., Momtaz, H., Soltani M., Pirali-Kheirabad, E. 2018. Effects of essential oils of *Satureja*

- bachtiarica and Nigella sativa on the efficacy of lactococcosis vaccine in rainbow trout (Oncorhynchus mykiss), Iranian Journal of Fisheries Sciences, 95-106.
- Ringo, E. and Gatesoupe, G.J., 1998. Lactic acid bacteria in fish: a review. *Aquaculture*, 160, 177-203.
- Safari, R., Adel, M., Lazado, C.C., Caipang, M.C.A. and Dadar, M., 2016. Host-derived probiotics Enterococcus casseliflavus improves resistance against Streptococcus iniae infection in rainbow trout (Oncorhynchus mykiss) via immunomodulation. Fishand Shellfish Immunology, 52, 198-205.
- Sheehan, B., Labrie, L., Lee, Y., Lim, W., Wong, F. and Chan, J., Komar C., Wendover N., Grisez L. 2009. Streptococcal diseases in farmed tilapia. *Aquaculture Asia Pacific*, 5, 26–9.
- Silva, B.C., Martins, M.L., Jatoba, A., Buglione-Neto, C.C., Vieira, F.N. and Pereira, G.V., Gabriela T. Jerônimo, Walter Q. Seiffert and José Luiz P. Mouriño 2009. Hematological and immunological responses of Nile tilapia after polyvalent vaccine administration by different routes. *Pesquisa Veterinária Brasileira*, 29(11), 874-880.
- Sivani, G., Bhaskar, M. and Sharma, G.R.K., 2016. Influence of probiotics on growth performance and digestive enzyme activities among common carps (*Cyprinus carpio*). International Journal of Science, Environment and Technology, 5, 564 574.

- Soltani, M., Jamshidi, Sh. and 2005. Sharifpour, I.. Streptococcosis caused by Streptococcus iniaein farmed rainbow (Oncorhynchus trout mykiss) in Iran: **Biophysical** characteristics and pathogenesis. Bulletin- European Association of Fish Pathologists, 25, 95-106.
- Soltani, M., Nikbakht, G., Ebrahimzadeh Moussavi, H.A. and Ahmadzadeh, N., 2008. Epizootic outbreaks of Lactococcosis caused by Lactoccoccus garviae in farmed rainbow trout (Onchorhynchus mykiss) in Iran. Bulletin of the European Association of Fish Pathologists, 28, 95-106.
- Soltani, M., Alishahi, M., Mirzargar, S. and Nikbakht, G., 2007. Vaccination of rainbow trout against *Streptococcus iniae* infection: comparison of different routes of administration and different vaccines. *Iranian Journal of Fisheries Sciences*, 7, 129–40.
- Soltani **Emami** M., **A.R.**, **Taherimrghaed** A., Moghani-Ghahremanlo M., Shahbazi M. 2017a. Effect of Macrogard as an immunstimulator on efficacy of bivalent streptococcosis/lactococcosis vaccine in rainbow trout (Onchorhynchus mykiss) fingerling. Journal of Aquaculture Development 11: 61-67.
- Soltani M., Pakzad K., Taheri-Mirghaed Mirzargar S., A., Shekarabi S.P.H., Yosefi P., Solymani N. 2017b. Dietary application of the probiotic

- Lactobacillus plantarum 426951 enhances immune status and growth of rainbow trout (*Oncorhynchus mykiss*) vaccinated against *Yersinia ruckeri*. *Probiotics and Antimicrobial*Proteins.DOI:10.1007/s12602-017-9376-5.
- Son, V.M., Chang, C.C., Wu, M.C., Guu, Y.K., Chiu, C.H. and Cheng, W., 2009. Dietary administration of the probiotic, *Lactobacillus plantarum*, enhanced the growth, innate immune responses, and disease resistance of the grouper *Epinephelus coioides*. *Fish and Shellfish Immunology*, 26, 691-698.
- Sun, Y., Hu, Y., Liu, C. and Sun, L., 2010. Construction and analysis of an experimental *Streptococcus iniae* DNA vaccine. *Vaccine*, 28, 3905–12.
- Sun, Y.Z., Yang, H.L., Ma, R.L., Song, K. and Li, J.S., 2012. Effect of Lactococcus lactis and Enterococcus faecium on growth performance, digestive enzymes and immune response of grouper Epinephelus coioides. Aquaculture Nutrition, 18, 281-289.
- Svobodova, Z., Pravda, D. and Palackova, J., 1991. Unified of methods haematological examination of fish. Research Institute of Fish Culture and Hydrobiology, Vodnany, Methods, 20, 31.
- Venkatalakshmi S., Ebanasar J. 2015. Immunostimulatory effect of *Lactobacillus* sporogenes on the nonspecific defense mechanisms of *Oreochromis mossambicus* (Peters).

International *Journal of Fisheries* and Aquatic Studies 2: 362–369.

Ward, P.D., Tatner, M.F. and Horne, M.T., 1985. Factors influencing the efficacy of vaccines against vibriosis caused by *Vibrio anguillarum*. In: Manning, MJ, Tatner, MF. ŽEds, Fish Immunology. Academic Press, London. pp. 221–229.

Yanbo, W. and Zirong, X., 2006. Effect of probiotics for common carp (*Cyprinus carpio*) based on growth performance and digestive enzyme activities. *Animal Feed Science and Technology*, 127, 283-292.

Yazici, I.S., Hisar, O., Yilmaz, S. and Yigit, M., 2015. Effects of different probiotic bacteria on growth, body composition, immune response and hematological parameters of rainbow trout (*Oncorhynchus mykiss*) under sublethal water temperature. *Marine Science and Technology Bulletin*, 4, 21-28.