Downloaded from jifro.ir on 2025-11-16

Research Article

Functional properties and antioxidant activities of protein hydrolysates from orangefin ponyfish (*Photopectoralis bindus*)

Ramezani Z.¹; Rajabzadeh Ghatarmi E.¹; Hosseini S.F.^{2*}; Regenstein J.M.³

Received: January 2020 Accepted: May 2020

Abstract

Amino acid profiles, functionality, and antioxidative potential of orangefin ponyfish (Photopectoralis bindus) protein hydrolysates, using 1% alcalase, with diverse hydrolysis times (1, 2, 3 and 4 h), were evaluated. The samples solubility in the pH range of 3-9, emulsifying and foaming properties at various protein contents (2.5, 5 and 10 mg/mL), as well as the scavenging of DPPH, ABTS free radicals and reducing power (at concentrations of 1-5 mg/mL), were studied. The amino acid profile of orangefin ponyfish was used to estimate nutritional value. A high amount of essential amino acids (35%) involved in protein hydrolysates and Glx, Asx, and Gly were the dominant amino acids. Solubility was above 90% in the pH range 3-9 and increased at all hydrolysis times by changing the pH from acidic to alkaline (p<0.05). The emulsifying activity (EAI) and stability (ESI) indices were reduced with increasing hydrolysis; at 2.5 mg protein/mL, 1 h hydrolysates displayed the highest EAI (87±2 m²/g) and ESI (22±0.4 min), when considered with those possessing higher hydrolysis time. On the other hand, the foam expansion and stability increased with increasing hydrolysate content. Antioxidant activity of hydrolyzed protein increased with increased hydrolysis (up to 3 mg/mL), especially DPPH and ABTS. However, the highest DPPH and ABTS scavenging activity of FPH was about 73% and 93% at 5 mg/mL hydrolysate concentration and hydrolysis time 4 h, respectively.

Keywords: Fish by-catch, Orangefin ponyfish, Fish hydrolysates, Amino acid composition, Emulsifying properties, Foaming attributes, Antioxidant activities

¹⁻Department of Fisheries, Faculty of Marine Natural Resource, Khorramshahr Marine Sciences and Technology University, Khorramshahr, Iran.

²⁻Department of Seafood Processing, Faculty of Marine Sciences, Tarbiat Modares University, Noor, Iran.

³⁻Department of Food Science, Cornell University, Ithaca, NY 14853-7201, USA.

^{*}Corresponding author's Emial: hosseinisf@modares.ac.ir

Introduction

Fish by-catch is the portion of total fish catch that is dumped back into the water (García-Moreno et al., 2014). Noncommercial fish species with low value and damaged fish that are not worth retaining comprise the marine discards. Discards cause environmental problems since they are normally dead when dumped back into the water. Therefore, it may be helpful to develop processes that will allow the development of products with added value from this underutilized material (García-Moreno et al., 2014). Enzymatic hydrolysis of proteins to produce fish protein hydrolysates (FPH) is one of these methods. The antioxidant activities. amino acid profile, and nutritional composition of FPH have been studied and they are being used in some industrial applications such cosmetics. pharmaceutical products. and human nutrition (Chalamaiah et al., 2012; Rabiei et al., 2019).

On the other hand, the food industry, as well as consumers, has a concern about the lipid oxidation. So, there is an escalating interest in finding natural antioxidants, which may have a less potential hazard in comparison to synthetic ones (You et al., 2009). Additionally, functional attributes of **FPH** including solubility, foam-forming emulsification, and ability have been reported (Hemker et al., 2020). FPH had better solubility at a high degree of hydrolysis, which, in affects the other functional turn,

attributes, like foaming and emulsifying (Ktari *et al.*, 2020).

Orangefin ponyfish (*Photopectoralis* bindus) is demersal and a amphidromous marine fish (Riede, 2004). This species is found in the Persian Gulf and the western part of the Oman Sea (Russell and Houston, 1989), and its maximum length is 11 cm. Orangefin ponyfish may be as much as 55% of the by-catch in the Persian Gulf (Nikoo et al., 2010). Therefore, the production of value-added products like FPH may allow greater use of this species. According to the author's knowledge, this study is the first report regarding functional properties and in vitro antioxidant characterization of protein hydrolysate from Orangefin ponyfish of the Persian Gulf. Thus, the present study aimed to make hydrolyzed proteins, and to (i) evaluate the influences of the hydrolysis time and hydrolysate concentrations on the functional attributes of FPH using solubility, foaming, and emulsifying tests, and (ii) to measure their in vitro antioxidant action by ABTS, DPPH, and reducing power assays.

Materials and methods

Materials

Orangefin ponyfish (8±1 cm length) were obtained from the fishing harbor of Khuzestan (Iran) and kept on ice (with a fish/ice ratio of 1:3 w/w) during transportation, which was about 3 h. Once they arrived in the laboratory, whole ungutted fish were minced by industrial mixer (ARSHIA MG135-

1409, Tehran, Iran), and frozen at -20 °C until use less than a week. Alcalase, DPPH (2,2-diphenyl-1-picrylhydrazyl), BSA (bovine serum albumin), ABTS (2,2-azino bis(3ethylbenzthiazoline)-6sulfonic acid), potassium persulfate, potassium ferricyanide, and ferrous chloride were obtained from Sigma-Aldrich (St. Louis, Mo., USA). Folin-Ciocalteu and SDS (sodium dodecyl sulfate) were gained from Merck Germany). **TCA** (Darmstadt, (trichloroacetic acid) and all other chemicals were obtained from Samchun Pure Chemical Co. Ltd. (Pveongtaek, Korea) as a reagent grade.

Production of orangefin ponyfish hydrolysates

Enzymatic hydrolysis of orangefin ponyfish was done according to the method of Ovissipour et al. (2013). Fifty g of the thawed (4°C, overnight) fish mince was heated at 85°C (for 20 min) for the inactivation of endogenous enzymes; then, the cooked mince was dispersed in 0.2 M SPB (sodium phosphate buffer, pH 8.5) at 1:2 (w:v). It was homogenized using a Heidolph DIAX 900 homogenizer (Kelheim, Germany) at 12000 rpm for about 2 min ambient temperature $(25\pm 2^{\circ}C)$. Alcalase (1%) was then added to the mixture and the reaction was done with constant agitation (200 rpm) at 55°C mixture (for 1-4 h). The was centrifuged at 8000×g (for 20 min at 4°C) utilizing a Universal 320R centrifuge (Hettich, Tuttlingen, Germany); supernatants the

(hydrolysates) were stored at -80°C for further assays, a maximum of one week.

Protein determination

Protein content was obtained based on the method of Lowry et al. (1951) using BSA assumed to be 100% pure as the standard in the range of 1-1000 µg/mL and assuming that BSA adequately represented the protein content of the fish samples. Using UV-vis spectroscopy (UV-1800, Shimadzu, Kyoto, Japan) the absorbance was quantified at 750 nm. The total protein content at 1, 2, 3 and 4 h was 32 ± 0.3 , 31.3 ± 0.5 , 31 ± 0.4 and 31 ± 0.2 mg/mL (n=3), respectively.

Hydrolysis degree (DH)

DH was determined according to the procedure described by Ovissipour *et al.* (2009). A sample volume of 500 μL was removed, mixed with 500 μL of 20% (w/v) TCA, and subsequently centrifuged at 6700×g (for 5 min at 4°C); to determine the soluble amino acids in the supernatant, recognizing that the loss of peptide bonds would decrease the apparent Lowry protein content. The DH was estimated as:

$$DH (\%) = \frac{\text{solubile protein in 10\% TCA}}{\text{total protein in sample}} \times 100$$

Amino acid composition

Detection and quantification of amino acids were measured by high-performance liquid chromatography (HPLC) according to the method of Girgih *et al.* (2015). The freeze-dried sample of hydrolysate (100 mg) was

dissolved in 7.5 mL of 6 mol/L HCl and the mixture was hydrolyzed in glass tubes under an atmosphere of nitrogen at 110°C for 24 h. Reverse-phase column (Sphere-Image 80-5 ODS 2, 25 cm, Knauer, Berlin, Germany) were applied for amino acids separation using methanol acetate buffer (as the mobile phase) and a fluorescence detector (Shimadzu, RF-20A). The 16 amino acids were determined from standard curves of each amino acid in the standard mixture based on peak area measurements assuming of response with concentration, using EZChrom Elite software using the Scientific Software Inc. module (SS420x) (Pleasanton, CA, USA), and expressed as mg amino acid/100 g of Lowry protein.

Functional attributes of protein hydrolysates

Solubility

For this, hydrolysates (200 mg) were dissolved in 20 mL of deionized water, and pH was adjusted to 3, 5, 7, and 9 with 1 M HCl or 1 M NaOH. The blend was agitated at ambient temperature and centrifuged at 8000×g (for 20 min at 4°C). Supernatant protein was determined and the solubility calculated (Lowry *et al.*, 1951):

Solubility (%) =

 $\frac{\text{Protein content in supernatant}}{\text{Protein content in sample}} \times 100$

Emulsifying attributes

The indices of emulsifying activity (EAI) and emulsion stability (ESI) were determined as previously demonstrated

by Giménez *et al.* (2009). Soybean oil (2 mL) and different concentrations (2.5, 5 and 10 mg/mL) of FPH solution (6 mL) were homogenized (~20000 rpm, 1 min); a sample volume of 50 µL was diluted by 0.1% SDS, and the absorbance was then quantified at 500 nm. The absorbance (A0 and A10) were utilized to compute the EAI and ESI as follows:

EAI
$$(m^2/g) = \frac{2 \times 2.303 \times 100 \times A}{c \times 0.25 \times 10000}$$

where A = absorbance at 500 nm; c = protein concentration (g/mL).

$$ESI (min) = \frac{A0 \times 10}{A0 - A10}$$

where A0 and A10 are the absorbance quantified immediately and 10 min after emulsion formation, respectively.

Foaming attributes

FE (foam expansion) and FS (foam stability) were ascertained as delineated by Nalinanon *et al.* (2011). FPH solutions with 2.5, 5, and 10 mg/mL protein were moved into a cylinder and homogenized at 14000 rpm for 1 min at ambient condition. FE and FS were then computed by these equations:

$$FE (\%) = (V_T/V_0) \times 100$$

FS (%) =
$$(V_t/V_0) \times 100$$

where V_T is total volume after whipping, V_0 is the initial volume before whipping, and V_t is total volume after 60 min.

Antioxidant activity determination DPPH assay

The DPPH radical-scavenging capacity was measured using the protocol of Bougatef *et al.* (2010), with some changes. An aliquot of 500 µL of each

sample (at 1-5 mg/mL) was mixed with $500 \mu L$ of 0.15 mM DPPH, and then vigorously mixed and kept in the dark (for 30 min) at room condition. The absorbance of the solution was quantified at 517 nm. The scavenging ability was:

DPPH radical-scavenging assay (%)
$$= \frac{\text{Ablank-Asample}}{\text{Ablank}} \times 100$$

ABTS assay

This assay was performed using the modified procedure of Giménez *et al.* (2009). Briefly, a sample volume of 20 µL was mixed with 980 mL of ABTS reagent and absorbance was quantified at 734 nm after 10 min incubation at 30°C. The scavenging capacity was:

ABTS radical-scavenging assay (%)
$$= \frac{\text{Ablank-Asample}}{\text{Ablank}} \times 100$$

Reducing power assay

This assay was done using the method of Bougatef *et al.* (2010). An aliquot of the sample (200 μ L) at 1-5 mg/mL was blended with the same quantity of sodium phosphate buffer and potassium ferricyanide solution (1% w/v). The mixture was incubated at 50°C for 30 min, pursued by the inclusion of 500 μ L of TCA (10% w/v); then the mixture was centrifuged at 1650×g (for 10 min at 4°C). Finally, 500 μ L of distilled water and 100 μ L of ferric chloride (0.1% w/v) were added to 500 μ L of supernatant solution. The solution was incubated at room condition for 10 min

for color development, and absorbance was read at 700 nm.

Statistical analysis

Statistical analyses of the data were evaluated using a one-way analysis of variance (ANOVA) followed by Duncan's multiple range tests. The threshold for statistical significance was set at p<0.05. All experiments were performed in triplicate and the results are presented as means \pm standard deviation.

Results

Degree of hydrolysis

The effect of time on the degree of hydrolysis (DH) of orangefin ponyfish protein is illustrated in figure 1. As can be seen from Figure 1, the lowest amount of hydrolysis was $25.61\pm0.61\%$ and the highest was $28.07\pm0.31\%$ for the first and fourth hours, respectively.

Amino acid composition

The amino acid composition of both protein hydrolysates, expressed as mg amino acid/100 g protein, is shown in Table 1. The total amino acids recovered from orangefin ponyfish protein hydrolysate was 97.7 mg/100 g (Table 1). Because amino acid residues add water on hydrolysis, the actual amount of amino acids that should be recovered is around 115 to 118 mg/100 g protein. Thus, errors in the protein determinations, the absence of hydroxyproline, tryptophan, and hydroxylysine along with the digestion destroying some amino acids and not hydrolyzing some peptide bonds would account for the lack of full recovery. As shown in Table 1, the orangefin ponyfish hydrolysate was rich in Glx (17.8 mg/100 g), Asx, Gly, Val, and Leu.




Figure 1: Degree of hydrolysis (DH) of orangefin ponyfish hydrolysates. Error bars represent standard deviation, n = 3.

Table 1: Amino acid composition of protein hydrolysates from orangefin ponyfish.

from orangetin ponytish.				
Amino acid	mg amino acid/100 g protein			
Aspartic acid (Asx)	11.8			
Glutamic acid (Glx)	17.8			
Serine (Ser)	7.3			
Histidine (His)	0.8			
Glycine (Gly)	11.1			
Threonine (Thr)	5.1			
Arginine (Arg)	3.2			
Alanine (Ala)	3.3			
Proline (Pro)	3.4			
Tyrosine (Tyr)	4.9			
Methionine (Met)	2.4			
Valine (Val)	9.2			
Phenylalanine (Phe)	3.3			
Isoleucine (Ile)	4.2			
Leucine (Leu)	8.8			
Lysine (Lys)	1.1			
HAA	50.6			
AAA	8.2			
EAA	34.9			
Total	97.7			
Total	* * * * * * * * * * * * * * * * * * * *			

Combined total of hydrophobic amino acids (HAA) = Ala, Val, Ile, Leu, Tyr, Phe, Pro, Met, Gly; romatic amino acids (AAA) = Phe, Tyr; essential amino acids (EAA) = Histidine, Ile, Leu, Lys, Met, Phe, Thr, Val.

Hydrolysates solubility

The solubility of orangefin ponyfish hydrolysates with various hydrolysis time at different pH values are tabulated in Table 2. All samples were soluble over a wide range of pH, in which more

than 90% solubility was acquired. Meanwhile, the solubility of hydrolysates at alkaline pH was higher than that of acidic one (p<0.05). The least solubility of FPH was in pH 5.

Table 2: Solubility (%) of hydrolysates from orangefin ponyfish at various pH.

Hydrolysis time (h)

pН	1	2	3	4
3	94 ± 0.1^{Cc}	$97 \pm 0.1^{\text{Bb}}$	$98 \pm 0.1^{\mathrm{Ab}}$	98 ± 0.1^{Ac}
5	92 ± 0.1^{Bd}	95 ± 0.2^{Ac}	95 ± 0.1^{Ac}	95 ± 0.1^{Ad}
7	94 ± 0.1^{Cb}	98 ± 0.2^{Ba}	98 ± 0.1^{ABa}	98 ± 0.1^{Ab}
9	96 ± 0.1^{Da}	98 ± 0.2^{Ca}	98 ± 0.1^{Ba}	99 ± 0.1^{Aa}

Values in the same column followed by different lower case letters and in the same row followed by different upper case letters, for each parameter, are significantly different (p<0.05).

Emulsifying properties

Emulsifying characteristics of orangefin ponyfish hydrolysates concerning the emulsion activity (EAI) and stability indices (ESI) with different hydrolysis time at various amounts (2.5, 5 and 10 mg/mL) are presented in Table 3. In ESI general, both **EAI** and of with raising hydrolysates reduced hydrolysis protein time and concentration, particularly regarding EAI when the protein content increased (p<0.05). At 2.5 mg protein/mL, 1 h hydrolysates displayed the highest EAI $(87\pm 2 \text{ m}^2/\text{g})$ and ESI $(22\pm 0.4 \text{ min})$, when considered with those possessing higher hydrolysis time (p < 0.05).

Foaming characteristics

Foam expansion (FE) and foam stability (FS) of orangefin ponyfish hydrolysates with various hydrolysis times at various

contents (2.5, 5 and 10 mg/mL) are shown in Table 3. At each hydrolysate concentration, insignificant decreases in FE and FS were noticed when hydrolysis time was raised (p<0.05). At 10 mg protein/mL, 1 h hydrolysates displayed the highest FE (75±1%) and FS (50±0.3%), when considered with those possessing higher hydrolysis time (p<0.05).

Antioxidant activities

DPPH-scavenging ability

DPPH-scavenging capacity of orangefin ponyfish hydrolysates with different hydrolysis time at various amounts (1-5 mg/mL) are presented in Table 4. At these concentrations, a significant increase in radical-scavenging capacity was observed when hydrolysis time increased from 1 to 4 h (p<0.05), except at concentration 1 (hydrolysis time 2-4

h) and 2 (hydrolysis time 2-3 h) mg/mL. As can be seen from Table 4, the highest DPPH scavenging activity

of FPH was about 73% at 5 mg/mL hydrolysate concentration and hydrolysis time 4 h.

Table 3: Emulsifying (emulsifying activity index, EAI; emulsifying stability index, ESI) and foaming properties (foam expansion, FE; foam stability, FS) of hydrolysates from orangefin ponyfish at various concentrations.

Protein oncentration	-	EAI (m^2/g)	ESI (min)	FE (%)	FS (%)
(mg/mL)	time (h)				
2.5	1	$87 \pm 2^{A*a**}$	22 ± 0.4^{Aa}	$29 \pm 0.2^{\text{Ca}}$	$18 \pm 0.4^{\text{Ca}}$
	2	80.3 ± 0.1^{Ab}	20 ± 0.3^{Ab}	25 ± 1^{Cb}	13 ± 0.4^{Cb}
	3	77.2 ± 0.2^{Ac}	19 ± 0.2^{Abc}	23 ± 1^{Ab}	9 ± 0.3^{Cb}
	4	72 ± 0.2^{Ad}	$18\pm0.2^{\text{Ac}}$	15 ± 0.3^{Cc}	3 ± 0.3^{Cc}
5	1	34.4 ± 0.3^{Bb}	19 ± 0.4^{Aa}	44 ± 1^{Ba}	31 ± 0.3^{Ba}
	2	37 ± 1^{Ba}	$19 \pm 0.1^{\text{Aa}}$	38 ± 2^{Bb}	25 ± 0.3^{Bb}
	3	28 ± 0.4^{Bc}	$18\pm0.5^{\mathrm{Aa}}$	38 ± 2^{Ab}	21 ± 1^{Bc}
	4	25 ± 0.4^{Bd}	$17 \pm 0.2^{\text{Bb}}$	34 ± 1^{Bc}	18 ± 0.3^{Bd}
10	1	18 ± 0.3^{Ca}	16 ± 0.3^{Ca}	$75\pm1^{\mathrm{Aa}}$	50 ± 0.3^{Aa}
	2	17 ± 0.2^{Cb}	$14 \pm 0.1^{\text{Bb}}$	$72\pm1^{\mathrm{Aa}}$	48 ± 0.3^{Aa}
	5	15 ± 0.2^{Cc}	14 ± 0.4^{Bbc}	63 ± 0.4^{Ab}	40 ± 0.4^{Ab}
	4	15 ± 0.3^{Cc}	$13 \pm 0.1^{\text{Cc}}$	60 ± 0.5^{Ab}	38 ± 0.4^{Ac}

^{*}Different capital letters in the same column within the same hydrolysate concentration indicate significant differences (p<0.05).

ABTS radical-scavenging capacity

The ABTS-scavenging ability of orangefin ponyfish hydrolysates with various hydrolysis times at different concentrations (1-5 mg/mL) is tabulated in Table 4. According to the results, with increasing concentrations, the

scavenging effects of hydrolysate on ABTS radical increased around 93% in 5 mg/mL in FPH concentration and hydrolysis time 4 h.

^{**}Different lower case letters in the same column within the same hydrolysis time indicate significant differences (*p*<0.05).

Reducing power activity

Table 4 shows the reducing power abilities of orangefin ponyfish hydrolysates at different concentrations (1-5 mg/mL) and different hydrolysis time. The reducing power was found to

be very low for all the fractions and exhibited a concentration dependency. Reducing power of hydrolysates significantly increased (p<0.05) with increasing concentrations (1 to 5 mg/mL).

Table 4: *In vitro* antioxidant activity of protein hydrolysates from orangefin ponyfish at various concentrations

concentrations.				
Protein concentration (mg/mL)	Hydrolysis time (h)	DPPH radical scavenging activity (%)	ABTS radical scavenging activity (%)	Ferric reducing power (OD at 700 nm)
1	1	39 ± 0.4 ^{B*e**}	45 ± 1 ^{Cc}	$0.1 \pm 0.01^{\text{Be}}$
	2	41 ± 0.3^{Ad}	$52\pm1^{\mathrm{Bd}}$	0.1 ± 0.01^{Be}
	3	$41 \pm 0.2^{\text{Ad}}$	52 ± 0.1^{Bd}	0.1 ± 0.001^{Be}
	4	42 ± 0.4^{Ae}	56 ± 1^{Ad}	$0.1\pm0.01^{\mathrm{Ae}}$
2	1	56 ± 1^{Cd}	80 ± 1^{Cb}	0.2 ± 0.002^{Cd}
	2	61 ± 0.2^{Bc}	82 ± 0.3^{Bc}	$0.2\pm0.002^{\mathrm{Bd}}$
	3	62 ± 1^{Bc}	82 ± 1^{Bc}	$0.2\pm0.002^{\mathrm{Bd}}$
	4	$65 \pm 0.3^{\text{Ad}}$	83 ± 0.2^{Ac}	$0.2\pm0.01^{\mathrm{Ad}}$
3	1	62 ± 0.4^{Cd}	$89 \pm 0.2^{\text{Ca}}$	0.2 ± 0.002^{Cc}
	2	64 ± 0.3^{Cb}	$90 \pm 0.2^{\text{Bb}}$	0.2 ± 0.003^{Cc}
	5	66 ± 1^{Bb}	90 ± 0.1^{ABb}	0.2 ± 0.002^{Bc}
	4	68 ± 0.1^{Ac}	91 ± 0.2^{Ab}	$0.2\pm0.003^{\mathrm{Ac}}$
4	1	63 ± 0.2^{Db}	90 ± 0.3^{Ca}	$0.2\pm0.003^{\mathrm{Bb}}$
	2	66 ± 1^{Cab}	91 ± 0.1^{Bab}	0.2 ± 0.01^{Bb}
	3	68 ± 0.4^{Ba}	91 ± 0.2^{Bab}	0.2 ± 0.001^{ABb}
	4	71 ± 0.3^{Ab}	92 ± 0.1^{Ab}	0.2 ± 0.003^{Ab}
5	1	65.3 ± 0.3^{Da}	90 ± 0.1^{Da}	0.3 ± 0.003^{Ca}
	2	67 ± 1^{Ca}	91 ± 0.2^{Ca}	0.3 ± 0.002^{Ba}
	3	69 ± 1^{Ba}	92 ± 0.1^{Ba}	0.3 ± 0.001^{Ba}
	4	73 ± 0.3^{Aa}	93 ± 0.1^{Aa}	0.3 ± 0.001^{Aa}

^{*}Different capital letters in the same column within the same hydrolysate concentration indicate significant differences (p<0.05).

^{**}Different lower case letters in the same column within the same hydrolysis time indicate significant differences (p<0.05).

Discussion

Degree of hydrolysis (DH), defined as the level of protein hydrolysis degradation, is the most extensively applied index for differentiation among various proteins (Bougatef et al., 2010; Raftani Amiri et al., 2016). The results revealed that the DH increased with increasing time of hydrolysis (Fig. 1); the hydrolysis curve acquired in the study is alike present those delineated earlier for hydrolysates from Chinese sturgeon (Acipenser sinensis) (Noman et al., 2019), hydrolysates obtained from discarded Mediterranean fish species (García-Moreno et al., 2014), and hydrolysates of defatted salmon backbones (Slizyte et al., 2016). Noman et al. (2018) reported that a longer incubation time would allow the enzyme to act more extensively on the protein, thus resulting in an increment in the DH.

The nutritional value of protein hydrolysates (as flavor enhancers and functional ingredients) relies on the composition of their small/large peptides and free amino acids (dos Santos et al., 2011). FPHs showed a difference in their amino acid composition that influenced by various elements e.g. source of the enzyme and fish species (Nasri et al., 2013). As shown in Table 1, the orangefin ponyfish hydrolysate was rich in Glx (17.8 mg/100 g). Determination of Glx as the most prevalent amino acid in fish protein hydrolysate has been addressed previous researches some (Ovissipour et al., 2009; Noman et al.,

2018). As shown in Table 1, the orangefin ponyfish protein had a high amount of HAA (hydrophobic amino acids) (50.6 mg/100 g), providing structural features that can enhance interactions with lipid foods (Girgih et al., 2015). Additionally, the protein hydrolysate had aromatic amino acids (AAA), such as Phe (3.3 mg/100 g) and Tyr (4.9 mg/100 g). Sarmadi and Ismail (2010) described that AAA can donate electrons to convert radicals to stable molecules while maintaining their own stability via resonance structures thus increasing the radical-scavenging activities of the amino acids (Rajapakse et al., 2005). Based on the amino acid profile, orangefin ponyfish hydrolysates showed a high nutritional value and, therefore. maybe a good dietary supplement to balance other dietary proteins.

Generally, the hydrolysates were more soluble in alkaline pH, in contrast with acidic one (Table 2). According to Ktari et al. (2020), the reduction of the secondary structure of FPH and the release of smaller polypeptide units from the protein resulted in an increase in solubility of hydrolysates. Similarly, stripe trevally vellow (Selaroides leptolepis) hydrolysates with various DHs (5-25%), also displayed high solubility (>85%) in the pH range of 2-12 (Klompong et al., 2007). In the all hydrolysis time, the lowest solubility was observed at pH 5, which is consistent with the results reported by Nalinanon et al. (2011) for hydrolysate obtained from ornate threadfin bream and Noman et al. (2019) for sturgeon hydrolysates. These authors demonstrated that high molecular weight (MW) proteins and/or peptides were precipitated at this pH, which was near the isoelectric points (pI) of myofibrillar proteins. Furthermore, since the net charge of the proteins is minimized near the pI, more proteinprotein interaction and less proteinwater interaction occur (Noman et al., 2019). Consequently, the protein solubility decreases, whenever, the pH moves away from this point, increases protein-water interactions At pH 9, the solubility of samples was increased with increasing hydrolysis time from 1 to 4 h. According to Nalinanon et al. (2011), the dissimilarities in FPH solubility with various DH could be ascertained by the peptides size, the peptides charge group, and the hydrophobic-hydrophilic balance.

Regarding emulsifying characteristics, both EAI and ESI of hydrolysates reduced with raising hydrolysis time and protein concentration (Table 3). Klompong et al. (2007) postulated that superfluous hydrolysis causes significant loss in the emulsifying attributes; this may be due to the low MW peptides which not being amphiphilic sufficient to display suitable emulsifying attributes (Nalinanon et al., 2011; Hemker et al., 2020). A decrease in emulsifying activity with increasing protein concentration had been described for other fish proteins such as ornate threadfin hydrolysates (Nalinanon et al., 2011) and sturgeon hydrolysates (Noman et al., 2019). Noman et al. (2019) demonstrated that this event may be also owing to a decrease in proteinprotein interactions, contributing a lower protein concentration at the interface of oil/water. At high protein concentration, the activation energy barrier does not allow protein migration to take place in a diffusion-dependent manner, leading to protein aggregation in the aqueous phase; while, at low concentrations, adsorption at the oil-water interface is diffusion-controlled (Nalinanon et al., 2011).

The feature of proteins to form foams is principal in the preparation of a variety of foods (Zayas, 1997); egg white, gelatin, casein, soy proteins, and gluten are the most commonly used protein as foaming agents. The foaming properties strongly depend on the transportation, penetration, and rearrangement molecules at the air-water interface (Noman et al., 2018). FPH has better foaming properties, thus it may be applied in the construction of foambased products (Barrow and Shahidi, 2007). According Table to insignificant decreases foaming in hydrolysate attributes at each concentration were noticed when hydrolysis time was raised. Similar findings were reported by Nalinanon et al. (2011) in ornate threadfin muscle protein hydrolysates and Liu et al. (2014) in hydrolysates from surimi processing by-products, which they

ascribed to a lesser alignment of small peptides at the air/water interface. The decrease in FE may also be related to the aggregation of proteins that interfere in the interactions between proteins and the water needed for the formation of foam (Noman et al., 2018). However, at same hydrolysis time. expansion of hydrolysates showed a significant increase with increasing concentration. The protein same tendency was observed by Thiansilakul et al. (2007) with round scad mince hydrolysates, where increased protein concentration led to higher foam expansion. Foam stability (FS) showed a slight decrease with increasing hydrolysis time (Table 3). A similar trend was also observed in ornate threadfin hydrolysates with 10-30% DH (Nalinanon et al., 2011). According to Nalinanon et al. (2011), peptides with low MW could not keep a well-ordered orientation of the molecule at the interface. Low FS possibly also arises from the formation of free amino acids during hydrolysis (Noman et al., 2018). Nevertheless, at the same hydrolysis time, FS of hydrolysate was enhanced by increasing content, resulting from the formation of stiffer foams (Lawal, 2004).

The ability of orangefin ponyfish hydrolysates to scavenge the free radicals was tested by their ability to scavenge the stable DPPH radical (Ramezanzade *et al.*, 2017). Generally, a significant increase in radical-scavenging capacity was observed when hydrolysis time increased from 1

to 4 h. Wu et al. (2003) showed that the scavenging effect of the hydrolysates slightly increased in the first 5 h, and remained constant throughout extended time of hydrolysis. At all hydrolysis times, **DPPH-scavenging** capacities increased with increasing FPH. Similarly, Noman et al. (2019) the **DPPH** showed that radical inhibitory percentage of sturgeon hydrolysates increased with increasing protein concentrations. Pires et al. (2013) also reported that there was a positive relationship between scavenging capacity and protein hydrolysate concentration of Cape hake (Merluccius capensis), in which the inhibiting percent increased increasing protein concentration. Also, the high level of DPPH radical scavenging ability of orangefin ponyfish hydrolysates is associated with a high amount of hydrophobic amino this acids (50.6% in study) (Ramezanzade et al., 2018).

To determine the antioxidant ability of hydrogen-donating constituents the ABTS assay is appropriate (Hosseini et al., 2017; Joghataei et al., 2019). Generally, the protein hydrolysates were able to quench ABTS radicals. Although the hydrolysis time had no significant influence **ABTS** on scavenging ability, the effect of protein concentrations from 1-3 mg/mL was significant, which is supported by previous reports (Ramezanzade et al., 2018; Ktari et al., 2020). The protein concentration can affect antioxidant activity, but excessive concentration has the opposite effect. ABTS results showed that orangefin ponyfish protein hydrolysates probably contain amino acids that act as electron donors, reacting with free radicals to give rise to more stable products (Musso *et al.*, 2019).

The reducing power assay was also applied evaluate the in vitro to antioxidant activities of protein hydrolysate since some research has indicated a direct correlation between antioxidant activities and the reducing power of the peptides (Bougatef et al., 2010; Hosseini et al., 2018). This assay is based on the ability of antioxidants to reduce ferricyanide (Fe⁺³) complex to the ferrous (Fe⁺²) form (Hosseini *et al.*. 2019). Generally, a significant increase in the reducing power was found with increasing protein concentration (Table 4), which is consistent with the report of Noman et al. (2019), who found a linear trend between the increases of the reducing power with increasing hydrolysates. sturgeon However, different hydrolysis times (1-4 h) had no significant influence on reducing power of orangefin ponyfish hydrolysates. Klompong et al. (2007) demonstrated that with increasing DH from 5 to 23%, no significant increase in the reducing power of the yellow stripe trevally hydrolysates observed. Contrary to the present study, Noman et al. (2019) demonstrated that reducing increased with power increased DH of sturgeon protein hydrolysates, which was attributed to a higher content of amino acids.

In conclusion, the protein hydrolysates produced from orangefin ponyfish maybe serve as a good source of beneficial peptides and amino acids. This study showed that it is possible to obtain fish hydrolysates having strong in vitro antioxidant capacities as well as good functionalities when using the discarded species (orangefin main ponyfish) from the Persian Gulf (~55%) raw material. The results a established that functional and antioxidant attributes of the hydrolysates were determined by the hydrolysis times as well as protein concentrations. The protein hydrolysates may be applied in food systems as a natural additive with emulsifying and foaming functionalities and antioxidant properties. They would probably inhibit lipid oxidation using radical scavenging activity. research should be performed to purify and identify the antioxidative peptides in orangefin ponyfish hydrolysate.

Acknowledgements

Financial support from Research Councils of Tarbiat Modares University and Khorramshahr University of Marine Science and Technology is gratefully acknowledged.

References

Barrow, C. and Shahidi, F., 2007.

Marine nutraceuticals and functional foods. CRC Press.

Bougatef, A., Nedjar-Arroume, N., Manni, L., Ravallec, R., Barkia, A., Guillochon, D. and Nasri, M., **2010.** Purification and identification of novel antioxidant peptides from enzymatic hydrolysates of sardinelle (*Sardinella aurita*) by-products proteins. *Food Chemistry*, 118(3), 559-565.

DOI:10.1016/j.foodchem.2009.05.02

- Chalamaiah, M., Hemalatha, R. and Jyothirmayi, T., 2012. Fish protein hydrolysates: proximate composition, amino acid composition, antioxidant activities and applications: a review. *Food Chemistry*, 135(4), 3020-3038. DOI:10.1016/j.foodchem.2012.06.10
- dos Santos, S.D.A., Martins, V.G., Salas-Mellado, M. and Prentice, C., 2011. Evaluation of functional properties in protein hydrolysates from bluewing searobin (*Prionotus punctatus*) obtained with different microbial enzymes. *Food and Bioprocess Technology*, 4(8), 1399-1406. DOI:10.1007/s11947-009 0301-0
- García-Moreno, P.J., Batista, I., Pires, C., Bandarra, N.M., Espejo-Carpio, F.J., Guadix, A. and Guadix, E.M., 2014. Antioxidant activity of protein hydrolysates obtained from discarded Mediterranean fish species. *Food Research International*, 65, 469-476. DOI:10.1016/j.foodres.2014.03.061
- Giménez, B., Alemán, A., Montero, P. and Gómez-Guillén, M.C., 2009.

 Antioxidant and functional properties of gelatin hydrolysates

obtained from skin of sole and squid. *Food Chemistry*, 114(**3**), 976-983. DOI:

10.1016/j.foodchem.2008.10.050

- Girgih, A.T., He, R., Hasan, F.M., Udenigwe, C.C., Gill, T.A. and Aluko, R.E., 2015. Evaluation of the in vitro antioxidant properties of a cod (*Gadus morhua*) protein hydrolysate and peptide fractions. *Food Chemistry*, 173, 652-659. DOI: 10.1016/j.foodchem.2014.10.079
- Hemker, A.K., Nguyen, L.T., Karwe, M. and Salvi, D., 2020. Effects of pressure-assisted enzymatic hydrolysis functional and bioactive properties of tilapia (Oreochromis niloticus) by-product protein hydrolysates. LWT, 122. 109003. DOI: 10.1016/j.lwt.2019.109003
- Hosseini, S.F., Ramezanzade, L. and Nikkhah, M., 2017. Nano-liposomal entrapment of bioactive peptidic fraction from fish gelatin hydrolysate. *International Journal of Biological Macromolecules*, 105, 1455-1463.

DOI:10.1016/j.ijbiomac.2017.05.141

Hosseini, S.F., Soleimani, M.R. and Nikkhah, M., 2018. Chitosan/sodium tripolyphosphate nanoparticles as efficient vehicles for antioxidant peptidic fraction from common kilka. *International Journal of Biological Macromolecules*, 111, 730-737.

DOI:10.1016/j.ijbiomac.2018.01.023

Hosseini, S.F., Nahvi, Z. and Zandi, M., 2019. Antioxidant peptide-

loaded electrospun chitosan/poly (vinyl alcohol) nanofibrous mat intended for food biopackaging purposes. *Food hydrocolloids*, 89, pp.637-648.

DOI:10.1016/j.foodhyd.2018.11.033

- Joghataei, M., Hosseini, S.F. and Arab-Tehrany, 2019. E., Cinnamaldehyde loaded chitosan/tripolyphosphate nanoassemblies: Fabrication, characterization, and in vitro evaluation of antioxidant activity. Journal of Food Processing and Preservation. 43(7). e13972. DOI:10.1111/jfpp.13972
- Klompong, V., Benjakul, S., Kantachote, D., and Shahidi, F. 2007. Antioxidative activity and functional properties of protein hydrolysate of yellow stripe trevally (*Selaroides leptolepis*) as influenced by the degree of hydrolysis and enzyme type. *Food Chemistry*, 102, 1317-1327.

DOI:10.1016/j.foodchem.2006.07.01

- Ktari, N., Salem, R.B.S.B., Bkhairia, I., Slima, S.B., Nasri, R., Salah, R.B. and Nasri, M., 2020. Functional properties and biological activities of peptides from zebra blenny protein hydrolysates fractionated using ultrafiltration. Bioscience, 34. 100539. DOI:10.1016/j.fbio.2020.100539
- Lawal, O.S., 2004. Functionality of African locust bean (*Parkia biglobossa*) protein isolate: effects of pH, ionic strength and various

- protein concentrations. *Food Chemistry*, 86(**3**), 345-355.

 DOI:10.1016/j.foodchem.2003.09.03
 6
- Liu, Y., Li, X., Chen, Z., Yu, J., Wang, F. and Wang, J., 2014. Characterization of structural and functional properties of fish protein hydrolysates from surimi processing by-products. *Food Chemistry*, 151, 459-465.

DOI:10.1016/j.foodchem.2013.11.08

- Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J., 1951. Protein measurement with the Folin phenol reagent. *Journal of Biological Chemistry*, 193, 265-275.
- Musso, Y.S., Salgado, P.R. and Mauri, A.N., 2019. Smart gelatin films prepared using red cabbage (*Brassica oleracea* L.) extracts as solvent. *Food Hydrocolloids*, 89, 674-681.

DOI:10.1016/j.foodhyd.2018.11.036

- Nalinanon, S., Benjakul, S., Kishimura, H. and Shahidi, F., 2011. Functionalities and antioxidant properties of protein hydrolysates from the muscle of ornate threadfin bream treated with pepsin from skipjack tuna. *Food Chemistry*, 124(4), 1354-1362. DOI:10.1016/j.foodchem.2010.07.08
- Nasri, R., Younes, I., Jridi, M., Trigui, M., Bougatef, A., Nedjar-Arroume, N., Dhulster, P., Nasri, M. and Karra-Châabouni, M., 2013. ACE inhibitory and

- antioxidative activities of Goby (*Zosterissessor ophiocephalus*) fish protein hydrolysates: effect on meat lipid oxidation. *Food Research International*, 54(1), 552-561. DOI:10.1016/j.foodres.2013.07.001
- Nikoo, S., Savari, A., Cochnian, P., Dehghan Medise, S. and Saki, S., 2010. Non-commercial fish species in by-catch composition of shrimp trawl landing from Mahshahr creeks. *Iranian Scientific Fisheries Journal*, 19(2), 149-153.
- Noman, A., Xu, Y., AL-Bukhaiti, W.Q., Abed, S.M., Ali, A.H., Ramadhan, A.H. and Xia, W., 2018. Influence enzymatic of hydrolysis conditions on the degree hydrolysis and functional properties of protein hydrolysate obtained from Chinese sturgeon (Acipenser sinensis) by using papain enzyme. Process Biochemistry, 67, 19-28.

DOI:10.1016/j.procbio.2018.01.009

Noman, A., Qixing, J., Xu, Y., Ali, A.H., Al-Bukhaiti, W.Q., Abed, S.M. and Xia, W., 2019. Influence of degree of hydrolysis on chemical composition, functional properties, and antioxidant activities of Chinese sturgeon (Acipenser sinensis) hydrolysates obtained by using alcalase 2.4 L. Journal of Aquatic Food Product Technology, 28(6), 583-597.

DOI:10.1080/10498850.2019.16265 23

Ovissipour, M., Abedian, A., Motamedzadegan, A., Rasco, B.,

- Safari, R. and Shahiri, H., 2009. The effect of enzymatic hydrolysis time and temperature on the properties of protein hydrolysates from Persian sturgeon (*Acipenser persicus*) viscera. *Food Chemistry*, 115(1), 238-242. DOI:10.1016/j.foodchem.2008.12.01
- Ovissipour, M., Rasco, B., Shiroodi, S.G., Modanlow, M., Gholami, S. and Nemati, M., 2013. Antioxidant activity of protein hydrolysates from whole anchovy sprat (*Clupeonella engrauliformis*) prepared using endogenous enzymes and commercial proteases. *Journal of the Science of Food and Agriculture*, 93(7), 1718-1726. DOI:10.1002/jsfa.5957
- Pires, C., Clemente, T. and Batista, I., 2013. Functional and antioxidative properties of protein hydrolysates from Cape hake by-products prepared by different three methodologies. Journal of the Science of Food and Agriculture, 771-780. 93(4), DOI:10.1002/jsfa.5796

Rabiei, S., Rezaei, M., Asgharzade, S., Nikoo, M. and Rafieia-kopai, M., 2019. Antioxidant and cytotoxic properties of protein hydrolysates obtained from enzymatic hydrolysis of Klunzinger's mullet (*Liza klunzingeri*) muscle. *Brazilian Journal of Pharmaceutical Sciences*, 55. DOI:10.1590/s2175-97902019000218304

Raftani Amiri, Z., Safari, R. and Bakhshandeh, T., 2016. Functional

- properties of fish protein hydrolysates from cuttlefish (*Sepia pharaonis*) muscle produced by two commercial enzymes. *Iranian Journal of Fisheries Sciences*, 15(4), 1485-1499.
- Rajapakse, N., Mendis, E., Byun, H.G. and Kim, S.K., 2005. Purification and in vitro antioxidative effects of giant squid muscle peptides on free radical-mediated oxidative systems. *The Journal of Nutritional Biochemistry*, 16(9), 562-569. DOI:10.1016/j.jnutbio.2005.02.005
- Ramezanzade, L., Hosseini, S.F. and Nikkhah, M., 2017. Biopolymer-coated nanoliposomes as carriers of rainbow trout skin-derived antioxidant peptides. *Food Chemistry*, 234, 220-229. DOI:10.1016/j.foodchem.2017.04.177
- Ramezanzade, L., Hosseini, S.F., Nikkhah, M. and Arab-Tehrany, E., 2018. Recovery of bioactive peptide fractions from rainbow trout (*Oncorhynchus mykiss*) processing waste hydrolysate. *Ecopersia*, 6(1), 31-40.
- **Riede, K., 2004.** Global register of migratory species: from global to regional scales: final report of the R&D-Projekt 808 05 081. Federal Agency for Nature Conservation.
- Russell, B. C., and Houston, W. 1989.

 Offshore fishes of the Arafura Sea.

 Beagle: Records of the Museums and

 Art Galleries of the Northern Territory,

 The, 6, 69-84.
- Sarmadi, B.H. and Ismail, A., 2010. Antioxidative peptides from food

- proteins: a review. *Peptides*, 31(**10**), 1949-1956.
- DOI:10.1016/j.peptides.2010.06.020
- Slizyte, R., Rommi, K., Mozuraityte, R., Eck, P., Five, K. and Rustad, T., 2016. Bioactivities of fish protein hydrolysates from defatted salmon backbones. *Biotechnology Reports*, 11, 99-109. DOI:10.1016/j.btre.2016.08.003
- Y., Thiansilakul, Benjakul, S. and F., 2007. Compositions, Shahidi, functional properties and antioxidative activity of protein hydrolysates prepared from round scad (Decapterus Food Chemistry, 103(4), maruadsi). 1385-1394.
 - DOI:10.1016/j.foodchem.2006.10.055
- Wu, H.C., Chen, H.M. and Shiau, C.Y., 2003. Free amino acids and peptides as related to antioxidant properties in protein hydrolysates of mackerel (*Scomber austriasicus*). Food Research International, 36(9-10), 949-957. DOI:10.1016/S0963-9969(03)00104-2
- You, L., Zhao, M., Cui, C., Zhao, H. and Yang, B., 2009. Effect of degree of hydrolysis on the antioxidant activity of loach (Misgurnus anguillicaudatus) protein hydrolysates. Innovative Food Science & Emerging Technologies, 10(2), 235-240.
 - DOI:10.1016/j.ifset.2008.08.007
- **Zayas, J.F., 1997.** Foaming properties of proteins. In Functionality of proteins in food. Springer, Berlin, Heidelberg. pp. 260-309. DOI:10.1007/978-3-642-59116-7_6