Research Article



Effect of dwarf elder (Sambucus ebulus) to improve hemato-immunological and biochemical parameters and resistance against Yersinia ruckeri in rainbow trout (Oncorhynchus mykiss)

Ghiasi M.1*; Binaii M.1; Mirmazlomi S.1; Adel M.2; Khara H.3

Received: January 2023 Accepted: June 2023

Abstract

The present study examined the effects of dietary dwarf elder (Sambucus ebulus) on hemato- immunological and biochemical parameters and protection against Yersinia ruckeri in rainbow trout. The fish were divided into four groups (in triplicate) and fed diets supplemented with 0, 2.5, 5, and 10% of the powdered leaf of dwarf elder for 8 weeks. The blood samples were taken on weeks 4 and 8 of the trial. The use of dietary dwarf elder did not significantly affect the red bloo cells (RBC), hematocrit, hemoglobin, mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHC), lymphocytes percentage, glucose, and alanine transaminase (ALT) levels of all the treated groups compared to the control group on weeks 4 and 8. At the same time, the total count of white blood cells (WBC) and the neutrophil percentage in the fish receiving the herb powder were significantly higher than the control group. The highest total protein, albumin, immunoglobulin levels, and respiratory burst activity were observed in 5% group after 4 and 8 weeks. On the 8th week, the lysozyme activity was significantly increased and the triglyceride was significantly decreased in 5 and 10% groups compared to the control group. After 4 and 8 weeks, the treated groups exhibited a significant decrease in the cholesterol and alanine aminotransferase levels compared to the control group. The lowest number of bacterial colonies in the serum antibacterial activity belonged to 5 %, 10 %, 2.5 %, and control groups, respectively. At the end of 8 weeks, the fish were challenged with Yersinia ruckeri and the survival rates of the control, 2.5 %, 5 %, and 10 % groups that were 22.2 %, 60.3 %, 79.5 %, and 68.5 %, respectively. The present findings indicate dwarf elder possesses beneficial dietary effects on immune responses and resistance against Y. ruckeri in rainbow trout.

Keywords: Dwarf elder, *Sambucus ebulus*, *Yersinia ruckeri*, Immunity, Total white blood cell, Lysozyme activity

¹⁻Caspian Sea Ecology Research Center, Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research, Education and Extension Organization (AREEO), Sari, Iran.

²⁻ Faculty of Veterinary Medicine, Shahrekord University, Shahrekord, Iran.

³⁻ Department of Fishery, Lahijan Branch, Islamic Azad University, Lahijan, Iran.

^{*}Corresponding author's Emial: E-mail address: ghiasimaryam4@gmail.com

Introduction

Rainbow trout (Oncorhynchus mykiss) is the second freshwater cultivated species in Iran, preceded by Carp, with total production increasing from 91519 metric tons in 2010 to 190287 metric tons in 2020 (FAO, 2020; IFO, 2021). The aquaculture production has shifted from extensive systems towards more intensified systems due to the increasing demand for seafood. However, the intensification of aquaculture poses many problems including stress in aquatic species, outbreaks of infectious diseases, and inappropriate applications of antibiotics (Moradyan et al., 2012; Yarahmadi et al., 2016). Unfortunately, various investigations have shown that the excessive use of antibiotics has led to a generation of bacteria that are resistant to drugs (Raissy and Moumeni 2016; Aflakian etal., 2022) and accumulation of antibiotics in rainbow trout muscle causes serious concern for public health in Iran (Soltani et al., 2014; Adel et al., 2017; Rafati et al., 2018). The enteric red-mouth disease (ERM) or Yersiniosis is one of the most important bacterial diseases of rainbow trout in Iran caused by Yersinia ruckeri (Tulaby Dezfuly et al., 2022). It causes subcutaneous hemorrhages the corners of the mouth, gums, and tongue, exophthalmia, skin darkness. inflammation, hemorrhage of the lower intestine, and splenomegaly (Kumar et al., 2015; Zorriehzahra et al., 2017). For the first time, versiniosis-like infection was reported from Iranian farmed rainbow trout in 1999 (Soltani et al., 1999), and then, Akhlaghi and Sharifi

Yazdi (2008), recognized *Y. ruckeri* by molecular techniques. In recent years, several studies have been conducted to produce a vaccine against yersiniosis by use of the native bacterial species (auto vaccine) in Iran (Soltani *et al.*, 2016; Erfanmanesh *et al.*, 2022), but this method is still expensive for farmers, and researchers are looking for cheaper and more appropriate ways to control the disease in Iran.

The use of herbal plants has recently deserved growing attention in the aquaculture industry due to their biological activities related to secondary metabolites, low costs, and minimal side effects compared to conventional antibiotics (Reverter et al., 2021). Plantenriched diets have been reported to increase disease resistance due to their phytochemical substances and immunomodulatory activities. Therefore, there has been increased interest in the possibility of using medicinal herbs as immunostimulants to enhance immune responses and disease resistance in cultured fish species (Ahmadifar et al., 2021; Alagawany et al., 2020; Elumalai et al., 2020; Firmino et al., 2021).

Several studies mentioned the positive effects of many Iranian medicinal plants such as Mentha piperita (Adel et al., 2016) Ducrosia anethifolia (Dehghan et al., 2016), Urtica dioica (Saeidi asl et al., 2017), Carthamus tinctorius (Zargari et al., 2018), Polygonum minus (Adel et al., 2019), Coriandrum sativum (Naderi Farsani et al., 2019), Zingiber officinale (Soltanian et al., 2019), Quercuse

brantii (Ghafarifarsani et al., 2020), Rhus coriaria (Gharaei et al., 2020), Mentha longifolia (Heydari et al., 2020), Malvae sylvestris (Rashidian et al., 2020), Ziziphora clinopodioides (Oroji et al., 2021), and Capsicum annuum (Firouzbakhsh et al., 2021) on hematobiochemical parameters, immune responses and immune-related gene expression, and growth performance as well as positive effect on oxidative stress resistance against *Y*. ruckeri infection in rainbow trout.

Elderberry or dwarf elder (Sambucus ebulus) is an herbaceous plant, wellknown in Iranian folk medicine and extensively grows in the northern regions of Iran (Aghajanzadeh et al., 2021). It has been reported to possess marked antiemetic and neuroprotective (Fathi et al., 2015), antiviral (Ghaffari et al., 2021), antibacterial (Mahboubi et al., 2012; Rodino et al., 2015; Hashemi al., 2022), antifungal (Rezaei-Moshaei et al., 2021; Mirmazloomi et al., 2022), anti-parasite (Heidari-Kharaji et al., 2019), antioxidant (Karami et al., 2015), anti-mutagenic and anticancer (Saeedi Saravi et al., 2013; Rezaei-Moshaei et al., 2021; Hashemi et al., 2022), anti-inflammatory and antinociceptive activities (Shokrzadeh et al., 2010; Schwaiger et al., 2011; Jabbari et al., 2016), and wound healing effects (Süntar et al., 2010; Babaei et al., 2017).

The most important phytochemical compounds have been found in dwarf elder including phytosterols, anthocyanins, quercetin, phenolic compounds, flavonoids, alkaloids, minerals (especially Ca, Mn, and Fe),

vitamin C, chlorogenic and ursolic acids, carotenoids, and chlorophylls (Ebadi and Hisoriev 2011; Fathi et al., 2015; Jabbari et al., 2017; Kaya et al., 2019; Tasinov et al., 2013, 2021). All these compounds made dwarf elder appropriate candidate for use in folk veterinary medicine. However, there are no reports on the dietary effect of dwarf elder on immune-hematological profiles in rainbow trout. Therefore, this study aimed to investigate the impact of dietary dwarf elder powder biochemical and hemato-immunological and disease resistance parameters against Y. ruckeri in rainbow trout.

Materials and methods

Diet preparation

Four experimental diets were obtained by adding the dwarf elder (dried powdered leaves and stem obtained from Sari, north of Iran) at 0, 2.5, 5 and 10% into a rainbow trout commercial basal diet (Beyza21TM Feed Com, Shiraz, Iran) containing 44.50% protein, 14.20% lipid, 7.62% ash, and 21.90 energy. MJ/kg gross The dietary ingredients were blended with water (100 mL of water per 1 Kg of diet) to form a stiff paste, which passed through a meat grinder (Techno Sanat, Iran) and pelleted to produce 5.0 mm pellets. The experimental diets were air-dried at room temperature (25°C) for 24 h and stored in plastic bags at -4°C until use.

Fish and experimental design

Rainbow trout (92.2±2.2 g initial weight) were obtained from a commercial fish farm in Babol

(Mazandaran province, north of Iran). The health of the fish (changes in behavior and physical appearance) was checked two weeks before starting the experiment. The fish were randomly distributed into 12 fiberglass tanks (1000 L) at 30 fish per tank density (3 tanks per treatment). During the acclimatization period (7 days), the fish were fed with the basal diet. During the experimental time (8 weeks), the feeding rate was 3.5-4% of the body weight and the fish were fed four times a day (07:00, 12:00, 17:00, 22:00). The and water temperature, dissolved oxygen, pH, and electrical conductivity were monitored daily and maintained at 14.4±1.3°C, 7.9±0.5 mg/L, 7.3, and 5733.1±137.1 respectively. MM/cm, Continuous aeration was provided to each tank through an air stone connected to a central air compressor.

Blood sampling

On weeks 4 and 8 after the feeding, 9 fish were randomly sampled from each treatment and were anesthetized by clove extract, a natural alcoholic extract of Eugenia aromatic, at 250 µL/L which was developed at Caspian Sea Ecology Research Center. The blood (~ 2 ml) was drawn from the caudal vein, using a nonheparinized syringe, after they were starved for 24 h. One half of each blood sample was transferred to microtube containing heparin as an anti-coagulant (50 µL) and immediately used for respiratory burst assav and hematological examinations, while the other half was transferred to nonheparinized microtube, placed at room

temperature and allowed to clot for 2 h. The sera were separated by centrifugation at $1500 \times g$ for 20 min and stored at -20°C until use.

Hematological and biochemical assays The red blood cells (RBC) and white blood cells (WBC) counts were determined using Neubauer hemocytometer. The hematocrit (Hct) was measured by microcentrifuge technique, using standard heparinized microhaematocrit capillary tubes (75 mm at $7000 \times g$ for 10 min). The haemoglobin level was analyzed spectrophotometrically at 540 nm by the cyamethaemoglobin method (Blaxhall and Daisley, 1973). The blood indices including mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), and corpuscular mean hemoglobin concentration (MCHC) were calculated according to Seiverd (1964)method. To estimate differential leukocyte percent (lymphocytes and neutrophils), blood smears were prepared, air-dried, fixed in methanol, and stained using May-Giemsa solution (Lee et al., 1998). Biochemical analyses were performed glucose, triglyceride, albumin, cholesterol, aspartate aminotransferase (AST), and alanine aminotransferase (ALT) by commercial kits (Pars Iran) and Azmoon, Tehran, biochemical auto analyzer instrument (Eurolyser, Belgium) (Binaii et al., 2014).

Immunological assays

Lysozyme activity

Serum lysozyme activity was measured using a modified turbidimetric method described by Ellis (1990). Briefly, aliquots (1.75 mL) of Micrococcus lysodeikticus (Sigma, USA) suspension (0.375 mg/mL, 0.05 M PBS, pH 6.2) were mixed with 250 mL of each sample and optical density was measured after 15 and 180 s by spectrophotometer (Biophotometer Eppendorf) at 670 nm. One unit of lysozyme activity was defined as a reduction in absorbance of 0.001/min. The units of lysozyme present in sera were obtained from a standard curve made with hen egg white lysozyme (Sigma, USA).

Respiratory burst activity

The oxidative burst produced by leukocytes of the blood samples was measured by Chemiluminescent assay (CL) (measuring of light emission) as described by Khoshbavar-Rostami *et al.* (2006) using an automated system for CL analysis (LUMI scan Ascent T392, Finland). The results of light emission are expressed in the form of relative light units per second (RLU/s) recorded by the CL assay. CL assays were performed in duplicate and the mean of replicate assays was used in subsequent evaluations.

Serum total protein and total immunoglobulin

The serum total immunoglobulin (Ig) levels were determined according to the method described by Siwicki and Anderson (1993). Briefly, the serum

total protein (TP) content was measured using a microprotein determination method (C-690; Sigma), before and after precipitating down the immunoglobulin molecules, using a 12% solution of polyethylene glycol (Sigma).

Serum antibacterial activity

In vitro bactericidal activity of the serum samples was examined against Y. ruckeri (KC291153, obtained from Faculty of Veterinary Medicine, University of Tehran, Iran) which was prepared as lyophilized stock. The isolate was cultured overnight in nutrient broth (Merck, Germany) and centrifuged at $1500 \times g$ for 20 min. The pellets were washed and re-suspended in PBS to an OD540 of 1.0 (5.32 \times 10⁷ CFU/mL). This bacterial suspension was serially diluted (1:10) with PBS five times. Serum bactericidal activity determined by incubating 2 mL of this diluted Y. ruckeri suspension with 20 ml of serum in a micro-vial for 1 h at 37 °C. In the bacterial control group, PBS replaced the serum. After incubation, the number ofviable bacteria was determined by counting the colonies grown on a nutrient agar plate (duplicate) for 24 h at 37°C (Rao et al., 2006).

Bacteria and challenge

Y. ruckeri (KC291153) was used for the challenge experiment. The bacteria were grown in LB-medium (Oxoid LP0042, Tryptone 10 g, Oxoid LP0021Yeast-extract 5 g, NaCl 5 g, H2O to 1000 ml, pH 7.4) at 20°C for 36 h and enumerated as colony forming units (CFU) by the

spread plate method on blood agar (40 g/L Oxoid blood agar base CM0055) with 5% bovine blood) (Raida and Buchmann, 2009). At the end of 8 weeks, eighty fish (ten fish in duplicate from each experimental group) was randomly selected and anesthetized by the clove extract (250 µL/L). The fish infected by intraperitoneal were injection with 6×10^7 CFU/fish in 100 μ L ringer. The mortalities were monitored for 14 days, and dead fish were dissected to confirm the presence of the pathogen through microbiological examination of the head kidney. The survival rate (SR) was calculated to evaluate the efficacy of the plant powder as $SR=(Nt / N0) \times 100$. Here, N0 is the initial number of fish and Nt is the final number of fish (Nya and Austin, 2009).

Statistical analysis

The effects of dietary dwarf elder on the hemato-biochemical parameters and immunity of rainbow trout were analyzed using SPSS (Ver. 20) and significant differences between the groups were determined by one-way analysis variance (ANOVA). of Duncan's multiple range tests were performed when the differences were significant. Mean values were considered significantly different at p<0.05. Data are expressed as mean values±SE.

Results

Hematological and biochemical profiles
The effect of dietary dwarf elder on
hematological and biochemical
parameters in weeks 4 and 8 is

summarized in Tables 1 and 2. The statistical analysis of data showed that there were no statistically significant changes in the RBC, Hb, Hct, MCV, MCH, MCHC, lymphocytes, glucose, and AST values in all treated groups compared to the control group on weeks 4 and 8 (p>0.05). At the end of week 4th, the WBC count significantly increased in the fish fed with 5 and 10 % dwarf elder when compared to the control and 2.5% groups (p<0.05). Also at the same time, there was a significant change in the neutrophil percentage of all treated groups compared to the control group (p<0.05). At the end of week 8th, the WBC counts of all herbal-enriched diets were significantly higher than the control group (p<0.05) and the highest WBC count was obtained in 5% group. At the same time, neutrophil percentage was also higher in the fish fed with the herbal enriched diets compared to the control group.

The highest levels of cholesterol and ALT were noticed in the control group at the end of weeks 4 and 8 (p<0.05). No significant difference was observed between the treated and control groups at the end of the 4 weeks in the triglyceride (p>0.05), whereas level it was significantly decreased in 5 and 10% herbal enriched diets on week 8th (p<0.05). Also, the TP and albumin levels were significantly increased in the fish fed with 5% dwarf elder when compared to the control group at the end of weeks 4 and 8 (p < 0.05).

[Downloaded from jifro.ir on 2025-11-26]

Immunological profile

The results showed that the serum total Ig value of 5% dwarf elder group was significantly higher than the control and 2.5% groups at the end of the week 4th. After 8 weeks, the serum total Ig value was increased in the fish fed with the

herbal-enriched diets compared to the control group, but just 5% group had a significant difference compared to the control group (Fig. 1).

Table 1: Average hematological parameters for rainbow trout fed with 0, 2.5,5 and 10% dwarf elder of feed for 4 and 8 weeks.

	Percent									
Sampling time	dwarf elder of feed	RBC (10 ⁶ mL ⁻¹)	WBC (10 ³ mL ⁻¹)	Htc (%)	$\mathbf{Hb}_{(gdL^{\cdot 1})}$	MCV (fL)	MCH (pg)	MCHC (gdL ⁻¹)	Lymphocytes (%)	Neutrophils (%)
After 4 weeks	10	1.11± 0.13 ^a	13.08± 0.41 ^b	36.44± 2.69 ^a	6.11± 0.60 ^a	343.92± 18.53 ^a	53.90± 1.95°	15.73± 0.53 ^a	93.66± 1.32°	6.33± 1.32 ^b
	5	1.08± 0.15 ^a	13.64± 0.62 ^b	37.01 ± 2.54^{a}	6.13± 0.36 ^a	350.83± 9.16 ^a	55.24± 1.23 ^a	15.20 ± 0.66^{a}	92.44 ± 1.28^{a}	$\substack{7.55\pm\\1.28^{\mathbf{b}}}$
	2.5	1.15± 0.9ª	10.42± 0.87 ^a	36.33± 2.95 ^a	6.06± 0.63 ^a	337.58± 8.41 ^a	56.11± 1.35 ^a	15.34 ± 0.43^{a}	94.33 ± 0.94^{a}	$\substack{5.66\pm\\0.94^{\text{b}}}$
	0	0.99± 0.1ª	9.56± 0.58 ^a	35.66± 1.87 ^a	5.87± 0.68 ^a	343.76± 8.06 ^a	57.81± 1.54 ^a	15.78 ± 0.75^{a}	98.22 ± 0.46^{a}	$1.77 \pm 0.46^{\mathbf{a}}$
After 8 weeks	10	1.17± 0.14 a	16.66± 0.85 ^{bc}	42.33± 4.12 ^a	6.58± 0.75 ^a	362.43± 11.41 ^a	60.30± 4.10 ^a	16.92± 1.13 ^a	92.66± 1.46 ^a	7.33± 1.46 ^b
	5	1.26± 0.24 ^a	18.02± 1.37°	42.44 ± 3.46^{a}	6.65± 0.62 ^a	364.36± 9.77 ^a	$\begin{array}{c} 58.01 \pm \\ 1.84^{a} \end{array}$	$16.53 \pm \\0.81^a$	91.6± 9.16 ^a	7.35± 1.05 ^b
	2.5	1.17± 0.21 ^a	11.54± 0.77 ^b	41.55± 2.87 ^a	6.45± 0.65°	366.36± 6.77 ^a	56.67± 1.09 ^a	$16.83 \pm \\ 0.46^{\mathrm{a}}$	93.66± 1.14 ^a	6.33± 1.12 ^b
	0	1.10± 0.14 ^a	0.95 ± 0.25^{a}	40.22 ± 3.59^{a}	6.17± 0.84 ^a	367.90± 14.96 ^a	57.08± 1.66°	16.60± 0.61 ^a	96.66± 4.6 ^a	3.33± 1.02 ^a

RBC, red blood cells; WBC, white blood cells; Ht, hematocrit; Hb, hemoglobin concentration; MCV, mean cell volume; MCH, mean cell hemoglobin; MCHC, mean cell hemoglobin concentration. Data represent as mean \pm SE. ata in the same columns with different superscript are significantly different (p<0.05).

Table 2: Biochemical indices of rainbow trout after feeding with different percent of dwarf elder for 4 and 8 weeks.

Sampling time	Percent dwarf elder of feed	Total protein (gdL ⁻¹)	Albumin (gdL ⁻¹)	Glucose (gdL ⁻¹)	Triglyceride (mgdL ⁻¹)	Cholesterol (mgdL ⁻¹)	ALT (IUdL ⁻¹)	AST (IUdL-1)
After 4 weeks	10	3.74 ± 0.77^{ab}	2.14± 0.13 ab	94.26± 14.01 ^a	243.62 ± 25.25^{a}	224.38± 10.15 ^a	9.27± 0.93 ^a	223.22 ± 21.16^{a}
	5	$\substack{4.15\pm\\1.07^{\textbf{b}}}$	$\substack{2.96 \pm \\ 0.11^{\text{b}}}$	95.02 ± 10.67^{a}	229.84± 17.16 ^a	221.84 ± 11.99^{a}	11.93± 1.24 ^a	223.97 ± 15.01^{a}
	2.5	$\begin{array}{c} 3.72 \pm \\ 0.78^{ab} \end{array}$	$\begin{array}{c} 2.23 \pm \\ 0.14^{ab} \end{array}$	95.84± 8.56 ^a	$209.77 \pm \\ 12.26^{a}$	237.14± 15.17 ^a	10.38± 1.32 ^a	226.85 ± 21.68^{a}
	0	3.47 ± 0.32^{a}	$\frac{2.02\pm}{0.18^{a}}$	100.08± 9.39 ^a	242.24± 17.41 ^a	283.93± 8.95 ^b	16.33± 1.81 ^b	258.12± 15.07 ^a

Table 2 cont	inued:							
	10	$\begin{array}{c} 4.31 \pm \\ 0.31^{ab} \end{array}$	$\begin{array}{c} 2.74 \pm \\ 0.33^{\text{b}} \end{array}$	92.35± 6.94 ^a	215.45± 16.05 ^a	234.02± 11.82 ^a	6.21± 1.43°	212.88± 15.19 ^a
After 8	5	$\substack{4.85\pm\\0.29^{\text{b}}}$	2.97± 0.31 ^b	93.33± 2.41 ^a	211.35 ± 13.08^{a}	258.61 ± 11.24^{a}	6.61 ± 0.46^{a}	221.71± 18.88 ^a
weeks	2.5	$\substack{4.22\pm\\0.35^{ab}}$	2.63 ± 0.42^{b}	94.06± 4.21 ^a	$282.05 \pm 8.42^{\mathbf{b}}$	243.82 ± 12.33^{a}	$6.38 \pm \\ 0.58^{a}$	229.92 ± 16.38^{a}
	0	$\begin{array}{c} 3.71 \pm \\ 0.12^{\mathbf{a}} \end{array}$	2.08 ± 0.45^{a}	101. 21± 4.24 ^a	294.81± 21.51 ^b	309.74± 7.78 ^b	16.75± 1.62 ^b	243.25± 11.93 ^a

ALT, alanine aminotransferase; AST, aspartate aminotransferase. Data represent as mean \pm SE. at a in the same columns with different superscript are significantly different (p<0.05).




Figure 1: Serum total immunoglobulin (Ig) levels of rainbow trout fed with 0, 2.5, 5 and 10% dwarf elder. Data represent the mean±S.D. Data in the same row with different superscript are significantly different (p < 0.05).

The lysozyme activity of rainbow trout was not affected by different doses of dietary dwarf elder powder until the week 4^{th} when compared to the control group (p>0.05), but it was significantly

enhanced in the fish fed with 5% and 10% dwarf elder powder on week 8^{th} when compared to the control group (p<0.05) (Fig. 2).




Figure 2: Lysozyme activity of rainbow trout fed with 0, 2.5, 5 and 10% dwarf elder. Data represent the mean \pm S.D. Data in the same row with different superscript are significantly different (p<0.05).

The respiratory burst activity was significantly enhanced in the fish fed with 5 and 10% dwarf elder powder on week 4^{th} when compared to the control and 2.5% groups (p<0.05), but it was

significantly enhanced in all treated groups on week 8^{th} when compared to the control group (p<0.05). The highest respiratory burst activity was found in 5% group (Fig. 3).




Figure 3: Respiratory burst activity of rainbow fed with 0, 2.5, 5 and 10% dwarf elder. Data represent the mean \pm S.D. Data in the same row with different superscript are significantly different (p<0.05).

The results of the serum bactericidal activity (based on colony count) showed that 5% group had the highest value

compared to the other groups and the control group had the lowest serum bactericidal activity (p<0.05) (Fig. 4).




Figure 4: Serum antibacterial activity of rainbow trout after 8 weeks feeding of dietary dwarf elder. Data represent the mean \pm S.D. Data in the same row with different superscript are significantly different (p<0.05).

Disease resistance

The use of dietary dwarf elder powder led to a marked reduction in the mortality rate after challenging with Y. survival ruckeri infection. The percentages were 22.2%, 60%, 79.5%, and 68.53% in the control group, 2.5, 5, dwarf 10% and elder powder, respectively. The survival rate of the treated groups did not show any sign of disease at the end of the feeding experiment.

Discussion

Immunostimulants have an important role to regulate fish health and prevent fish losses due to infectious diseases and medicinal herbs have been extensively used as an immunostimulant to enhance immunity and fish increase resistance against infectious diseases (Liu *et al.*, 2022). In this study, we determined the effect of dwarf elder (*Sambucus ebulus*) powdered leaf supplementation of

rainbow trout diet on immuno hematological responses and resistance against Yersinia ruckeri. In our study, the results showed that WBC levels in the 5% and 10% dwarf elder enriched diet treated group significantly increased in rainbow trout on week 4th, but at the end ofweeks. **WBC** levels significantly rose in all groups treated by dwarf elder. WBCs monitoring can reflect the general immune system status in fish and many investigations have shown that medicinal plants could improve and increase the total WBC of rainbow trout (Adel et al., 2016, 2019; Farsani et al., 2019; Gharaei et al., 2020; Heydari et al., 2020; Rashidian et al., 2020; Firouzbakhsh et al., probably as a result of the stimulation of erythropoiesis and leukopoiesis by plant active compounds (Ahmed, 2020).

Teleost neutrophils are professional phagocytes and the key components of the innate immune response. They ingest pathogens and produce reactive oxygen species and release toxic substances intracellular granules, becoming very potent killers in fish blood (Kordon et al., 2018). The use of dwarf elder enriched diet significantly rose the neutrophil percentage of all the treatment groups on weeks 4th and 8th. Joshi et al. (2022) found that phytosterol components (especially β-sitosterol) the neutrophils increase count in humans. Shokrzadeh and Saeedi Saravi (2010) described that β -sitosterol is one of the dwarf elder leaf's phytosterol components and it could be explained by the rise of neutrophil count in our investigation.

Lysozyme is normally used as an indicator of fish's non-specific immune response in defense against bacterial pathogens. Its bactericidal action is due to destroy of Gram-positive and some Gram-negative bacteria. Moreover, it promotes phagocytosis of macrophages and polymorphonuclear leucocytes (by acting as an opsonin) and complement system in marine and freshwater fish (Uribe et al., 2011; Alagawany et al., 2020). In the present study, the lysozyme activity of fish was not changed after feeding with different doses of dwarf elder at the end of week 4th, but it was significantly increased in the 5% and 10% groups after 8 weeks when compared to the control. The increase in lysozyme activity within the dwarf elder groups of rainbow trout had been confirmed in previous studies with other Iranian herbs including Mentha piperita, Polygonum minus (Adel et al., 2016, 2019), Capsicum annuum

(Firouzbakhsh et al., 2021), Aloe vera (Alishahi et al., 2017), Nigella sativa, Echinacea angustifolia (Fadeifard et al., 2018), Zingiber officinale (Fadeifard et al., 2018; Soltanian et al., 2019) Verbascum speciosum (Nofouzi et al., 2017), Urtica dioica (Saeidi asl et al., 2017), Echinacea purpurea (Sharif Rohani et al., 2016) Allium sativum (Zaefarian et al., 2017). Various studies on other Iranian medicinal plants have shown that the flavonoid compounds are effective in elevating of lysozyme activity of rainbow trout (Firouzbakhsh et al., 2021; Hosseini Shekarabi et al., 2021) and there is information that dwarf elder leaves are rich in these compounds (Fathi et al., 2015).

The respiratory burst is a potent antimicrobial defense mechanism of phagocytes. It is activated after the stimulation of the plasma membrane during phagocytosis and triggers the production of reactive oxygen species (Bulfon et al., 2018). The results showed that the respiratory burst activity in fish fed by the 5% and 10% dwarf elder diet significantly rose compared to two other groups at the end of 4 weeks, but at the end of 8 weeks, the respiratory burst activity significantly rose in all groups treated by dwarf elder and the highest respiratory burst activity were observed in 5% group at both times. The same results were reported in rainbow trout after receiving the other Iranian herbs (Adel et al., 2016, 2019; Saeidi asl et al., 2017; Heydari et al., 2020). Akbay et al., (2003) have shown that the flavonoid glycosides of nettle can cause degranulation modulate the and

oxidative burst of appropriately stimulated human neutrophils. Five flavonoid glycosides were recognized in dwarf elder (Zahmanov *et al.*, 2015) and it seems that the same mode of action could happen in the respiratory burst activity of rainbow trout after receiving dwarf elder.

The components of TP (globulin and albumin) are mainly secreted by the liver and can be used as an indicator of fish health to determine physiological changes following different feeding herbal plants (Hosseini Shekarabi et al., 2021; Farsani et al., 2019; Adel et al., 2016, 2019). At the end of the 4 and 8 weeks, the total serum protein and albumin concentrations were increased in all treatment groups, but this increase was only significant in the 5% group compared to the control. The total Ig of the fish after receiving dwarf elder was significantly increased in the 5% group compared to the 2.5% group and control on the week 4th and just control on the week 8th. Albumin is the most prevalent protein in the blood circulation of fish and functions as a carrier protein for a variety of nutrients, metabolites, and xenobiotics. Moreover, the increase in the levels of globulins in fish is thought to be associated with a stronger immune response and it is among the most frequently tested immune parameters in herbal-supplemented diets in fish (Wang et al., 2017; Awad and Awaad, 2017).

The glucose and AST of rainbow trout were not affected by different levels of dwarf elder, but its usage significantly decreased the cholesterol, triglyceride, and ALT levels in the treatment groups compared to the control. Various studies have shown that dwarf elder is rich in phenolic compounds (Jabbari *et al.*, 2017; Tasinov *et al.*, 2013, 2021). Phenols and polyphenolic compounds can serve as antioxidants (Fathi *et al.*, 2015) and improve human blood lipid profile (Ivanova *et al.*, 2014). In our study, it seems that the phenols and polyphenolic compounds have affected and changed cholesterol, triglyceride, and ALT in the treated groups.

Serum bactericidal activity introduced as an indicator of innate immunity in teleost fish. The lowest number of bacterial colonies indicated the efficiency of immune cells in serum to kill the pathogen (Uribe et al., 2011). The results showed that the serum bactericidal activity against Y. ruckeri was significantly increased in the group supplemented with dwarf elder. The enhancement of serum bactericidal activity has been reported previously with other herbs in rainbow trout (Awad et al., 2013; Alishahi et al., 2017). These findings indicate that the bioactive compounds of medicinal herbs may provide some non-specific and specific immune responses in fish.

The results of the survival rate of fish fed with dwarf elder supplementation feeds were higher than those of the control after infection with *Y. ruckeri* and the maximum protection was detected in the group fed with 5% of dwarf elder. Previous works have shown that rainbow trout which received dietary herbs revealed protection against *Y. ruckeri* (Adel *et al.*, 2016, 2019;

Dehghan *et al.*, 2016; Soltanian *et al.*, 2019; Bilen *et al.*, 2020; Ghafarifarsani *et al.*, 2020; Heydari *et al.*, 2020; Firouzbakhsh *et al.*, 2021). The improved immune indices that were tested in the current research could be correlated with the improved resistance against *Y. ruckeri*.

In conclusion, the present results suggested that dwarf elder was found beneficial for rainbow trout when applied as a feed additive. Several hemato-serological and immunological parameters were enhanced in fish that received diets containing dwarf elder at different levels and the best results were found in 5% group compared to the other groups. Thus, it seems that dwarf elder could be introduced as an immunomodulator and improver of hemato-biochemical parameters to aquaculture. But there is still limited information available regarding the possible effects of dwarf elder on fish, and therefore, further research is required.

Acknowledgments

The authors would like to thank the staff of Aquatic Animal Health of Caspian Sea Ecology Research Center for their kind help during the experiment.

References

Adel, M., Pourgholam, R., Zorriehzahra, J. and Ghiasi, M., 2016. Hemato - Immunological and biochemical parameters, skin antibacterial activity, and survival in rainbow trout (*Oncorhynchus mykiss*) following the diet supplemented with

Mentha piperita against *Yersinia* ruckeri. Fish and Shellfish Immunology, 55, 267-273. DOI:10.1016/j.fsi.2016.05.040.

Adel, M., Dadar, M. and Conti, G.O., 2017. Antibiotics and malachite green residues in farmed rainbow trout (*Oncorhynchus mykiss*) from the Iranian markets: A risk assessment. *International Journal of Food Properties*, 20(2), 402–408. DOI:10.1080/10942912.2016.11635 77.

Adel, M., Dawood, M.A.O, Shafiei, S., Sakhaie, F. Hosseini and S.P., 2019. Shekarabi, Dietary Polygonum minus extract ameliorated the growth performance, humoral immune parameters, immune-related expression and resistance against Yersinia ruckeri in rainbow trout (Oncorhynchus mykiss). Aquaculture, 519. 734738. DOI:10.1016/j.aquaculture.2019.734 738.

Aflakian, F., Nikoueian, H., Salimizand, H., Rad, M. and Rafati Zomorodi, M., 2022. Heterogeneity of *Yersinia ruckeri* isolated from rainbow trout (*Oncorhynchus mykiss*) infected with enteric red mouth disease. World's Veterinary Journal, 12(1), 19-27.

Aghajanzadeh, H., Abdolmaleki, M., Ebrahimzadeh, M.A., Mojtabavi, N., Mousavi, T. and Izad, M., 2021. Methanolic extract of *Sambucus ebulus* ameliorates clinical symptoms in experimental type 1 diabetes through anti-inflammatory and immunomodulatory actions. *Cell*

- Journal, 23(4), 465-473. DOI:10.22074/cellj.2021.7287.
- Ahmadifar, E., Pourmohammadi Fallah, H., Yousefi, M., Dawood, M.A.O., Hoseinifar. S.H., Adineh, H., Yilmaz. S., Paolucci, M. and Van Doan, H., 2021. The gene regulatory roles of herbal extracts on the growth, immune system, and reproduction of fish. *Animals*, 11, 2167. DOI:10.3390/ani11082167.
- Ahmed, I., Reshi, Q.M. and Fazio, F., 2020. The influence of the endogenous and exogenous factors on hematological parameters in different fish species: a review. *Aquaculture International*, 28(3), 869-899. DOI:10.1007/s10499-019-00501-3.
- Akbay, P., Basaran, A.A., Undeger, U. and Basaran, N., 2003. In vitro immunomodulatory activity of flavonoid glycosides from *Urtica dioica* L. *Phytotherapy Research*, 17(1), 34-37. DOI:10.1002/ptr.1068.
- Akhlaghi, M. and Sharifi, Y.H., 2008.

 Detection and identification of virulent *Yersinia ruckeri*: the causative agent of enteric redmouth disease in rainbow trout (*Oncorhynchus mykiss*) cultured in Fars province, Iran. *Iran Journal Veterinary Research*, 9(4), 347-352. DOI:10.22099/ijvr.2008.2617.
- Alagawany, M., Farag, M.R., Salah, A.S. and Mahmoud, M.A., 2020. The role of oregano herb and its derivatives as immunomodulators in fish. *Reviews in Aquaculture*, 1–12. DOI:10.1111/raq.12453.

- Alishahi, M., Tulaby Dezfuly, Z., Mesbah, M. and Mohammadian, T., 2017. Effects of Aloe vera crude extract on growth performance and some hemato-immunological indices of *Oncorhynchus mykiss* in farm scale. *Iranian Journal of Veterinary Medicine*, 11, 383-393. DOI:10.22059/ijvm.2017.231790.10 04806.
- Awad, E., Austin, D. and Lyndon, A.L., 2013. Effect of black cumin seed oil (*Nigella sativa*) and nettle extract (Quercetin) on enhancement of immunity in rainbow trout *Oncorhynchus mykiss* (Walbaum). *Aquaculture*, 193-197. DOI:10.1016/j.aquaculture.2013.01. 008.
- **Awad, E. and Awaad, A., 2017.** Role of medicinal plants on growth performance and immune status in fish. *Fish Shellfish Immunology*, 67, 40-54.
 - DOI:10.1016/j.fsi.2017.05.034.
- Babaei, E., Asghari, M.H., Mehdikhani, F., Moloudizargari, M., Ghobadi, E. and Pouya, S.R.H., 2017. The healing effects of herbal preparations from *Sambucus ebulus* and *Urtica dioica* in full-thickness wound models. *Asian Pacific Journal of Tropical*, 7(5), 421-427. DOI:10.1016/j.apjtb.2017.01.013.
- Bilen, S., Ispir, S., Kenanoglu, O.N., Taştan, Y., Güney, K. and Terzi, E., 2020. Effects of Greek juniper (*Juniperus excelsa*) extract on immune responses and disease resistance against *Yersinia ruckeri* in rainbow trout (*Oncorhynchus*)

Downloaded from jifro.ir on 2025-11-26]

- *mykiss*). *Journal of Fish Diseases*, 44(**6**), 729-738. DOI:10.1111/jfd.13293.
- Binaii, M., Ghiasi, M., Farabi, S.M.V., Pourgholam, R., Fazli, H., Safari, R., Alavi, S.E., Taghavi, M.J. and Bankehsaz, Z., 2014. Biochemical hemato-immunological and parameters in juvenile beluga (Huso following the diet huso) supplemented with nettle (Urtica dioica). Fish and Shellfish Immunology, 36(1), 46-51. DOI:10.1016/j.fsi.2013.10.001.
- Blaxhall, P.C. and Daisley, W., 1973.

 Routine haematological methods for use with fish blood. *Journal of Fish Biology*, 5, 771- 781.

 DOI:10.1111/j.1095-8649.1973.tb04510.x.
- **Bulfon, C., Galeotti, M. and Volpatti, D., 2018.** Medicinal plant extracts modulate respiratory burst and proliferation activity of rainbow trout (*Oncorhynchus mykiss*) leukocytes. *Fish Physiology and Biochemistry*, 44, 109–117. DOI:10.1007/s10695-017-0417-5.
- Dehghan, F., Vazirzadeh, A., Soltanian, S., Karami, A. and Akhlaghi, M., 2016. Mortality rate and immune responses of rainbow trout (*Oncorhynchus mykiss*) infected with *Yersinia ruckeri* subsequent to feeding on diet supplemented with *Ducrosia anethifolia* essential oil. *International Journal of Aquatic Biology*, 4(5), 340-344.
- Ebadi, A.G. and Hisoriev, H., 2011.

 Review on distribution of *Sambucus*ebulus L. in the north of Iran.

- American-Eurasian Journal of Agricultural & Environmental Sciences, 10(3), 351-353.
- Ellis, A.E., 1990. In: Stolen JS, Fletcher TC, Anderson DP, Robertson BS, Van Muisvinkel WR, (ed) Lysozyme assay in techniques in fish immunology, Fair Haven, SOS Publications, New Jersey, USA, pp. 101 103.
- Elumalai, P., Kurian, A., Lakshmi, S., Faggio, C., Esteban, M.A. and Ringø, E., 2020. Herbal immunomodulators in aquaculture. Reviews in Fisheries Science and Aquaculture, 29(1), 33-57. DOI:10.1080/23308249.2020.17796 51.
- Erfanmanesh, A., Mohajerfar, T., Nikaein, D., Mokhtari, A. and Beikzadeh, B., 2022. Comparative protection of two antigens (whole-cell and outer membrane vesicle) of Yersinia ruckeri in rainbow trout (Oncorhynchus mykiss). Iranian Journal of Fisheries Sciences, 21(1), 187-201.

DOI:10.22092/ijfs.2022.125884.

- Fadeifard, F., Raissy, M., Jafarian, M., Rouhi Boroujeni, H., Rahimi, M. and Faghani, M., 2018. Effects of black seed (Nigella sativa), ginger (Zingiber officinale) and cone flower (Echinacea angustifolia) on the immune system of rainbow trout, **Oncorhynchus** mykiss. Arauivo Brasileiro de Medicina Veterinaria e Zootecnia. 70 **(1)**. 199-204. DOI:10.1590/1678-4162-8489.
- FAO (Food and Agriculture Organization of the United

- Nations), 2020. The State of World Fisheries and Aquaculture. Sustainability in Action; Food and Agriculture Organization of the United Nations, Rome, Italy.
- Farsani, M.N., Hoseinifar, S.H.. Rashidian, G., Farsani, H.G.. Ashouri, G. and Van Doan, H., 2019. Dietary effects of Coriandrum sativum extract on growth performance, physiological and innate immune responses and resistance of rainbow trout (Oncorhynchus mykiss) against Yersinia ruckeri. Fish and Shellfish 91. 233-240. Immunology, DOI:10.1016/j.fsi.2019.05.031
- Fathi, H., Ebrahimzadeh, M.A., Ziar, A. and Mohammadi, H., 2015. Oxidative damage induced by retching; antiemetic and neuroprotective role of Sambucus Cell ebulus L. Biology and DOI:10.1007/s10565-Toxicology, 015-9307-8.
- Firmino, J.P., Galindo-Villegas, J., Reyes-Lo' pez, F.E. and Gisbert, E., 2021. Phytogenic bioactive compounds shape fish mucosal immunity. *Frontiers in Immunology*, 12, 695973. DOI:10.3389/fimmu.2021.695973.
- Firouzbakhsh, F., Haghparast, S. and Memarzadeh, M.R., 2021. Study on the effects of red pepper (Capsicum annuum) extract on immune responses and resistance of rainbow trout (Oncorhynchus mykiss) juveniles against Yersinia ruckeri. Iranian Journal of Fisheries

- *Sciences*, 20(**6**), 1573-1588. DOI:10.22092/ijfs.2021.350686.0.
- Ghafarifarsani, H., Rashidian, G., Sheikhlar, A., Naderi Farsani, M., Hoseinifar, S.H. and Van Doan, H., 2020. The use of dietary oak acorn extract to improve haematological parameters, mucosal and serum immunity, skin mucus bactericidal activity, and disease resistance in rainbow trout (Oncorhynchus mykiss). *Aquaculture* Research, 2518-2527. 52(6), DOI:10.1111/are.15101.
- Ghaffari, H., Ataei-Pirkooh, A., Mirghazanfari, S.M. and Barati, M., 2021. Inhibition of herpes simplex virus type 1 infection by Sambucus ebulus extract in vitro. Medical Journal of the Islamic Republic of Iran.35, 9. DOI:10.47176/mjiri.35.9.
- Gharaei, A., Shafiei, M., Mirdar Harijani, J., Hassanein, P. and Arshadi, A., 2020. **Immune** and haematological responses parameters changes of rainbow trout (Oncorhynchus mykiss) under effects of dietary administration of sumac (Rhus coriaria L.). Journal of Science Agricultural and Technology, 22(1), 173-186.
- Mizwari, **Z.M.**, Hashemi, Z., Mohammadi-Aghdam, S., Mortazavi-Derazkola, S. and Ebrahimzadeh. M.A., 2022. Sustainable green synthesis of silver nanoparticles using Sambucus ebulus phenolic extract (AgNPs@SEE): optimization and assessment photocatalytic degradation of methyl

orange and them in vitro antibacterial and anticancer activity. *Arabian Journal of Chemistry*, 15, 103525. DOI:10.1016/j.arabjc.2021.103525.

- Heidari-Kharaji, M., Fallah-Omrani, V., Badirzadeh, A., Mohammadi-Ghalehbin, B., Nilforoushzadeh, M.A., Masoori, L., Montakhab-Yeganeh, H. and Zare, M., 2019. Sambucus ebulus extract stimulates cellular responses in cutaneous leishmaniasis. Parasite Immunology, 41(1), e12605. DOI:10.1111/pim.12605.
- Heydari, M., Firouzbakhsh, F. and Paknejad, H., 2020. Effects of *Mentha longifolia* extract on some blood and immune parameters, and disease resistance against yersiniosis in rainbow trout. *Aquaculture*, 15, 734586.

DOI:10.1016/j.aquaculture.2019.734 586.

- Hosseini Shekarabi, S.P., Mostafavi, **Z.S.**, Shamsaie Mehrgan, M., Rajabi Islami, H., 2021. Dietary supplementation with dandelion (Taraxacum officinale) flower extract provides immunostimulation resistance against Streptococcus iniae rainbow infection in trout (Oncorhynchus mykiss). Fish and Shellfish Immunology, 118, 180–187. DOI:10.1016/j.fsi.2021.09.004.
- **IFO, 2021.** Fisheries statistic yearbook. Planning and development office, Fisheries public relation department, Tehran Publications, Iran. pp 34–35. [In Persian].
- Ivanova, D., Tasinov, O. and Kiselova-Kaneva, Y., 2014. Improved lipid

profile and increased serum antioxidant capacity in healthy volunteers after *Sambucus ebulus* L. fruit infusion consumption, *International Journal of Food Science and Nutrition*, 65(6), 740–744.

DOI:10.3109/09637486.2014.89825

- Jabbari, M., Hashempur, M.H., **Emami** Razavi, S.Z., Raeisi Shahraki, H., Kamalinejad, M. and Emtiazy, M., 2016. Efficacy and short-term safety of topical Dwarf elder (Sambucus ebulus L.) versus diclofenac for knee osteoarthritis: a randomized, double-blind, activetrial. J. controlled Ethnopharmacology, 80-86. 188. DOI:10.1016/j.jep.2016.04.035.
- Jabbari, M., Daneshfard, B., Emtiazy, M., Khiveh, A. and Hashempur, M.H., 2017. Biological effects and clinical applications of dwarf elder (Sambucus ebulus L): A Review. Journal of Evidence-Based Complementary & Alternative Medicine. 22(4), 996-1001. DOI: 10.1177/2156587217701322.
- Joshi, T., Patni, K., Belwal, T., Maurya, H. and Kumar, A., 2022. Sterols: beneficial or detrimental for human health. In: Antioxidants Effects in Health, Elsevier pp. 475-484.
- Karami, M., Ale-Nabi, S.S., Nosrati, A. and Naimifar, A., 2015. The protective effect of *Sambucus ebulus* against lung toxicity induced by gamma irradiation in mice.

- Pharmaceutical and Biomedical Research, 1, 48-54.
- Kaya, Y., Kelil Haji, E., Arvas, Y.E. and Aksoy, H.M., 2019. Sambucus ebulus L.: past, present and future. Proceedings of the 2nd International Conference on Biosciences and Medical Engineering (ICBME2019). 2155, 020030-1–020030-7. DOI:10.1063/1.5125534.
- Khoshbavar-Rostami, H.A., Soltani, M. and Hassan, H.M.D., 2006. Immune response of great sturgeon (*Huso huso*) subjected to long-term exposure to sub lethal concentration of organophosphate, diazinion. *Aquaculture*, 256, 88-94. DOI:10.1016/j.aquaculture.2006.02. 041
- Kordon, A.O., Karsi, A. and Pinchuk, L., 2018. Innate immune responses in fish: antigen presenting cells and professional phagocytes. *Turkish Journal of Fisheries and Aquatic Sciences*, 18, 1123-1139. DOI:10.4194/1303-2712-v18_9_11.
- Kumar, G., Menanteau-Ledouble, S., Saleh, M and El-Matbouli., M., 2015, *Yersinia ruckeri*, the causative agent of enteric redmouth disease in fish. *Veterinary Research* 46:103. DOI 10.1186/s13567-015-0238-4.
- Lee, R.G., Foerster, J., Jukens, J., Paraskevas, F., Greer, J.P. and Rodgers, G.M., 1998. Wintrobe's clinical hematology. Lippincott Williams & Wilkins, New York.
- Liu, J., Zhang, P., Wang, B., Lu, Y., Li, L., Li, Y. and Liu, S., 2022. Evaluation of the effects of Astragalus polysaccharides as

- immunostimulants on the immune response of crucian carp and against SVCV in vitro and in vivo. Comparative Biochemistry & Physiology Part C: Toxicology and Pharmacology, 253, 109249. DOI:10.1016/j.cbpc.
- Mahboubi, A., Kamallnejad, M., Shalviri, M., Karbasi, Z., Jafariazar, Z. and Asgharian, R., 2012. Evaluation of antibacterial activity of three Iranian medicinal plants. *African Journal of. Microbiology Research*, 6(9), 2048-2052. DOI:10.5897/AJMR11.1281.
- Mirmazloomi, S., Ghiasi, M. and Khosravi, A.R., 2022. Chemical composition and in vitro antifungal activity of *Sambucus ebulus* and *Actinidia deliciosa* on the fish pathogenic fungus, *Saprolegnia parasitica*. *Aquaculture International*, 30(2), 1037-1046. DOI: 10.1007/s10499-022-00835-5
- Moradvan, H., Karimi, Gandomkar, H., Sahraeian, M.R., Ertefaat, S. and Hosseinzadeh Sahafi, H., 2012. The effect of density stocking on growth parameters and survival rate of rainbow trout alevins (Oncorhynchus mykiss). World Journal of Fisheries, 4(5), 480-485. DOI:5829/idosi.wjfms.2012.04.05.6 4143.
- Naderi Farsani, M., Hoseinifar, S.H., Rashidian, G., Ghafari Farsani, H., Ashouri, G. and Van Doan, H., 2019. Dietary effects of *Coriandrum sativum* extract on growth performance, physiological and

- innate immune responses and resistance of rainbow trout (Oncorhynchus mykiss) against Yersinia ruckeri. Fish and Shellfish Immunology, 91, 233-240.
- Nofouzi, K., Aghapour, M., Ezazi, A., Sheikhzadeh, N., Tukmechi, A., Khordadmehr, M., Akbari, M., Tahapour, K. and Mousavi, M., 2017. **Effects** of Verbascum speciosum on growth performance, intestinal histology, immune system biochemical parameters and rainbow (Oncorhynchus trout mykiss). Turkish Journal of Fisheries and Aquatic Sciences, 17, 145-152. DOI:10.4194/1303-2712-v17_1_17.
- Nya, E.J. and Austin, B., 2009. Use of garlic, *Allium sativum*, to control *Aeromonas hydrophila* infection in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases*, 32, 963–970. DOI:10.1111/j.1365-2761.2009.01100.x
- Oroji, E., Shamsaie Mehrgan, M., Rajabi Islami, H. and Sharifpour, I., 2021. Dietary effect of Ziziphora clinopodioides extract on zootechnical performance, immune and disease resistance response, against Yersinia ruckeri in Oncorhynchus mykiss. Aquaculture 21, Reports, 100827. DOI:10.1016/j.agrep.2021.100827
- Rafati, L., Ehrampoush, M.H., Mokhtari, M., Sohrabi, A., Shirazi, S., Mahvi, A.H. and Momtaz, S.M., 2018. The analysis of oxytetracycline residue in tissues of cultured rainbow trout (*Oncorhynchus Mykiss*). *Health*

- *Scope*, 7(**2**), e57495. DOI:10.5812/jhealthscope.57495.
- Raida, M.K. and Buchmann, K., 2009.
 Innate immune response in rainbow trout (*Oncorhynchus mykiss*) against primary and secondary infections with *Yersinia ruckeri* O1.

 Developmental and Comparative Immunology, 33, 35–45.
 DOI:10.1016/j.dci.2008.07.001.
- Raissy, M. and Moumeni, M., 2016.

 Detection of antibiotic resistance genes in some *Lactococcus garvieae* strains isolated from infected rainbow trout. *Iranian Journal of Fisheries Sciences*, 15(1), 221-229.
- Rao, Y.V., Das, B.K., Jyotyrmayee, P. and Chakrabarti, R., 2006. Effect of *Achyranthes aspera* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. *Fish and Shellfish Immunology*, 20, 263-273. DOI:10.1016/j.fsi.2005.04.006
- Rashidian, G., Kajbaf, K., Prokić, M.D. and Faggio, C., 2020. Extract common mallow (Malvae sylvestris) enhances growth, immunity, and resistance of rainbow (Oncorhynchus trout mykiss) fingerlings against Yersinia ruckeri Fish infection. and Shellfish Immunology, 96, 254 261. DOI:10.1016/j.fsi.2019.12.018.
- Reverter, M., Tapissier-Bontemps, N., Sarter, S., Sasal, P. and Caruso, D., 2021. Moving towards more sustainable aquaculture practices: a meta-analysis on the potential of plant-enriched diets to improve fish growth, immunity and disease

- resistance. *Reviews in Aquaculture*, 13(1), 537-555.
- Rezaei-Moshaei, M., Dehestani, A., Bandehagh, A., Pakdin-Parizi, A., Golkar, M. and Heidari-Japelaghi, Recombinant pebulin R., 2021. protein, type 2 ribosomeinactivating protein isolated from dwarf elder (Sambucus ebulus L.) shows anticancer and antifungal activities in vitro. *International* Journal of Biological Macromolecules, 174, 352-36. DOI:10.1016/j.ijbiomac.2021.01.129
- Rodino, S., Butu, A., Petrache, P., Butu, M., Dinu-Pîrvu, C. and Cornea, C.P., 2015. Evaluation of the antimicrobial and antioxidant activity of *Sambucus ebulus* extract. *Farmacia*, 63(5), 751-754.
- Saeedi Saravi, S. S., Shokrzadeh, M. and Hosseini Shirazi, F., 2013. Cytotoxicity of *Sambucus ebulus* on cancer cell lines and protective effects of vitamins C and E against its cytotoxicity on normal cell lines. *African Journal of Biotechnology*, 12(21), 3360-3365. DOI:10.5897/AJB09.1577.
- Saeidi asl, M.R., Adel, M., Caipang, C.M.A. and Dawood, M.A.O., 2017. Immunological responses and disease resistance of rainbow trout (*Oncorhynchus mykiss*) juveniles following dietary administration of stinging nettle (*Urtica dioica*), *Fish and Shellfish Immunology*, 71, 230-238. DOI:10.1016/j.fsi.2017.10.016.
- Schwaiger, S., Zeller, I., Pölzelbauer, P., Frotschnig, S., Laufer, G.,

- Messner, B., Pieri, V., Stuppner, H. and Bernhard, D., 2011. Identification and pharmacological characterization of the anti-inflammatory principal of the leaves of dwarf elder (*Sambucus ebulus* L.). J. *Ethnopharmacology*, 133(2), 704-709. DOI:10.1016/j.jep.2010.10.049.
- **Seiverd, C.E., 1964.** Hematology for Medical Technologists. Journal of Medical Education 39(9), p 867.
- **Seiverd, C.E., 1964.** Hematology for medical technologists. PA: Lea and Febiger, Philadelphia, US.
- Sharif Rohani, M, Pourgholam, R. and Haghighi, M., 2016. Evaluation the effects of different levels of *Echinacea purpurea* extract on the immunity responses, biochemical and hematological indicesand disease resistance against *Streptococcus iniae* in rainbow trout (*Oncorhynchus mykiss*). *Iranian Journal of Aquatic Animal Health*, 2(2), 1-13.
- Shokrzadeh, M. and Saeedi Saravi, S.S., 2010. The chemistry, pharmacology and clinical properties of *Sambucus ebulus*: A review. *Journal of Medicinal Plants*, 4(2), 95-103. DOI: 10.5897/JMPR09.026.
- Siwicki, A.K. and Anderson, D.P., 1993. Nonspecific defense mechanisms assay fish: II. in killing activity **Potential** of neutrophils and macrophages, lysozyme activity in serum and organs and total immunoglobulin level in serum. In: Siwicki AK, Anderson DP, Waluga J (ed). Fish disease diagnosis and prevention

- methods, Olsztyn, Poland, pp 105-112.
- Soltani, M., Fadaii Fard, F. and Mehrabi, M.R., 1999. First report of a Yersiniosis-like infection in Iranian Farmed Rainbow Trout. Bulletin of the European Association of Fish Pathologists, 19(4), 173.
- Soltani, M., Pirali, E., Rasoli, A., Shafiei, S., Kakoolaki, S. and Shams, G., 2014. Antibiotic residuals in some farmed rainbow trout (*Oncorhynchus mykiss*) of market size in Iran. *Iranian Journal of Aquatic Animal Health*, 1(1), 71-77.
- Soltani, M., Mokhtari, A., Mirzargar, S.S., Taherimirghaed, A., Zargar, A., Shafiei, S. and Hosseini-Shekarabi, S.P., 2016. Efficacy and immune response of intraperitoneal vaccination of rainbow trout (Oncorhynchus mykiss) with Yersinia ruckeri bacterin formulated with MontanideTM ISA 763 AVG adjuvant. Bulletin of the European Association of Fish Pathologists, 36(6), 225-236.
- Soltanian, M., Faghani Langrodi, H. and Mohammad Nejad, M., 2019. The use of Zingiber officinale extract against Yersinia ruckeri and its effects on the antioxidant status and immune response in Oncorhynchus mykiss. International Journal of Aquatic Biology, 7(5), 301-314. DOI:10.22034/ijab.v7i5.630.
- Süntar, I.P., Küpeli Akkol, E., Yalc, F.N., Koca, U., Keleş, H. and Yesiladad, E., 2010. Wound healing potential of *Sambucus ebulus* L.

- leaves and isolation of an active component, quercetin 3-O-glucoside. *Ethnopharmacology*. 129, 106–114. DOI:10.1016/j.jep.2010.01.051.
- Tasinov, O., Kiselova-Kaneva, Y. and Ivanova, D., 2013. Sambucus ebulus
 from traditional.
 DOI:10.14748/ssm. v45i2.319.
- Tasinov, O., Dincheva, I., Badjakov, I., Kiselova-Kaneva, Y., Galunska, B., Nogueiras, R. and Ivanova, D., 2021. Phytochemical composition, anti-inflammatory and ER stress-reducing potential of *Sambucus ebulus* L. fruit extract. *Plants*, 10, 2446. DOI:10.3390/plants10112446.
- Tulaby Dezfuly Z, Alishahi M, Ghorbanpoor M, Tabandeh M, Mesbah M., 2022. Effect of extracellular proteins (ECP) on protective efficacy of Yersiniosis vaccine in juvenile rainbow trout (Oncorhynchus mykiss). Iranian Journal of Fisheries Sciences, 21(5), 1166-1179.

DOI: 10.22092/ijfs.2022.127765

- **Uribe, C., Folch, H., Enriquez, R. and Moran, G., 2011.** Innate and adaptive immunity in teleost fish: a review. *Veterinarni Medicina*, 56(10), 486–503.
- Wang, W., Sun, J., Liu, C. and Xue, Z., 2017. Application of immunostimulants in aquaculture: current knowledge and future perspectives. *Aquaculture Research*, 48(1), 1–23. DOI:10.1111/are.13161.
- Yarahmadi, P., Kolangi Miandare, H., Fayaz, S. and Caipang, C.M.A., 2016. Increased stocking density causes changes in expression of

selected stress and immune related innate immune genes, humoral parameters and stress responses of Rainbow (Oncorhynchus trout mykiss). Fish and Shellfish 43-53. Immunology, 48. DOI:10.1016/j.fsi.2015.11.007.

Zaefarian, A., Yeganeh, S. and Adhami, B., 2017. Dietary effects of garlic powder (*Allium sativum*) on growth, blood indices, carcass composition, and lysozyme activity in brown trout (*Salmo caspius*) and resistance against *Yersinia ruckeri* infection. *Aquaculture International*, 25(6), 1987-1996. DOI:10.1007/s10499-017-0169-3.

Zahmanov, G., Alipieva, K., Denev, P., Todorov, D., Hinkov, A., Shishkov, S., Simova, S. and Georgiev, M.I., 2015. Flavonoid glycosides profiling in dwarf elder

fruits (*Sambucus ebulus* L.) and evaluation of their antioxidant and anti-herpes simplex activities. *Industrial Crops and Products*, 63, 58-64.

DOI:10.1016/j.indcrop.2014.10.053.

Zargari, A., Mazandarani, M. and Hoseini, S.M., 2018. Effects of safflower (Carthamus tinctorius) extract on serum antibacterial activity of rainbow trout (Oncorhynchus mykiss) against Aeromonas hydrophila, Streptococcus iniae and Yersinia ruckeri. International Journal of Aquatic Biology, 6(1), 1-7. DOI:10.22034/ijab.y6i1.392.

Zorriehzahra, M.J., Adel, M. and Torabi Delshad, S., 2017. Enteric redmouth disease: Past, present and future: A review. *Iranian Journal of Fisheries Sciences*, 16(4), 1135-1156.