

Research Article



Comparative effects of Ovaprim™ and Ovapass™ hormones on some reproductive characteristics of *Schizothorax zarudnyi*

Mohammadi Avizi B.¹; Arshadi A.^{1*}; Rahdari A.²

Received: January 2023

Accepted: April 2023

Abstract

This study was conducted to compare the effectiveness of Ovaprim™ and Ovapass™ hormones to induce spawning in *Schizothorax zarudnyi*. The female (n=48; 1535.12±180.09 g) and male broodstocks (n=53; 752.00±48.30 g) were randomly allocated in 4 experimental groups including T1: females which received 0.2, 0.5 and 0.5 mL/kg body weight (BW) Ovaprim with 24-h time intervals, T2: females which treated with 0.5 and 0.5 mL/kg BW Ovapass with 12-h time intervals, T3: females were injected with 0.2, 0.5 and 0.5 mL/kg BW Ovapass with 24-h time intervals, and T4: females were injected with a combination of Ovaprim (0.2 mL/kg) and Ovapass (0.5 and 0.5 mL/kg BW) with 24-h time intervals. The amount of injection in the first stage was 10%, the second stage was 90%, and the male fish were simultaneously injected with the second stage of the females at 0.3 mL/kg. The results showed that the highest rate of fertilization was detected in treatments 2 and 4 (93.33% and 92.33%, respectively), which had a significant difference with other groups ($p<0.05$). No significant differences were observed between the treatments in the indices of relative and working fecundity and dry egg diameter ($p>0.05$). The lowest and highest latency period was observed in treatments 2 (27.33± 0.66 h) and 1 (36.20±3.77 h), respectively ($p<0.05$). The results of this study clearly suggested that the Iranian Ovapass hormone is highly effective for *S. zarudnyi* reproduction and might be successfully replace with Ovaprim.

Keywords: *Schizothorax zarudnyi*, Artificial reproduction, Ovaprim, Ovapass, Fertilization

1-Department of Fisheries, Faculty of Natural Resources, University of Zabol, Zabol, Iran.

2-Department of Aquatic Sciences, Hamoun International Wetland Institute, Research Institute of Zabol, Zabol, Iran.

*Corresponding author's Email: arshadi@uoz.ac.ir

Introduction

The Hamun mahi, *Schizothorax zarudnyi*, is native to the Sistan basin that is located in the east of Iran (Oveisi and Kavosh 2021) and the species exists only in this region in Iran. In fact, this fish belongs to the Hamun international wetland, which is shared between Iran and Afghanistan (Vekerdy *et al.*, 2006). Several factors such as drought, man-made actions and the introduction of non-native species caused a decline in the population of this fish and it was in danger of extinction (Gharaei *et al.*, 2019). In addition to the biological and ecological importance of *S. zarudnyi*, it is delicious and in high demand in the seafood domestic market (Rahdari *et al.*, 2013). Therefore, the species could be a potential candidate in the aquaculture industry (Gharaei *et al.*, 2010). The introduction of new species for aquaculture is rapidly expanding due to the development of new technologies for fish breeding. On the other hand, the number of threatened or endangered species is increasing (Arthington *et al.*, 2016). There are many ways to protection of native species. Although artificial breeding is not the best solution, there is no other solution under the current conditions when other solutions such as restoring natural spawning grounds or removing non-native fish are not possible (Azari Takami, 2014). In sustainable aquaculture, artificial propagation is the most promising and reliable method to produce high egg quality throughout the year. It involves the use of natural (hypophysation) or synthetic hormones

to induce ovulation and spawning in farmed fish (Apollos, 2020; Asadi Edivand *et. al.*, 2022). In captivity, many commercial fish of aquaculture industry exhibit reproductive dysfunctions (Zohar and Mylonas, 2001). So, in these fish hormone therapy is the only way to achieve controlled and reliable reproduction, and not only spawning should be induced in this way, but it is necessary for having good quality gametes and followed having good larvae and fry (Podhorec and Kouril, 2009; Kerdgari *et. al.*, 2022). The reason for the lack of spawning in these conditions is the cessation of the release of LH (luteinizing hormone) from the pituitary (Mylonas *et al.*, 1998). The use of hormone therapy causes the release of LH stored in the pituitary and, as a result, advances the final maturation of the oocyte and ovulation (Zohar and Mylonas, 2001). The use of GnRH analogs has been used for hormone therapy of many fish. Of course, in some fish such as carp, catfish, salmon, and some perciforms, there is a strong dopamine system that stops ovulation and sperm emission in captivity (Cabrita *et al.*, 2008).

In breeding sectors, aquaculture species are artificially propagated using different hormones, which are produced by synthetic and recombinant methods. There are several reasons for the production of recombinant hormones such as the small number of desired peptides in the production tissues available, and relationship between the structure and function of the hormones is dedicated to removing or replacing one

or more amino acids (Yeganeh *et al.*, 2022). For artificial reproduction, synthetic hormones are cheaper and more stable than carp pituitary extract (CPE), which has been used in the past for the reproduction of carp, and therefore have a longer shelf life. Also, these hormones are mostly available in the market in pure form without any reproductive inhibiting agents (Yousefian and Mousavi, 2011). Ovaprim contains combinations of salmon gonadotropin-releasing hormone (sGnRH) and domperidone. The Ovaprim hormone has been used for the artificial breeding of many fish species, such as Indian carps (Cyprinidae) (Mijkherjee *et al.*, 2002; Sharma and Singh, 2002; Sarkar *et al.*, 2004; Naeem *et al.*, 2005 a, b), pike, *Esox lucius* (Khaval *et al.*, 2015) and Longspine scraper, *Capoeta trutta* (Zadmajid, 2016). In Danube bleak (*Alburnus chalcoides*), higher efficiency of Ovaprim than pituitary extract and combination of HCG (Human chorionic gonadotropin) with metoclopramide has been observed (Nosrati *et al.*, 2019). Induction of ovulation with Ovaprim in catfish, *Clarias gariepinus* species has been successfully performed (Musa

Ahmed and Hagar Talib, 2018). However, there is currently no published data on the use of two types of recombinant hormones (Iranian-made hormone: Ovapass with foreign-made hormone: Ovaprim) in the artificial reproduction of *S. zarudnyi*, but total reliance on Ovaprim hormone means that reproduction of this fish will be disrupted if unavailable. Therefore, the aim of this study was to compare the effect of the hormones OvapassTM and OvaprimTM on spawning induction in *S. zarudnyi*. So far, no study has been published about the effects of this hormone on fish.

Materials and methods

Broodstock selection and maintenance

In this experiment, the fish specimens were obtained from healthy broodstocks of Zahak Aquatic Restoration and Genetic Conservation Center (Sistan and Baluchestan province, Iran, 89°30'N, 67°61'E). Before the experiment, 48 female broodstocks (1535.12 ± 180.09 g) and 60 male broods (752.00 ± 48.30 g) were placed separately in rectangular-shaped concrete tanks of running water (3 m³ volume) of 15-17°C for 22-24 h (Table 1).

Table 1: Weights and lengths of injected groups (mean \pm SD).

Parameter	Groups				
	1	2	3	4	
Female	Weight (g)	1601.33 ± 315.15	1483.33 ± 56.86	1550.00 ± 245.37	1505.83 ± 102.97
	Length (cm)	52.20 ± 0.97	52.60 ± 1.16	49.40 ± 1.24	49.60 ± 1.58
Male	Weight (g)	761.12 ± 29.41	783.59 ± 60.11	725.17 ± 67.12	738.12 ± 36.57
	Length (cm)	24.85 ± 0.63	27.86 ± 1.29	23.12 ± 0.81	24.35 ± 0.57

Females were characterized with soft, swollen belly and pink-red genital. Also, males were characterized by releasing

milt upon slight pressure to the abdomen were selected. Prior to injections, the fish were anesthetized with a clove

solution bath at 0.05-0.07 mg/L for 2-3 min (Rahdari *et al.*, 2018). The weighing (g), labeling, length measurement (cm), and hormone injection were performed while the fish were unconscious. Following the hormone injection, the specimens (12 females and 15 males) were kept together in circular concrete tanks with a flow-through circuit. Tanks were filled with filtered freshwater and 50% daily water exchanged. During the experiment, the water temperature was $16.4 \pm 1.1^\circ\text{C}$ and values for dissolved O₂ and pH were 6.9 ± 0.3 mg/L and 7.5 ± 0.4 , respectively. The photoperiod was kept natural (10 h light and 14 h darkness).

Hormones and experimental design

Ovapass™ was purchased from Protein Array Saman (PAS) Company (Tehran,

Iran). Its composition is declared similar to Ovaprim™. Ovaprim (each milliliter contains 20 µg Salmon gonadotropin-releasing hormone analog (GnRH) and 10 mg domperidone), purchased from Syndel Laboratories, Ltd., Vancouver, Canada.

Females (n=48) were allocated into four experimental groups. 24 h after adaptation at $15-17^\circ\text{C}$, four treated groups received hormones via intraperitoneal injections at the base of the pectoral fins as shown in Table 2. The males (n=60) were allocated into four groups and received hormones synchronized to the 2nd female's injection for inducing spermiation (Table 3).

Table 2: The doses of hormones used to induce spawning in female *S. zarudnyi*.

Treatment	Fish No.	Injections dosage/kg BW			Time interval	Reference	
		First	Second	Third			
T1	Ovaprim (mL/kg)	12	0.2	0.5	0.5	24h	Rahdari <i>et al.</i> , 2013
T2	Ovapass (mL/kg)	12	0.5	0.5	-	12h	Recommended by the manufacturer
T3	Ovapass (mL/kg)	12	0.2	0.5	0.5	24h	
T4	Ovaprim (mL/kg) + Ovapass (mL/kg)	12	0.2 Ovaprim	0.5 Ovapass	0.5 Ovapass	24h	

Table 3: The doses of hormones used to induce spawning in male *S. zarudnyi*.

Groups	Treatment	Fish No.	Injection's dosage/kg BW.
1	Ovaprim (mL/kg)	15	0.3
2	Ovapass (mL/kg)	15	0.3
3	Ovapass (mL/kg)	15	0.3
4	Ovaprim (mL/kg) + Ovapass (mL/kg)	15	0.3

To avoid repeated anesthesia and stress to the brood stocks, an arrangement was made to inject the first stage of the

female broodstocks immediately after labeling and weighing. The injections were done at the base of the pectoral fin

(Rahdari *et al.*, 2013). To minimize stress in the broodstocks, insulin syringes were used for injection. The male and female brood stocks, which were kept separately in rectangular ponds before the injection, were transferred together to the round pond in the indoor facility, after the injection, for further stimulation. After the second

injection, the females were checked every 6 h. Therefore, the fish were anesthetized and the eggs were stripped manually, when ovulation was observed. The spawning rate index (%) was calculated with the following relationship (Billard, 1990):

$$\text{Spawning rate} = (\text{Number of ovulated fish} / \text{Total number of injected fish}) \times 100$$

Collection of Gametes, fertilization, and incubation

During the examination, ripe gamete donors were anaesthetized in a solution containing 0.05-0.07 mg/L clove powder. The females were checked each 8 to 10 h after the second injection. The milt was collected from the male donors by stripping. The ovum was collected from individual female donor by abdominal gently stripped method. Eggs were stripped into a plastic vessel and were fertilized using a "dry method" (Kucharczyk *et al.*, 1997). The eggs were placed on trays and the milt and eggs were mixed slowly with a turkey feather for one min. In general, the ratio

of males to females was 3:1 (Gharaei *et al.*, 2019). All spawners were kept one week after gamete collection and all fish were closely monitored to record survival rate.

The fecundity rate was calculated using volumetric technique. This technique uses simple proportionality to estimate total fecundity from a given number of eggs in a known subsample volume and total sample volume value, and then calculates the total number of eggs in the ovary. To determine the relative fecundity, the ratio of the number of extracted eggs to the weight of the fish was calculated (Billard, 1990):

Working fecundity=The total volume of stripped eggs \times The average number of eggs in the samples

Relative fecundity=Working fecundity/ total body weight (kg)

For fertilization, a very small amount of water was added so that the sperms become active and fertilization takes place. Then, the adhesion was removed gradually with the water of the hatchery facility. Egg diameter (mm) was

measured from 10 eggs from each fish, using an eyepiece micrometer fitted to a dissection microscope. The water was changed over time and after half an hour at most, the adhesion of the eggs was completely removed then left to water

harden for 30 min. Fertilized eggs were incubated in jar incubators (Vase) to start the incubation stage. To determine the percentage of fertilization about 24 h after transferring the eggs to jar incubators, about 100 eggs of each brooder were sampled and after clarification in the clarification solution Fertilization rate (FR) (%)=Number of fertilized egg/ total eggs×100

Two types of incubators, jar Vase and troughs California, were used for the incubation stages of *S. zarudnyi* eggs. In this way, from the beginning to the egg-hatching stage, a jar Vase was used, and from hatching to post-hatching, and after that, California incubators were used. In each jar incubator, 300 mL of water-

$$\text{Hatching rate} (\%) = (\text{Number of viable larvae} / \text{total number of eyed eggs}) \times 100$$

Larvae Samples were fixed in formaldehyde and external condition were checked by a loop.

Statistical Analysis

Data analysis was done with SPSS 16 software (Chicago, IL, USA). First, their normality was checked by the Kolmogorov-Smirnov test and the homogeneity of variances was checked by Levene's test. Then, differences among the experimental groups were analyzed using One-way variance followed by Duncan's test at a 95% confidence level. Data are represented as mean±standard deviation.

Results

According to our results, ovulation and spawning occurred between 85% and

(1:1:1 methanol+acetic acid+distilled water (Zarei *et al.*, 2019)), the stages of embryo growth and development was observed under the loop. Samples with a neural streak were counted to determine the fertilization rate (Pankhurst *et al.*, 1996):

hardened eggs were poured and water flow was established. After all the eggs in each jar incubator reached the hatching time, they were transferred to the trough's Californian incubator. The percentage of egg-to-larvae conversion was calculated using the following relationships (Pankhurst *et al.*, 1996):

100% in all groups after hormone injection. There were significant differences between experimental groups in terms of fertilization rate (FR) (Fig. 1), the number of dry eggs per milliliter, spawning success (Table 4), latency period, eyed eggs, and hatching rate ($p<0.05$; Table 5). In this regard, the highest values of FR were observed in T2 ($p<0.05$). Fish of T2 showed the highest spawning rate among all experimental groups, also, there were no significant differences between T2 and T4 ($p>0.05$). As well as, there were no significant differences between experimental groups in terms of working fecundity and relative fecundity (Table 4), the diameter of dry egg and total hormone injected ($p>0.05$; Table 5). Also, the lowest working fecundity and

relative fecundity were observed in T1 ($p>0.05$; Table 4). Also, the lowest and highest values of total hormone injected were observed in T2 and T1, respectively. The lowest and highest values of the latency period, which is the time interval between the first injection and the time of ovulation, were observed

in T1 and T2, respectively ($p>0.05$; Table 5). The results of the incubation period showed that the time to reach the hatching stage and from hatching to post-hatching was influenced by the water temperature of the incubators.




Figure 1: Fertilization rate of different groups injected with Ovaprim and Ovapass hormones (T1: Fish that received 0.2, 0.5 and 0.5 mL/kg body Ovaprim with 24-h time intervals, T2: females treated with 0.5 and 0.5 mL/kg BW Ovapass with 12-h time intervals, T3: females were injected with 0.2, 0.5 and 0.5 mL/kg BW Ovapass with 24-h time intervals, and T4: females were injected with a combination of Ovaprim (0.2 mL/kg) and Ovapass (0.5 and 0.5 mL/kg BW) with 24-h time intervals).

Table 4: The effect of different doses of two types of hormones, Ovaprim and Ovapass, on different parameters of *S. zarudnyi* eggs and larvae (mean \pm standard deviation; n=12).

Group	Female brooder weight (g)	Spawning success (%)	Working fecundity	Relative fecundity	The volume of dry eggs per female fish (mL)	The number of dry eggs/mL
1	1601.33 \pm 315.15	85 \pm 00.00 ^a	42258.33 \pm 10651.89	26504.79 \pm 2821.39	153.67 \pm 38.73 ^b	275.00 \pm 1.00 ^a
2	1483.33 \pm 56.86	100 \pm 00.00 ^c	45070 \pm 10203.03	30236.04 \pm 5803.13	160.00 \pm 36.06 ^a	281.67 \pm 1.53 ^b
3	1550.00 \pm 245.37	85 \pm 00.00 ^a	52977.22 \pm 6191.04	34522.62 \pm 4946.76	191.67 \pm 23.63 ^b	278.00 \pm 1.00 ^a
4	1505.83 \pm 102.97	90 \pm 00.00 ^b	50746.67 \pm 5306.65	33785.46 \pm 3229.86	210.00 \pm 55.68 ^a	273.00 \pm 1.65 ^a

^{a,b} Different superscripts within a column indicate significant differences ($p<0.05$). (T1: Fish that received 0.2, 0.5 and 0.5 mL/kg body Ovaprim with 24-h time intervals, T2: females treated with 0.5 and 0.5 mL/kg BW Ovapass with 12-h time intervals, T3: females were injected with 0.2, 0.5 and 0.5 mL/kg BW Ovapass with 24-h time intervals, and T4: females were injected with a combination of Ovaprim (0.2 mL/kg) and Ovapass (0.5 and 0.5 mL/kg BW) with 24-h time intervals).

As the temperature increased, the duration of these periods became shorter and counter. The lowest values of completion of hatching (h) obtained in fish administrated with Ovapass (35 ± 3.2 h; T2) compared to other experimental

groups ($p < 0.05$; Table 6). No abnormality was observed in the offsprings of the experimental groups after hormone therapy and spawning.

Table 5: Effect of different doses of two types of hormones, Ovaprim and Ovapass, on different parameters of *S. zarudnyi* eggs and larvae (mean \pm standard deviation; n=12).

Group	Diameter of dry egg (mm)	Total hormone injected (mL)	Latency period (h)	Eyed egg (%)	Hatching rate (%)
1	1.67 ± 0.25	1.93 ± 0.36	36.1 ± 3.8^c	87.27 ± 0.23^a	80.67 ± 0.12^a
2	1.48 ± 0.012	1.52 ± 0.98	27.33 ± 0.66^a	93.33 ± 0.58^d	87.67 ± 4.93^b
3	1.61 ± 0.12	1.70 ± 0.19	32.38 ± 2.20^b	90.33 ± 0.58^b	82.00 ± 1.00^a
4	1.69 ± 0.03	1.80 ± 0.15	31.8 ± 1.30^b	91.33 ± 0.58^c	89.33 ± 0.58^b

^{a,b} Different superscripts within a column indicate significant differences ($p < 0.05$). (T1: Fish that received 0.2, 0.5 and 0.5 mL/kg body Ovaprim with 24-h time intervals, T2: females treated with 0.5 and 0.5 mL/kg BW Ovapass with 12-h time intervals, T3: females were injected with 0.2, 0.5 and 0.5 mL/kg BW Ovapass with 24-h time intervals, and T4: females were injected with a combination of Ovaprim (0.2 mL/kg) and Ovapass (0.5 and 0.5 mL/kg BW) with 24-h time intervals).

Table 6: Characteristics of water temperature and period of eyed and hatching eggs.

	Group 1	Group 2	Group 3	Group 4
Water temperature until eyed egg period (°C)	16.0 ± 1.3	16.0 ± 1.1	16.8 ± 1.2	17.2 ± 1.0
Eyed egg period (h)	170.0 ± 10.0^b	165.0 ± 12.5^b	158.5 ± 11.5^a	152.6 ± 8.5^a
Water temperature from eyed egg stage to hatching (°C)	18.4 ± 1.5	19.5 ± 1.9	17.3 ± 1.1	17.3 ± 1.0
Completion of hatching (h)	48.0 ± 3.5^b	35.0 ± 3.2^a	52.0 ± 4.3^b	52.0 ± 4.1^b

^{a,b} Different superscripts within a line indicate significant differences ($p < 0.05$).

Discussion

In the present study, a combination of Ovaprim and Ovapass was applied for the first time for inducing spawning in *S. zarudnyi*. *S. zarudnyi* belongs to the Cyprinidae family that has a strong dopamine system (Peter *et al.*, 1986). One of the strong types of evidence for this claim is that the use of pituitary extract and HCG never induces ovulation in the female fish. Therefore, to weaken and eliminate the dopamine system, an anti-dopamine agent (e.g. domperidone, reserpine, metoclopramide or pimozide) must be used (Brzuska, 2021). So far, the

Ovaprim hormone has been used for the artificial breeding of this fish (Rahdari *et al.*, 2013; Gharaei *et al.*, 2019). However, pituitary extract has been effective for inducing spermatogenesis in male *S. zarudnyi*, and even some of its sperm cognitive parameters have been better than Ovaprim. For example, the highest levels of sperm volume, spermatoctrit, and sperm density were observed in the treatment that used the pituitary extract (Arabnejad *et al.*, 2013).

The efficacy of two synthetic (Ovaprim and Ovatide) for the induced breeding of Indian major carps were compared. The results revealed that

Ovaprim is more effective than ovatide for the breeding of *Catla catla* (Dhawan and Kaur, 2004). Akbari Nargesi *et al.* (2022) showed that two injections of Ovaprim and/or a priming dose of Ovopel (mammalian GnRH analog+metoclopramide, at 18-20 µg and 9-10 mg, respectively) with a resolving dose of Ovaprim were suitable for the artificial reproduction of Rudd (*Scardinius erythrophthalmus*) female breeders, however two injections of Ovaprim recommended for Rudd spawning due to the proper effects and easy application.

The reason for the success of Ovaprim in inducing the spawning of *S. zarudnyi* should be found in the composition of this hormone, which contains dopamine antagonists. In other words, each milliliter contains 20 micrograms of sGnRH (Salmons GnRH) and 10 mg of the anti-dopamine domperidone (Ovaprim™ www.syndel.com). In this study, the effectiveness of Ovapass in inducing spawning of *S. zarudnyi* was compared to Ovaprim. The composition of both hormones is similar, but there are two non-structural differences between them. First of all, Ovapass is an available preparation for fish fry producers. Secondly, the protocol provided for the use of these two hormones by the manufacturing companies is different from each other. This difference becomes important when the duration of hormone therapy until spawning and the amount of hormone used decrease (T2). The frequency of injection is one of the parameters that is important in terms of time, cost and stress to the brooders. In

the case of *S. zarudnyi*, at least three consecutive injections are needed for ovulation to occur, while according to the brochure of the manufacturer of Ovaprim, in most carps, only one injection of Ovaprim at the rate of 0.5 mg/kg is enough for ovulation to occur (www.syndel.com). The issue of the necessity of multiple injections in white fish is true for both hormones, Ovaprim and Ovapass, with the difference that the interval between injections for Ovaprim was 24 h, but for Ovapass, it was 12 h, which reduces the time by half, at least in terms of time and cost. The need for one or more injections is related to the characteristics of the fish, such as, in contrast to *S. zarudnyi*, in *Capoeta razii*, the weight of its brooders was less than 100 g (Abdolahpour *et al.*, 2021) and/or in *Cyprinus rubrofuscus* with an average weight of 536.7 ± 6 g (Malik *et al.*, 2014), spawning occurred only with a single injection of 0.5 mL of Ovaprim per kilogram of body weight. Also, in Indian cyprinid, a single injection of 0.5-2.5 mg/kg of Ovaprim has caused the final treatment (More *et al.*, 2010). On the other hand, the total amount of hormone injected into the reproductive system until the induction of ovulation, in treatment 2 (Ovapass), although it did not have a statistically significant difference with other treatments, the amount was 0.4 mL less than Ovaprim, which is practically and economically very important.

Spawning ratio or Spawning success is one of the good indicators to evaluate hormonal effects on ovulation (Szabo *et al.*, 2002). In this research, fish of T2

showed the highest spawning rate (100 %) among all experimental groups, which is a very acceptable ratio in fish reproduction.

The observation of high FR in treatments 2 and 4 (93.33 and 90.33 percent) compared to the other two treatments cannot be attributed to the type of hormone, because the composition of Ovaprim and Ovapass is similar and probably other factors such as the type of brooders, sperm quality and inoculation operation have been involved. The fact that the percentage of fertilization did not decrease with Ovapass compared to Ovaprim, indicates the absence of a negative effect of Ovapass. In *Barbus grypus*, the percentage of fertilization in fish injected with Ovaprim was 75-90%, while it was 65-80% in pituitary extract treatment (Ghanemi and Khodadadi, 2017).

In the present study, there was no significant difference in some reproductive parameters, including total hormone injected, fecundity (working and relative), and egg diameter among different treatments, which indicates the comparable efficiency of Ovapass with Ovaprim. Egg quality is defined based on its ability to become an embryo (Bobe and Labb  , 2010). The substances used to induce spawning affect not only the time of spawning but also on the number of fish ready to spawn and the quantity and quality of the obtained gametes (Kucharczyk *et al.*, 2007). In perch, *Sander lucioperca*, the quality of eggs collected after induction of ovulation by GnRHs was better than in fish induced

with gonadotropic compounds (such as HCG) (Ljubobratovi   *et al.*, 2019). In the present study, Ovaprim and Ovapass were GnRH hormones, therefore it cannot be concluded that hormone therapy had an effect on egg quality indicators such as egg diameter.

The lowest latency period in the present study was related to T2 (27.33  0.66), which used Ovapass. In fish, the latency period depends on various parameters such as water temperature, biological characteristics (species, age and weight), hormone type and injection frequency (Billard, 1990; Yaron, 1995). In grass carp *Ctenopharyngodon idella*, there was no significant difference in the latency period between the pituitary extract injected group and Ovaprim group (Khodabandeh Shelamani *et al.*, 2012). In *Aspius aspius*, the latency period in the group that received Ovopel (mammalian GnRH+anti-dopamine metoclopramide) or Ovopel and Ovaprim was 40 h, but in the group that received only Ovaprim it was longer (42-44 h) (Targo  ska *et al.*, 2011). In *Abramis brama*, the shortest latency period was also observed when the pituitary extract was used together with HCG hormone, and the duration of the period was increased with Ovopel (Kucharczyk *et al.*, 2007). Since in our study, all treatments were performed on the same species and weight, age and hormone composition were the same. So, the difference between the latency periods of the treatments may be related to the dose and frequency of hormone injections (Rahdari *et al.*, 2014).

In conclusion, the artificial spawning of *S. zarudnyi* was enhanced by using stimulating hormones, but Ovapass gave the best results followed by Ovaprim. These stimulating hormones have enhanced reproductive performances. The data of the present study showed that the effectiveness of Iranian Ovapass hormone for inducing spawning of *S. zarudnyi* for reasons such as good performance, cheapness, and easy availability is very suitable and it can be replaced Ovaprim hormone.

Acknowledgment

The authors would like to thank the University of Zabol (Zabol, Sistan and Baluchestan Province, Iran) for providing the funds and facilities for this study.

References

Abdolahpour, H., Jafari, N., Karimzadeh, M. and Falahatkar, B., 2021. Artificial breeding of *Capoeta razii* using OvaprimTM. *Journal of Aquaculture Sciences*, 9, 15-26.

Akbari Nargesi, E., Falahatkar, B. and Źarski, D. 2022. Artificial reproduction of Caspian roach, *Rutilus caspicus* following stimulating ovulation with Ovaprim, Ovopel, and their combinations under controlled conditions. *Animal Reproduction Science*, 238. DOI:10.1016/j.anireprosci.2022.106932.

Apollos, T.G., 2020. Effect of synthetic hormone (Overprim) on some haematological parameters of African catfish *Clarias gariepinus* (Burchell, 1822). *Journal of Agriculture and Aquaculture*, 2(1), 1-5.

Arabnejad, S., Gharaei, A., Ghaffari, M. and Rahdari, A., 2013. Investigating the qualitative changes of Snow trout *Schizothorax zarudnyi* sperm in response to hormonal induction. *Journal of Cell and Molecular Research*, 27, 617-611. DOI:20.1001.1.23832738.1393.27.4. 13.7

Arthington, A.H., Dulvy, N.K., Gladstone, W. and Winfield, I.J., 2016. Fish conservation in freshwater and marine realms: status, threats and management. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 26, 838-857. DOI:10.1002/aqc.2712

Asadi Edivand A., Mousavi S., Fatemi Tabatabaei S., Zakeri M. and Zanguee N., 2022. Determining the optimal time for artificial propagation in Hilsa shad (*Tenualosa ilisha*) based on thyroid and steroid hormones levels. *Iranian Journal of Fisheries Sciences*, 21, 741-757. DOI:10.22092/ijfs.2022.127015

Azari Takami, Q., 2014. Breeding and Cultivation of Sturgeon Fish. Tehran University Press, Iran. 406 P. [In Persian]

Billard, R., 1990. The major carps and other cyprinids. In: Nash, C.E. and Novotny, A.J. (eds) Production of Aquatic Animals (fishes). Elsevier, Amsterdam, pp. 21-55.

Bobe, J. and Labb  , C., 2010. Egg and sperm quality in fish. *General and Comparative Endocrinology* 165, 535-548.

Brzuska, E., 2021. Reproduction effectiveness of carp (*Cyprinus carpio* L.) from the Hungarian W breeding line after stimulating ovulation with spawning inducing agents of natural (CPH, hCG, PMSG) and/or synthetic origin (Ovopel, Digin, Ovaprim, mGnRH-a). *Aquaculture*, 532. DOI:10.1016/j.aquaculture.2020.736 023.

Cabrita, E., Robles, V. and Paz Herráez, P., 2008. Methods in reproductive aquaculture: marine and freshwater species. Taylor and Francis Group, 574 P. DOI:10.1201/9780849380549

Dhawan, A. and Kaur, K., 2004. Comparative efficacy of ovaprim and ovatide in carp breeding. *Indian Journal of Fisheries*, 51, 227-228.

Ghanemi, M. and Khodadadi, M., 2017. Effects of ovaprim administration on reproductive parameters of Shirbot, *Barbus grypus*, Cyprinidae. *Turkish Journal of Fisheries and Aquatic Sciences*, 17, 1025-1030. DOI:10.4194/1303-2712-v17_5_18

Gharaei, A., Rahdari, A. and Ghaffari, M., 2010. *Schizothorax zarudnyi* as a potential species for aquaculture. Farming the waters for people and food Conference. 22-25 September 2010, Thailand. 3 P.

Gharaei, A., Rahdari, A. and Ghaffari, M., 2019. Artificial reproduction of snow trout *Schizothorax zarudnyi* using synthetic hormones. *Journal of Marine Science and Technology*, 10, 1-11.

DOI:20.1001.1.20088965.1390.10.1. 1.1.

Kerdgari M., Afkhami M., Ehsanpour M. and Ghanbarzadeh M., 2022. Monitoring anthropogenic pollutants in northern coasts of Hormuz Strait using blood indices and thyroid hormone levels in *Periophthalmus argentilineatus* (Pisces: Gobiidae). *Iranian Journal of Fisheries Sciences*, 21 (2), 431-444. DOI:10.22092/ijfs.2022.350699.0

Khaval, A., Dezhandiyani, S., Mahisefat, F., Amiri, A. and Sharifian, M., 2015. The effect of optimum dosage of Ovaprim injection on artificial spawning of *Esox lucius*. *Iranian Scientific Fisheries Journal*, 23, 1-15. DOI:10.22092/isfj.2015.103165 [In Persian]

Khodabandeh Shalmani, L., Maboudi, H., Askari Sari, A. and Basak Kahkesh, F., 2012. Comparison of the effect of Ovaprim hormone along with antidopamine domperidone with carp pituitary gland extract on latency period, larval survival percentage and absolute fecundity of *Ctenopharyngodon idella*. National Conference of non-operating defense in the agricultural sector, Pishtaz Iranian Science Cooperative, Qeshm, Iran. 10 P.

Kucharczyk, D., Kujawa, R., Luczynski, M., Glogowski, J., Babiak, I. and Wyszomirska, E., 1997. Induced spawning in bream *Abramis brama* (L.), using carp pituitary extract and HCG.

Aquaculture Research, 28, 139-144. DOI:10.1046/j.1365-2109.1997.t01-1-00840.x

Kucharczyk, D., Kestemont, P. and Mamcarz, A., 2007. Artificial reproduction of pikeperch. Olsztyn: EC project (Luciperceimprove, CT 2005-17646), pp. 52-54. DOI:10.13140/2.1.3621.4560

Ljubobratović, U., Péter, G., Sándor, Z., Kugyela, N. and Rónyai, A., 2019. The effect of hormonal preparation (gonadotropins vs. gonadoliberins) on pre-seasonally obtained eggs and larvae quality in pikeperch (*Sander lucioperca* L.). *Aquaculture International*, 27: 1009-1024. DOI:10.1007/s10499-019-00368-4

Malik, A., Abbasi, A.R., Kalhoro, I.B., Shah, S.A., Narejo, N.T. and Kalhoro, H., 2014. Effect of Ovaprim hormone (Syndel Laboratory, Canada) on spawning of koi carp at fish hatchery Chilya Thatta, Sindh, Pakistan. *Sindh University Research Journal*, 46, 273-276.

Mijkherjee, M., Praharaj, A. and Das, S., 2002. Conservation of endangered fish stocks through artificial propagation and larval rearing technique in West Bengal, India. *Aquaculture Asia*, 7, 8-14.

More, P.R., Bhandare, R.Y., Shinde, S.E., Pathan, T.S. and Sonawane, D.L., 2010. Comparative study of synthetic hormones Ovaprim and carp pituitary extract used in Induced breeding of Indian major carps. *Libyan Agriculture Research Center Journal International*, 1, 288-295.

Musa Ahmed, A.M. and Hagar Talib, E.A., 2018. Effects of Ovaprim Hormone on Induced Breeding of *Clarias gariepinus*. *Agricultural Extension Journal*, 2(2), 75-77.

Mylonas, C., Woods, L.C. and Zohar, Y., 1998. Hormone profiles of captive Striped Bass (*Morone saxatilis*) during spermiation and long-term enhancement of milt production. *Journal of the World Aquaculture Society*, 29, 379-392. DOI:10.1111/j.1749-7345.1998.tb00662.x

Naeem, M., Salam, A. and Jafar, A., 2005a. Induced spawning of major carp *Catla catla* by a single intramuscular injection of Ovaprim-C and fecundity at fish hatchery Islamabad, Pakistan. *Journal of Biological Sciences*, 5, 776-780. DOI:10.3923/pjbs.2005.1126.1130

Naeem, M., Salam, F., Diba, A. and Saghir, S., 2005b. Fecundity and Induced Spawning of Silver carp, *Hypophthalmichthys molitrix* by using a Single Intramuscular Injection of Ovaprim-C at Fish Hatchery Islamabad, Pakistan. *Pakistan Journal of Biological Sciences*, 8, 26-30. DOI:10.3923/pjbs.2005.1126.1130

Nosrati, M., Khara, H., Vahabzadeh, H. and Mirhashemi Nasab, S.F., 2019. Effect of pituitary extract, Ovaprim and combination of human chorionic gonadotropin and metoclopramide on reproductive performance of Caspian shemaya,

Alburnus chalcoides (Guldenstadt, 1772). *Indian Journal of Fisheries*, 66, 51-55. DOI:10.21077/ijf.2019.66.1.67434-07

Oveisi, Z. and Kavosh, H., 2021. Evolution of patterns in architecture of Sistan, east of Iran. *Journal of Sistan and Baluchistan Studies*, 1, 87-93.

Pankhurst, N., Purser, G., Van Der Kraak, G., Thomas, P. and Forteath, G., 1996. Effect of holding temperature on ovulation, egg fertility, plasma levels of reproductive hormones and in vitro ovarian steroidogenesis in the rainbow trout *Oncorhynchus mykiss*. *Aquaculture*, 146, 277-290. DOI:10.1016/S0044-8486(96)01374-9

Peter, E.R., Chang John, P., Nahorniak Carol, S., Omeljaniuk R.J., Sokolowska, M., Shih, S.H. and Billard, R., 1986. Interactions of Catecholamines and GnRH in Regulation of Gonadotropin Secretion in Teleost Fish, Editor(s): ROY O. GREEP, In Recent Progress in Hormone Research, Proceedings of the 1985 Laurentian Hormone Conference, Academic Press, 42, 513-548. DOI:10.1016/B978-0-12-571142-5.50016-1.

Podhorec, P. and Kouril, J., 2009. Induction of final oocyte maturation in cyprinidae fish by hypothalamic factors: a review. *Veterinarni Medicina*, 54, 97-110. DOI:10.17221/50/2009-VETMED

Rahdari, A., Gharaei, A., Ghaffari, M. and Najafi, M., 2013. Biotechnics of artificial propagation of *Schizothorax zarudnyi* in Sistan water basin. *Journal of Aquaculture Sciences*, 1, 82-73. [In Persian]

Rahdari A., Gharaei A. and Ghaffari M. 2014. Spawning Latency Period in Hormonal Induced Reproduction of Snow trout (*Schizothorax zarudnyi* (Nikolskii, 1897)). *Iranian Journal of Biotechnology*, 12, 1-5.

Rahdari, A., Khosravanizadeh, A., Dahmardeh, H., Gharaei, A. and Mirdar Harijani, J., 2018. Anesthetic effects and biochemical changes of peppermint essence (*Mentha spicata*) in snow trout (*Schizothorax zarudnyi*). *Aquatic Animals Nutrition* 3, 35-46. [In Persian]

Sarkar, U.K., Negi, R.S., Deepak, P.K., Singh, S.P., Srivastava, S.M. and Roy, D., 2004. Captive breeding of vulnerable Indian carp *Cirrhinus reba* with Ovaprim for conservation of wild populations. *Aquaculture Asia*, 9, 5-7. DOI: 10.14798/73.1.786.

Sharma A.P. and Singh, V.K., 2002. Induced breeding response of Indian major carps, viz. *Labeo rohita*, *Catla catla* and *Cirrhinus mrigala* using Ovaprim and carp pituitary extract. *The Indian Journal of Animal Sciences*, 72, 35-354.

Szabo, T., Medgyasszay, C. and Horváth, L., 2002. Ovulation induction in nase (*Chondrostoma nasus*, Cyprinidae) using pituitary extract or GnRH analogue combined with domperidone. *Aquaculture*, 203,

389-395. DOI:10.1016/S0044-8486(01)00635-4.

Targońska, K., Kucharczyk, D., Żarski, D., Mamcarz, A. and Falahatkar, B., 2011. Optimization of artificial reproduction of Asp, *Aspius aspius* (L.) under controlled conditions. *Polish Journal of Natural Sciences*, 26, 151-157.

Vekerdy, Z., Dost, R.J.J., Reinink, G. and Partow, H. 2006. History of environmental change in the Sistan Basin based on satellite image analysis: 1976-2005. United Nations Environmental Programme (UNEP): Geneva, Switzerland. 60 P.

Yaron, Z., 1995. Endocrine control of gametogenesis and spawning induction in the carp. *Aquaculture*, 129, 49-73. DOI: 10.1016/0044-8486(94)00229-H.

Yeganeh, S., Mohammadzadeh, S., Moradian, F. and Milla, S., 2022. The effects of recombinant GnRH with dopamine antagonist on reproduction performance, sex steroid levels, and stress response in female koi carp (*Cyprinus carpio*), *Aquaculture Reports*, 22. DOI:10.1016/j.aqrep.2021.101001.

Yousefian, M. and Mousavi, S.E., 2011. The mechanism of reproduction and hormonal function in finfish species. *Scientific Research and Essays*, 6, 3561-3570.

Zadmajid, V., 2016. Comparative effects of human chorionic gonadotropin (hCG) and Ovaprim™ (sGnRHa+ domperidone) on the reproductive characteristics of wild-caught male Longspine scraper, *Capoeta trutta* (Heckel, 1843). *Aquaculture*, 463, 7-15. <http://dx.doi.org/10.1016/j.aquaculture.2016.05.029>

Zarei, S., Dorafshan, S. and Paykan Heyrati, F., 2019. First cleavage interval determination in rainbow trout (*Oncorhynchus mykiss*) and the Caspian salmon (*Salmo trutta caspius*). *Journal of Animal Environment*, 11, 253-256. [In Persian]

Zohar, Y. and Mylonas, C.C., 2001. Endocrine manipulations of spawning in cultured fish: from hormones to genes. In: Lee, C.S. and Donaldson, E.M., (eds). *Reproductive Biotechnology in Finfish Aquaculture*. Elsevier, Netherlands. Pp. 99-136. DOI:10.1016/S0044-8486(01)00584-1

