

## Research Article

# Microplastic contamination in *Nerita albicilla*: Implications for marine ecosystem health along Karachi coast

Tariq H.<sup>1</sup>, Arshad N.<sup>1\*</sup>, Taha M.<sup>1</sup>, Su'ud M.B.<sup>2</sup>, Alam M.M.<sup>3</sup>, Muhammad A.A.<sup>4</sup><sup>1</sup>Institute of Marine Science, University of Karachi, Karachi, Pakistan<sup>2</sup>Multimedia University, Cyberjaye Campus, Malaysia<sup>3</sup>Riphah International University, Islamabad campus, Pakistan<sup>4</sup>Coastal Development and Fisheries Department, Government of Balochistan, Balochistan, Pakistan

\*Correspondence: narshad@uok.edu.pk

**Keywords**

Microplastics,  
*Nerita albicilla*,  
Hawksbay,  
Buleji,  
Morphometric characteristics,  
Plastic pollution,  
Marine gastropods,  
Karachi coast

**Abstract**

Microplastics (MPs) have become a significant environmental concern globally, with their pervasive presence in various ecosystems posing threats to marine life and human health. This study investigates the MPs contamination in a gastropod species *Nerita albicilla* found commonly along the Karachi coast. A total of 60 individuals of *N. albicilla*, 5 specimens from each site monthly were handpicked from the intertidal area of Hawksbay and Buleji rocky shore from November 2022-April 2023. The shells were noted for morphometrics and visceral tissues were digested in 10% KOH solution in a ratio of 3:1. Each digested suspension was filtered using 0.45  $\mu\text{m}$  organic filter paper and observed with the help of a compound microscope. A total of 64 MPs ( $1.4 \pm 0.244 - 3.6 \pm 0.678$  item  $\text{ind.}^{-1}$ ) were recorded from 30 *N. albicilla* specimens collected from Hawksbay and 59 MPs ( $1.4 \pm 0.4 - 3 \pm 0.632$  item  $\text{ind.}^{-1}$ ) from Buleji. The most dominant MPs types were microfibers, which made up to 84% and 88 % of total MPs at Hawksaby and Buleji, respectively whereas the most of MPs ranged in size range of 0-25  $\mu\text{m}$  indicating anthropogenic pollution sources such as textiles, fishing gears, and synthetic materials commonly found in marine environments. The study underscores the ecological significance of gastropods in monitoring marine pollution and highlights the urgent need for mitigation measures to safeguard coastal ecosystems.

**Article info**

Received: April 2024

Accepted: June 2024

Published: November 2024



Copyright: © 2023 by the authors.  
Licensee MDPI, Basel, Switzerland.  
This article is an open access article  
distributed under the terms and  
conditions of the Creative Commons  
Attribution (CC BY) license  
(<https://creativecommons.org/licenses/by/4.0/>).

## Introduction

Microplastics (MPs) have become a major environmental concern due to their widespread presence in various ecosystems, including oceans, rivers, lakes, and even the air (Akdogan and Guven, 2019). MPs are small plastic particles that are less than 5 millimeters in size (Lim, 2021). They can be classified into two categories: primary microplastics, which are intentionally manufactured at small sizes for specific purposes (*e.g.*, microbeads in cosmetics), and secondary microplastics, which result from the breakdown of larger plastic items over time (Lots *et al.*, 2017, Arshad *et al.*, 2023). MPs are commonly transported through rivers and estuaries from terrestrial to aquatic environments, where they persist for extended durations, making them accessible to living organisms (Wright *et al.*, 2013). MPs are widely found in marine environments and can be ingested by organisms with filter-feeding strategies, such as marine gastropods. This ingestion can lead to various detrimental effects, such as reduced feeding, internal injuries, blocked digestive tracts, and impaired reproduction (Ding *et al.*, 2022). MPs particles accumulate in the digestive tract, potentially causing obstructions that disrupt normal digestive processes and adversely affect the health and well-being of the gastropods (Li *et al.*, 2021; Jeyavani *et al.*, 2022). It can also disrupt the entire food chain (Gola *et al.*, 2021) as MPs can be passed along from smaller organisms to larger predators. They contribute to nutrient cycling and are economically significant in the field of fisheries and aquaculture. Additionally, some species are known for

their intricate and beautiful shells, making them popular in the shell trade and as collectibles (Moretzsohn, 2023).

The majority of gastropods species either exhibit limited mobility or are completely sessile as adults. These species, therefore, are ideal for studying contamination levels of their habitat (Lester *et al.*, 2009). Bioaccumulation of contaminants by shellfish is a human health issue and is indicative of poor water quality in the ecosystem (Larsen, 1979; Metian *et al.*, 2009; Kumar *et al.*, 2021).

*Nerita albicilla* is an essential part of the food web and is consumed by other organisms, including humans. Examining the levels of MPs in gastropods offers valuable insights into their potential effects on higher trophic levels, including human health (Pastorino *et al.*, 2021; Multisanti *et al.*, 2022). *N. albicilla* has been chosen for this investigation owing to its ecological significance and abundance along the Karachi coast. This study aims to explore the contamination of microplastics in *N. albicilla* sampled from Hawksbay and Buleji beaches along the Karachi coast, thereby contributing to the ongoing monitoring efforts of MPs levels in gastropods.


## Material and methods

### *Study location*

Pakistan has a vast coastline that stretches from the province of Sindh in the south, to the province of Balochistan in the west. The largest and most populous city in Pakistan, Karachi, is endowed with a long beachfront along the Arabian Sea that stretches for around 64 kilometers (Arshad and Farooq, 2018). Because of the Port of Karachi,

which is strategically located along this coast and is one of the busiest and most important ports in the area, a large amount of Pakistan's trade is facilitated. The Karachi coastline is a center of economic activity and trade (Hameed *et al.*, 2012). The nation's maritime prominence has also been further elevated by the emergence of Gwadar Port in Balochistan as a hub for international trade and a key component of the China-Pakistan Economic Corridor (CPEC) initiatives (Khetran *et al.*,

2017). The coastal area is particularly well-known for its marine life and provides chances for fishing, which contributes significantly to the livelihoods of the local populace. The coastline is distinguished by a blend of rocky outcrops, sandy beaches, and a busy urban setting (Inam *et al.*, 2007). Two nearby rocky beaches on the Karachi coast, Hawksbay and Buleji, were chosen for the present study are shown in Figure 1.



**Figure 1:** Study area map showing studied stations at Buleji (B) and Hawksbay (H) on the Karachi coast.

#### *HawksBay*

Hawksbay approximately covers an area of 990 Km and is characterized by a

combination of rocky and sandy beaches. The rocky regions sustain a varied marine ecology by offering special habitats for a

range of marine organisms. Crevices and tide pools are frequently found in the rocky terrain, where marine life like crustaceans, mollusks, and tiny fish can flourish. The sandy beaches, on the other hand, are crucial sea turtle breeding grounds and offer a natural backdrop for beachgoers and visitors looking to engage in leisure activities like beach sports and picnics.

### *Buleiji*

Buleiji is an 800-meter-long rocky beach. The intertidal zone is made up of sporadic, slightly elevated, and depressed parts. Large and small boulders are mainly found in the high tidal zone. The rocks in the mid- and low-tidal zones are comparably smaller and more level. Mid- and low-tidal zones have rock ponds of various sizes (Zafar *et al.*, 2018). The majority of the organisms in the high tidal zone are barnacles and gastropods. Gastropods, crustaceans (crabs, pistol shrimp, and amphipods), echinoderms (sea cucumber and sea urchin), bivalves (mussels and oysters), etc. live in the mid- and low-tidal zones. The algal vegetation on the shore is abundant.

### *Faunal sampling and MPs analysis*

A total of 60 individuals of *N. albicilla* were randomly handpicked from the intertidal area of Hawksbay and Buleiji rocky ledge along the coast of Karachi. Five individuals of *N. albicilla* were collected from each station every month from both locations from November 2022 to April 2023. The months between November to February are referred to northeast monsoon (winter) whereas, March and April are referred to spring inter-monsoon. The samples were placed in aluminum foil to avoid

contamination and brought to the laboratory, placed in the freezer at  $-20^{\circ}\text{C}$  to prevent deterioration till further laboratory procedures. The collected specimens were defrosted at room temperature in the clean laboratory, and analyzed using Bio protocol by Lusher *et al.* (2015, 2018). The molluscan species were identified using Shells, Dance (2002) and available literature. Digital weight balance (Akira, Tx-200) was used to calculate the weight whereas a Vernier caliper was used to measure the length and width of molluscan shells. After morphometrics for each sample, an iron hammer covered with clean sterilized aluminum foil was used to crack the shells, the visceral tissues were extracted carefully and weighed to get wet weight of the samples. The whole visceral tissue of every specimen was placed into pre-cleaned vials and digested with 10% KOH solution in a ratio of 3:1 (KOH: visceral tissue) as described by Lusher *et al.* (2018) and Arshad *et al.* (2023). The samples were kept in 10% KOH for 3 weeks to break down the living residues without heat (Foekema *et al.*, 2013). Some of the samples which were turbid after 3 weeks of KOH digestion, were heated on a magnetic stirrer at 125 rpm till the suspension became transparent. Each digested suspension was filtered using  $1.6\ \mu\text{m}$  (pore size) GF/A filter papers. Filter papers were placed and covered with Petri dishes washed with Milli-Q water. After filtration, stereomicroscope (Micros Austria) was used to study the total number of MPs and their characteristics in each sample. MPs were evaluated according to size, shape, color, and abundance. The morphotypes of MPs were categorized into

fibers and fragments (including irregular particles) and confirmed according to their physical characteristics. MPs were categorized into four size ranges in this study, including 0-25 $\mu$ m, 26-50 $\mu$ m, 151-175 and 176-200 $\mu$ m, respectively and eight different colors i.e. black, blue, brown, green, orange, pink, purple and red. Regression analysis was performed to find out the significant relation between the morphometrics and MPs count in the samples. Statistical analyses were performed using Statistical Package for the Social Sciences (SPSS) program for windows, version 26.0 (IBM, America).

## Results

### *Monthly microplastic abundance in molluscan Species at Hawksbay*

A total of 64 MPs were recorded from 30 *N. albicilla* collected from Hawksbay from November 2022 to April 2023. The morphometric characteristics of all specimens are described in Table 1. MPs abundance differed significantly among months and ranged between 1.400 $\pm$ 0.244-3.600 $\pm$ 0.678 item ind.<sup>-1</sup>. The highest number of MPs (3.600 $\pm$ 0.678 item ind.<sup>-1</sup>, 2.420 $\pm$ 1.489 MPs/g) was recorded in November with an average wet weight of 0.52  $\pm$  0.062 g. The lowest number of MPs (1.4 $\pm$ 0.244 item ind.<sup>-1</sup>, 0.422 $\pm$ 0.088 Mps/g) was recorded in March with an average wet weight of 0.508 $\pm$ 0.041 g.

**Table 1: Morphometric characteristics and microplastic (Mps) abundance in *Nerita albicilla* species collected from Hawksbay beach monthly (November 2022- April 2023).**

S.No.	Months	Number	Avg. width	Avg. Height	Avg. wt. with shell	Avg. wt. without shell	Total MPs	Total MPs /g
1	Nov.	5	1.98 $\pm$ 0.037	1.66 $\pm$ 0.081	3.31 $\pm$ 0.208	0.52 $\pm$ 0.062	3.60 $\pm$ 0.678	2.420 $\pm$ 1.489
2	Dec.	5	1.78 $\pm$ 0.156	1.36 $\pm$ 0.067	2.78 $\pm$ 0.172	0.58 $\pm$ 0.054	1.40 $\pm$ 0.600	0.891 $\pm$ 0.288
3	Jan.	5	2.34 $\pm$ 0.112	3.48 $\pm$ 0.152	9.29 $\pm$ 1.052	1.71 $\pm$ 0.214	2.60 $\pm$ 0.927	0.933 $\pm$ 0.267
4	Feb.	5	2.00 $\pm$ 0.044	1.00 $\pm$ 0.031	2.53 $\pm$ 0.145	0.45 $\pm$ 0.038	2.60 $\pm$ 1.077	1.710 $\pm$ 0.315
5	Mar.	5	1.46 $\pm$ 0.040	1.08 $\pm$ 0.058	2.67 $\pm$ 0.108	0.50 $\pm$ 0.041	1.40 $\pm$ 0.244	0.422 $\pm$ 0.088
6	Apr.	5	1.50 $\pm$ 0.031	1.08 $\pm$ 0.020	2.87 $\pm$ 0.115	0.60 $\pm$ 0.020	1.80 $\pm$ 1.067	0.383 $\pm$ 0.139

Avg. = average; MPs = microplastics

### *Monthly Microplastic Abundance in Molluscan Species at Buleiji*


A total of 59 MPs were recorded from 30 samples of *N. albicilla* collected from Buleiji from November 2022 to April 2023. Whereas, the morphometrics of all specimens are described in Table 2. MPs abundance differed significantly among months and ranged between 1.4 $\pm$ 0.4 - 3 $\pm$ 0.632 item ind.<sup>-1</sup> The highest number of

MPs (3 $\pm$ 0.632 item ind.<sup>-1</sup>, 0.448 $\pm$ 0.095 MPs/g) were recorded in November with an average wet weight of 1.108 $\pm$ 0.082 g. However, the lowest number of MPs (1.4 $\pm$ 0.4 item ind.<sup>-1</sup>, 0.456 $\pm$ 0.144 MPs/g) was recorded in March with an average wet weight of 0.865 $\pm$ 0.123 g. The results of regression analysis do not indicate any significant relationship among height, weight, and number of MPs in studied samples (Fig. 2).

**Table 2: Morphometric characteristics and microplastic (Mps) abundance in *Nerita albicilla* recorded monthly (November 2022 - April 2023) from Buleiji beach.**

S.No.	Months	Number	Avg. width	Avg.Height	Avg. wt. with shell	Avg. wt. without shell	Total MPs	Total MPs/g
1	Nov.	5	1.54 ± 0.04	1.86±0.067	2.99±0.384	0.48±0.038	2.20±0.374	0.25±0.049
2	Dec.	5	2.14±0.092	1.34±0.024	4.13±0.268	0.48±0.039	1.80±0.860	0.37±0.088
3	Jan.	5	1.98±0.109	1.30±0.054	3.69±0.208	0.65±0.034	2.40±0.509	0.34±0.087
4	Feb.	5	2.72±0.048	1.32±0.058	5.56±0.281	1.10±0.082	3.00±0.632	0.45±0.095
5	Mar.	5	1.84±0.024	1.28±0.048	5.39±0.447	0.86±0.123	1.40±0.400	0.46±0.144
6	Apr.	5	1.60±0.137	1.14±0.074	4.09±0.468	0.66±0.100	2.20±0.374	0.34±0.062

Avg. = average; MPs = microplastics

**Figure 2: Regression analysis between height (A, C), wet weight without shell (B, D) of *Nerita albicilla* and number of microplastics (MPs) recorded from Hawksbay (A, C) and Buleiji (B, D), respectively.**

Identified MPs were classified into eight size ranges, i.e., <0-25  $\mu\text{m}$  >151-175 $\mu\text{m}$ . The most observed MPs size ranged from 0-25 $\mu\text{m}$  while the lowest particle sizes ranged from 76-100  $\mu\text{m}$ , respectively. A negligible difference has been recorded in both sites. The highest number of MPs was recorded in November (18, 27%) followed by February (14, 23%), December (9, 19%) and January (7, 15%) at Hawksbay. At Buleiji, the highest Mps count was recorded

in January (12, 31%) followed by November (11, 17%), December (10, 17%), February (9, 13%) and March (7, 9%) (Fig. 3A and B). A total of 129 MPs were examined from both sites, however, a negligible difference has been recorded the highest number of MPs (66, 51%) were recorded from Hawksbay whereas, (63, 49%) MPs were recorded from Buleiji. Black and Blue color MPs were dominant,

followed by red, green, and orange from both sites (Fig. 3C and D).

Overall, two different morphologies of MPs were recorded. Fibers were the most dominant type in all samples (84%)

followed by fragments (16%) from Hawksbay while 88% of fibers were recorded from Buleiji followed by 12% fragments (Fig. 4).




Figure 3: Size range of MPs particle (A, B), colors of MPs (C, D) in *Nerita albicilla* collected from Hawksbay (A, C) and Buleiji (B, D), respectively.




Figure 4: Types of MPs recorded from Hawksbay (A) and Buleiji (B) sites

## Discussion

This is the first study investigating and confirming microplastics (MPs) contamination in *Nerita albicilla* from Karachi coast. Our findings indicate a lower number of MPs compared to other gastropod species from various regions, such as *Siphonalia subdilatata* from the Bohai Sea in China (Zhao *et al.*, 2024), *Thais mutabilis* and *Cerithidea cingulata* from the northern Persian Gulf (Naji *et al.*, 2018), *Babylonia spirata* from the southwest coast of India (Abisha *et al.*, 2024), and *Telescopium telescopium* from East Java, Indonesia (Vidayanti and Retnaningdyah, 2024). Comparing the MPs abundance in mollusk species from different regions worldwide is difficult due to the lack of standardized methods (KOH digestion, H<sub>2</sub>O<sub>2</sub> treatment, acid destruction) and units (microplastics per individual or gram tissue) to quantify MPs (Vandermeersch *et al.*, 2015).

MPs contamination comparatively our finding has less number of MPs than in other gastropods species from different regions of the world such as *Thais mutabilis* from north part of the Persian Gulf (Naji *et al.*, 2018), *Siphonalia subdilatata* from Bohai sea China (Zhao *et al.*, 2024), *Thais mutabilis* and *Cerithidea cingulate* from north part of the Persian Gulf (Naji *et al.*, 2018), *Babylonia spirata* southwest coast of India (Abisha *et al.*, 2024) and *Telescopium telescopium* East java Indonesia (Vidayanti and Retnaningdyah, 2024)

Comparing these results with other studies helps to highlight the global nature of MP pollution and the need for comprehensive strategies to mitigate its

impact on marine ecosystems. The most dominant fraction of MPs were microfibers which were observed in an alarming number in all gastropod samples, made up 84% from Hawksbay and 88% from Buleiji, respectively. Similar results have been documented by many researchers. *i.e.*, 66.66% by Wang *et al.* (2021), 68.72%, Patria *et al.* (2020), 50.2%, and Nikhil *et al.* (2024). For microfibers with different colors, black stands out as the most prevalent hue, primarily attributed to its larger proportion within marine environments as reported by Abidli *et al.* (2019) at 99%, Expósito *et al.* (2022) at 74%, and Akindele *et al.* (2019) at 100%. Additionally, it has been suggested by Akindele *et al.* (2019) that near-shore ecosystems are predominantly characterized by these fibers. Microfibers, being degradable plastic waste in yarn or fiber form, typically originate from sources such as fishing lines, ship mines, and synthetic fibers (Pirc *et al.*, 2016). Hawksbay and Buleiji serve as an ecotourism spot for boating and fishing. Moreover, the fragments type of microplastics found in the sample could be from degradation and fragmented plastic debris such as plastic bottles, plastic bags, and plastic bowls. The fiber type of microplastic known could be released from textiles and garments such as clothes (Napper and Thompson, 2016). The fibers are known to remain for a longer period of time on the surface of the water because of their relatively low densities while fragments and granules with higher densities tend to sink (Priscilla and Patria, 2020). Ecologically gastropods are considered a valuable model for



environmental biomonitoring of genotoxic pollutants (Radwan *et al.*, 2020). They have been extensively studied for possessing therapeutically significant proteins and peptides which exhibit antiviral, antimicrobial, and antifungal activities (Cheung *et al.*, 2014). This study underscores the ecological significance of gastropods in monitoring marine pollution and highlights the urgent need for mitigation measures to safeguard the coastal ecosystem.

### Conclusion

This study found considerable microplastic (MPs) contamination in the gastropod species *Nerita albicilla* along the Karachi coast, with microfibers accounting for up to 84-88% of total MPs with the majority ranging in size from 0-25  $\mu\text{m}$ . These findings provide valuable insights into the extent of MPs pollution and its potential impacts on marine life. The results highlight the critical need for actions to safeguard coastal ecosystems and the function of gastropods in monitoring marine pollution.

### Conflicts of interest

The authors have no conflicts of interest to declare. All co-authors have seen and agree with the contents of the manuscript and there is no financial interest to report.

### References

**Abidli, S., Lahbib, Y. and El Menif, N.T., 2019.** Microplastics in commercial molluscs from the lagoon of Bizerte (Northern Tunisia). *Marine Pollution Bulletin*, 142, 243-252. DOI:10.1016/j.marpolbul.2019.03.048

**Abisha, C., Kutty, R., Gurjar, U.R., Jaiswar, A.K., Deshmuke, G., Sasidharan, A. and Xavier, K.M., 2024.** Microplastic prevalence, diversity and characteristics in commercially important edible bivalves and gastropods in relation to environmental matrices. *Journal of Hazardous Materials Advances*, 13, 100392. DOI:10.1016/j.hazadv.2023.100392

**Akdogan, Z. and Guven, B., 2019.** Microplastics in the environment: A critical review of current understanding and identification of future research needs. *Environmental Pollution*, 254, 113011. DOI:10.1016/j.envpol.2019.113011

**Akindele, E.O., Ehlers, S.M. and Koop, J.H., 2019.** First empirical study of freshwater microplastics in West Africa using gastropods from Nigeria as bioindicators. *Limnologia*, 78, 125708. DOI:10.1016/j.limno.2019.125708

**Arshad, N. and Farooq, S., 2018.** Evaluation of Clifton beach by using macrobenthic assemblages for beach management. *Ocean & coastal management*, 163, 30-36. DOI:10.1016/j.ocecoaman.2018.06.002

**Arshad, N., Alam, M.M., Su'ud, M.B.M., Imran, S., Siddiqui, T., Saleem, K., Bashir, A. and Batool, A., 2023.** Microplastic contamination from surface waters and commercially valuable fishes of Karachi Coast, Pakistan. *Regional Studies in Marine Science*, 62, 102955. DOI:10.1016/j.rsma.2023.102955

**Cheung, G.Y., Joo, H.S., Chatterjee, S.S. and Otto, M., 2014.** Phenol-soluble modulins—critical determinants of

- staphylococcal virulence. *FEMS Microbiology Reviews*, 38(4), 698-719. DOI:10.1111/1574-6976.12057
- Dance, S.P., 2002.** Shells. The photographic recognition guide to sea shells of the world. Dorling Kindersley Ltd. England.
- Ding, J., Sun, Y., He, C., Li, J. and Li, F., 2022.** Towards risk assessments of microplastics in bivalve mollusks globally. *Journal of Marine Science and Engineering*, 10(2), 288. DOI:10.3390/jmse10020288
- Expósito, N., Rovira, J., Sierra, J., Gimenez, G., Domingo, J. L. and Schuhmacher, M., 2022.** Levels of microplastics and their characteristics in molluscs from North-West Mediterranean Sea: Human intake. *Marine Pollution Bulletin*, 181, 113843. DOI:10.1016/j.marpolbul.2022.113843
- Foekema, E.M., Gruijter, C., Mergia, M.T., van Franeker, J.A., Murk, A.J. and Koelmans, A.A., 2013.** Plastic in North Sea fish. *Environmental Science and Technology*, 47, 15. DOI:10.1021/es400931b
- Gola, D., Tyagi, P. K., Arya, A., Chauhan, N., Agarwal, M., Singh, S. K. and Gola, S., 2021.** The impact of microplastics on marine environment: A review. *Environmental Nanotechnology, Monitoring and Management*, 16, 100552.
- Hameed, I., Soomro, Y., Butt, A.S. and Shakoor, R., 2012.** Coastline pollution problems of Karachi. *International Journal of Emerging trends in Engineering and Development*, 4(2), 528-537.
- Inam, A., Clift, P.D., Giosan, L., Tabrez, A.R., Tahir, M., Rabbani, M.M. and Danish, M., 2007.** The geographic, geological and oceanographic setting of the Indus River. *Large rivers: Geomorphology and management*, 1, 333-346.
- Jeyavani, J., Sibiya, A., Bhavaniramy, S., Mahboob, S., Al-Ghanim, K.A., Nisa, Z. U. and Vaseeharan, B., 2022.** Toxicity evaluation of polypropylene microplastic on marine microcrustacean *Artemia salina*: An analysis of implications and vulnerability. *Chemosphere*, 296, 133990. DOI:10.1016/j.chemosphere.2022.133990
- Khетran, M.S.B. and Saeed, M.A., 2017.** The CPEC and China-Pakistan relations: a case study on Balochistan. *China Quarterly of International Strategic Studies*, 3(03), 447-461. DOI:10.1142/S2377740017500191
- Kumar, P., Sivaperumal, P., Manigandan, V., Rajaram, R. and Hussain, M., 2021.** Assessment of potential human health risk due to heavy metal contamination in edible finfish and shellfish collected around Ennore coast, India. *Environmental Science and Pollution Research*, 28, 8151-8167. DOI: 10.1007/s11356-020-10764-6
- Larsen, P.F., 1979.** The distribution of heavy metals in the hard clam, *Mercenaria mercenaria*, in the Lower Chesapeake Bay Region. *Estuaries*, 2(1): 1-8.
- Lester, S.E., Halpern, B.S., Grorud-Colvert, K., Lubchenco, J., Ruttenberg, B.I., Gaines, S.D., Airamé, S. and Warner, R.R., 2009.**

- Biological effects within no-take marine reserves: a global synthesis. *Marine Ecology Progress Series*, 384, 33-46.
- Li, M., Wu, D., Wu, D., Guo, H. and Han, S., 2021.** Influence of polyethylene-microplastic on environmental behaviors of metals in soil. *Environmental Science and Pollution Research*, 28, 28329-28336. DOI:10.1007/s11356-021-12718-y
- Lim, X., 2021.** Microplastics are everywhere-but are they harmful?. *Nature*, 593(7857), 22-25. DOI:10.1038/d41586-021-01143-3
- Lots, F.A., Behrens, P., Vijver, M.G., Horton, A.A. and Bosker, T., 2017.** A large-scale investigation of microplastic contamination: abundance and characteristics of microplastics in European beach sediment. *Marine Pollution Bulletin*, 123(1-2), 219-226. DOI:10.1016/j.marpolbul.2017.08.057
- Lusher, A.L., Tirelli, V., O'Connor, I. and Officer, R., 2015.** Microplastics in Arctic polar waters: the first reported values of particles in surface and sub-surface samples. *Scientific reports*, 5(1), 14947. DOI:10.1038/srep14947
- Lusher, A.L., Hernandez-Milian, G., Berrow, S., Rogan, E. and O'Connor, I., 2018.** Incidence of marine debris in cetaceans stranded and bycaught in Ireland: Recent findings and a review of historical knowledge. *Environmental Pollution*, 232, 467-476. DOI:10.1016/j.envpol.2017.09.07
- Metian, M., Warnau, M. and He'douin, L., 2009.** Bioaccumulation of essential metals (Co, Mn and Zn) in the king scallop *Pecten maximus*: seawater, food and sediment exposures. *Marine Biology*, 156, 2063-2075.
- Moretzsohn, F., 2023.** *A Natural and Cultural History. In: Shells.* Reaktion Books, London, 2023 P.
- Multisanti, C.R., Merola, C., Perugini, M., Aliko, V. and Faggio, C., 2022.** Sentinel species selection for monitoring microplastic pollution: A review on one health approach. *Ecological Indicators*, 145, 109587. DOI:10.1016/j.ecolind.2022.109587
- Naji, A., Nuri, M. and Vethaak, A.D., 2018.** Microplastics contamination in molluscs from the northern part of the Persian Gulf. *Environmental pollution*, 235, 113-120.
- Napper, I.E. and Thompson, R.C., 2016.** Release of synthetic microplastic plastic fibres from domestic washing machines: Effects of fabric type and washing conditions. *Marine pollution bulletin*, 112(1-2), 39-45. DOI:10.1016/j.marpolbul.2016.09.025
- Nikhil, V.G., Amritha, G.G., Ranjeet, K. and Varghese, G.K., 2024.** Distribution of microplastics in seafloor sediments and their differential assimilation in nearshore benthic molluscs along the south-west coast of India. *Environmental Pollution*, 344, 123350. DOI:10.1016/j.envpol.2024.123350
- Pastorino, P., Nocita, A., Ciccotelli, V., Zaccaroni, A., Anselmi, S., Giugliano, R. and Prearo, M., 2021.** Health risk assessment of potentially toxic elements, persistence of NDL-PCB, PAHs, and microplastics in the translocated edible freshwater *Sinotaia quadrata* (Gasteropoda, Viviparidae): a case study from the Arno River Basin (Central

- Italy). *Exposure and Health*, 13(4), 583-596. DOI: 10.1007/s12403-021-00404-w
- Patria, M.P., Santoso, C.A. and Tsabita, N., 2020.** Microplastic ingestion by periwinkle snail *littoraria scabra* and mangrove crab *metopograpsus quadridentata* in Pramuka Island, Jakarta Bay, Indonesia. *Sains Malaysiana*, 49(9), 2151-2158. DOI: 10.17576/jsm-2020-4909-13
- Pirc, U., Vidmar, M., Mozer, A. and Kržan, A., 2016.** Emissions of microplastic fibers from microfiber fleece during domestic washing. *Environmental Science and Pollution Research*, 23, 22206-22211. DOI:10.1007/s11356-016-7703-0
- Priscilla, V. and Patria, M.P., 2020.** Comparison of microplastic abundance in aquaculture ponds of milkfish *Chanos chanos* (Forsskål, 1775) at Muara Kamal and Marunda, Jakarta Bay. In: IOP conference series, Earth and environmental science. IOP, England, 012027 P.
- Radwan, M.A., El-Gendy, K.S. and Gad, A.F., 2020.** Biomarker responses in terrestrial gastropods exposed to pollutants: A comprehensive review. *Chemosphere*, 257, 127218. DOI:10.1016/j.chemosphere.2020.127218
- Vandermeersch, G., Lourenço, H.M., Alvarez-Muñoz, D., Cunha, S., Diogène, J., Cano-Sancho, G., Sloth, J.J., Kwadijk, C., Barcelo, D., Allegaert, W. and Bekaert, K., 2015.** Environmental contaminants of emerging concern in seafood—European database on contaminant levels. *Environmental Research*, 143, 29-45.
- Vidayanti, V. and Retnaningdyah, C., 2024.** Microplastic pollution in the surface waters, sediments, and wild crabs of mangrove ecosystems of East Java, Indonesia. *Emerging Contaminants*, 10(4), 100343. DOI:10.1016/j.emcon.2024.100343
- Wang, F., Wu, H., Wu, W., Wang, L., Liu, J., An, L. and Xu, Q., 2021.** Microplastic characteristics in organisms of different trophic levels from Liaohe Estuary, China. *Science of The Total Environment*, 789, 148027. DOI:10.1016/j.scitotenv.2021.148027
- Wright, S.L., Thompson, R.C. and Galloway, T.S., 2013.** The physical impacts of microplastics on marine organisms: a review. *Environmental Pollution*, 178, 483-492. DOI:org/10.1016/j.envpol.2013.02.031
- Zafar, F.H.S., Zarrien, A., Karim, A., Zahid, M. and Levent, B., 2018.** Seasonal variations in physico-chemical parameters of Buleji and Paradise Point rocky shores at Karachi coast. *International Journal of Environment and Geoinformatics*, 5(2), 154-168. DOI:0.30897/ijegeo.418709
- Zhao, S., Liu, Y., Sun, C., Wang, X., Hou, C., Teng, J., Zhao, J., Fang, Y. and Wang, Q., 2024.** The pollution characteristics and risk assessment of microplastics in mollusks collected from the Bohai Sea. *Science of The Total Environment*, 913, 169739.