

Research Article

Feeding habits of the marbled flounder *Pseudopleuronectes yokohamae* (Günther, 1877) in the coastal waters off Pohang, East Sea of Korea

Cho J.I.¹, Kim M.S.¹, Kim D.J.¹, Son J.Y.¹, Choi H.Y.², Moon S.Y.², Park J.H.³, Baeck G.W.^{1*}

1 Department of Marine Biology and Aquaculture Science, Institute of Marine Industry, Marine Bio-Education and Research Center, College of Marine Science, Gyeongsang National University, Tongyeong-si 53064, Republic of Korea

2 South Sea Fisheries Research Institute, National Institute of Fisheries Science, Yeosu 59780, Republic of Korea

3 Distant Water Fisheries Resources Division, National Institute of Fisheries Science, Busan 46083, Republic of Korea

Correspondence: gwbaeck@gnu.ac.kr

Keywords

Pseudopleuronectes yokohamae,
East Sea of Korea,
Feeding habits,
Trophic level

Abstract

The diet composition of the marbled flounder *Pseudopleuronectes yokohamae* was examined using 975 specimens collected from January to December 2022 in the coastal waters off Pohang, East Sea of Korea. The range of total length was 14.4–47.6 cm. Polychaetes were the most important prey item, accounting for 91.9% of the index of relative importance, with the Lumbrineridae family being the most dominant, followed by Ampharetidae and Terebellidae. The trophic level of *P. yokohamae* was 3.08. Polychaetes showed no significant differences in dietary composition among size classes (<25.0 cm, 25.0–30.0 cm, 30.0–35.0 cm, and ≥35.0 cm), although the mean prey weight per stomach differed significantly with size. Polychaetes were the dominant prey item across all seasons. Both the mean prey weight per stomach and mean number of prey items per stomach varied significantly by season. A two-way PERMANOVA revealed that both size class and season had significant effects on dietary composition ($p > 0.05$). To assess the nutritional status of individuals, the condition factor was calculated. It was lowest in March and highest in November. Therefore, increase in feeding activity and prey intake during spring is likely associated with post-spawning energy recovery.

Article info

Received: June 2025

Accepted: August 2025

Published: January 2026



Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Introduction

The marbled flounder *Pseudopleuronectes yokohamae*, belonging to the family Pleuronectidae within the order Pleuronectiformes, is a coastal benthic fish species distributed from the East China Sea to southern Hokkaido, including the coastal waters of Korea (NIFS, 2014). Moreover, *P. yokohamae* is the most dominant pleuronectid species inhabiting the coastal waters off Pohang in the East Sea of Korea, migrating to nearshore areas shallower than 40 m during its spawning season between December and February (Hong *et al.*, 2008; Kim *et al.*, 2016; Tomiyama *et al.*, 2021).

Owing to its high commercial value as a seafood resource, *P. yokohamae* has been subject to resource conservation measures in Korea, including the establishment of a closed fishing season from December 1 to January 31 and minimum legal catch size (Huh *et al.*, 2012; Kim *et al.*, 2016). Although the catch of *P. yokohamae* is aggregated under pleuronectid fishes in Korean fisheries statistics, making it difficult to determine species-specific catch data, the annual average catch of pleuronectid fishes, including *P. yokohamae*, in Gyeongsangbuk-do Province was 4,344 tons in 2011, declined to 2,772 tons in 2016, and subsequently increased to 5,254 tons in 2024 (KOSIS, 2025). Previous studies have shown that *P. yokohamae* reaches 50% of its population maturity size (50% maturity size) at approximately three years of age (Kim *et al.*, 2016; Yang *et al.*, 2017). Given this relatively long maturation period, fishing activities are expected to have a considerable impact on the population dynamics of *P. yokohamae*. Therefore,

biological studies of the species are essential for effective monitoring and sustainable management.

Previous studies on the feeding habits of *P. yokohamae* have been conducted in various regions, including the central Yellow Sea (Park *et al.*, 2016; Roh *et al.*, 2022) and Gwangyang Bay (Kwak and Huh, 2003) and Tongyeong in the South Sea (Huh *et al.*, 2012) in Korea, as well as in the Seto Inland Sea (Hata *et al.*, 2016), Sendai Bay (Takahashi *et al.*, 2018), and Tokyo Bay (Lee *et al.*, 2019) in Japan. However, to date, no studies have investigated the feeding habits of *P. yokohamae* inhabiting the coastal waters off Pohang in the East Sea, despite existing research on its reproductive biology and growth (Kim *et al.*, 2016; Yang *et al.*, 2017).

As feeding habits can vary by habitat and environmental conditions, the analysis of stomach contents is fundamental to understanding the ecological role of a species in different regions (Zhang, 2006; Park and Gwak, 2009; Kim *et al.*, 2022). Because *P. yokohamae* is known to occur year-round in the coastal waters off Pohang (Hong *et al.*, 2008), and resource enhancement programs such as juvenile releases are actively underway, it is crucial to investigate its prey composition to support successful resource enhancement efforts.

Therefore, the objective of this study was to analyze the stomach content composition of *P. yokohamae* inhabiting the coastal waters off Pohang in the East Sea of Korea, identify its primary prey items, and assess variations of prey composition by size and season. The

findings provide essential ecological information for the ecosystem-based management of *P. yokohamae*.

Materials and methods

Sample collection and stomach content analysis

Specimens of *P. yokohamae* used in this study were collected monthly from January to December 2022 by gillnet fisheries in the coastal waters off Pohang, East Sea of Korea. Each gill net was 600 meters in length with a mesh size of less than 20 cm. The nets were soaked for approximately three days and deployed more than once per month until the target sample size was obtained. After capture, the total length (TL, cm) and body weight (BW, g) of each specimen were measured at the South Sea Fisheries Research Institute, Korea National Institute of Fisheries Science (NIFS). Subsequently, the stomachs were excised, preserved in 10% neutral formalin, and transported to the Marine Animal Resources Laboratory at Gyeongsang National University for analysis.

Stomach contents were analyzed under a dissection microscope (LEICA 12; LEICA, Wetzlar, Germany), and prey items were identified to the lowest possible taxonomic level. The wet weight of each prey item was measured to the nearest 0.001 g using a precision analytical balance (ME204TE/00; Mettler Toledo, Greifensee, Switzerland).

The stomach content analysis results were expressed as the frequency of occurrence (%F), numerical proportion (%N), and weight proportion (%W), calculated according to the following formulas:

$$\%F = A_i/N \times 100$$

$$\%N = N_i/N_{total} \times 100$$

$$\%W = W_i/W_{total} \times 100$$

Where, A_i is the number of fish containing prey item i , N is the total number of fish with nonempty stomachs, N_i is the number of prey item i , N_{total} is the total number of prey items, W_i is the wet weight of prey item i , and W_{total} is the total wet weight of all prey items.

The index of relative importance (IRI) for each prey category was calculated following the method described by Pinkas *et al.* (1971):

$$IRI = \%F \times (\%N + \%W)$$

The IRI was then standardized to a percentage (%IRI) as:

$$\%IRI = \frac{IRI_I}{\sum_{i=1}^n IRI} \times 100$$

Trophic level estimation

The trophic level (TL_k) of *P. yokohamae*, indicating its ecological position, was estimated using TrophLab (Pauly *et al.*, 2000), based on the following formula:

$$TL_k = 1 + \sum_j^n |P_j \times TL_j|$$

Where, P_j is the relative importance (%IRI) of the prey category j and TL_j is the trophic level of the prey item j . The trophic levels of the prey taxa were referenced from Pauly *et al.* (1998), Cortés (1999), and Ebert and Bizzarro (2007).

Size and seasonal grouping

To examine size-related differences in diet, specimens with nonempty stomachs were categorized into four size classes: <25.0 cm, 25.0–30.0 cm, 30.0–35.0 cm, and ≥35.0 cm. For seasonal comparisons, samples were grouped into spring (March–May; $n =$

73), summer (June–August; $n = 62$), autumn (September–November; $n = 70$), and winter (December–February; $n=42$).

To examine the dietary differences of *P. yokohamae* by size class and season, the data were grouped into subgroups consisting of 3 to 5 individuals for each size class and season. Since the number of individuals in the largest size class was small, the 30.0–35.0 cm and ≥ 35.0 cm groups were combined into a single ≥ 30.0 cm category for analysis. The resulting matrix was then subjected to two-way PERMANOVA to identify significant effects of three size classes (<25.0 cm, 25.0–30.0 cm, and ≥ 30.0 cm) and four seasons.

For each size class and season, the mean number of prey items per stomach (mN/ST) and the mean prey weight per stomach (mW/ST) were calculated. Statistical differences were analyzed using one-way

analysis of variance (ANOVA) performed with Microsoft Excel 365.

Condition factor

To assess the nutritional condition of *P. yokohamae* before and after the spawning season, the monthly condition factor (CF) was calculated using the following formula:

$$CF = BW(g)/TL(cm)^3 \times 10^2$$

$$CF = BW(g)/TL(cm)^3 \times 10^2$$

Where, BW is the body weight (g) and TL is the total length (cm).

Results

Total length distribution

In total, 975 specimens of *P. yokohamae* were collected during the study period, exhibiting a total length (TL) range of 14.4–47.6 cm (Fig. 1). The size class of 20.0–25.0 cm accounted for the highest proportion, representing 40.9% of the total collected specimens.




Figure 1: Location of sampling area in the coastal waters off Pohang, East Sea of Korea.

Stomach content composition and trophic level

The stomach contents of *P. yokohamae* were analyzed (Table 1). Of the 975 specimens examined, 728 individuals had empty stomachs, yielding an empty stomach rate of 74.7%. Of the 247

individuals with prey, polychaetes were the predominant prey item, with a frequency of occurrence of 77.7%, numerical proportion of 50.1%, and weight proportion of 61.6%. The relative importance index (%IRI) of polychaetas was 91.9%.

Table 1: Composition of the stomach contents of *P. yokohamae* collected from January to December 2022 in the coastal waters off Pohang, East Sea of Korea, showing frequency of occurrence (%F), number (%N), weight (%W), and index of relative importance (%IRI).

	Prey organism	%F	% N	% W	%IRI
Amphipoda		28.3	12.2	2.5	4.4
	<i>Ampelisca</i> sp.	9.3	2.6	0.3	
	<i>Byblis</i> sp.	10.5	4.7	0.4	
	Caprellidae	4.0	1.8	1.4	
	<i>Melita</i> sp.	6.5	1.1	0.2	
	<i>Themisto</i> sp.	0.4	+	+	
	Unidentified Amphipoda	13.4	2.0	0.2	
Brachyura		3.6	0.4	0.5	+
Crustacea		1.6	0.1	+	+
Cumacea		0.4	0.1	+	+
Euphausiacea		2.8	29.4	17.7	1.4
	<i>Euphausia</i> spp.	2.8	29.4	17.7	
Isopoda		1.2	0.1	+	+
Macrura		1.6	0.1	0.3	+
	Alpheidae	0.4	+	+	
	Unidentified Macrura	1.2	0.1	0.3	
Pycnogonida		1.6	0.2	+	+
Stomatopoda		1.2	0.2	0.3	+
Tanaidacea		1.6	0.2	+	+
Anthozoa		13.4	1.5	4.3	0.8
Aplysiidae		4.0	0.4	4.6	0.2
Astroidea		0.4	+	+	+
Bivalvia		3.6	0.3	1.0	+
Desmospongia		0.4	0.1	1.0	+
Gastropoda		1.6	0.1	0.1	+
Hydrozoa		3.6	0.3	0.3	+
Ophiuroidea		19.0	1.7	3.0	1.0
Ostracoda		1.6	0.2	+	+
Polychaeta		77.7	50.1	61.6	91.9
	Ampharetidae	14.2	7.4	3.4	
	<i>Amage</i> sp.	1.6	0.2	0.1	
	Aphroditidae	0.8	0.1	1.1	
	Cirratulidae	0.8	0.1	0.2	
	<i>Cirratulus</i> sp.	1.2	0.1	0.1	
	Eunicidae	9.7	3.0	3.0	
	Glyceridae	5.7	1.0	0.7	
	<i>Goniada</i> sp.	0.8	0.2	0.2	

Table 1 (continued):

	Prey organism	%F	%N	%W	%IRI
	Lumbrineridae	19.4	7.1	3.3	
	Nereididae	6.9	1.3	5.2	
	<i>Ophioglycera</i> sp.	0.4	0.1	+	
	Orbiniidae	0.4	+	0.1	
	Syllidae	1.2	0.1	0.1	
	Terebellidae	10.5	3.4	4.3	
	Unidentified Polychaeta	64.8	26.1	39.6	
Polyplacophora		0.4	+	+	+
	Chitonidae	0.4	+	+	
Sipunculida		3.6	0.3	2.6	0.1
Eggs		1.6	1.6	+	+
Seaweeds		2.0	0.2	+	+
Vinyls		0.4	+	+	+
	Total		100.0	100.0	100.0

Among the polychaetes, the Lumbrineridae family was the most dominant, followed by Ampharetidae and Terebellidae. Amphipods (Amphipoda) were the second most important prey group, showing a %IRI of 4.4%, based on a frequency of occurrence of 28.3%, numerical proportion of 12.2%, and weight proportion of 2.5%. Other prey items such as euphausiids (Euphausiacea), brittle stars (Ophiuroidea), and anthozoans (Anthozoa) were also found, but each accounted for less than 1.4% of the total %IRI.

The trophic level of *P. yokohamae* inhabiting the coastal waters off Pohang was estimated to be 3.08.

Ontogenetic changes in diet composition
To assess ontogenetic changes in the diet, the specimens were categorized into four size classes (Fig. 2). In all size classes, polychaetes were identified as the primary prey group (PERMANOVA, $p>0.05$).

Mean number and weight of prey items per stomach by size class

The mean number of prey items per stomach (mN/ST) and the mean prey weight per stomach (mW/ST) for each size class were analyzed (Fig. 3). The mN/ST values were 3.74 (± 0.50), 2.91 (± 0.26), 9.52 (± 4.41), and 2.61 (± 0.67) for the <25.0 cm, 25.0–30.0 cm, 30.0–35.0 cm, and ≥ 35.0 cm size classes, respectively, showing no statistically significant differences (one-way ANOVA, $p>0.05$).

The mW/ST values were 0.15 (± 0.02), 0.21 (± 0.03), 0.44 (± 0.15), and 0.24 (± 0.10) g for the <25.0 cm, 25.0–30.0 cm, 30.0–35.0 cm, and ≥ 35.0 cm size classes, respectively. The mean prey weight increased up to the 30.0–35.0 cm class and then decreased, indicating a statistically significant difference among size classes (one-way ANOVA, $p<0.05$).




Figure 2: Total length frequency distribution of *P. yokohamae* collected from January to December 2022 in the coastal waters off Pohang, East Sea of Korea.




Figure 3: Ontogenetic changes in the composition of stomach contents by the index of relative importance (%IRI) of *P. yokohamae* collected from January to December 2022 in the coastal waters off Pohang, East Sea of Korea.

Seasonal changes in diet composition

Seasonal variations in diet composition were analyzed by dividing the specimens

into spring, summer, autumn, and winter groups (Fig. 4). In all seasons, the polychaetes were the predominant prey

group. The %IRI of the polychaetes was highest in autumn (94.7%) and lowest in winter (82.5%) (PERMANOVA, $p>0.05$).




Figure 4: Variations in the mean (\pm standard error) number of prey items per stomach (mN/ST) and the prey weight per stomach (mW/ST) among the size classes of *P. yokohamae* collected from January to December 2022 in the coastal waters off Pohang, East Sea of Korea (<25.0 cm, $n = 104$; 25.0–30.0 cm, $n = 85$; ≥ 30.0 cm, $n = 58$).

Mean number and weight of prey items per stomach by season

Seasonal differences in the mean number and weight of prey items per stomach were examined (Fig. 5). The mN/ST values were 9.99 (± 4.21), 3.63 (± 0.43), 2.78 (± 0.35), and 2.59 (± 0.33) for spring, summer, autumn, and winter, respectively, showing significant seasonal variation (one-way ANOVA, $p<0.05$).

The mW/ST values were 0.54 (± 0.15) g, 0.13 (± 0.02) g, 0.14 (± 0.02) g, and 0.18 (± 0.03) g for spring, summer, autumn, and winter, respectively, also showing significant seasonal differences (one-way ANOVA, $p<0.05$).

Monthly changes in the condition factor

The monthly variation in the condition factor (CF) of *P. yokohamae* was analyzed (Fig. 6). The CF was lowest in March (mean: 1.22), increased to 1.30 in April, and then gradually increased to reach the highest mean value of 1.52 in November (Fig. 7).




Figure 5: Seasonal variations in the composition of stomach contents by the index of relative importance (%IRI) of *P. yokohamae* collected from January to December 2022 in the coastal waters off Pohang, East Sea of Korea.




Figure 6: Seasonal variations in the mean (\pm standard error) number of prey items per stomach (mN/ST) and the prey weight per stomach (mW/ST) of *P. yokohamae* collected from January to December 2022 in the coastal waters off Pohang, East Sea of Korea (Spring, $n = 73$; Summer, $n = 62$; Autumn, $n = 70$; Winter, $n = 42$).




Figure 7: Monthly changes in the mean (\pm standard error) value of the condition factor (CF) for *P. yokohamae* collected from January to December 2022 in the coastal waters off Pohang, East Sea of Korea.

Discussion

In this study, *P. yokohamae* exhibited a high empty stomach rate, with approximately three-quarters of individuals showing no stomach contents. High empty stomach rates in fish have been attributed to digestion during capture or transportation, piscivorous feeding habits, or a mismatch between feeding and capture times (Huh *et al.*, 2006; Choi *et al.*, 2019). Given that the specimens were captured using gillnets, it is likely that digestion occurred either within the fishing gear or during transportation, resulting in the observed high empty stomach rate.

The present results indicate that polychaetes are the most important prey group for *P. yokohamae*. Polychaetes, being less mobile, are considered good indicators of environmental changes and habitat characteristics. They primarily feed on organic matter in sediments or the water column and play a key role in energy

transfer to higher trophic levels (Koo *et al.*, 2008; Lee *et al.*, 2022). Moreover, previous studies have reported that *P. yokohamae* inhabiting the central Yellow Sea, South Sea of Korea, and the Seto Inland Sea and Sendai Bay of Japan primarily feed on polychaetes (Kwak and Huh, 2003; Hata *et al.*, 2016; Park *et al.*, 2016; Takahashi *et al.*, 2018; Roh *et al.*, 2022). However, in the coastal waters off Tongyeong, *P. yokohamae* primarily consumed bivalves, likely reflecting the dense distribution of bivalve aquaculture farms in that region (Huh *et al.*, 2012; Roh *et al.*, 2022). These findings suggest that the main prey of *P. yokohamae* may vary depending on local environmental conditions. Nonetheless, the dominance of polychaetes observed in this study is consistent with most previous research (Shin and Koh, 1992; Kim *et al.*, 2022; Lee *et al.*, 2023).

Among polychaetes, members of the Lumbrineridae family were the most

dominant. Shin and Koh (1992) and Han *et al.* (2015) reported that lumbrinerid polychaetes are abundant in the coastal waters off Pohang. Moreover, members of the Lumbrineridae family are recognized as indicator species for polluted environments, suggesting that their high abundance in this region may be associated with environmental degradation due to port facilities and wastewater inflow (Lee *et al.*, 2003; Lee *et al.*, 2023). Consequently, *P. yokohamae* is characterized as an opportunistic benthophagous carnivore primarily feeding on the abundant lumbrinerid polychaetes within its habitat.

The trophic level of *P. yokohamae* in this study was estimated at 3.08. In comparison, previous studies reported higher trophic levels for *P. yokohamae*, 3.33 in the Yellow Sea and 3.14 in Gwangyang Bay in the South Sea (Kwak and Huh, 2003; Roh *et al.*, 2022). This difference likely reflects the dominance of polychaetes (91.9% of %IRI) in the present study, whereas a greater diversity of prey such as gastropods, fish, and amphipods was reported in previous studies, leading to higher trophic levels. Thus, *P. yokohamae* in the coastal waters off Pohang functions as a key intermediary in the local food web, primarily consuming low-trophic benthic organisms such as polychaetes and serving as prey for higher predators such as Pacific cod (*Gadus macrocephalus*) and Black edged sculpin (*Gymnoanthus herzensteini*) (Yoon *et al.*, 2012; Yang *et al.*, 2013).

Regarding ontogenetic changes, *P. yokohamae* across all size classes primarily consumed polychaetes, and no evident dietary shift was observed with growth. However, *P. yokohamae* exhibited a dietary

shift from amphipods to polychaetes at a body length of 4.0 cm in Gwangyang Bay (Kwak and Huh, 2003) and at 8.0 cm in the Seto Inland Sea (Hata *et al.*, 2016). In the Yellow Sea, the importance of polychaetes decreased while that of gastropods increased with growth (Roh *et al.*, 2022). Because the specimens collected in the present study ranged from 14.4 to 47.6 cm in TL, smaller individuals (<14.0 cm) were not sampled. Juvenile *P. yokohamae* (≤ 10.0 cm) typically inhabits shallow seagrass beds or estuarine areas and migrate offshore as they grow (Kwak and Huh, 2003; Able *et al.*, 2005), which may explain the absence of smaller individuals in this study. Therefore, it is presumed that dietary shifts occur at sizes below 10.0 cm, as suggested by the findings of Kwak and Huh (2003) and Hata *et al.* (2016), with juveniles feeding more heavily on small invertebrates such as amphipods.

Analysis of the mean number and weight of prey items per stomach by size class revealed that although the mean number of prey items per stomach did not differ significantly, the mean prey weight showed statistically significant differences among size classes. Previous studies have reported varying patterns: *P. yokohamae* from Tongyeong exhibited significant differences in both mean prey number and weight with growth (Huh *et al.*, 2012), whereas individuals from the Yellow Sea exhibited significant differences in mean prey weight but not in prey number, consistent with the present study findings (Roh *et al.*, 2022). Such regional differences are likely attributable to variations in dominant prey types, with

bivalves dominating in Tongyeong and polychaetes in the Yellow Sea and Pohang.

Seasonal analysis indicated that polychaetes were the predominant prey across all seasons. Given their significant pollution in the coastal waters of Pohang, polychaetes, including lumbrinerids, which are known as pollution indicator species, are abundant year-round (Lee *et al.*, 2003; Lee *et al.*, 2023). The high abundance of polychaetes, coupled with limited escape ability compared to more mobile prey such as fish or cephalopods, suggests that *P. yokohamae* adopts an optimal foraging strategy with a high success rate.

Seasonal variations were also observed in the mean number and weight of prey items per stomach, with the highest values recorded in spring. The CF of *P. yokohamae* was lowest in March and increased thereafter until June. Considering that the spawning season for *P. yokohamae* in the East Sea occurs from December to February (Kim *et al.*, 2016), the observed increase in feeding activity and prey intake during spring is likely associated with post-spawning energy recovery (Bond, 1979; Balanov *et al.*, 2006; Baeck *et al.*, 2010). The results of this study are expected to serve as fundamental data for research on the stock structure of *P. yokohamae*, as well as studies on interspecific and intraspecific food competition and food web dynamics.

Conclusion

The coastal waters off Pohang in the East Sea are continuously affected by the inflow of industrial effluents and domestic wastewater from nearby industrial complexes (Lee *et al.*, 2003; Lee *et al.*, 2023). Due to the bay-shaped geography of

the region, these pollutants tend to accumulate in the benthic environment rather than being discharged outward. In this study, the marbled goby (*P. yokohamae*) was found to feed primarily on benthic prey organisms. These findings suggest that future changes in the benthic environment may directly affect the feeding ecology of *P. yokohamae*. In particular, the accumulation of pollutants and shifts in benthic community composition could alter the diet composition and feeding patterns of this species. This study may serve as fundamental data for ecological restoration and resource management efforts such as juvenile release programs and sea forest creation aimed at enhancing the stocks of *P. yokohamae*. In such efforts, the importance of ensuring a stable food supply and improving benthic habitat conditions should be emphasized. Moreover, the results provide valuable baseline information not only for understanding the stock structure of *P. yokohamae*, but also for future studies on interspecific and intraspecific food competition and food web dynamics.

Acknowledgements

This research was a part of the project titled 'Survey of coastal fisheries resources and marine environmental ecology in the South Sea' (grant number R2025011), funded by the National Institute of Fisheries Science (NIFS), Korea.

References

Able, K.W., Neuman, M.J. and Wennhage, H., 2005. Ecology of juvenile and adult stages of flatfishes: Distribution and dynamics of habitat

associations. *Flatfishes: Biology and Exploitation*, pp. 165–184. London: Blackwell Science. <https://doi.org/10.1002/9780470995259.ch8>

Baeck, G.W., Park, C.I., Jeong, J.M., Kim, M.C., Huh, S.H. and Park, J.M., 2010. Feeding habits of *Chaenogobius gulosus* in the coastal waters of Tongyeong, Korea. *Korean Journal of Ichthyology* 22, 41-48.

Balanov, A.A., Badaev, O.Z., Napazakov, V.V. and Chuchukalo, V.I., 2006. Distribution and some biological features of *Lycodes raridens* (Zoarcidae) in the western part of the Bering Sea. *Journal of Ichthyology*, 46, 148-155. <https://doi.org/10.1134/S0032945206020020>

Bond CE. 1979. Biology of Fishes. W.B., Saunders Co Philadelphia, 514.

Choi, H.C., Cho, J.H., Huh, S.H. and Park, J.M., 2019. Feeding habits of larval *Acanthogobius flavimanus* from the Nakdong River estuary, Korea. *Korean Journal of Ichthyology*, 31, 150-158. <https://doi.org/10.35399/ISK.31.3.4>

Cortés, E., 1999. Standardized diet compositions and trophic levels of sharks. *ICES Journal of Marine Science*, 56, 707-717. <https://doi.org/10.1006/jmsc.1999.0489>

Ebert, D.A. and Bizzarro, J.J., 2007. Standardized diet compositons and trophic levels of skates (Chondrichthyes: Rajiformes: Rajoidei). *Environmental Biology of Fishes*, 80, 221-237.

<https://doi.org/10.1007/s10641-007-9227-4>

Han, H.S., Ma, C.W. and Choi, M.S., 2015. Ecological variations of macrobenthos in subtidal at Chonsu bay in Chungcheongnam-do, Korea. *Korean Journal of Environmental Ecology*, 29, 3, 421-430. <https://doi.org/10.13047/KJEE.2015.29.3.421>

Hata, M., Sugimoto, R., Hori, M., Tomiyama, T. and Shoji, J., 2016. Occurrence, distribution and prey items of juvenile marbled sole *Pseudopleuronectes yokohamae* around a submarine groundwater seepage on a tidal flat in southwestern Japan. *Journal of Sea Research*, 111, 47-53. <https://doi.org/10.1016/j.seares.2016.01.009>

Hong, B.K., Kim, J.K., Park, K.D., Jeon, Y.Y., Hwang, K.S., Kim, Y.S. and Park, K.Y., 2008. Species composition of fish collected in gill nets from Youngil Bay, East Sea of Korea. *Journal of Korean Fisheries Society*, 41, 353-362.

Huh, S.H., Park, J.M. and Baeck, G.W., 2006. Feeding habits of John Dory *Zeus faber* in the coastal waters off Gori, Korea. *Korean Journal of Fisheries & Aquatic Sciences*, 39, 357-362. <https://doi.org/10.5657/kfas.2006.39.4.357>

Huh, S.H., Nam, K.M., Park, J.M., Jeong, J.M. and Baeck, G.W., 2012. Feeding habits of the Marbled flounder, *Pleuronectes yokohamae* in the coastal waters of Tongyeong, Korea. *Korean Journal of Ichthyology*, 24, 77-83.

Kim, D.G., Jeong, J.M., Lee, S.J., Kang, S.K., Seong, G.C., Kang, D.Y., Jin, S.Y. and Baeck, G.W., 2022. Comparison of stomach contents of yellow goosefish *Lophius litulon*, in the South Sea and Yellow Sea, Korea. *Korean Journal of Fisheries and Aquatic Sciences*, 55, 714-720.

Kim, S.R., Cha, H.K., Lee, J.B., Lee, H.W., Yang, J.H., Baek, H.J. and Kim, S.T., 2016. Maturity and spawning of the Marbled flounder *Pseudopleuronectes yokohamae* off the coast of Pohang, East Sea. *Korean Journal of Fisheries & Aquatic Sciences*, 49, 367-375. <https://doi.org/10.5657/KFAS.2016.0367>

Koo, B.J., Shin, S.H., Woo, H.J., Kim, E.S. and Je, J.G., 2008. Changes in macrobenthic community structure on Gunsan tidal flat after the closing of the Saemangeum 4th dyke. *Ocean & Polar Research*, 30, 497-507. <https://doi.org/10.4217/OPR.2008.30.4.497>

KOSIS (Korean Statistical Information Service), 2022. *Statistic databased for fishery production survey*. Retrieved from <http://kosis.kr/> on Oct. 31, 2023.

Kwak, S.N. and Huh, S.H., 2003. Feeding habit of *Limanda yokohamae* in the eelgrass (*Zostera marina*) bed in Kwangyang Bay. *Korean Journal of Fisheries & Aquatic Sciences*, 36, 522-527. <https://doi.org/10.5657/KFAS.2003.36.5.522>

Lee, M.K., Lim, D.I., Um, I.K., Shin, E.B. and Jung, H.S., 2003. Seasonal variation and spatial distribution of water qualities in Youngil bay, southeast coast of Korea. *JKSEE*, 25, 898-908.

Lee, J.H., Kodama, K., Shiraishi, H. and Horiguchi, T., 2019. Qualitative and quantitative changes in dietary composition of marbled flounder *Pseudopleuronectes yokohamae* between periods of different stock sizes in Tokyo Bay, Japan. *Aquatic Animals*, AA2019-2.

Lee, J.H., Lim, H.S., Shin, H.C. and Ryu, J.S., 2022. Spatial distribution of macrozoobenthic organisms along the Korean coasts in summer season. *Journal of Korean Society and Oceanography*, 27, 87-102. <https://doi.org/10.7850/jkso.2022.27.2.087>

Lee, M.O., Kim, J.K. and Kim, B.K., 2023. Marine environmental characteristics of Yeongil bay, Korea. *Journal of the Korean Society for Marine Environment & Energy*, 26, 11-33. <https://doi.org/10.7846/JKOSMEE.2023.26.1.11>

National Institute of Fisheries Science (NIFS), 2014. Technical report of national fisheries research and development institute. Busan, Korea: NIFS.

Park, C.Y. and Gwak, W.S., 2009. Comparison of stomach contents of pacific cod (*Gadus macrocephalus*) in Korean coastal waters. *Korean Journal of Ichthyology*, 21, 28-37.

Park, J.M., Kwak, S.N., Choi, H.C., Jawad, L.A. and Riedel, R., 2016. Diet patterns of the marbled flounder, *Pseudopleuronectes yokohamae*, in the mid-western coast of Korea. *Science*

International, 4, 94-100. <https://doi.org/10.17311/sciintl.2016.94100>

Pauly, D., Froese, R., Saa, P., Palomares, M.L., Christensen, V., and Rius, J., 2000. TrophLab in MS Access.

Pauly, D.V., Christense, J.D., Froese, R. and Torres, F., 1998. Fishing down marine food webs. *Science*, 279, 860-863. <https://doi.org/10.1126/science.279.5352.860>

Pinkas, L., Oliphant, M.S. and Iverson, I.L.K., 1971. Food habits of albacore, bluefin tuna, and bonito in California waters. *California Department of Fish and Game's Fish Bulletin*, 152, 1-105.

Roh, T.H., Choi, D.H., Lee, S.H., Kwon, D.H., Han, K.H. and Kim, M.J., 2022. Feeding habits of the marbled flounder *Pseudopleuronectes yokohamae* in the West Sea in Korea. *Journal of Fisheries & Aquatic Science*, 55, 583-589. <https://doi.org/10.5657/KFAS.2022.0583>

Shin, H.C. and Koh, C.H., 1992. Seasonal and spatial variation of polychaetous community in Youngil bay, southeastern Korea. *The Korean Society of Oceanography*, 27, 46-54.

Takahashi, S., Kurita, Y., Togashi, H. and Tomiyama, T., 2018. Diel feeding rhythms, daily ration, and seasonal changes thereof in marbled flounder *Pseudopleuronectes yokohamae*. *Journal of Applied Ichthyology*, 34, 888-895. <https://doi.org/10.1111/jai.13719>

Tomiyama, T., Yamada, M., Yamanobe, A. and Kurita, Y., 2021. Seasonal bathymetric distributions of three coastal flatfishes: Estimation from logbook data for trawl and gillnet fisheries. *Fisheries Research*, 233, 105733. <https://doi.org/10.1016/j.fishres.2020.105733>

Yang, J.H., Yoon, S.C., Park, J.H., Yoon, B.S., Choi, Y.M. and Lee, J.B., 2013. Feeding habits of the Black-edged sculpin *Gymnothorax herzensteini* in the coastal waters off Mukho, Gangwondo of Korea. *Korean Journal of Fisheries & Aquatic Science*, 45, 379-386. <https://doi.org/10.5657/KFAS.2012.0379>

Yang, J.H. and Lee, H.W., Lee, J.B., Cha, H.K., Kim, S.T. and Yoon, B.S., 2017. The age and growth of marbled flounder *Pseudopleuronectes yokohamae* in the coastal waters of East Sea off Pohang. *Journal of Fisheries & Aquatic Science*, 50, 278-286. <https://doi.org/10.5657/KFAS.2017.0278>

Yoon, S.C., Yang, J.H., Park, J.H., Choi, Y.M., Park, J.H. and Lee, D.W., 2012. Feeding habits of the Pacific cod *Gadus macrocephalus* in the coastal waters off Jumunjin, Gangwondo of Korea. *Korean Journal of Fisheries and Aquatic Sciences*, 45, 379-386

Zhang, C.I., 2006. A study on the ecosystem-based management system for fisheries resources in Korea. *Journal of Korean Society of Fisheries and Ocean Technology*, 42, 240-258. <https://doi.org/10.1016/j.fishres.2020.105733>