

Survey on the metazoan parasites in *Neogobius* fishes from Southeastern part of the Caspian Sea

Pazooki, J.^{1*}; Mansouri-Habibabadi; Z.¹; Masoumian, M.²;
Aghae-Moghadam, A.³

Received: December 2009

Accepted: April 2010

Abstract

In this study a parasitological investigation on 238 samples of *Neogobius* species from the south-east of Caspian Sea was carried out. The samples including *Neogobius melanostomus* (35), *N. fluviatilis pallasii* (103), *N. kessleri gorlap* (70) and *N. bathybius* (30) were seasonally seined by beach net during one year (summer 2006-spring 2007), fixed in 10 % buffer formalin and transferred to the laboratory. From 231 of examined fishes 235 specimens were infected (97/6 %). The collected parasites were as follows: *Hysterothylacium aduncum*, *Raphidascaris acus*, *Cucullanus sphaerocephalus*, *Dichelyne minutus*, *Raphidascaroides sp.*, *Cystidicola sp.*, *Gyrodactylus sp.*, and *Corynosoma strumosum*. Hereby, *N. melanostomus* is introduced as a new host for *Raphidascaroides sp.*, *C. sphaerocephalus* and *Cystidicola sp.*. *N. fluviatilis pallasii* and *N. kessleri gorlap* for *C. sphaerocephalus*, *Cystidicola sp.* and *H. aduncum*. *N. bathybius* for *C. sphaerocephalus*, *Cystidicola sp.*, *H. aduncum*, *E. excisus*, *R. acus* and *C. strumosum*. *Cystidicola sp.* and *H. aduncum* were reported for the first time from Iran. Among the investigated fishes *N. fluviatilis pallasii* possessed the highest and *N. bathybius* the lowest variety in infection with various species of parasites. The results of this study indicated that there are significant relations between total length (T.L.) and weight in *N. fluviatilis pallasii* with the number of *C. sphaerocephalus*, *D. minutus* and *E. excisus*. Also there are significant relations between total length and weight in *N. melanostomus* and between weights in *N. kessleri gorlap* with the number of *C. sphaerocephalus*.

Keywords: Neogobiidea, Nematoda, Acanthocephala, Monogenea, Caspian Sea

1-Department of Marine Biology, Faculty of Biological Sciences, Shahid Beheshti University, G. C., P. O. Box: 19834, Tehran, Iran.

2-Department of Fish Diseases, Iranian Fisheries Research Organization, P.O.Box:14155-6116, Tehran, Iran.

3-Asipenseridae Hatchery Center of Golestan Province, Gorgan, Iran.

*Corresponding author's email: pazooki2001@yahoo.com

Introduction

Many studies have been conducted on parasite fishes in the Northern part of the Caspian Sea; Bykhovskaya-Pavlovskaya et al. (1962), Lomaki (1970), Gibson (1972), Mikailov (1975), Gussev (1985) and Bauer et al. (2002). In Iran, also, many researches have been done on parasite fishes of the Southern part of the Caspian Sea by Eslami et al. (1972), Mokhayer and Anwaar (1973), Eslami and Kohneshahri (1978), Mokhayer (1983), Ghorogi and Pourgholam (1996), Pazooki and Masoumian (2002), Sattari et al. (2002, 2003), Khara et al. (2005) and Nik-Syrat et al. (2006). Altogether 129 species of parasites were reported from Neogobiidae fishes in the Northern and Southern part of the Caspian Sea; 126 species from *N. melanostomus*, 17 from *N. kessleri* and 13 from *N. fluviatilis*. Acipenseridae fishes of the Caspian Sea are very important as commercial fishes. The role of Neogobiidae fishes in the food chain of Acipenseridae, are emphasized. The aims of the research are to investigate parasitic fauna of Neogobiidae fishes in the Southern part of the Caspian Sea, to find out the relations between parasitic infections and length and weight of the examined fishes and also to survey on intensity and prevalence of the infections. The relations between the seasonality and the sex of hosts with infections are discussed as well.

Materials and methods

This study has been done in four seasons from summer 2006 to spring 2007. A total of 238 samples of four different *Neogobius* species including *Neogobius melanostomus* (35), *N. fluviatilis pallasi*

(103), *N. kessleri gorlap* (70) and *N. bathybius* (30) were seined by beach nets. The samples were collected from Southern part of the Caspian Sea, North of Golestan Province, fixed in 10 % buffer formalin, transferred to the laboratory and weighted and measured. Then, they were examined for different parasites. The collected parasites were identified according to Bykhovskaya et al., 1964, Moravec, 1994 and Jalali, 1999. Data analyses were done according to SPSS15 and SPSS9. The percentage of prevalence infections in each fish and season were done as well. Table 1 indicates the sex, length and weight of the examined fishes in each season.

Results

During this study 9 different genera of parasites were detected from 231 fish (97.6%); *Gyrodactylus*, *Cucullanus*, *Dichelyne*, *Cystidicola*, *Eustrongylides*, *Hysterothylacium*, *Raphidascaris*, *Raphidascoroides* and *Corynosoma* spp. were detected and only 7 fish specimens (2.4%) were not infected. Table 2 indicates the names of different parasites in their hosts.

Gyrodactylus sp.

Gyrodactylus sp. was found in the gill of *N. fluviatilis pallasi* only in the winter. The dimensions of the parasite were: length 0.36 mm, width 0.102 mm, ir (Innerroot) 0.0243 mm, vb (Ventral anchor) 0.019 and pr (Point recure) 0.016 mm. The prevalence was 10 % (2) (Fig. 1; A, B, C). *Dichelyne minutes* (Rudolphi, 1819).

This parasite was only collected from the intestine of the *N. fluviatilis pallasi* in the summer. The body length was 1.76-3.17 mm, width was 0.29-0.51

mm, length of the pharynx was 0.44-0.64 mm, and the conical tail was 0.071-0.16 mm. The prevalence was 36% (11), the

mean intensity 3.5 and the range was 1-7 (Fig. 1; D, E, F). Table 3 indicates infections in different seasons.

Table 1: The sex, length and weights of the examined fishes in each season

Fishes	Sex	No.	Summer	Autumn	Winter	Spring	Total	Length	Weigh
			2006	2006	2006	2007	No.	Cm	gr
<i>N. melanostomus</i>	♂	10	35	-	-	-	35	4.8-8.9	1.5-10.8
<i>N. fluviatilis</i>	♀	25							
<i>N. fluviatilis</i>	♂	53	30	32	20	21	103	6.9-25.3	4.73-268.8
<i>N. pallasii</i>	♀	50							
<i>N. kessleri</i>	♂	43	-	33	19	18	70	16.4-27.9	88-292.6
<i>N. gorlap</i>	♀	27							
<i>N. bathybius</i>	♂	25	-	-	10	20	30	20.7-30	101.2-296.2
	♀	5							

Table 2: The different parasites in 4 *Neogobius* fishes

parasites	<i>N. melanostomus</i>	<i>N. fluviatilis pallasii</i>	<i>N. kessleri gorlap</i>	<i>N. bathybius</i>
	<i>Gyrodactylus sp.</i>	-	+	-
<i>C. sphaerocephalus</i>	+	+	+	+
<i>Cystidicola sp.</i>	+	+	+	-
<i>D. minutus</i>	-	+	-	-
<i>E. excisus</i>	-	+	+	+
<i>H. aduncum</i>	+	+	+	-
<i>R. acus</i>	+	+	+	+
<i>Raphidascares sp.</i>	+	-	-	-
<i>Corynosoma strusoma</i>	-	+	+	+

***Cucullanus sphaerocephalus* (Rudolphi, 1809)**

This parasite was collected in four seasons and in the intestine of four hosts. The body length was 3.3-20.6 mm and the maximum width was 0.1-0.4 mm. The length of the pharynx and spicules were 0.44-0.64 mm and 0.3-0.5 mm respectively. The genital pole of the females was 6.5-10.3 mm from the posterior end. The conical tail was 0.2-

0.5 mm. The prevalence was 36% (11), the mean intensity 3.5 and the range was 1-7

***Hysterothylacium adunacum* (Rudolphi, 1802)**

In this research 14 of these parasites were collected from the intestine of *N. fluviatilis pallasii*, (in summer and winter) and *N. kessleri gorlap* (in autumn, winter and spring). The cysts of parasites were in the intestinal wall. The body length and

maximum width were 1.7-2.9 mm and 0.07-0.3 mm respectively. The posterior part of the conical tail was covered with spin cuticle and the length was 0.2 mm. The length of the pharynx was 0.1-0.7 mm. The parasites have downward secum, the length of it was 1/3 of the pharynx length. The intestine was dark and direct. (Fig. 2; A, B). Table 3 indicates infections in different seasons.

***Raphidascaris acus* (Bloch, 1770)**

In this research 6 of the parasites were collected from the intestinal wall of *N. melanostomus* and *N. fluviatilis pallasi*, (in autumn), *N. kessleri gorlap* (in autumn and winter) and *N. bathybius* (in winter). The body length was 3.29 mm, the maximum width 0.2 mm, the length of the downward secum 0.45 mm and the pharynx length was 0.357 mm. The nerve ring was 0.153 mm from the interior end of the body. The tail was conical and short (Fig.2; C, D). Table 3 indicates infections in different seasons.

***Raphidascaroides* sp.**

This parasite was found only in the abdominal cavity of two *N. melanostomus* in the summer. The body length and maximum width were 22.5-24 mm and 0.51 mm respectively. The body surface was covered with very thick cuticle. The pharynx length was 3.14 mm and the conical tail 0.188 mm. (Fig. 2; E, F). The prevalence of infections was 5.2% and the mean intensity was 1.

***Eustrongylides excises* (Jägerskiöld, 1909)**

The third larvae stage of this parasite was found only in *N. melanostomus* and *N. fluviatilis pallasi* in autumn, winter and spring. The body length and maximum width were 20-30 mm and 0.3-0.8 mm

respectively. The apical end of the body was round with 12 papillae in two row circles. The cuticle was rather thick and the genital pore was opened in the end of the body. The color of these parasites was dark red. The mesentery muscles, intestine and abdominal cavity of fishes were infected by cysts and the larvae of parasites (Fig. 3; A, B). Table 3 indicates infections in different seasons.

***Cystidicola* sp.**

In this study 8 specimens of these parasites were found in the intestine of *N. melanostomus*, *N. fluviatilis pallasi* and *N. kessleri gorlap* in autumn. The body length, maximum width and the pharynx length were 0.31-0.46 mm, 2.75 mm and 0.15-0.2 mm respectively. The tail was round and the length was 0.105-0.16 mm. The abdomen was elongated and narrow. The nerve ring up to the anterior end was 0.18 mm. The prevalence of infections in *N. melanostomus* (in summer), *N. fluviatilis pallasi* and *N. kessleri gorlap* (in autumn) were 2.8%, 3.1% and 3% respectively (Fig. 3; C, D).

***Corynosoma strumosum* (Rudolphi, 1802)**

This parasite was collected from *N. fluviatilis pallasi*, *N. kessleri gorlap* and *N. bathybius* in winter and spring. The body length was small, males and females were the same size; 75 mm. The rostrum had 18 torn rows, in each 10 to 12 torn. The fifth and seventh rows were big (Fig.3; E, F). Table 4 indicates infections in winter and spring.

The relation between length and weight of fishes with infection

The results show that there is a significant direct positive relation, between length and weight of *N. melanostomus*, *N. fluviatilis*

pallasi and *N. kessleri gorlap* with infections of *Cucullanus sphaerocephalus* (Figs. 4-10). There is also a significant direct negative relation, between length and weight of *N. fluviatilis pallasi* with

infections of *Dichelyne minutus* (Figure 10-11). The results also show that there is a significant relation between infections of *C. sphaerocephalus*, *E. excisus*, and *D. minutus* in different seasons (Tables 5, 6).

Table 3: Infections of Nematoda in different seasons*

Seasons	Fs	IF	%	NP	MI	IF	%	NP	MI	IF	%	NP	MI	IF	%	NP	MI
		<i>Cs</i>	<i>Cs</i>	<i>Cs</i>	<i>Cs</i>	<i>Ha</i>	<i>Ha</i>	<i>Ha</i>	<i>Ha</i>	<i>Ra</i>	<i>Ra</i>	<i>Ra</i>	<i>Ee</i>	<i>Ee</i>	<i>Ee</i>	<i>Ee</i>	<i>Ee</i>
Summer	<i>Nm</i>	29	83	357	12	1	3	1	1	1	3	1	1	-	-	-	-
	2006 <i>Nfp</i>	18	60	72	4	4	13	8	2	1	3	2	2	-	-	-	-
Autumn	<i>Nkg</i>	23	70	121	5	2	6	2	1	1	3	1	1	25	75	80	3
	2006 <i>Nfp</i>	28	88	155	6	-	-	-	-	-	-	-	-	20	63	37	2
Winter	<i>Nkg</i>	18	95	227	13	1	5	1	1	1	5	1	1	13	68	54	4
	2006 <i>Nfp</i>	14	70	229	16	1	5	1	1	-	-	-	-	11	55	36	3
	<i>Nb</i>	9	90	136	15	-	-	-	-	1	10	1	1	5	50	5	1
Spring	<i>Nkg</i>	12	67	93	8	1	6	1	1	-	-	-	-	11	61	45	4
	2007 <i>Nfp</i>	10	48	18	2	-	-	-	-	-	-	-	-	10	68	24	2
	<i>Nb</i>	11	55	77	7	-	-	-	-	-	-	-	-	9	45	40	4

*Fs: Fish species, IF: Infected fishes, *Cs*: *Cucullanus sphaerocephalus*, *Nm*: *N. melanostomus*, %: Prevalence of infections, *Ha*: *Hysterothylacium aduncum*, *NfP*: *N. fluviatilis pallasi*, NP: Number of parasites, *Ra*: *Raphidascaris acus*, *Nkg*: *N. kessleri gorlap*, MI: Mean intensity, *Ee*: *Eustrongylides excisus*, *Nb*: *N. bathybius*

Table 4: Infections of *Corynosoma strumosum* in different seasons*

Seasons	Fs	IF	%	NP	MI
Winter	<i>N. k. g</i>	5	26.3	7	1.4
	2006 <i>N. f. P</i>	2	10	3	1.5
	<i>N. b</i>	1	10	1	1
Spring	<i>N. k. g</i>	3	16.7	4	1.3
	2006 <i>N. f. P</i>	2	9.5	3	1.5
	<i>N. b</i>	4	20	5	1.2

*Refer to the abbreviations of table 3

Table 5: Difference of infections in different season*

Seasons	Fs	IF	%	NP	MI
Winter 2006	<i>N. k. g.</i>	5	26.3	7	1.4
	<i>N. f. P</i>	2	10	3	1.5
	<i>N. b</i>	1	10	1	1
Spring 2006	<i>N. k. g.</i>	3	16.7	4	1.3
	<i>N. f. P</i>	2	9.5	3	1.5
	<i>N. b</i>	4	20	5	1.2

*Refer to the abbreviations of table 3

Table 6: Differences between infections in different hosts' sex*

Parasites	Chi-square			df			Asymp.sig		
	<i>N.f.p.</i>	<i>N.k.g.</i>	<i>N.b</i>	<i>N.f.p.</i>	<i>N.k.g.</i>	<i>N.b</i>	<i>N.f.p.</i>	<i>N.k.g.</i>	<i>N.b</i>
<i>C. sphaerocephalus</i>	1.586	0.102	0.480	1	1	1	0.208	0.750	0.488
<i>E. excisus</i>	0.713	0.347	0.107	1	1	1	0.398	0.556	0.743
<i>D. minutus</i>	0.178	---	---	1	---	---	0.673	---	---
<i>Corynosoma strumosum</i>	0.004	0.498	1.500	1	1	1	0.953	0.480	0.221

*Refer to the abbreviations of table 3




Figure 1: A; *Gyrodactylus* sp., B; Opisthaptor of *Gyrodactylus* sp., C; The anterior of *Gyrodactylus* sp.; D; The anterior of *Dichelyne Minutus*, E; The posterior of *D. Minutus*, F; *Cucullanus sphaerocephalus*




Figure 2: A; The posterior of *Hysterothylacium aduncum*, B; The anterior of *H. aduncum*, C; The anterior of *Raphidascaris acus*, D; The posterior of *R. acus*., E; The anterior of *Raphidascaroides* sp., F; The posterior of *Raphidascaroides* sp.,




Figure 3: A; The anterior of *Eustrongylides excisus*, B; The posterior of *E. excisus*, C: The anterior of *Cystidicola* sp. D: The posterior of *Cystidicola* sp., E; *Corynosoma strumosum*, F; The rostrum of *Corynosoma strumosum*




Figure 4: Relationship between weight and infection of *C. sphaerocephalus* in *N. melanostomus*




Figure 5: Relationship between length and infection of *C. sphaerocephalus* in *N. fluviatilis pallasii*




Figure 6: Relationship between weight and infection of *C. sphaerocephalus* in *N. fluviatilis pallasii*




Figure 7: Relationship between weight and infection of *C. sphaerocephalus* in *N. kessleri gorlap*




Figure 8: Relationship between length and infection of *E. excisus* in *N. fluviatilis pallasii*




Figure 9: Relationship between weight and infection of *E. excisus* in *N. fluviatilis pallasii*




Figure 10: Relationship between length and infection of *D. minutus* in *N. fluviatilis pallasii*




Figure 11: Relationship between weight and infection of *D. minutus* in *N. fluviatilis pallasii*

Discussion

The fish sampling area is located in the Sarmatian Fauna Region, which is one of the three faunal regions of freshwater fishes of Iran. Neogobiidae fishes are spread through this region. It is expected that the parasitic fauna in the sampling area are the same as the neighboring territories. In Iran only a few researches were done on Neogobiidae parasites (Pazooki and Aghmandi, 1999, Daghigh-Rouhi and Sattari, 2004). The collected parasites found in this study are the same as the ones found in other examinations made in the basin of Northern parts of the Caspian Sea, Black Sea, Azov and other rivers. During this study, *Gyrodactylus* sp. was reported for the first time from the gills of *N. fluviatilis pallasii* in Iran. Before this study only a single *Gyrodactylus* sp., *G. protesrohini* was reported from the Northern part of the Caspian Sea; from *N. melanostomus* in the Russian and Turkish part of the Black Sea and Azov, Danube River, and from *N. kessleri gorlap*, in Danube River (Naydenova, 1974; Gaevskaya et al., 1975; Ondrachova et al., 2005; Oumlzer 2007).

Another parasite *Cucullanus sphaerosephalus*, was collected from 4 species of Neogobiidae. Up to this time only *C. heterochrous* has been reported from *N. melanostomus* from the Black Sea (Naydenova, 1974; Gaevskaya, et al., 1975). This parasite is specific for Acipenseridae and was reported from Europe (Bauer et al., 2002) and from the Iranian part of the Caspian Sea (Mokhayer and Anwar, 1973; Ghoroghi and Pourgholam, 1995; Sattari et al., 2002). Finding this parasite from both Neogobiidae and Acipenseridae shows

that, the first fish group has an important role of transferring the parasite to the second one. Because of a significantly positive relation between the length and weight by infections, the bigger infected fishes are more sensitive than the smaller ones.

The next parasite, *Cystidicola* sp. was found in the intestine of *N. melanostomus*, *N. fluviatilis pallasii* and *N. kessleri gorlap*. This parasite was reported from Salmonidae fishes in Europe, North America and Asia (Bykhovskaya et al., 1964). During this study *N. melanostomus*, *N. fluviatilis pallasii* and *N. kessleri gorlap* were reported as new hosts. The only species reported from *N. melanostomus* was *Cystidicola epherneridarum* (Rolbiecki, 2006).

D. minutus has already been reported from different parts of Europe (Moravec, 1994) and Southern parts of the Caspian Sea (Pazooki and Aghmandi, 1999; Daghigh-Roihi and Sattari, 2004). Markowski in 1996 assumed that *Polychaeta nereis diversicolour* is the intermediate host for the parasites, if this theory is accepted the reason for more infections in the younger (smaller size) hosts is due to the their nutrition diet changes during growth. In this study there were no infections in the large aged fishes and there was a significant negative direction between length and infections (Figs. 7, 8).

Eustrongylides excisus had already been reported from the *N. fluviatilis pallasii* in the Danube River (Ondraková et al., 2005) and from *N. melanostomus* in the Black Sea (Naydenova, 1974). In Iran this parasite was reported from Acipenseridae,

Cyprinidae and some Neogobiidae fishes (Sattari et al., 2003; Daghigh-Rouhi and Sattari, 2004). In this study *Eustrongylides excisus* is reported for the first time from *N. bathybius*.

Hysterothylacium aduncum was reported from *N. melanostomus* in Poland, Gdansk Gulf and the Black Sea (Rolbiecki, 2006; Kvach and Skora, 2007; Oumlzer, 2007). During this study, *N. fluviatilis pallasi*, *N. kessleri gorlap* and *N. bathybius* were presented as new hosts and this was the first time they were reported from Iran. *Raphidascaris acus* has already been reported from many fish hosts throughout the world (Moravec, 1994) and from Iran (Eslami et al., 1978; Pazooki and Masoumian 2002; Khara et al., 2005). This parasite was reported from *N. fluviatilis pallasi*, *N. kessleri gorlap* and *N. melanostomus* (Ondrackva et al., 2005) but in this study it was also reported from *N. bathybius* for the first time.

N. melanostomus is introduced as a new host for *Raphidascaroides sp.* This parasite was reported from fishes of different parts of the world and Iran (Pazooki and Masoumian 2002, Sattari et al., 2002).

Corynosoma strumosum has already been reported from Acipenseridae, Neogobiidae and Clupeidae in Iran (Sattari et al., 2002; Daghigh-Rouhi and Sattari 2004; Nik-Sirat et al., 2006). Earlier it was presumed that only Clupeidae could transfer this parasite to Acipenseridae fishes but later on it was detected that Neogobiidae fishes can also do so. This parasite was reported for the first time from *N. bathybius* in Iran.

This study revealed that there is a positive direct significant relation between

length and weight of *N. fluviatilis pallasi*, *N. kessleri gorlap* and *N. melanostomus* with the number of *C. sphaerocephalus*, (Figs. 4-7). The same correlation is between length and weight of *N. fluviatilis pallasi* with the number of *Eustrongylides excisus* (Figs. 8-11). There is a significant difference between infections of *C. sphaerocephalus*, *Eustrongylides excisus* and *D. minutus* in different seasons (Table 3). The reason was due to the ecological accepts and nutritional behavior in different seasons (Reimchen and Nosil, 2001).

References

- Bykhovskaya – Pavlovskaya, I. E., Gussev, A. V., Dubinina, M. N., Izyumova, N. A., Smirnova, T. S., Sokolovskaya, A. L. et al., 1962.** Key to parasites of freshwater fishes of the U.S.S.R. Publication House of the V.S.S.R. Academy of Sciences. Moscow- Leningrad, pp: 776. (In Russian)
- Bauer, O. N., Pugachev, O. N., Voronin, V., 2002.** Study of parasites and diseases of sturgeon in Russia: a review. *Journal of Applied Ichthyology*. XI, 420-429.
- Daghigh-Rouhi, J. and Sattari, M., 2004.** Survey on incidence of parasitic infection in the Gobiidae fishes from South-Western part of the Caspian Sea. *Journal of Veterinary Medicine Faculty*, 59 (1), 22-17. (In Persian)
- Eslami, A. and Kohneshahri, M., 1978.** Study on the helminthiasis of *Rutilus frisii katum* from south Caspian Sea. *Acta Zoologica Et Pathologica Antverpiensia*, 70, 153-155.

- Eslami, A., Anwar, M. and Khatibi, Sh., 1972.** Incidence and intensity of helminthiasis in pike (*Esox lucius*) of the Caspian Sea (Northern Iran). *Rivista Italiana Di Pisciocoltura Ed Ittiopatologia*, 1, 11-14.
- Gaevskaya, A. V., Gusev, A. V., and Delyamure, S. L., 1975.** Key to the Parasites of vertebrates of the Black Sea and Sea of Azov. Parasitic invertebrates of fishes, fish-eating birds and marine animals, Nauk. Dumka, Kiev, 552 pp. (In Russian)
- Ghoroghi, A., and Pougholam, R., 1996.** Identification the parasitic diseases of *Huso huso*. *Journal of Pajouhesh & Sazandegi*, 28 (In Persian).
- Gibson, D. I., 1972.** Contributions to the life – histories and development of *Cucullanus minutus* Rudolphi, 1819 and *C. heterochrous* Rudolph: 1802 (Nematoda :Ascaridata). *Bulletin of British Museum Natural History. Zoology*, 22, 153-170.
- Golovin, J. V., and Mokhayer, B., 1973.** Acanthocephales des Estur geo ns de la mer Caspiane . *Annales De Institut Parasitologie Humaine Et Comparee*, 4, 597-602.
- Golovin, A. V., 1985.** Parasitic Metazoans, (In Russian), In Bauer, O.N (Ed): key to parasites of freshwater fishes in U.S.S.R., vol.3 .Nauka, Leningrad.
- Gushev, A. V., 1985.** Parasitic Metazoan, Class-Monogenidae (In Russian), in Bauer, O. N. (Ed): Key to the parasites of freshwater fish of the USSR, Vol., 2. Nauka, Leningrad.
- Jalali, B., 1999.** Parasites and parasitic Diseases in the freshwater fishes of Iran. Iranian Fisheries Company, Aquaculture Department, 652pp. (in Persian).
- Khara, H., Sattari, M., Nezami, Sh. A., Mosawi, S. A. and Jaafarzadeh, E., 2005.** Survey on prevalence and intensity parasitic infection *Esox lucius* from Amir-Kolateh, Lahijan. *Journal of Veterinary Faculty*. 57(4), 333-340. (In Persian)
- Kvach, Y. and Skora, K., 2007.** Metazoan parasites of the invasive round goby *Apollonia melanostoma* (*Neogobius melanostomus*) (Pallas) (Gobiidae: Osteichthyes) in the Gulf of Gdańsk, Baltic Sea, Poland: a comparison with the Black Sea. *Parasitology Research*, 100, N. 4. pp. 767-774(8)
- Lomaki, V. V., 1970.** Distribution and some question of the biology of *Cucullanllus minutus* (Rud, 1819) (Nematode: Camallanata) in fishes of the Caspian Sea. *Voprosy morskoy parazitologi.*, Izd. Naukova Dumka, kiev, pp: 68 - 69. (In Russian).
- Markowski, S., 1966.** The diet and infection of fishes in Cavendish Dock, Barrow-in-Furness. *journal of zoology London*, 150, 183-197
- Mikhailov, T. K., 1975.** Parasite fauna of fishes of the basins of Azerbaidzhan. *Izd. Elm, Baku*, pp: 299. (In Russian)

- Mokhayer, B. and Anwar, M., 1973.** Effect pathogen des parasites de l'esturgeon dans le milieu naturel et artificiel. *Rivista Italiana Di Piscicoltura Ed Ittiopatologia*, 4, 111-115.
- Mokhayer, B., 1983.** Parasites and parasitic diseases of fish. The Fifth International Symposium of Ichthyoparasitology, *Ceske Budejovice*, 8-13 Aug.
- Moravec, F., 1994.** Parasitic Nematodes of Freshwater Fishes of Europe, Kluwer Academic publisher.
- Naydenova, N. N., 1974.** Parasite fauna of fishes of the goby family from the Black Sea and Sea of Azov, *Nauk. Dumka, Kiev*, 182 pp. (In Russian)
- Nik-Syrat, H., Hatf, A., Hajmoradlou, A., M., Ghorbani, R., and Nikoo, M., 2006.** Infection of *Gasterosteus aculeatus* to *Corynosoma strumosum* in the Goumishan Lagoon. *Journal of Iranian Fisheries Sciences*, 15(2), 155-159. (In Persian)
- Ondrackova, M., Davidova, M., Pecinkova, M., Blazek, R., Gelnar, M., Valova, Z., Cernv, J., and Jurajda, P., 2005.** Metazoan parasites of *Neogobius* fishes in the Slovak section of the River Danube *Journal of Applied Ichthyology*, 21 (4), 345-349
- Oumlzer, A., 2007.** Metazoan parasite fauna of the round goby *Neogobius melanostomus* Pallas, 1811 (Perciformes: Gobiidae) collected from the Black Sea coast at Sinop, Turkey, *Journal of Natural History*, 41(9-12), 483-492.
- Pazooki, J., and Aghlmandi, F., 1999.** Infections in two species of *Neogobius fluviatilis* and *Neogobius kessleri* to the *Dichelyne minutus Rudolphi*, 1819. *Journal of Iranian Fisheries Sciences*. (7) 2. (In Persian)
- Pazooki, J., and Masoumian, M., 2002.** Nematode Parasites from some freshwater fishes of Gyilan Mazandaran Provinces. *Journal of Pajouhesh and Sazandegi*, 51. (In Persian)
- Reimchen, T. E., and Nosil, P., 2001.** Ecological causes of sex-biased parasitism in three spine stickleback, *Biology Journal of Linnaeus Society*, 73,51-63.
- Rokicki, J., and Rolbiecki, L., 2002.** Colonization of the round goby, *Neogobius melanostomus* (Gobiidae) by parasites in the new environment of the Gulf of Gdańsk (Southern Baltic), *Wiadomosci Parazytologiczne*, 48 (2), 197-200.
- Rolbiecki, L., 2006.** Parasites of the round goby, *Neogobius melanostomus* (Pallas, 1811), an invasive species in the Polish fauna of the Vistula Lagoon ecosystem. *Journal of Oceanologia*, 48(4), 545-561.
- Sattari, M., Roustaii-Alimehr, M., and Shafii, Sh., 2002.** Survey on intensity to the Raphiascarid in some fishes of

Anzali Lagoon. *Journal of Pajouhesh and Sazandegi*,. 52. (In Persian)
Sattari, M., Shafii, Sh., Daghigh-Rouhi, J., Bourbiria, H., and Najest, N.,

2003. Survey on Eustrongilides larvae in some fishes of the Caspian Sea. *Journal of Veterinary Medicine Faculty*, (57)1. (In Persian)