

**The first report of *Amphipholis squamata* (Delle Chiaje, 1829)  
(Echinodermata: Ophiuroidea) from Chabahar  
Bay – northern Oman Sea**

**Attaran-Fariman G. \* ; Beygmoradi A.**

Received: December 2014

Accepted: April 2016

Chabahar Maritime University, Faculty of Marine Sciences, Department of Marine Biology,  
Daneshgah Avenue, 99717-56499, Chabahar, Iran.

\*Corresponding author's email: [gilan.attaran@gmail.com](mailto:gilan.attaran@gmail.com)

**Keywords:** Echinoderms, Amphiuroidae, Morphology, Taxonomy, Chabahar Bay;  
Oman Sea

### **Introduction**

*Amphipholis squamata* is an important Ophiuroid species belonging to the family Amphiuroidae which is widely used in biotechnological and molecular studies. It is a cosmopolitan species and capable to inhabit a wide variety of habitats except the polar regions, from subtidal zone to the depth of 2000 meters (Hendler, 1995). According to Fell (1962) its widespread distribution all over the world is the result of its coastal migration. *A. squamata* is characterized by its small body size, hermaphroditic reproduction, lack of larval phase, (Nisolle, 1990) and brood protection properties. It is an omnivorous species filtering sediment food particles and planktons (Emson and Whitfield, 1989). Variety in coloration patterns are reported among

the cognates of this species (Deheyn and Jangoux, 1999). Also, this species is one of the most important echinoderms in terms of bioluminescence (Deheyn *et al.*, 1997). Bioluminescence echinoderms were identified about two centuries ago (Viviani, 1805), consisting 4 out of 5 class of Echinodermata (Herring, 1987). The only class without bioluminescence ability is Echinoidea (Herring, 1987).

In the present study, *Amphipholis squamata* was reported for the first time from the subtidal zone of Chabahar Bay in northern part of the Oman Sea. This paper also provides information about the taxonomy and morphology of this group of aquatic taxa dwelling in the marine waters of the Iranian coasts of Chabahar Bay and the Oman Sea.

## Materials and methods

This study was carried out in March 2014. Sediment samples were collected by grab from the subtidal zone at Shahid Beheshti Port in Chabahar Bay (Fig. 1). Brittle stars were sorted in situ and put in a jar including sea water. Specimens were transferred to Laboratory of Chabahar Maritime University and living specimens were maintained in the well aerated aquariums. Dimensions, radial shields, form and color of disc and arms' length were measured in all of the specimens. Photographs were taken by stereo microscope (model C-DS ) equipped with the camera (model T4AL250 V (21 V 150 W)). Identification keys used in this study were from Clark and Rowe (1971); Price (1983) and Pomory (2007). Specimens (n=3) are currently fixed and preserved in 4% formaline and maintained in the same laboratory and are ready for scientific demands.

## Results and discussion

*A. squamata* was recorded for the first time during this study.

### Systematic

Order OPHIURIDA Müller and Troschel, 1840

Suborder OPHIURINA Müller and Troschel, 1840

Family AMPHIURIDAE Ljungman, 1867

Genus *Amphipholis* Ljungman, 1866

*Amphipholis squamata* (DelleChiaje, 1829)

### Redescription

Generally trunk brownish gray carrying small circular disk with black speckles (ca. 5 mm) and arms (6 cm). Dorsally, disk covered by small and irregular scales overlapping partially. Radial shields with elongated, thin, connective, rounded outer margins and straight inner margin (Fig. 2A). Coloration and coverage in ventral part similar to dorsal side. Bursa slits elongated. Oral shield pentagonal with an enlarged distal margin and convex, somewhat longer than wide. Dorsal shield large and united proximally. Two oral papilla present on each corner of jaw. infradental papillae present in pair (Fig. 2B). Dorsal arm plate oval with smooth distal margin (Fig. 2C). Ventral arm plate pentagonal (Fig. 2D). 3arm spines conic, transparent, erected, serrated at tip, 2 small tentacle scales, thin and elongated (Fig. 2E).

Ophiuroidea have a widespread distribution in the tropics. Totally 651 Echinodermata species have been reported from the Indian Ocean so far, 152 of which are belonging to Ophiuroidea. Chabahar Bay includes a variety of substrate such as, sandy, muddy and rocks. This geomorphological variation makes this region qualified to be habitable for various groups of animals (Sharifi, 2004).




Figure 1: Geographical location of the study area at Shahid Beheshti Port in Chabahar Bay, and its position at the Iranian coast of the Oman Sea.




Figure 2: Species of the family Amphipuridae. *Amphipholis squamata* A dorsal view, detail of the radial shields B ventral view C jaw D dorsal view of the arms E ventral view of the arms. Scale bar = A and B 2 mm; C, D and E 100  $\mu$ m.

However, the Ophiuroid species which were reported from this area so far are: *Macrophiothrix elongata* (H.L. Clark, 1938) and *Ophiiothrix savigny* (Müller and Troschel, 1842 ) belonging to family Ophiotrichidae (Ljungman, 1867) (Khaleghi and Owfi, 2011) and *Ophionereis dubia* (Müller and Troschel, 1842) belonging to the family Ophionereididae Ljungman, 1867 (Attaran-Fariman *et al.*, 2014). During the present study, *A. squamata* was reported for the first time. Specimens were collected from the depth of 10 m with a muddy substrate. *A. squamata* have a diversity of genetic, color and bioluminescence features through the world (Sponer *et al.*, 1999). For instance, 11 different color patterns of this species were reported till now (Mallefet and Jangoux, 2000). Therefore, variation in color and brightness could proof the presence of correlation between polychromatism and genetics of native populations. Because of its brightening feature, *A. squamata* had previously been nominated as *Asterias noctiluca* (Viviani, 1805). Presence of dark and light pigments on the body give it a noctilucous feature while bioluminescence is controlled by nervous system (Deheyn *et al.*, 1994). Bioluminescence mechanism in these taxa is utilized for avoiding predation and also, scaring the predators (Grober, 1990). Tortonesi (1932) showed the presence of various coloration patterns in this species. Orange, beige, dark brown, gray, black polka dots are the

colours that were reported between various populations of this species collected from inter tidal zone at 1971 by Binaux and Bocquet. Whereas, our species was brownish gray with polka dots. However, Clark (1987) proposed and fixed the name of this species as *Amphipholis squamata* instead of its previous name, *Ophiura elegans* (Leach, 1815). *A. squamata* is a very complicating species with very divergent mitochondrial lineages (Sponer, 2002). Unpublished data E. Boissinon mitochondrial DNA successive determination show that this species has a common lineage in equatorial regions (Stohr *et al.*, 2008). This species has a spherical disc with 1.08 – 2.47 mm diameter. The disc is covered with enlarged scales irregularly. In some cases the central plate is appeared. The length of shield is longer than its width, connective with rounded external edge and smooth internal edge. Spaces between radials are covered by small scales. Bursa slits elongated and vast. Oral shield fan-shaped with an enlarged and convex distal margin, its length is longer than width. Dorsal shield large, standing in equal distance from its proximal edge. Two oral papilla with a long distal part present on both apical sides of jaw. Pair of infradental papilla is present as well. Ventral arm plate pentagonal, elongated with 3 conical straight arm spine and denticulate apex. There are 2 small, thin and elongated tentacle scale (Gondim *et al.*, 2013). Our specimen is small with spherical disc 5mm in diameter. The

lengths of arms are as long as 6 times of the disc diameter bearing some short spines. The disc is covered by small scales while radial shields are visible. There are a colored area with different colors (such as, gray, white and Bluish gray) in different individuals. Present type has Gray polka dots. This species has recently been reported by Pomory (2007) with 5 mm disc diameter with arms as long as 3-4 times of disc diameter. However, we measured this rate 6 times as long as disc's diameter. Dorsal plate of arm is lunate with smooth distal edge. Radial shield is long, thin and is half time of radius of disc (Pomory, 2007). Although in some references the diameter of disc was reported longer than 5 mm its average length is 0.7 – 3 mm (Mortensen, 1927). Radial shields are small and connected to each other. Big species has 2 tentacle scales on proximal extremity of arm while small species normally has only 1 (Bartsch, 2008). Finding of this species from Chabahar Bay for the first time is not surprising because of its wide distribution through the world. According to the present literatures related to morphology and identification of *A. squamata* from different area our species demonstrates very close features with them. Therefore, due to the obtained results from this study and previously published data it would be concluded that this species has a invariable and monotonic morphological features all ove the world (Sponer *et al.*, 1999).

*A. squamata* dwells in various substrates such as, under stones, btween algae and bryozoas, rock pools, sandy bottom (Rodrigues *et al.*, 2011) and also sea grasses, mangrove, estuaries and water with high salinity (Gondium *et al.*, 2013). It is mostly in relation with Corallina alge and bryozoas (Fish and Fish, 2011). This cosmopolitan species could be found from intertidal zone to 1300 m depth of sea (Rodrigues *et al.*, 2011).

*A. squamata* is distributed all over the world except to arctic and antarctic regions. This species has already been reported from west Alantic, United States of America, Mexico, Netherlands Antilles, Belize, Costa Rica, Panama, Colombia, Brazil, Uruguay, and Argentina's Santa Cruz Province (Benavides-Serrato *et al.*, 2011), South Carolina, Florida, Islands of South Florida, Texas, Antilles, Belize, Panama, Brazil (Alvarado *et al.*, 2008), Persian Gulf (Price, 1983), eastern Atlantic along the periphery of Europe, the Mediterranean, the west coast of Africa, the eastern coast of South Africa and Madagascar (Alva and Vadon, 1989). This is the first report of *A. squamata* species from Chabahar Bay located at the North part of the Oman Sea.

#### **Acknowledgements**

The present work is a part of the MSc work of A.Beygmoradi at Chabahar Maritime University (CMU) and working space and facilities granted by the university are highly acknowledged.

The authors are grateful to Professor Andrew Price from University of Warwick for his help with the identification of *Amphipholis squamata*.

## References

- Alva, V. and Vadon, C., 1989.** Ophiuroids from the western coast of Africa (Namibia and Guinea-Bissau). *Scientia Marina*, 53(4), 827–845.
- Alvarado, J.J., Solís-Marín, F.A. and Ahearn, Y.C., 2008.** Equinodermos (Echinodermata) del Caribe Centroamericano. *Revista Biología Tropical / Internacional Journal of Tropical Biology and Conservation*, 56 (Supl. 3), 37-55.
- Attaran-Fariman, G., Beygmoradi, A. and Boos, K., 2014.** First record of *Ophionereis dubia* (Echinodermata: Ophiuroidea) from Chabahar Bay (Oman Sea, Iran). *Marine Biodiversity Records*.7(2),1-5.
- Bartsch, I., 2008.** Notes on ophiuroids from the Great Meteor Seamount (Northeastern Atlantic). *Spixiana*, 31 (2), 233-239.
- Benavides-Serrato, M., Borrero-Perez, GH. and Dias Sanchez, CM., 2011.** Equinodermos del Caribe colombiano I: Crinoidea, Asteroidea y Ophiuroidea. Serie de Publicaciones Especiales de Invemar, 22. Santa Marta, 384P.
- Binaux, R. and Bocquet, C., 1971.** Sur le polychromatisme de l'ophiure *Amphipholis squamata* (Delle Chiaje). *Comptes Rendus Académie Des Sciences Paris*. 273, 1618-1619.
- Clark, A.M. and Rowe, F.W., 1971.** Monograph of Shallow-water Indo-west Pacific Echinoderms. Trustees of the British Museum (Natural History), London. 234P.
- Clark, A.M., 1987.** *Asterias squamata* Delle Chiaje, 1828 (currently *Amphipholis squamata*; Echinodermata, Ophiuroidea): proposed conservation of the specific name. *Bulletin of Zoological Nomenclature*, 52, 246–247.
- Deheyn, D., Jangoux, M., 1999.** Colour varieties as sibling species in the polychromatic ophiuroid *Amphipholis squamata* (Echinodermata): evidence from inheritance of body colour and luminescence characters. *Journal of Experimental Marine Biology and Ecology*, 234(2), 219-234.
- Deheyn, D., Mallefet, J. and Jangoux, M., 1997.** Intraspecific variations of bioluminescence in a polychromatic population of *Amphipholis squamata* (Echinodermata: Ophiuroidea). *Journal of the Marine Biological Association UK*, 77, 1213-1222.
- Deheyn, D., Mallefet, J. and Jangoux, M., 1994.** Variations of bioluminescence intensity in the ophiuroid *Amphipholis squamata* (Delle Chiaje, 1828). In *Echinodermata* (ed. B. David *et al.*), 409P.
- Emson, R.H. and Whitfield, P.J., 1989.** Aspects of the life history of a tide pool population of *Amphipholis*

- squamata* (Ophiuroidea) from south Devon. *Journal of the Marine Biological Association of the United Kingdom*, 69, 27-41.
- Fell, H.B., 1962.** Evidence for the validity of Matsumoto's classification of the Ophiuroidea. *Publications of the Seto Marine Biological Laboratory* 10(2), 145-152.
- Fish, J. D. and Fish, S., 2011.** A student's guide to the seashore. United States of America by Cambridge University Press, New York.
- Gondim, A.I., Alonso, C., Dias, T.L., Manso, C.L. and Christoffersen, M. L., 2013.** A taxonomic guide to the brittle-stars (Echinodermata, Ophiuroidea) from the State of Paraíba continental shelf, Northeastern Brazil. *ZooKeys*, 307, 45-96.
- Grober, M.S., 1990.** Aposematism and bioluminescence in coastal marine communities. In *Adaptative coloration in invertebrates* (ed. M. Wicksten), pp. 77-87.
- Hendler, G., 1995.** Echinoderms of Florida and the Caribbean, Sea stars, sea urchins and allies Class Ophiuroidea: 89-195, Smithsonian Institution Press, Washington.
- Herring, P.J., 1987.** Systematic distribution of bioluminescence in living organisms. *Journal of Bioluminescence and Chemiluminescence*, 1, 147-163.
- Khaleghi, M. and Owfi, F., 2011.** Identification of Echinoidea species in the intertidal zones of Chabahar Bay. *Journal of Animal Environment*, 4, 31-36.
- Mortensen, T., 1927.** Handbook of the Echinoderms of the British Isles. London. 471P.
- Nisolle, V., 1990.** Modalités de l'incubation chez l'ophiure *Amphipholis squamata* (Delle Chiaje, 1829) (Echinodermata). Mémoire de licence. Université de Mons- Hainaut.
- Rodrigues, C.F., Paterson, G.L.J., Cabrinovic, A. and Cunha M.R., 2011b.** Deep-sea ophiuroids (Echinodermata) from mud volcanoes in the Gulf of Cadiz (NE Atlantic). *Zootaxa*, 2754,1-26.
- Pomory, M.C., 2007.** Key to the common shallow-water brittle stars (Echinodermata:Ophiuroidea) of the Gulf of Mexico and Caribbean Sea. *Caribbean Journal of Science* 1-42.
- Price, A.R.G., 1983.** Echinoderms of Saudi Arabia. Echinoderms of the Arabian Gulf coast of Saudi Arabia. *Fauna of Saudi Arabia*, 5, 28-108.
- Sponer, R., 2002.** Phylogeography and evolutionary history of the cosmopolitan, brooding brittle star *Amphipholis squamata* (Delle Chiaje, 1828; Echinodermata: Ophiuroidea). PhD. thesis, University of Otago.
- Sponer, R., Roy, M.S. and Mladenov, P.V., 1999.** Molecular evolution of *Amphipholis squamata* (Echinodermata: Ophiuroidea): High genetic differences among populations. In: M.D. Candia Carnevali and F. Bonasoro (Eds):

Echinoderm Research 1998: 405-408.

**Stohr, S., Conad, C. and Boissin, E., 2008.** Brittle stars (Echinodermata: Ophiuroidea) from La Réunion and the systematic position of *Ophiocanops* Koehler, 1922. *Zoological Journal of the Linnean Society*, 153, 545–560.

**Tortonese, E., 1932.** Osservazioni sul colore di alcuni Echinodermi. *Natura (Milano)*, 23, 160-164.

**Viviani, D., 1805.** Phosphorescencia maris quattuordecim lucescentium animalculorum novis speciebus illustrata. Genova.