

The effects of different levels of *Aloe vera* extract on some of the hematological and non-specific immune parameters in Siberian sturgeon (*Acipenser baerii*)

Bazari Moghaddam S.¹; Haghghi M.^{2*}; Sharif Rohani M.³;
Hamidi M.⁴; Ghasemi M.¹

Received: February 2016

Accepted: April 2016

Abstract

In this study, *Aloe vera* extract was used to evaluate its effects on hematological and immune parameters of Siberian sturgeon (*Acipenser baerii*). A total of 360 Siberian sturgeon weighing on average 10.95 ± 0.04 g were randomly distributed in four treatments including a control group and three experimental groups (each with three replications). *A. vera* extract powder at the rate of 0.5%, 1% and 1.5% were added to the food. The fish were fed for 60 days in fiberglass tanks. At the end of the first and second months of breeding, the necessary samplings for hematological studies were done. The physicochemical parameters of water were recorded every day. The results showed significant differences ($p < 0.05$) in the amount of RBC, hemoglobin, MCV, WBC, lysozyme and ACH50 in the treatments compared to the control group at the end of the breeding period. The results showed that *A. vera* extract has the effect of a non-specific immune system booster and using the extract in particular at the level of 1.5% as an immune stimulant in the diet of Siberian sturgeon can be effective in preventing disease in this species.

Keywords: *Aloe vera* extract, Hematological, Non-specific immune, Siberian sturgeon

1- Inland Waters Aquaculture Research Center, Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research Education and Extension Organization (AREEO), Bandar Anzali, Iran

2- Cold-water Fishes Research Center, Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research Education and Extension Organization (AREEO), Tonekabon, Iran

3-Iranian Fisheries Science Research Institute, Agricultural Research Education and Extension Organization (AREEO), Tehran, Iran

4-School of Pharmacy, Zanzan University of Medical Sciences, Zanzan, Iran

*Corresponding author's Email: masoud126@yahoo.com

Introduction

Along with the increasing development of the aquaculture industry in the world, emphasizing the principles of operation according to the economic and health aspects of aquaculture is of particular importance. In this regard, ensuring proper growth and preventing disease in aquatic animals are among the important issues that have attracted the attention of researchers in this industry more and more. Siberian sturgeon (*Acipenser baerii*) is a species of commercial value that easily adapts to the conditions of breeding and is resistant to changes in environmental conditions (Pyka and Kolman, 2003). Because of its fast growth, early sexual maturity and caviar production in a short time, as well as its diverse diet, this species was introduced as a candidate for freshwater culture for the production of sturgeon meat (Adamek *et al.*, 2007). In recent years, in order to treat and the fight against bacterial infections in fish, chemical drugs such as antibiotics, have been used. However, in addition to the high costs involved, the administration of antibiotics has resulted in the development of drug resistance, destruction of natural bacterial flora, emergence and spread of bacterial resistance towards antimicrobial agents, and the transfer of antibiotic residues from aquatic animals to humans. Antibiotics also weaken the immune system against viral diseases and parasites.

The use of herbal immunostimulants in the aquaculture industry, in particular to strengthen the immune system and increase fish resistance against a variety of infectious diseases has spread worldwide. Many studies have shown that immunostimulants of plants are capable of enhancing specific and non-specific immune defense mechanisms, as well as increase or reduce fish losses against a variety of viral and bacterial or parasitic infections (Kim *et al.*, 1999; Dügenci *et al.*, 2003; Yin *et al.*, 2006; Nya and Austin, 2009; Yin *et al.*, 2009; Bilen and Bulut, 2010; Harikrishnan *et al.*, 2010; Bilen *et al.*, 2011; Begum and Navaraj, 2012; Ardo, 2013; Govind, 2013; Haghghi and Sharif Rohani, 2013; Sharif Rohani *et al.*, 2013).

Aloe vera is one of the most valuable plants in the Liliaceae family, which is native to tropical regions. *A. vera* contains more than 75 nutrients, 200 active compounds, 20 minerals, 18 amino acids and 12 vitamins and compounds such as Aloin, famodin, Antrokinon and barbaloin (Shelton, 1991; Atherton, 1998; Mandrioli *et al.*, 2011). This plant has a wide variety of pharmacological effects such as skin lesions and wound healing effects, anti-viral, antibacterial and other effects have been attributed to this plant. However, the effects of immune stimulation and growth of warm-blooded animals has not been proven. In addition, little information about the effects of immunogenicity, anti-toxicity and the effects on growth performance by *A. vera* in different fish species have

been found (Alishahi *et al.*, 2010; Zodape, 2010; Wang *et al.*, 2011; Alishahi and Abdi, 2013; Haghighi *et al.*, 2014).

According to the characteristics and properties of this medicinal plant, this study was performed to evaluate the effects of three doses of *A. vera* on hematological and immune parameters of this valuable species (Siberian sturgeon).

Materials and methods

Preparation of A. vera extract

The plant of *A. vera* was procured from Medicinal Plants Production Cooperation of Havin and the plant species was identified and confirmed by a botanist. The leaves were collected and washed in sterile distilled water and evacuated from gel. The leaves were separately shade-dried for 10 day till

weight constancy was achieved. The sample was powdered in an electric blender. The extract was prepared with the standard method of percolation. To do this, chopped dried plant leaves in 80% ethanol were percolated for 72 hours. Then, the slurry was filtered with Whatman No. 1 filter paper and centrifuged for 5 min at 5000 rpm. The filtrate obtained from ethanol using a rotary device, the excess solvent was separated from the extract. These crude extract was stored at -18°C until use (Ozakan *et al.*, 2007; Haghighi *et al.*, 2014). At the end of the extraction process, to analyze compounds in *A. vera* extract gas chromatography (GC-MS) and liquid chromatography (HPLC) (Rouessac and Rouessac, 2007; Lakshmi and Rajalakshmi, 2011) was used (Table 1).

Table 1: Types and amounts of compounds in *Aloe vera* extract.

Compounds	Amount (%)	Compounds	Amount (%)
Aloin	28.81	Comaric acid	7.62
Oleic acid	6.23	Squalene	13.8
B-Sitostrol	1.41	Limoene	10.26
Lupeol	4.7	n- Hexadecanoic acid	10.24
Campestrol	2.18	other components	6.3
Carvone	8.43		

Supplementation of the normal diet with dried A. vera extract

In this study, in order to produce food containing different percentages of *A. Vera* extract powder (0.5%, 1% and 1.5%), Biomar commercial feed containing 49% crude protein, 15% crude lipid, 2.5% cellulose, 1.24% phosphorus, 8.7% ash, 1.85% calcium,

0.55% sodium, Cu 1.5 mg kg⁻¹, Mn 12 mg kg⁻¹, Zn 75 mg kg⁻¹ and I1.8 mg kg⁻¹ was used. In order to add extract powder and prepare food diets containing different amounts of *A. vera* (0.5%, 1% and 1.5%), the the the required amounts of the extract powder was dissolved in water and sprayed on the food and then stirred. Feeds were

spread on separate trays (for 24 hours) and dried at room temperature (Noga, 2000). Feeds were stored in special containers with lids at a temperature of 4-6°C.

Fish and experimental conditions

A total number of 360 Siberian sturgeon (with average weight of 10.95 ± 0.04 g), were randomly divided into 12 fiberglass tanks of 500 liter (dewatering volume 350 liters, at 3 liters per minute water flow and permanent aeration) placed at the International Sturgeon Research Institute (Rasht, Iran) and cultured for two months.

Physico-chemical parameters of water including dissolved oxygen (WTW Oxi 330i digital devices made in Germany), pH and water temperature (WTW pH 330i digital devices made in Germany) were measured on a daily basis. Average water temperature (22.8 ± 0.88 °C), dissolved oxygen (6.74 ± 0.42 mg/L⁻¹) and 6.8 ± 0.19 of pH were determined. Fish adaptation for 14 days was carried out in the fiberglass tanks. During the two months breeding, fish were fed the prepared feeds containing *A. vera* extract (0.5%, 1% and 1.5%) and Biomar basic diet (for control group) at a rate of 3% of body weight at 8:00, 15:00 and 22:00 hours.

Blood sampling

At the end of each month, to evaluate the effects of *A. vera* extract on hematological and immune parameters of Siberian sturgeon and to compare

between different treatments, blood samples from the caudal vein of 6 fish from each group (18 samples in total) were taken. 0.5 cc of blood with anticoagulant heparin and 1.5 cc of blood without anticoagulant heparin were taken to measure immune parameters.

It should be noted that anesthetics were not used in the samplings because they probably have an impact on the blood indices (Torrecillas *et al.*, 2007 and 2011). The samples were centrifuged using centrifugation at 3000 rpm for 10 minutes. They were then transferred using a sampler to a new vial, and stored at -80°C. Preparation and measurement of all hematological and serological parameters were carried out at the International sturgeon research Institute and Viomed laboratory.

Haematological assay

Blood samples were analyzed following routine methods adopted in fish hematology (Blaxhall and Daisley, 1973; Klontz, 1994). The total red blood cell counts (RBC $\times 10^6$ /mL) were determined in a 1:200 dilution of the blood sample in Hayem's solution and total white blood cell counts (WBC $\times 10^3$ /mL) in a 1:20 dilution of the blood sample with a Neubauer hemocytometer. The hematocrit (Hct) and leucocrit percentages were determined in duplicate using micro hematocrit-heparinized capillary tubes of 75 μ L volume and a micro hematocrit centrifuge at 15000 rpm for 5 min

(Houston, 1990; Klontz, 1994). The percentages of erythrocyte (hematocrit) and leucocyte (leucocrit) volumes were calculated by overlaying the tubes on a sliding scale hematocrit reader. The hemoglobin (Hb, g dL⁻¹) concentrations were determined by the cyanomethaemoglobin method (Valery *et al.*, 1991; Klontz, 1994) using a haemoglobin reagent set (Pars Azmun Diagnostics). All the values of red blood cell indices, the mean values of cell haemoglobin (MCH pg), cell haemoglobin concentration (MCHC %), and cell haemoglobin volume (MCV fl) were calculated according to Wintrobe formulae (Anderson and Klontz, 1965).

The differential leukocytes count was carried out using blood smears stained with Wright-Giemsa. The percentage composition of leukocytes was determined based on their identification characters listed (Ivanova, 1983).

Immunological assay

Alternative complement activity

Alternative complement activity (ACH50) was evaluated following the procedure of Yano (1992) by using rabbit red blood cells (RaRBC). Briefly, RaRBC were washed and adjusted to 2×10^8 cell/mL in EDTA-magnesium-gelatin veronal buffer (0.01 M). Precisely 100 μ L of the RaRBC suspension was lysed with 3.4 mL of distilled water and the absorbance of the haemolysate was measured at 414 nm against distilled water to acquire the 100% lysis value.

The test plasma was appropriately diluted, and different volumes ranging from 0.1 to 0.25 mL were made up to 0.25 mL total volume before being allowed to react with 0.1 mL of RaRBC in test tubes. After incubation at 20°C for 90 min with occasional shaking, 3.15 mL of a 0.9% (w/v) saline solution was added to each tube with centrifugation at 1600 rpm for 10 min at 4°C. The absorbance (A) of the supernatant was measured using a spectrophotometer at 414 nm. A lysis curve was obtained by plotting the percentage of haemolysis against the volume of plasma added. The volume of plasma producing 50% haemolysis (ACH50) was determined and the number of ACH50 units/mL was obtained for each fish.

Lysozyme activity

The turbidimetric assay for lysozyme was carried out according to Sahoo *et al.* (2008) with minor modifications. Thus, plasma (50 μ L) was added to 2 mL of a suspension of *Micrococcus lysodeikticus* (0.2 mg/mL⁻¹) in a 0.05 M sodium phosphate buffer (pH 6.2). The reaction was carried out at 25°C and absorbance was measured spectrophotometrically at 570 nm after 0.5 min and 4.5 min. PBS was used as a blank. A unit of lysozyme activity was defined as the sample amount causing a decrease in absorbance of 0.001 min⁻¹. Lysozyme of sample calibrated using a standard curve determined with hen's egg white lysozyme (Sigma) in PBS.

Statistical analysis

Statistical analyses were performed using SPSS (version 20) software. Data are presented as mean±SD. All the data were tested for normality (Kolmogorov-Smirnov test). Data were analyzed by one-way of variance analysis (ANOVA). The significant means were compared by Duncan test and a $p<0.05$ was considered statistically significant. Also using independent samples T-Test compares the average blood sampling between the two stages.

Results

The results showed significant differences ($p<0.05$) in the hematological parameters including Hct, Hb, MCH, MCV and WBC in fish fed *A. vera* compared with the control group (Table 2) in both samples (the first and second months of breeding). However no significant differences ($p>0.05$) were observed in the results of the first and second months of training in any of the hematological parameters.

Table 2: The haematological parameters of Siberian sturgeon fed with different levels of *Aloe vera* extract for two months (mean ±SD).

Parameters	<i>Aloe vera</i> extract levels (%)							
	First month				Second month			
	Control (0)	0.5	1	1.5	Control (0)	0.5	1	1.5
RBC (10^6 mL^{-1})	0.77±0.06 ^a	0.78±0.05 ^a	0.8±0.06 ^a	0.78±0.06 ^a	0.85±0.07 ^a	0.86±0.04 ^a	0.88±0.05 ^a	0.92±0.07 ^b
Hct (%)	26±0.26 ^b	25±0.52 ^a	25.17±0.4 ^a	24.83±0.17 ^a	30.67±0.2 ^{ab}	29.17±0.6 ^a	29.83±0.48 ^a	31.5±0.22 ^b
Hb (g dL ⁻¹)	5.1±0.04 ^a	7.6±0.07 ^b	5.98±0.06 ^b	5.8±0.07 ^b	6.68±0.03 ^a	5.87±0.07 ^b	7.75±0.04 ^b	8.1±0.07 ^c
MCH (pg)	65.33±0.56 ^a	75.17±0.54 ^b	75.17±0.6 ^b	74.67±0.33 ^b	78.17±0.31 ^a	88.18±0.65 ^b	88.17±0.4 ^b	87.67±0.22 ^b
MCV (fl)	338.7±2.56 ^b	321±5.53 ^{ab}	316.8±3.18 ^a	320.5±0.99 ^{ab}	360±2.67 ^b	338.5±5.49 ^a	339.7±3.22 ^a	342.2±1.11 ^{ab}
MCHC (%)	21.5±0.25	23.5±0.5	23.67±0.49	23.33±0.21	21.8±0.14	26.05±0.32	25.97±0.25	25.7±0.11
WBC (10^3 mL^{-1})	9.2±0.8 ^a	11.6±0.94 ^b	13.23±1.12 ^c	13.51±1.1 ^c	12.1±0.93 ^a	15.78±1.32 ^b	16.6±1.41 ^c	16.68±1.32 ^c
Neut (%)	9.83±0.98	10.5±1.05	14.33±0.82	15±1.89	10.17±1.12	10.83±0.98	14.33±1.21	14.33±1.21
Mon (%)	1.83±0.41	1.5±0.55	1.33±0.52	1.17±0.75	2.33±0.52	2±0.63	1.83±0.41	1.33±0.52
Lymp (%)	85.83±1.17	85.17±0.75	82.1±1.21	82±1.26	86.83±1.17	86±0.89	82.5±1.05	82.83±1.62
Eos (%)	1.5±0.55	2±0.69	1.83±0.41	1.67±0.52	1.67±0.52	2±0.69	1.83±0.41	1.67±0.52

Data are expressed as mean±SD (n=72). Neut: neutrophil; Mon: Monocyte; Lymp: Lymphocyte; Eos: Eosinophil.

Differences in Latin letters in each row shows a significant difference ($p<0.05$).

In this study, the results of the immune parameters (Table 3) showed that serum lysozyme levels and ACH50 in different *A. vera* treatments compared to the control in the tanks during the

two months were significantly different ($p<0.05$). At the end of the breeding period, the highest values in the immune parameters were recorded in the 1.5% *A. vera* extract group.

Table 3: The immunological parameters of Siberian sturgeon fed with different levels of *Aloe vera* extract for two months (mean \pm SD).

Parameters	<i>Aloe vera</i> extract levels (%)							
	First month				Second month			
	Control	0.5	1	1.5	Control	0.5	1	1.5
Lysozyme Activity (u mL ⁻¹)	8.67 \pm 0.33 ^a	17.67 \pm 0.33 ^b	19.33 \pm 0.33 ^b	19 \pm 0.58 ^b	11 \pm 0.58 ^a	21.33 \pm 0.67 ^b	25.33 \pm 0.88 ^c	28 \pm 0.58 ^d
ACH50 (u mL ⁻¹)	171 \pm 1 ^a	175 \pm 0.58 ^a	177.67 \pm 0.88 ^b	180.67 \pm 0.88 ^b	191.67 \pm 1.45 ^a	217.67 \pm 1.2 ^b	220 \pm 1 ^b	224 \pm 2.3 ^c

Differences in Latin letters in each row shows a significant difference ($p < 0.05$; $n = 72$).

Discussion

Attention to the prevention of disease and increase in survival rate can affect the security of investments and the development of sturgeon breeding. In this regard, medicinal plants have attracted much attention due to their therapeutic effects. Moreover they are also known to cause no or less biological and environmental risks and cost less compared to other antibiotics (Christyapita *et al.*, 2007; Alishahi *et al.*, 2010; Watanuki *et al.*, 2010). Important indicators on health status of fish in intensive culture conditions can be non-specific resistance of different fish species, assessment of nutritional status to evaluate the impact of fish food additives, and the measurement hematologic parameters noted (Cataldi *et al.*, 1998).

A number of studies have shown that herbal immune stimulants did not have a significant effect on red blood cell counts (Shalaby *et al.*, 2006; Kumar Jha *et al.*, 2007; Sahu *et al.*, 2007; Harikrishnan *et al.*, 2010). The study showed that the amount of hemoglobin in sampling stages (the first and second months of breeding) in the control group was significantly lower than that in all treatments. The amounts of hemoglobin in the second stage of

sampling, in treatments and control, increased. Studies have shown that levels of hemoglobin, hematocrit and red blood cells are under the influence of different factors such as age, sex, race and environment (Ameri mahabadi, 1999).

Hematocrit values change in the presence of pathogens or imbalance in mineral density in blood. In addition, the very high volume of hematocrit may indicate the possibility of unfavorable environmental conditions associated with stress (Parker, 1974). The results of this study are consistent with that of some other studies regarding changes in the levels of red blood cells, hemoglobin, hematocrit and blood indices (Iwama and Nakanishi, 1996; Shalaby *et al.*, 2006; Nya and Austin, 2009). Haghghi *et al.* (2014) observed significant changes in blood in levels of red blood cells and other indices at the end of the two months period in rainbow trout fed a diet containing *A. vera* extract. However, Alishahi *et al.* (2010) did not see any differences in the amount of red blood cells and related indices. Generally, an increase in white blood cells in the second month of breeding was evident. Increased levels of white blood cells in the study conducted by Pourgholam *et al.* (2013)

using *Echinacea purpurea* extract on immune indices and survival of rainbow trout corresponded with the results of this study.

Hajibeglou and Sudagar (2010) reported an increase in the number of white blood cells in common carp fish with the use of dietary supplements containing herbal immune stimulants. Also, the results of Sahu *et al.* (2007) indicated an increase in the rate of WBC. Gopalakannan and Arul (2006) reported that WBC significantly increased after feeding carp with herbal stimulants such as chitin. Oskoi *et al.* (2012) also found that the use of herbal stimulants on blood indices, including white blood cells in rainbow trout was effective. The WBC increased in common carp blood by adding *Stachys lavandulifolia* (2% and 8%) to the diet of the fish (Bahrami Babahydari *et al.*, 2015). In research conducted by Alishahi and Abdi (2013) amounts of WBC in the different treatments of *A. vera* rainbow trout showed statistically significant differences ($p < 0.05$). However no significant differences ($p > 0.05$) were observed in the lymphocytes, monocytes, neutrophils and eosinophils, which confirms the results of the present study.

Increase in the number of neutrophils after taking herbal stimulants, can be related to beta-glucan that are able to recognize specific receptors in their white blood cells. (Andrews *et al.*, 2009). When these receptors are occupied by glucan, active white blood cells surround, kill and

digest pathogenic bacteria more so that all these factors improve the host immune system (Andrews *et al.*, 2009).

Lysozyme is an important component of non-specific immunity in fish that causes destruction of bacterial wall, enhances phagocytic activity and activates both of the complement pathways involved in stimulating the immune response of the fish. The level of serum lysozyme shows improvement in the immune status of the fish and an increase in the immune system of the fish to better fight against infectious agents and stress reduction. Lysozyme activity increases immune stimulants following administration of some vaccines and some prebiotics in fish (Alishahi *et al.*, 2010). Results of this study showed that adding different levels of *A. vera* extract in lysozyme activity had significant effects ($p < 0.05$) and the highest lysozyme activity was observed with 1.5% *A. vera*.

In this study, an increasing trend in lysozyme activity has been shown which is in agreement with several reports indicating the role of herbal immunostimulants in enhancing lysozyme activity (Rao *et al.*, 2006; Choi *et al.*, 2008; Haghighi *et al.*, 2014). Lysozyme is a humoral component of the non-specific defense mechanism which has the ability to prevent the growth of bacteria by splitting β -1, 4 glycosidic bonds in the peptidoglycan of bacterial cell walls (Secombes, 1996; Choi *et al.*, 2008).

In both sampling stages, ACH50 amounts between experimental and

control groups were statistically significant ($p < 0.05$). Minimum and maximum ACH50 were observed in the control and 1.5% *A. vera* extract group, respectively. Awad *et al.* (2010) reported that the use of *Lupinus perennis*, *Urtica dioica* and *Managifera indica*, especially at a concentration of 1% and 2% in the diet of rainbow trout caused a significant increase in the complement activity. Feeding tilapia fish within two weeks using *Eclipta alba* extract could create a significant increase in complement activity (Christybapita *et al.*, 2007). Alishahi and Abdi (2013) reported that the use of *A. Vera* extract (1%) in common carp diet is able to significantly increase ACH50 amounts in comparison with the control group and 0.5% *A. vera* extract group. Alternative complement pathway activity is a strong non-specific defense mechanism to protect fish against a wide range of pathogenic organisms such as bacteria, fungi, viruses and parasites (Chiu *et al.*, 2010). In general, the results of this study indicate that the use of *A. Vera* extract (especially 1.5%) was more effective on some hematological and immune parameters in Siberian sturgeon.

Acknowledgements

We are grateful to Dr. Mohammad Ali Yazdani, Director of the International Sturgeon Research Institute and Dr. Shahram Abdolmaleki, Research Deputy of this institute for supporting us during this study. Also, sincere

thanks to colleagues in Dept. of Fish Diseases of this institute. This article is part of the project No. 92026531 by the Iranian National Science Foundation (INSF) that was implemented with financial support from INSF.

References

- Adamek, Z., Prokes, M.M., Barus, V. and Sukop, I., 2007.** Diet and growth of seberian, *Acipenser baerii* on alternative pond culture. *Turkish Journal of Fisheries and Aquatic Sciences*, 7, 153-160.
- Alishahi, M., Ranjbar, M., Ghorbanpour, M., Peyghan, R., Mesbah, M. and Razi jalali, M., 2010.** Effects of dietary *Aloe vera* on specific and nonspecific immunity of common carp (*Cyprinus carpio*). *Journal of Veterinary Research*, 4(3), 85-91.
- Alishahi, M. and Abdi, E., 2013.** Effects of different levels of *Aloe vera* L. Extract on growth performance, hemato-immunological indices of *Cyprinus carpio* L. *Iranian Journal of Veterinary Science and Technology*, 5(2), 33-44.
- Ameri Mahabadi, M., 1999.** Laboratory methods of veterinary hematology. Tehran University Press. 126P. (In Persian).
- Anderson, D. and Klontz, G.W., 1965.** Basic haematology for the fish culturist. Northwest Fish Culture Conference, 16, 38-41.
- Andrews, S.R., Sahu, N.P., Pal, A.K. and Kumar, S., 2009.**

- Hematological modulation and growth of *Labeo rohita* fingerlings: effect of dietary mannan oligosaccharide, yeast extract, protein hydrolysate and chlorella. *Aquaculture Research*, 41, 61-69.
- Ardó, L., 2013.** Application of herbal extracts to improve health status and innate immune response of fish. Doctoral PhD Thesis. University of Debrecen. Doctoral School of Animal Husbandry.
- Atherton, P., 1998.** *Aloe vera* revised. *The British journal of phytotherapy*, 4, 176-183.
- Awad, E.S., 2010.** Studies on plant based dietary supplements for control of *Aeromonas hydrophila* infections in rainbow trout (*Oncorhynchus mykiss* Walbaum). PhD Thesis, University of Heriot Watt.
- Bahrami Babaheydari, S., Paykan Heyrati, F., Dorafshan, S., Mahboobi Soofiani, N. and Vahabi, M.R., 2015.** Effect of dietary wood betony, *Stachys lavandulifolia* extract on growth performance, haematological and biochemical parameters of common carp, *Cyprinus carpio*. *Iranian Journal of Fisheries Sciences*, 14(4), 805-817.
- Begum, S.S. and Navaraj, P.S., 2012.** Synergistic effect of plant extracts supplemented diets on immunity and resistance to *Aeromonas hydrophila* in *Mystus keletius*. *IOSR Journal of Pharmacy and Biological Sciences*, 2(4), 30-36.
- Bilen, S. and Bulut, M., 2010.** Effects of Laurel (*Laurus nobilis*) on the non-specific immune responses of rainbow trout (*Oncorhynchus mykiss*, Walbaum). *Journal of Animal and Veterinary Advances*, 9(8), 1275-1279.
- Bilen, S., Bulut, M. and Bilen, A.M., 2011.** Immunostimulant effects of *Coggyria coggyria* on rainbow trout (*Oncorhynchus mykiss*). *Fish and Shellfish Immunology*, 30, 451-455.
- Blaxhall, P.C. and Daisley, K.W., 1973.** Routine haematological methods for use with fish blood. *Journal of Fish Biology*, 5, 771-781.
- Cataldi, E., Di Marco, P., Mandich, A. and Cataudella, S., 1998.** Serum parameters of Adriatic sturgeon *Acipenser naccarii* (Pisces: Acipenseriformes); effects of temperature and stress. *Comparative Biochemistry and Physiology*, 121A, 351-354.
- Chiu, C.H., Cheng, C.H., Gua, W.R., Guu, Y.K. and Cheng, W., 2010.** Dietary administration of the probiotic, *Saccharomyces cerevisiae* P13, enhanced the growth, innate immune responses, and disease resistance of the grouper, *Epinephelus coioides*. *Fish and Shellfish Immunology*, 29, 1053-1059.
- Choi, S.H., Park, K.H., Yoon, T.J., Kim, J.B., Jang, Y.S. and Choe, C.H., 2008.** Dietary Korean mistletoe enhances cellular non-specific immune responses and survival of Japanese eel (*Anguilla*

- japonica). *Fish and Shellfish Immunology*, 24, 67-73.
- Christyapita, D., Divyagnaneswari, M. and Dinakaran, M.R., 2007.** Oral administration of *Eclipta alba* leaf aqueous extract enhances the non-specific immune responses and disease resistance of *Oreochromis mossambicus*. *Fish and Shellfish Immunology*, 23, 840-852.
- Düğenci, S.K., Arda, N. and Candan, A., 2003.** Some medicinal plants as immunostimulant for fish. *Journal of Ethnopharmacology*, 88, 99-106.
- Gopalakannan, A. and Arul, V., 2006.** Immunomodulatory effects of dietary intake of chitin, chitosan and levamisole on the immune system of *Cyprinus carpio* and control of *Aeromonas hydrophila* infection in ponds. *Aquaculture*, 255, 179-187.
- Govind, P., 2013.** Some medicinal plants to treat fish ectoparasitic infections. *International Journal of Pharmaceutical Sciences and Research*, 2(2), 532-538.
- Haghighi, M. and Sharif Rohani, M., 2013.** The effects of powdered ginger (*Zingiber officinale*) on the haematological and immunological parameters of rainbow trout *Oncorhynchus mykiss*. *Journal of Medicinal Plants and Herbal Therapy Research*, 1, 8-12.
- Haghighi, M., Sharif Rohani, M., Samadi, M., Tavoli, M., Eslami, M. and Yusefi, R., 2014.** Study of effects *Aloe vera* extract supplemented feed on hematological and immunological indices of rainbow trout (*Oncorhynchus mykiss*). *International Journal of Advanced Biological and Biomedical Research*, 2(6), 2143-2154.
- Hajibeglu, A. and Sudagar, M., 2010.** Immune response of common carp (*Cyprinus carpio*) fed with herbal immunostimulants diets. *Journal of Animal and Veterinary Advances*, 9(13), 1839-1897.
- Harikrishnan, R., Balasundaramb, C. and Heo, M.S., 2010.** Herbal supplementation diets on hematology an innate immunity in goldfish against *Aeromonas hydrophila*. *Fish and Shellfish Immunology*, 28, 354-361.
- Houston, A. H., 1990.** Blood and circulation. In: Moyle (ed) Methods for fish biology. *American Fisheries Society*, 273-334.
- Ivanova, N.T., 1983.** Atlas of fish blood cells. *LPP Moscow, Russia* (In Russian).
- Iwama, G. and Nakanishi, T., 1996.** Innate immunity in fish. In: The fish immune system. Academic Press, London, UK. pp. 73-114.
- Kim, K.H., Hwang, Y.J. and Bai, S.C., 1999.** Resistance to *Vibrio alginolyticus* in juvenile rockfish (*Sebastes schlegel*) fed diets containing different doses of aloe. *Aquaculture*, 180, 13-21.
- Klontz, G.W., 1994.** Fish hematology. In: Techniques in fish immunology. Stolen, J.S., T.C. Flecher, A.F. Rowely, T.C. Zelikoff, S.L. Kaattari and S.A. Smith (Eds.). Vol. 2, SOS

- Publications, USA., ISBN: 0962550582, 121-132.
- Kumar Jha, A., Pal, A.K., Sahu, N.P., Kumar, S. and Mukherjee, S.C., 2007.** Haemato-immunological responses to dietary yeast RNA, ω -3 fatty acid and β -carotene in *Calta calta* juveniles. *Fish and Shellfish Immunology*, 23, 917-927.
- Lakshmi, P.T.V. and Rajalakshmi, P., 2011.** Identification of phyto-components and its biological activities of *Aloe vera* (L.) through Gas Chromatography-Mass Spectrometry. *International Research Journal of Pharmacy*, 2(5), 247-249.
- Mandrioli, R., Mercolini, L., Ferranti, A., Fanali, S. and Raggi, M.R., 2011.** Determination of aloe emodin in *Aloe vera* extracts and commercial formulations by HPLC with tandem UV absorption and fluorescence detection. *Food Chemistry*. 126, 387–393.
- Noga, E., 2000.** Fish Diseases: diagnosis and treatment. Wiley-Blackwell press. 366-367.
- Nya, E.J. and Austin, B., 2009.** Use of garlic, *Alium sativum*, to control *Aeromonas hydrophila* infection in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases*, 32, 963-970.
- Oskoi, S.B., Kohyani, A.T., Parseh, A., Salati, A.P. and Sadeghi, E., 2012.** Effects of dietary administration of *Echinacea purpurea* on growth indices and biochemical and hematological indices in rainbow trout (*Oncorhynchus mykiss*) fingerlings. *Fish Physiology and Biochemistry*, 38(4), 1029-1034.
- Ozakan, G., Simsek, B., and Kuleasan, H., 2007.** Antioxidant activities of *Satureja cilicica* essential oil in butter and in vitro. *Journal of Food Engineering*, 79, 1391-1396.
- Parker, R.B., 1974.** Probiotics. The other half of the antibiotics story. *Animal Nutrition Health*, 29, 4–8.
- Pourgholam, R., Sharif Rohani, M., Safari, R., Saeedi, A. A., Binaeei, M., Najafeyan, R., Bankehsaz, Z., Taghavi, M.J. and Sepahdari, A., 2013.** Effect of *Echinacea purpurea* extract on the immune system of rainbow trout (*Oncorhynchus mykiss*) and its resistance to Streptococcus. *Iranian Scientific Fisheries Journal*, 22(3), 1-12. (In Persian).
- Pyka, J. and Kolman, R., 2003.** Feeding intensity and growth of Siberian sturgeon (*Acipenser baerii* Brandt), under pond conditions. *Archives of Polish Fisheries Journal*, 5, 267-277.
- Rao, Y.V., Das, B.K., Pradhan, J. and Chakrabarti, R., 2006.** Effect of *Achyranthes aspera* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. *Fish and Shellfish Immunology*, 20, 263-273.
- Rouessac, F. and Rouessac, A., 2007.** Chemical analysis modern instrumentation methods and

- techniques. 2nd Edition, England, John Wiley and Sons Ltd. pp. 98-115.
- Sahoo, P.K., Mahapatra, K. D., Saha, J.N., Barat, A., Sahoo, M., Mohanty, B.R., Gjerde, B., Ødegard, J., Rye, M. and Salte, R., 2008.** Family association between immune parameters and resistance to *Aeromonas hydrophila* infection in the Indian major carp, *Labeo rohita*. *Fish and Shellfish Immunology*, 25, 163-169.
- Sahu, S., Das, B.K., Pradhan, J., Mohapatra, B.C., Mishra, B.K. and Sarangi, N., 2007.** Effect of *Magnifera indica* Kernel as a food additive on immunity and resistance to *Aeromonas hydrophila* in *Labeo rohita* fingerlings. *Fish and Shellfish Immunology*, 23, 109-118.
- Sharif Rohani, M., Masoumzadeh, M., Haghighi, M., Jalilpoor, J., Pourdehghani, M., Shenavar Masouleh, A., Alizadeh, M. and Bazari Moghaddam, S., 2013.** Effects of oral administration of *Zataria multiflora* essential oil on some blood and serum parameters in *Acipenser persicus*. *Iranian Journal of Fisheries Sciences*, 12(4), 908-915.
- Secombes, C. J., 1996.** The non-specific immune system: cellular defenses. In: Iwama G, Nakanishi T, editors. *The fish immune system: organism, pathogens and environment*. CA Academic Press San Diego. pp. 63-103.
- Shalaby, A.M., Khattab, Y.A. and Abdel Rahman, A. M., 2006.** Effects of garlic (*Allium sativum*) and chloramphenicol on growth performance, physiological parameters and survival of Nile tilapia (*Oreochromis niloticus*). *Journal of Venomous Animals and Toxins including Tropical Diseases*, 12, 172-201.
- Shelton, R.M., 1991.** *Aloe vera*: its chemical and therapeutic properties. *International Journal of Dermatology*, 30, 679-683.
- Torrecillas, S., Makol, A., Caballero, M.J., Montero, D., Robaina, L., Real, F., Sweetman, J., Tort, L. and Izquierdo, M.S., 2007.** Immune stimulation and improved infection resistance in European sea bass (*Dicentrarchus labrax*) fed mannan oligosaccharides (MOS). *Fish and Shellfish Immunology*, 23, 969-981.
- Torrecillas, S., Makol, A., Caballero, M.J., Montero, D., Gines, R., Sweetman, J. and Izquierdo, M.S., 2011.** Improved feed utilization, intestinal mucus production and immune parameters in sea bass (*Dicentrarchus labrax*) fed mannan oligosaccharides (MOS). *Aquaculture Nutrition*, 17(2), 223-233.
- Valery, H., Gowenblock, A.H. and Bell, M., 1991.** Practical clinical biochemistry. 5th ed. CBS Publication and distributors, Dehli, India: 479-480.
- Wang, C., Xu, Q.Y., Xu, H., Zhu, Q., Zheng, Q.S. and Sun, D.J., 2011.**

Effects of aloe powder on the growth performance and plasma indices of sturgeon (*Acipenser baerii* Brandt). *Journal of Shanghai Ocean University*, 4, 123-132.

Watanuki, H., Ota, K., Malina, A.C. and Tassakka, A.R., 2010.

Immunostimulant effects of dietary *Spirulina platensis* on carp, *Cyprinus carpio*. *Aquaculture*, 258, 157–163.

Yano, T., 1992. Assay of hemolytic complement activity. In: Techniques in fish immunology (ed. by J.S. Stolen, T.C. Fletcher, D.P. Anderson, S.C. Hattari and A.F. Rowley), pp. 131–141. SOS Publications, Fair Haven, NJ.

Yin, G., Jeney, G., Racz, T., Xu, P., Jun, X. and Jeney, Z., 2006. Effect of two Chinese herbs (*Astragalus radix* and *Scutellaria radix*) on non-specific immune response of tilapia, *Oreochromis niloticus*. *Aquaculture*, 253, 39-47.

Yin, G., Ardó, L., Thompson, K.D., Adams, A., Jeney, Z. and Jeney, G., 2009. Chinese herbs (*Astragalus radix* and *Ganoderma lucidum*) enhance immune response of carp, *Cyprinus carpio* and protection against *Aeromonas hydrophila*. *Fish and Shellfish Immunology*, 26(1), 140-145.

Zodape, G.V., 2010. Effect of *Aloe vera* Juice on toxicity induced by metal (chromium) in *Labeo Rohita* (Hamilton). *Journal of Applied Sciences Research*, 6, 1788-1793.