

Morphometric and molecular characterization of *Dactylogyrus lamellatus* isolated from farmed Grass carp, *Ctenopharyngodon idella* (Valenciennes, 1844), in Guilan Province, Iran

Daghigh Roohi J.¹; Dalimi A.^{2*}; Pourkazemi M.³; Ghasemi M.¹; Shamsi S.⁴

Received: July 2018

Accepted: April 2019

Abstract

Grass carp is an herbivorous species, which is actually popular in Iran. Monogenean parasites, particularly members of the family Dactylogyridae, have been one of the main causes of mortalities in Iranian fish farms and therefore they have been subject of many studies in Iran. The main aim of the present study was to describe morphological and molecular characteristics of monogenean parasites recovered from the farmed grass carp (*Ctenopharyngodon idella*) in the Guilan Province during 2015 to 2016. A total of 80 grass carps have been examined for infestation with monogenean parasites. 10923 *Dactylogyrus* species were recovered from 95% of the farmed grass carp. The parasite was first classified based on their morphological characteristics and identified as *Dactylogyrus lamellatus*. The parasite was then subjected to PCR and sequencing of the 28S rDNA gene. The phylogenetic tree showed that *D. lamellatus* genetically was similar to those previously reported from other countries and Mashhad farms in Iran.

Keywords: Grass carp, *Dactylogyrus lamellatus*, 28S rDNA, Guilan, Iran.

1-Inland water Aquaculture Research Center, Iranian Fisheries Sciences Research Institute, Agricultural Research Education and Extension Organization (AREEO), Bandar Anzali, Iran

2-Department of Parasitology, Faculty of Medical Sciences, Tarbiat Modares University, P.O.Box: 14115-331, Tehran, Iran

3-Iranian fisheries Science Research Institute, Agricultural Research Education and Extension Organization (AREEO), Tehran, Iran

4-School of Animal and Veterinary Sciences; Graham Centre for Agricultural Innovations, Charles Sturt University, NSW 2650, Australia

*Corresponding author's Email: dalimi_a@modares.ac.ir

Introduction

Grass carp is a large member of the carp and minnow family (Cyprinidae) and is native to south eastern Russia and north western China. This herbivorous species has been deliberately introduced into many countries for vegetation control purposes (Cudmore and Mandrak, 2004) and also for aquaculture due to rapid growth rate (Zolfinejad *et al.*, 2017). In Iran, this species was intentionally released for vegetation control in Anzali lagoon and introduced to the Guilan province for polyculture fish farming about 40 years ago. Today, this fish is artificially propagated in fish hatcheries of Guilan province and was sent to the whole of the country for cultivation.

Class Monogenea is a group of parasitic worms commonly found in fishes. These parasites feed of mucus and epithelial cells of the skin and gills and sometimes of the blood (Chaudhary *et al.*, 2013). Infestation with monogenean parasites may lead to serious hyperplasia of the gill filaments epithelium, impairs respiratory function, negatively affects growth, and even associates with high mortalities, especially in small carps (Lu *et al.*, 2012; Jiang *et al.*, 2013; Tu *et al.*, 2015). *Dactylogyrus* represents the most dominant genus among the Monogenea. These are common ectoparasites usually attached to the gills of freshwater fish of the family Cyprinidae (Gibson *et al.*, 1996; Woo 2006). The fish heavily infected with *Dactylogyrus* spp. is susceptible to secondary infestations that can play a

significant role in the pathogenicity of these parasites, which result in considerable economic losses (Dove and Ernst, 1998; Woo *et al.*, 2002; Reed *et al.*, 2009; Tu *et al.*, 2015).

Thus far, grass carp is known to be parasitized by at least two species of *Dactylogyrus* (Shamsi *et al.*, 2009; Daghigh Roohi *et al.*, 2016) in both pond and natural lake habitats in Iran. Identification of the Monogenea in Iran has often been based on a morphological basis. Monogenean identification solely based on morphological features can be problematic and ambiguous for closely related species. For effective diagnosis, treatment and control of parasitic diseases, the accurate identification of parasites is necessary (Mcmanus and Bowles, 1996). Recently, molecular markers have provided a useful alternative to the identification and validation of monogenean species (Cunningham *et al.*, 1995; Tu *et al.*, 2015). Our knowledge on genetic character of Iranian monogenea is poor and little is known about molecular aspects of these parasites (Mozhdeganlou *et al.*, 2011; Omidzahir *et al.*, 2012; Mousavi *et al.*, 2013; Omidzahir *et al.*, 2015). In the present study, *Dactylogyrus* that recovered from cultured grass carp in the Caspian Sea region are described morphologically, then their genetic characterization was done based on sequences of 28SrDNA gene.

Materials and methods

Fish collection

This work was carried out from April 2015 to November 2016 in Guilan Province, North of Iran (Fig. 1). A total of 80 grass carp (*Ctenopharyngodon idella*) were collected from 10 fish farms. Fish were one summer old with the average total length of 11.38 ± 2.84 cm and an average weight of 15.89 ± 12.69 g. The fish were placed in plastic tanks with water obtained from the collection sites and transferred to the Parasitology Laboratory of National Inland Water Aquaculture Research Center in Anzali where they were randomly distributed into several 100-L aquariums containing water from the fish farm and investigated within 72 hours. Fish were examined for the presence of Dactylogyrids parasites under a dissection microscope (4 to 40 magnifications). Live parasites were picked up from the gills scraping by using small needles. Then mounted in glycerine-jelly for morphological investigations (Gusse, 1983).



Figure 1: Distribution of studied ponds in Guilan Province, Iran.

Parasites collection and morphological identification

Monogenean parasites were recovered from gill filaments of *Ctenopharyngodon idella*. Parasite identification was performed based on the character of copulatory organ, the haptor sclerites like anchors, bars and hooks. Photographs of mounted parasites were captured at magnifications of 20X and 40X using a Nikon Eclipse 50i compound microscope with Nikon Digital Sight DS-L1 image analysis software and Nikon Digital Sight DS-SM camera. For morphometric analysis, a total of 7 point to point measurements were made on *Dactylogyrus* specimen photographs, using Image J software. Drawings were made with the aid of a drawing tube Nikon Y-IDT Japan. Finally, drawing of specimens and their measurement data were compared by parasites identification keys (Bykhovskaya-Pavlovskaya *et al.*, 1962; Gussev, 1985). After species identification, the same parasite specimen was stored in 75% ethanol tube for molecular analysis.

Molecular Studies

DNA extraction and PCR

The genomic DNA was extracted from one specimen using the YTA Genomic DNA Extraction Mini Kit (Yekta Tajhiz Azma, Iran) according to the manufacturer's manual. The 28S rDNA region of specimens was amplified using primers (forward, 5'TCTAGTAACGGCGAGTGAACG-3') (Chiary *et al.*, 2014) and the modified reverse primer (5'-

GTGGGAAGGTCTACCTCAGC-3').

Each amplification reaction was performed in a final volume of 25 µl containing 12.5 µl 2× Taq Mastermix, 1 µl of each primer (Macrogen, South Korea), and 2 µl of genomic template DNA. The amplification was carried out in a thermocycler (BIO-RAD, USA) under the following conditions: after an initial denaturation at 94 °C for 3 min, 35 cycles at 94 °C for 30 sec (denaturation), 59 °C for 30 sec (annealing), 72 °C for 1 min (extension) with a final extension at 72 °C for 10 min. The PCR products were analysed on 1.5% agarose gel and visualized under a UV illuminator, and the results were recorded.

DNA Sequencing

Purified fragments of PCR products were sequenced from both forward and reverse sites of each PCR product. Sequencing was carried out using the same primers as used for PCR amplification (Macrogen, South Korea).

Phylogenetic analysis

28SrDNA sequences of the *Dactylogyrus lamellatus* specimens in the present study were aligned separately using the Clustal W software and then manually adjusted to perform the phylogenetic analysis. Gaps and ambiguously aligned regions were removed. The phylogenetic tree was built using Mega 6.0 by the UPGMA method.

Statistical calculation and analysis

In this study, all calculations were performed using the following formulas (Bush *et al.*, 1997):

Prevalence = Number of infected hosts / Total of hosts × 100

Mean intensity = Total number of parasites / Number of infected hosts

Mean abundance = Total number of a particular parasite / Total number of infected and uninfected hosts

Dominance = Total number of a particular parasite / Total number of parasites × 100

The mean intensity of infestation and abundances of Monogenean among different seasons was tested by the Kruskal-Wallis test (KW, multiple comparisons) and Mann-Whitney test. The results were considered significant at the 95% level ($p < 0.05$). Computations were performed using the SPSS.15 programme.

Results

Totally, 10923 *Dactylogyrus* parasites were collected from 80 pieces of grass carp. Morphological examination of taxonomically important features such as the shape of the anchors, connective bar and the shape of a copulatory organ (Fig. 2) as well as detailed morphometric (Table 1) revealed that the only one species is identified as *D. lamellatus* in the present study.

The selected rDNA region was successfully amplified and sequenced. The sequence was deposited in GenBank with accession number MG657261. The 28SrDNA sequence size for *D. lamellatus* was 624 base pairs (bp) (Fig. 3). To build a

phylogenetic tree we used of different 28SrDNA sequences, which are brought in Table 2.

Table 1: Morphometrics (μm) of *Dactylogyrus lamellatus* recovered from *Ctenopharyngodon idella*.

Characters	Species	<i>Dactylogyrus lamellatus</i> (n=10)	
		Present Study	Bykhovskaya Pavlovskaya (1962)
Body length		320.55 \pm 80.92 (221.48-444.63)	Up to 480
Body width		71.57 \pm 21.23 (44.58-94.76)	Up to 110
Length of marginal hooks		27.12 \pm 3.16 (20.62-32.47)	21-31
Length of median hooks		35.59 \pm 2.02 (32.08-37.64)	35-41
Connecting bar		26.95 \pm 1.91 (24.04-29.11)	28-30
Length			
Connecting bar		3.07 \pm 0.42 (2.57-3.82)	Up to 4
Width			
The total length of the copulatory organ		42.68 \pm 3.77 (35.78-46.86)	About 50

n = number of specimens measured and range in size given in brackets.




Figure 2: Drawing of *Dactylogyrus lamellatus*: the opisthohapthoral central hook complex (A); copulatory organ (B).




Figure 3: Agarose gel electrophoresis of PCR amplification products from *Dactylogyrus lamellatus* in the present study following exposure to UV light.

Table 2: List of *Dactylogyrus* species used in this study with their host species, Genbank accession numbers and locality to build a phylogenetic tree.

Taxa	Host name	GenBank Accession number	Locality of collection
<i>Dactylogyrus extensus</i>	<i>Cyprinus carpio</i>	AJ969944	Zzech Republic
<i>Dactylogyrus nminutus</i>	<i>Cyprinus carpio</i>	MF926269	Guilan, Iran
<i>Dactylogyrus nachmerowi</i>	<i>Cyprinus carpio</i>	MF979966	Guilan, Iran
<i>Dactylogyrus vastator</i>	<i>Cyprinus carpio</i>	MF928712	Guilan, Iran
<i>Dactylogyrus aristichthys</i>	<i>Aristichthys nobilis</i>	MH023397	Guilan, Iran
<i>Dactylogyrus nobilis</i>	<i>Aristichthys nobilis</i>	MH023399	Guilan, Iran
<i>Dactylogyrus hypophthalmichthys</i>	<i>Aristichthys molitrix</i>	EF100532	China
<i>Dactylogyrus suchengtaii</i>	<i>Aristichthys molitrix</i>	MG825765	Guilan, Iran

Table 2 continued:

<i>Dactylogyrus anchoratus</i>	<i>Cyprinus carpio</i>	JX524546	Mashhad, Iran
<i>Dactylogyrus inexpectatus</i>	<i>Carassius auratus</i>	AJ969945	Zzech Republic
<i>Dactylogyrus quanfami</i>	<i>Cirrhinus moliorella</i>	EF100536	China
<i>Dactylogyrus lahei</i>	<i>Catla catla</i>	JX566720	India
<i>Dactylogyrus sphyrna</i>	<i>Rutilus rutilus</i>	AJ969943	Zzech Republic
<i>Dactylogyrus cryptomeres</i>	<i>Gobio gobio</i>	AJ969947	Zzech Republic
<i>Dactylogyrus lamellatus</i>	<i>Ctenopharyngodon idella</i>	AY307019	China
<i>Dactylogyrus lamellatus</i>	<i>Ctenopharyngodon idella</i>	MG657261	Guilan, Iran •
<i>Dactylogyrus lamellatus</i>	<i>Ctenopharyngodon idella</i>	KY039154	Mashhad, Iran
<i>Dactylogyrus lamellatus</i>	<i>Ctenopharyngodon idella</i>	EF100533	China
<i>Dactylogyrus lamellatus</i>	<i>Ctenopharyngodon idella</i>	AG969948	Zzech Republic
<i>Dactylogyrus pekinensis</i>	<i>Megalobrama amblycephala</i>	EF100535	China
<i>Dactylogyrus petruschewskyi</i>	<i>Megalobrama amblycephala</i>	AY548927	China
<i>Tetraonchus monenteron</i>	<i>Esox lucius</i>	AJ969953	Zzech Republic

(• This study)

In the present study, 95% of the fish were found to be infected with *D. lamellatus*. The highest rate of infestation has been observed with an average of 505 parasites per fish in summer. In Table 3 the prevalence, mean intensity, range and total of *D. lamellatus* in grass carp in different seasons is compared (Mann-Whitney test, $p<0.05$). As shown in Table 3 this parasite was present during all seasons

and cultivation period, however, the mean intensity was significantly higher in summer than other seasons ($p<0.05$). The low intensity of this parasite was observed in autumn.

Phylogenetic trees showed 100% similarity between our isolates and other *Dactylogyrus lamellatus* specimens registered in GenBank (Fig. 4).

Table 3: The prevalence, mean intensity, range and total number of *Dactylogyrus lamellatus* in *Ctenopharyngodon idella*, in the present study and different seasons.

Parameters Season	Prevalence (%)	Mean±SE	Range	Total
Spring N=47	100	98.17±66.04 ^a	3-836	4614
Summer N=12	100	505±146.51 ^c	61-1527	6060
Autumn N=21	80.95	14.64±5.92 ^b	1-101	249

N: number of examined fish, SE: Standard Error, * Data with different letters in a column are statistically significant at the $p<0.05$ level




Figure 4: Phylogenetic tree constructed by UPGMA analysis based on 28S rDNA for selected species of *Dactylogyrus*, with *Tetraonchus monenteron* used as the outgroup.

Discussion

Result indicated that only one species of monogenean, *D. lamellatus*, was recovered from *Ctenopharyngodon idella* in the Guilan province fish farms. The parasite identification was performed by morphometric and molecular analysis.

Grass carp is readily infested by several monogeneans, among them *D. lamellatus* is the most harmful parasite, which is spread throughout the country and causes high mortality in fingerlings (Jalali and Molnar, 1990). This survey demonstrated, *D. ctenopharyngodonis* as another *Dactylogyrus* on grass carp which introduced from China in 1991 and has been reported from some of the water resources in the country (Shamsi *et al.*, 2009; Daghigh Roohi *et al.*, 2016) has not been spread out in fish ponds yet. The study of grass carp parasites in Zarrineh rud River showed that 11.87% of the fish were infested with *D. lamellatus*, whereas in this study, the rate of fish infestation was

determined about 95% due to the high density of host fish in these ponds. As mentioned in table 3 the mean intensity of this parasite in Guilan fish ponds was also higher than natural waters like Anzali wetland in Guilan province or Khandaghlou Dam Lake in Zanjan province (Pazooki *et al.*, 2006; Daghigh Roohi *et al.*, 2016). Phylogenetic tree showed that *D. lamellatus* is genetically identical to those previously reported from other countries and also other parts of the country like Mashhad region which reported by Ahmadi *et al.* (2017). On the other hand, this phylogenetic tree shows that *D. lamellatus* is closely related to parasites of other Chinese carp like big head and silver carp. Because of grass carp is an alien fish, which introduced to Iran for cultivation so, we can conclude its private parasite, *D. lamellatus* probably has introduced to Iran due to lack of proper quarantine for their host. Therefore, as suggested by other authors a widely distributed species has

the ability to adapt itself to a broad range of climates.

As shown in Table 3, this parasite infestation was significantly higher in summer than other seasons. It seems that the parasite intensity significantly depends on the water temperature. According to the previous studies, the time of *D. lamellatus* sexual maturation depends on the temperature of water. It is indicated that the egg production in *D. lamellatus* began 8, 6 and 4 days after infection at temperatures of 17-19 °C , 20-24 °C and 22-26 °C respectively (Molnar, 1971).

Therefore, due to the temperature conditions in Guilan province, the different seasonal infection intensity, which obtained in our study, is justifiable.

This study provides further insights into the biodiversity and taxonomic status of monogenean parasites in Iran. Iran is a country with various zoogeographical regions, resulting in a high diversity of monogenean parasites reported from the country. Therefore, there is a huge potential for further work on the genetic character of monogenean fauna in the region and to understand the taxonomic relationship between Iranian monogenea with those from the rest of the world.

Acknowledgements

The present work is part of PhD. thesis, supported financially by Iranian fisheries Science Research Institute, Agricultural Research Education and Extension Organization (AREEO), Tehran, Iran. The author wishes to thank all the personnel of the

Parasitology Department of Medical Sciences Faculty of Tarbiat Modares University for their kind assistance.

Conflict of interest statement

We declare that we have no conflict of interest.

Reference

- Ahmadi, A., Borji, H., Naghibi, A., Nasiri, M.R. and Sharifiyazdi, H., 2017.** Morphologic and molecular (28SrDNA) characterization of *Dactylogyrus* spp. in *Cyprinus carpio* and *Ctenopharyngodon idella* in Mashhad, Iran. *Canadian Journal of Comparative Medicine*, 81(4), 280-284.
- Bush, A.O., Lafferty, K.D., Lotz, J.M. and Shostak, A.W., 1997.** Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *Journal of Parasitology*, 83(4), 575-583.
- Bykhovskaya-Pavlovskaya, I.E., Gussev, A.V., Dubinina, M.N., Izyumova, N.A., Smirnova, T.S., Sokolovskaya, A.L., Schtein, G.A., Shulman, S.S. and Epshtein, V.M., 1962.** Key to parasites of freshwater fishes of the USSR. Academy of Science of the USSR, Zoological Institute. 919 P.
- Chaudhary, A., Verma, C., Shobhna, M.V. and Singh, H.S., 2013.** A review of Monogenean diversity in India: Pathogens of fish diseases. *Journal of Coastal Life Medicine*, 1, 151-168.
DOI:10.12980/JCLM.1.20133D130

- Chiary, H.R., Chaudhary, A., Singh, H.S. and Goswami, U.C., 2014.** Molecular characterization of two non-native species of *Dactylogyrus* (Monogenea: Dactylogyridae) recovered from introduced hosts in India. *BioInvasions Record*, 3, 297-300. DOI:10.3391/bir.2014.3.4.12
- Cudmore, B. and Mandrak, N.E., 2004.** Biological Synopsis of Grass Carp (*Ctenopharyngodon idella*). Canadian Manuscript Report of Fisheries and Aquatic Sciences 2705. 44 P.
- Cunningham, C.O., McGillivray, D.M. and MacKenzie, K., 1995.** Phylogenetic analysis of *Gyrodactylus salaris* Malmberg, 1957 based on the small subunit (18S) ribosomal RNA gene. *Molecular and Biochemical Parasitology*, 71, 139-142
- Daghigh Roohi, J., Sattari, M., Masoumian, M., Nezamabadi, H., Mirhasheminasab, S. F., Asgharnia, M., Ghorbanpour, N., Nahrvar, M., Rufchaei, R., Ramezani, B., Musavi, A., Abbasi, K., Rastin, R., Mahisefat, F. and Syaddokht, J., 2016.** Study of parasites occurrence and intensity in fishes of Anzali lagoon. Iranian Fisheries Science Research Institute. Final report of the project No. 48174. 36 P (In Persian).
- Dove, A.D. and Ernst, I., 1998.** Concurrent invaders—four exotic species of Monogenea now established on exotic freshwater fishes in Australia. *International Journal for Parasitology*, 28, 1755-1764.
- Gibson, D.I., Timofeeva, T.A. and Gerasev, P.I., 1996.** A catalogue of the nominal species of the monogenean genus *Dactylogyrus* Diesing, 1850 and their host genera. *Systematic Parasitology*, 35, 3-48.
- GusseV, A.V., 1983.** Methods for the collection and processing fish parasites monogenean material Nauka, Leningrad, USSR, 48 P. (In Russian).
- GusseV, A.V., 1985.** Identification key to parasites of freshwater fish of USSR vol. II. Nauka, Leningrad, (In Russian). pp.10-424.
- Jalali, B. and Molnar, K., 1990.** Occurrence of monogeneans on freshwater fishes in Iran. *Dactylogyrus* spp. on cultured Iranian fishes. *Acta Veterinaria Hungarica*, 38, 239-242.
- Jiang, B., Chi, C., Fu, Y.W., Zhang, Q.Z. and Wang, G.X., 2013.** In vivo anthelmintic effect of flavonol rhamnosides from *Dryopteris crassirhizoma* against *Dactylogyrus intermedius* in goldfish (*Carassius auratus*). *Parasitology Research*, 112, 4097-4104. DOI:10.1007/s00436-013-3600-3
- Lu, C., Zhang, H.Y., Ji, J. and Wang, G.X., 2012.** In vivo anthelmintic activity of *Dryopteris crassirhizoma*, *Kochia scoparia*, and *Polygala tenuifolia* against *Dactylogyrus intermedius* (Monogenea) in goldfish (*Carassius auratus*). *Parasitology Research*, 110, 1085-1090. DOI:10.1007/s00436-011-2592-0
- Mcmanus, D.P. and Bowles, J., 1996.** Molecular genetic approaches to parasite identification: their value in

- diagnostic parasitology and systematic. *International Journal for Parasitology*, 96(7), 687-704.
- Molnar, K., 1971.** Studies on gill parasitosis of the grass carp (*Ctenopharyngodon idella*) caused by *Dactylogyrus lamellatus* Achmerow, 1952. 1. Morphology and Biology of *Dactylogyrus lamellatus*. *Acta Veterinaria Academiae Scientiarum Hungaricae, Tomus*, 21(2-3), 267-289.
- Mousavi, H.E., Omidzahir, S., Soltani, M., Shayan, P., Ebrahimzadeh, E., Mousavi, S. and Hoseini. M., 2013.** Morphometrical and molecular characterization of *Gyrodactylus cichlidarum* (Gyrodactylidae) from *Astronotus ocellatus* (Cichlidae) in Iran. *Comparative Clinical Pathology*, 22, 1093-1097. DOI:10.1007/s00580-012-1534-2
- Mozhdeganlou, Z., Ebrahimzadeh, M., Shayan, P., Soltani, M., Ebrahimzadeh, E. and Rostami, M., 2011.** Detection of single *Dactylogyrus* spp. in DNA extracted from infected gill tissue of fishes using polymerase chain reaction. *Iranian Journal of Veterinary Research*, 5, 77-144.
- Omidzahir, S., Ebrahimzadeh Mousavi, H.A., Soltani, M., Shayan, P., Ebrahimzadeh, E. and Hoseini, M., 2012.** Identification of *Gyrodactylus gurleyi* in *Carassius auratus* using morphometric and molecular characterization. *Iranian Journal of Veterinary Medicine*, 6, 41-46.
- Omidzahir, S., Mousavi, H., Shayan, P., Abkooh, E. and Mahmoodzadeh, H., 2015.** Morphometric and molecular analysis of *Gyrodactylus kobayashi* in *Carassius auratus* (Linnaeus, 1758). *Journal of Veterinary Research*, 70, 425-432.
- Pazooki, J., Jalali Jafari, B. and Ghobadian, M., 2006.** Monogenean species from freshwater fishes of Zanjan province, Iran. *Iranian Journal of Fisheries Sciences*, 6 (1), 103-112.
- Reed, P., Francis-Floyd, R., Klinger, R. and Petty, D., 2009.** Monogenean parasites of fish. Fisheries and Aquatic Sciences University of Florida UF, FA28, USA. pp. 1-4.
- Shamsi, S., Jalali, B. and Aghazadeh Meshgi, M., 2009.** Infection with *Dactylogyrus* spp. among introduced cyprinid fishes and their geographical distribution in Iran. *Iranian Journal of Veterinary Research*, 10, 70-74.
- Tu, X., Ling, F., Huang, A. and Wang, G., 2015.** The first report of *Dactylogyrus formosus* Kulwiec, 1927 (Monogenea: Dactylogyridae) from goldfish (*Carassius auratus*) in central China. *Parasitology Research*, DOI: 10.1007/s00436-015-4474-3
- Woo, P., Bruno, D. and Lim, L., 2002.** Diseases and disorders of finfish in cage culture. 365 hal. CABI, Malaysia. 353 P.
- Woo, P., 2006.** Fish diseases and disorders: protozoan and metazoan

infections. vol 1. 2th edition. CABI, UK. 791 P.

Zolfinejad, K., Khara, H. and Filizadeh, Y., 2017. Food preference and growth of grass carp, *Ctenopharyngodon idella* (Cuvier and Valenciennes, 1844) fed some aquatic and terrestrial plants. *Iranian Journal of Fisheries Sciences*, 16(4), 1278-1286.