

Research Article

Effect of monsoon variations on the antioxidant contents and antioxidant activity of *Sargassum illicifolium*

 Mahdi Abkenar A.^{1*}

1Department of Fisheries, Chabahar Branch, Islamic Azad University, Chabahar, Iran

 * Correspondence: aliabkenar@gmail.com
Keywords

Sargassum illicifolium,
Phaeophyta,
Antioxidant activity,
Monsoon,
Oman Sea

Abstract

Algae has recently drawn attention due to its strong potential for biological activities as a promising natural antioxidant resource. In this regard, seasonal variations of *Sargassum illicifolium* antioxidant activities were evaluated in the present study. *S. illicifolium* was collected from three coastal sites of the Oman Sea from April to May (spring) 2020 and October to November (autumn) 2020, representing pre- and post-monsoon, respectively. The methanolic extract was used to evaluate its antioxidant activities and properties. Regarding antioxidant activity (DPPH), there is a significant difference for the ferric reducing antioxidant power (FRAP), total flavonoids, total anthocyanin, and β -carotene variables analyzed between different sites during pre- and post-monsoon ($p > 0.05$). Meanwhile, IC₅₀ in pre-monsoon was significantly higher than in post-monsoon ($p < 0.05$). However, statistical analysis revealed a strong positive correlation between IC₅₀ and FRAP content in pre-monsoon; total phenolic and anthocyanin contents positively correlated to β -carotene post-monsoon. This study revealed that the seasonal variations lead to alter the biological activities regarding antioxidant contents in *S. illicifolium*, which need to be considered when considering the seaweed as a commercial antioxidant resource.

Article info

Received: November 2022

Accepted: December 2023

Published: July 2024



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Introduction

The distribution and diversity of marine macroalgae on the Iranian coast of the Oman Sea are prosperous (Mahdi Abkenar *et al.*, 2021). More than 83 dominant marine macroalgal species, such as Rhodophyta, Phaeophyta, and Chlorophyta have been recorded in this area (Gharanjik and Mahdi Abkenar, 1999).

Sargassum illicifolium is known as a tropical and sub-tropical species from Phaeophyta (family Sargassaceae and order Fucales), which is found abundantly in the coastal intertidal area of the Oman Sea, Iran (Rohani-Ghadikolaei *et al.*, 2012). *S. illicifolium* has several functional bioactive compounds that have created an imminent consequence in the biomedical fields (Marinho *et al.*, 2019). In addition, the *S. illicifolium* extract is well known for antioxidant, anti-inflammatory, antibacterial, and anticancer activities (Taheri *et al.*, 2018; Aghajanpoor *et al.*, 2020; Sahragard *et al.*, 2021; Yende *et al.*, 2021).

Global climate change has an extreme effect on macroalgae species that form habitats, including the extinction of algae populations in marine ecosystems (Wang *et al.*, 2021). According to Ramage (1971), a monsoon is traditionally a seasonal reverse wind along with the appropriate change in precipitation. However, it is now used to describe seasonal changes in atmospheric circulation and precipitation associated with the annual latitude oscillation of the intertropical convergence zone between the northern and southern equators (Wang *et al.*, 2019).

Algae present seasonal variations in their composition, e.g., proteins and lipids

(Marinho *et al.*, 2015a, b; Marinho *et al.*, 2019), which may also be the case of antioxidants, and this needs to be considered when evaluating their potential as a source of antioxidant compounds and capacity. Meanwhile, it is reported that the seasonal pattern of the pigment concentration generally is lower in summer and higher in winter; it has been reported for many seaweed species and is related to changes in the availability of light and nutritional concentration in seawater (Stengel and Dring, 1998; Schmid *et al.*, 2017).

The present study aimed to evaluate seasonal variation in the content of antioxidant activity and properties based on standard assays of *S. illicifolium*. It was collected at three stations at Tang, Chabahar, and Gowatr sites on the Oman Sea coast (Iran) from April to May (spring) and October to November (autumn) 2020, each representing pre- and post-monsoon, respectively.

Materials and methods

Seaweeds collection

Samples of *S. illicifolium* specimens were collected during low tide times (depending on the tide table obtained from www.tideforecast.com) by 2020 for two seasons: April to May (pre-monsoon) and October to November (post-monsoon), representing the autumn and spring, respectively, from three different locations on the northern coasts of the Oman Sea, Iran (Tang: latitude, 25° 14' 49" N; longitude: 59° 48' 02" E, Chabahar Bay: latitude, 25° 14' 17" N; longitude: 60° 38' 57" E; and Gowatr Bay: latitude, 25° 12' 59" N; longitude: 61° 29' 22" E). The Offshore

Fisheries Research Center determined taxonomic algae. The algae were identified by Aleem (1993); they belonged to the

family of *S. illicifolium* (Turner) from Phaeophyta (Fig. 1).




Figure 1: Sampling sites of *Sargassum illicifolium* in different locations of the Oman Sea, Iran.

Preparation of algae methanolic extract

After collection, the algae were taken to the laboratory in plastic bags containing seawater to avoid evaporation. Epiphytic and foreign materials were removed by washing first with seawater and then with distilled water to separate possible contaminants. Dry samples have been prepared by drying fresh algae in the air at room temperature for seven days and storing them in plastic bags for further analysis (González del Val *et al.*, 2001).

Sargassum illicifolium (≈ 1 kg) was washed with water to remove suspended materials, salts, epiphytes, and microorganisms and dried at room temperature, kept in a dark room and then, ground into a fine powder. Then 20 g of powdered samples were added to 10 mL of 80% methanol and stored at 37°C for 6 h in an orbital shaker incubator (AXYOS, Australia) and centrifugation (Kubota, KN-70, HD8292, Japan) at 1500×RPM for 15 minutes. After 72 h, they were getting

centrifuged agent extract. Finally, the extract obtained was filtered with WHATMAN No. 1 filter paper and concentrated to dry to produce a rough extract residue. The extract was diluted with solvents (50 mg/mL) and stored at 4°C for further analysis (Chidambararajan *et al.*, 2019) (Fig. 2).



Figure 2: Brown macroalgae *Sargassum illicifolium* collected from different locations in the Oman Sea, Iran.

Antioxidant activity (AOA)

The AOA of *S. illicifolium* extract was determined by hydrogen atom transfer

(HAD) (DDPH) and electron transfer (ET) (FRAB) methods (Heavisides *et al.*, 2018).

2, 2-diphenyl-1-picrylhydrazyl (DPPH %) radical scavenging ability

The DPPH scavenging activity assay of *S. illicifolium* extract was studied following Duan *et al.* (2006). In short, 1 mL of 0.15 mM DPPH was added to a different extract

$$\text{Scavenging activity (\%)} = \frac{A_{\text{Sample}} - A_{\text{Control}}}{A_{\text{Control}}} \times 100$$

The half-maximal inhibitory concentration (IC₅₀) was calculated by linear regression analysis and expressed as a mean of three determinations. IC₅₀ has been used as an index to compare the AOA of individuals. Ascorbic acid has been used as a positive control (1, 10, 100, and 1000 µg/mL), and reducing power has been expressed as equivalents of mg ascorbic acid (AAE) per mL of dry weight of the sample (DW).

Ferric reducing antioxidant power (FRAP) assay

The FRAP assay was determined using the Oyazu (1986) modification method. Briefly, 1 mL of *S. illicifolium* extract in different concentrations (0.1, 0.5, 1 and 2 mg/mL) was added to 1 mL of 0.2 M phosphate buffer (PSB, pH=6.6) and 1 mL of 1% potassium ferricyanide. The mixture was incubated at 50°C for 20 minutes, and 1 mL of 10% TCA was added to this reaction mixture. An aliquot of 1 mL from the incubation mixture was mixed with 1 mL of distilled water and 0.2 mL of 0.1% ferric chloride in the test tube. Absorbance was measured at 700 nm. Gallic acid was used as a standard power and reduction was

dilution (1, 10, 100, and 1000 µg/mL). Then, the reaction mixture was incubated in a dark room for 30 min. Its absorbance was measured at 517 nm using a spectrophotometer (ELISA Reader Epock, USA). DPPH (%) free radical referral activity is calculated using the following equation (Chidambarajan *et al.*, 2019):

expressed as equivalent to mg of gallic acid (GAE) per g of dry weight sample (DW).

Total phenolic and flavonoid content

The total phenolic content (TPC) of *S. illicifolium* extract was assessed by the Folin-Ciocalteu method (Cagalj *et al.*, 2021) using a UV spectrophotometer (Shimadzu, UV-160A, Japan) with minor modification. Briefly, 20 µL of *S. illicifolium* extract (1 mg/mL) reacted with 1.50 mL of distilled water and 125 µL of 10% Folin-Ciocalteu reagent. The solution was mixed, and after one minute, 375 µL of 20% sodium carbonate solution and 475 µL of distilled water were added. The mixture was left in the dark for 2 h at room temperature. Absorbance was read at 720 nm using a spectrophotometer. The standard curve (0-500 mg/l) was prepared using gallic acid for the calculation of phenolic content ($y = 0.004x$, $r^2 = 0.9999$). The TPC was expressed as mg of gallic acid equivalents (GAE) per g of dry weight sample (DW).

The total content of flavonoids (TFC) in *S. illicifolium* extract was assessed by the aluminum chloride colorimetric method described by Cagalj *et al.* (2021). Briefly, 250 µL of *S. illicifolium* extract (1 mg/mL)

was mixed with 750 μL of 96% ethanol, 50 μL of 10% (w/v) aluminum chloride, 50 μL 1M sodium acetate, and 1400 μL distilled water. The mixture was stored for 30 minutes at room temperature. Absorbance was read at 415 nm using a spectrophotometer. The standard calibration curve (0-100 $\mu\text{g}/\text{mL}$) was plotted using a quercetin solution ($Y=0.0076-0.0042x$, $r^2=0.9412$). TFC is the equivalent of mg quercetin (QE) per g of dry weight sample (DW).

Total anthocyanin content

The total anthocyanin content has been measured using a differential protocol spectrophotometric pH, as Heavisides *et al.* (2018) described, with minor

modifications. Briefly, 0.5 mL of *S. illicifolium* extract mixed with 3.5 mL of 0.025 M potassium chloride buffer; pH=1. The mixture was stirred and left at room temperature for 15 minutes. Then, the absorbance was measured at 515 and 700 nm against distilled water as a blank.

Then, the extract was combined with 3.5 mL of 0.025 M sodium acetate buffer (pH=4.5). Once again, the mixture was stirred and left at room temperature for 15 minutes. The absorbance of the mixture was measured at the same wavelengths (512 and 700 nm). The total anthocyanin content has been calculated using the following equation:

$$TAC (mgC - 3 - GE/100g DW) = (A \times MW \times 1000 / (\epsilon \times C))$$

Where, A is absorbance= $(A_{\lambda 515} - A_{\lambda 700})$ pH 1.0 - $(A_{\lambda 515} - A_{\lambda 700})$ pH 4.5; MW is molecular weight for cyanidin-3-glucoside=484.2; DF is the factor of the extract; ϵ is the molar absorptivity of cyanidin-3-glucoside=24825; C is the concentration of the buffer in mg/mL=0.025. The result was expressed as milligrams of cyanidin-3-glucoside equivalents (mg C-3-GE) per 100 grams of the dry weight of the sample (DW).

β -carotene content

The quantification of the β -carotene was carried out following the methods of Eijkelhoff and Dekker (1995) with minor modifications. Briefly, 20 g of *S. illicifolium* crude extract was dissolved in 10 mL of 85% acetone-hexane and centrifuged at 3000 rpm for 15 minutes at 4°C. Then, the absorbance was measured at 453, 505, 645, and 663 nm using a UV spectrophotometer (Shimadzu, UV -160A, Japan). The contents of the β -carotene have been calculated according to the following equations and expressed as mg per g of dry weight (DW):

$$\beta - Carotene (mg/100g DW) = 0,216A_{663} - 1,22A_{645} - 0,304A_{505} + 0,452A_{453}$$

Statistical analysis

All statistical analyses were conducted by SPSS Software (New York, USA) version

16.0 in triplicate. Shapiro-Wilk and Levene's tests were used to check the normality and homogeneity of variance,

respectively. To determine whether there are differences between means, ANOVA analysis and Duncan's multiple range test (DMRT) as a post hoc test are applied to correlate the potential of antioxidants in the samples. In addition, the comparison between monitored factors and locations in pre and post-monsoon with independent sample T-tests.

Results

DPPH (%) scavenging activity

Table 1 indicates no significant variation in the scavenging DPPH (%) radicals of methanolic *S. illicifolium* extract tested at 1, 10, 100, and 1000 µg/mL harvested in pre- and post-monsoon among three locations of Tang, Chabahar, and Gowatr gulf ($p>0.05$).

Table 1: DPPH radical scavenging activity (%DPPH) of 1, 10, 100, and 1000 µg/ml of *Sargassum illicifolium* extracts in different seasons.

Seasons	Concentration (µg/mL)	Stations		
		Tang	Chabahar	Gowatr
Pre- (Spring)	1	48.46±1.37	41.72±1.74	44.9±1.91
	10	50.19±83	58.80±2.01	53.15±0.45
	100	66.28±3.34	57.20±1.15	59.79±0.74
	1000	76.43±1.19	57.96±1.23	61.90±3.01
Post- (autumn)	1	56.19±0.85	53.76±0.99	50.38±1.09
	10	63.27±3.45	61.96±1.36	56.76±2.59
	100	76.98±2.36	66.80±2.78	60.26±2.34
	1000	82.50±2.54	71.12±2.46	61.81±2.19
	Ascorbic acid	55.1±1.00	55.1±1.00	55.1±1.00

*Data were presented as the average of four replicates standard deviation (±SD).

IC50

As shown in Figure 3, The IC50 content of ascorbic acid as a standard antioxidant has shown a significant difference ($p<0.05$) between methanolic *S. illicifolium* extract which was harvested in pre-monsoon (85.80±11.40 mg AAE/mL) compared to post-monsoon (46.63±4.65 mg AAE/mL) (sig=0.020; t=9.53; DF=16). Based on the results, the IC50 content between the three

locations of Tang, Chabahar, and Gowatr in pre-monsoon was significantly higher than post-monsoon ($p>0.05$). However, there were no significant differences between samples harvested from studied locations in pre- and post-monsoon, separately ($p>0.05$). The IC50 content in *S. illicifolium* extract at pre-monsoon ranges from 67.71 to 101.31 mg AAE/mL, whereas in post-monsoon it is found at 41.71 to 54.17 mg AAE/mL.




Figure 3: Comparison of the IC50 content of *Sargassum illicifolium* extract between three different locations (Left) and monsoon conditions (Right) on the northern coast of the Oman Sea ($p>0.05$).

Ferric-reducing power (FRAP) assay

Figure 4 presents the comparison of the FRAP content in *S. illicifolium* extract among three locations and seasonal variations on the northern coast of the Oman Sea. According to the results, there was no significant difference found between *S. illicifolium* extract which was harvested in pre- (0.62 ± 0.15 mg GAE/g DW) compared to post- (1.21 ± 0.11 mg GAE/g DW) monsoon (sig=0.28; $t=-9.45$;




Figure 4: Comparison of the FRAP content in *Sargassum illicifolium* extract between three different locations (Left) and monsoon conditions (Right) on the northern coast of the Oman Sea ($p>0.05$).

The FRAP content in *S. illicifolium* extract at pre-monsoon ranged from 0.43 to 0.81 mg GAE/g DW, whereas at post-monsoon, it was found to be 1.11 to 1.39 mg GAE /g DW.

Total phenolic content (TPC)

Compression of The TPC content in *S. illicifolium* extract is presented in Figure 5. Based on the results for both the pre- (3.42 ± 0.38 mg GEA/g DW) and post- (3.77 ± 0.65 mg GAE/ g DW) monsoon, no significant differences were observed (sig=0.12; $t=-1.41$; DF=12.87). However, no significant differences were observed between the three locations of Tang, Chabahar, and Gowatr at pre- and post-

DF=14.65). Based on the results, the FRAP content between the three locations of Tang, Chabahar, and Gowatr in pre-monsoon was significantly lower than at post-monsoon ($p> 0.05$). However, there were no significant differences between samples harvested from studied locations at pre- and post-monsoon, separately ($p> 0.05$).

monsoon, separately ($p>0.05$). The TPC compounds in the extract harvested in the pre-monsoon ranged from 3.12 to 4.08 mg GEA/g DW, whereas the foe sample harvested in the post-monsoon was found at 2.79 to 4.91 mg GEA/g DW.

Total flavonoid content (TFC)

As shown in Figure 6, there was no significant difference between TFC content measured in samples harvested from the pre- (12.45 ± 2.31 mg QE/g DW) and post- (28.14 ± 4.56 mg QE/g DW) monsoon for Methanolic *S. illicifolium* extract (Sig=0.07; $T=- 9.19$; DF=11.85). However, the TFC content between the three locations of Tang, Chabahar, and Gowatr at pre-

monsoon was significantly lower than at post-monsoon ($p>0.05$). However, there were no significant differences between samples harvested from all studied locations at pre- and post-monsoon, separately ($p>0.05$). The content of total

flavonoid compounds in the pre-monsoon ranges from 8.38 to 16.07 mg QE/g DW, while in the post-monsoon it is found at 21.81 to 32.17 mg QE/g DW.




Figure 5: Comparison of the TPC content in *S. i.* extract between three different locations (Left) and monsoon conditions (Right) on the northern coast of the Oman Sea ($p>0.05$).




Figure 6: Comparison of the TFC content in *Sargassum illicifolium* extract between three different locations (Left) and monsoon conditions (Right) on the northern coast of the Oman Sea ($p>0.05$).

Total anthocyanin content

Figure 7 shows no significant ($p>0.05$) seasonal variation for the anthocyanin content of *S. illicifolium* extract harvested in pre (3.97 ± 0.68 mg C-3-GE/100g DW) and post (8.23 ± 1.04 mg C-3-GE/100g DW) monsoon (Sig=0.24; T=- 10.28; DF=13.77). However, there was no significant difference found in the three locations of samples harvested pre-monsoon ($p>0.05$); Meanwhile, we observed a statistically significant difference for samples harvested post-

monsoon between different locations ($p<0.05$). Based on the results, at the post-monsoon, the anthocyanin content in samples harvested from Gowatr (7.00 ± 0.62 mg C-3-GE/100g DW) was significantly lower than in Tang (9.10 ± 0.43 mg C-3-GE/100g DW) and Chabahar (8.59 ± 0.42 mg C-3-GE/100g DW) ($p<0.05$). The content of anthocyanin compounds in the pre-monsoon ranged from 3.15 to 5.11 mg C-3-GE/100g DW, while in the post-monsoon it was found at 6.28 to 9.59 mg C-3-GE/100g DW.




Figure 7: Comparison of the anthocyanin content in *Sargassum illicifolium* extract between three different locations (Left) and monsoon conditions (Right) on the northern coast of the Oman Sea ($p > 0.05$).

β -carotene content

Figure 8 shows no significant ($p > 0.05$) seasonal variation in the contents of β -carotene in *S. illicifolium* extract in the pre (0.076 ± 0.015 mg/g DW) and the post (0.079 ± 0.010 mg/g DW) monsoon (Sig=0.42; T= -0.57; DF=13.81). The β -carotene content between all studied locations showed a significant difference during pre- and post-monsoon ($p < 0.05$). The highest and lowest levels of β -carotene at pre-monsoon were observed in Chabahar Bay (0.088 ± 0.006 mg/g DW) and Tang (0.060 ± 0.016 mg/g DW) locations,

respectively ($p < 0.05$). The β -carotene content measurement in the sample harvested from Gowatr at pre-monsoon was 0.079 ± 0.006 mg/g DW. For post-monsoon, the β -carotene content in the sample harvested from Gowatr (0.068 ± 0.003 mg/g DW) was significantly lower than the Tang (0.085 ± 0.006 mg/g DW) and Chabahar Bay (0.086 ± 0.008 mg/g DW) samples ($p < 0.05$). The β -carotene content in the methanolic extract of *S. illicifolium* harvested in pre-monsoon ranges from 0.042 to 0.093 mg/g DW, while in post-monsoon it is found at 0.066 to 0.095 mg/g DW.




Figure 8: Comparison of the β -carotene content in *Sargassum illicifolium* extract between three different locations (Left) and monsoon conditions (Right) on the northern coast of the Oman Sea ($p > 0.05$).

Correlation analysis

The Pearson correlation analysis was carried out to determine the relationship

between each antioxidant activity index and the properties of *S. illicifolium* extract in different seasonal samplings. As shown in

Table 2, there was a significant strong positive correlation between IC₅₀ with FAR content ($r=0.83$; $p<0.001$) in pre-monsoon.

However, at post-monsoon (Table 3), there was a significant negative correlation

between IC₅₀ and DPPH content ($r=-0.67$; $p<0.05$). However, we observed a significantly positive correlation between TPC ($r=0.68$; $p<0.05$) and anthocyanin ($r=0.77$; $p<0.05$) with β -carotene content.

Table 2: Pearson's correlation coefficients between the variables of pre-monsoon.

Indices	DPPH	IC50	FRAP	TPC	TFC	Anthocyanin	β -Carotene
DPPH	1	-0.24	-0.27	-0.17	-0.59	-0.58	0.35
IC50	-0.24	1	0.83**	0.47	0.26	-0.21	-0.19
FRAB	-0.27	0.83**	1	0.20	0.36	0.13	-0.19
TPC	-0.17	0.47	0.20	1	-0.23	0.02	0.15
TFC	-0.59	0.26	0.36	-0.23	1	0.40	-0.46
Anthocyanin	-0.58	-0.21	0.13	0.02	0.40	1	0.18
β -carotene	0.35	-0.19	-0.19	0.15	-0.46	0.18	1

Table 3: Pearson's correlation coefficients between the variables to post-monsoon.

Variables	DPPH	IC50	FRAP	TPC	TFC	Anthocyanin	β -carotene
DPPH	1	-0.67*	-0.63	-0.20	-0.22	-0.47	-0.21
IC50	-0.67*	1	0.38	0.23	0.62	0.45	0.55
FRAP	-0.63	0.38	1	-0.13	0.16	0.54	0.05
TPC	-0.20	0.23	-0.13	1	0.52	0.42	0.68*
TFC	-0.22	0.62	0.16	0.52	1	0.40	0.61
Anthocyanin.	-0.47	0.45	0.54	0.42	0.40	1	0.77*
β -carotene	-0.21	0.55	0.05	0.68*	0.61	0.77*	1

Discussion

S. illicifolium is a good source of bioactive constituents that cope with various applications in different industries (Yende *et al.*, 2021). To our knowledge, this is the first report on the antioxidant content and activities of *S. illicifolium* in different seasons of the Iranian coast of the Oman Sea. The results indicated the difference in antioxidant activity, FRAP, TPC, TFC, anthocyanin, and β -carotene from seaweed based on different geographical locations. Based on the results, seasonal changes in DPPH radical scavenging activity showed no statistical significance between geographical locations. Nevertheless, the lowest IC₅₀ values, corresponding to the highest radical scavenging activity, were found from October to November (post-

monsoon), matching the period in which the maximum concentrations of FRAP, TFC, and anthocyanin were found, respectively. This data suggests a possible correlation between DPPH scavenging activity and TFC and TPC. This assumption was further supported by the partial least square coefficient plot revealing that both FRAP were positively correlated with IC₅₀ at pre-monsoon and DPPH with IC₅₀ at post-monsoon, suggesting that these variables had a positive effect on the radical scavenging activity. These results follow previous studies, which directly correlated DPPH scavenging activity with the IC₅₀, FRAP, and TFC found in the extracts of both seaweeds (Palanisamy *et al.*, 2017; Sharifian *et al.*, 2019; Nosrati *et al.*, 2021; Sahragard *et al.*, 2021). Nosrati *et al.*

(2021) DPPH activity of brown algae of *Sargassum glaucescens* was reported to be between 16.1-31.9%, which was lower than our results. Palanisamy *et al.* (2017) studied the antioxidant effect of *Sargassum polycystum* and reported that DPPH and FRAP contents were 61.2% and 67.59%, respectively. The DPPH radical scavenging activity (12.08 ± 0.28 and 7.24 ± 0.18 mg GAE/100g) was high in *Zonaria tournefortii* and *Sphaerococcus coronopifolius* during summer. In comparison, for *Halopteris scoparia* (11.12 ± 0.19 mg GAE/100g), the highest DPPH activity was observed in autumn. Likewise, the ferric reducing power for *H. scoparia* (41.93 ± 1.08 mg GAE/100g), while the highest level of FRAP was observed at autumn in *S. coronopifolius* and *Z. tournefortii* (54.69 ± 0.75 and 31.02 ± 1.45 mg GAE/100g, respectively) (Fellah *et al.*, 2017).

On the other hand, the analysis revealed a positive correlation between TPC and anthocyanin with β -carotene, suggesting a negative impact of β -carotene on the radical scavenging activity. However, in several studies, pigments have been reported to have vigorous radical scavenging activity (Jensen, 1966; Stengel and Dring, 1998; Petrushkina *et al.*, 2017; Marinho *et al.*, 2019).

Our results demonstrate that the total phenolic content was not significantly different between autumn and spring in *S. illicifolium* samples. The TPC found in this study was previously reported for water ($209 \mu\text{g GAE/g DM}$) and ethanol extracts ($354 \mu\text{g GAE/g DM}$) of *Saccharina latissima* (Farvin and Jacobsen, 2015). This difference may likely be related to the

solvents' polarity (Marinho *et al.*, 2019). On the other hand, the concentrations of TPC reported here for *S. illicifolium* (2.79-4.91 mg GAE/g DM) are similar to those reported for *L. digitata* harvested monthly on a sheltered rocky shore in Brittany (France), ranging seasonally from 0.09 to 0.19% DM (using phloroglucinol as standard; Connan *et al.*, 2004). Likewise, the seasonal variation in the TPC followed the pattern reported for *S. latissima* in the same study, with the highest values found in winter and the lowest in summer/autumn (Marinho *et al.*, 2019). However, Fellah *et al.* (2017) reported that TPC was higher in *S. coronopifolius* (144.33 ± 1.76 mg GAE/100g) in autumn. The TPC content in *Zonaria tournefortii* in the summer (12.08 ± 0.28 mg GAE/100g) was significantly higher than in the spring (6.61 ± 0.14 mg GAE/100g) and autumn (3.73 ± 0.28 mg GAE/100g). It could be that the annual variation in phenolic content is dependent on diverse species composition and the different environmental conditions obtained in each season. The phenolic compound content of macroalgae varies with the reproductive stage of algae and with physical factors such as light density and quality, photoperiod, and temperature (Fellah *et al.*, 2017). Gobbo-Neto and Lopes (2007) report that these factors have correlations and do not act in isolation; they may jointly influence the secondary metabolism.

To our knowledge, the amount of TFC did not show significant differences between the samples harvested in spring (pre-monsoon) and autumn (post-monsoon). In comparison, samples harvested in spring showed a higher level of

TFC compared to samples harvested in autumn. TFC varied seasonally, reaching a maximum in summer and a minimum in winter. Flavonoids are UV-absorbing compounds that function as UV screens in terrestrial plants, among many other vital functions. For instance, UV-B can induce quercetin, which is primarily found in epidermal cells. This role is assumed to be played by simpler UV-absorbing compounds, such as mycosporine-like amino acids, in the aquatic environment (Rozema *et al.*, 2002). Nevertheless, structural identification of flavonoids in red and brown seaweeds, and microalgae, has recently been reported, contradicting the general assumption that these compounds could be exclusively present in terrestrial plants (Klejdus *et al.*, 2010; Rodríguez-Bernaldo de Quirós *et al.*, 2010; Goiris *et al.*, 2014). The TFC found in the present study (8.38–32.17 mg QE/g DM over all seasons) is much higher than in several brown and green seaweed species originating from the Black Sea (6.5–229 µg RE/g DM).

The qualitative pigment composition did not change seasonally, whereas the number of specific pigments changed seasonally and with β-carotene concentrations that could be of commercial interest (Marinho *et al.*, 2019).

β-carotene is the first and most crucial carotenoid (Butler *et al.*, 2018). Our results reveal a seasonal pattern in the concentration of β-carotene, which ranged from summer and autumn. However, the samples harvested in the Tang location post-monsoon were significantly higher than those harvested pre-monsoon. Furthermore, the increased β-carotene

content observed in this location in autumn is supported by studies showing that this increase can be a response to, i.e., shade and depth adaptation to increase light-harvesting efficiency (Ramus *et al.*, 1977; Fortes and Lüning, 1980). These results can be correlated with the lower irradiance and short day length observed during autumn and winter in Iran. On the other hand, several specific pigments changed seasonally, and with β-carotene concentrations that could be of commercial interest, varying seasonally from 5.29 to 23.46 µg/g DM of *S. latissima* in September, November, and January by Hjørnø Havbrug A/S in the vicinity of Horsens Fjord, in the inner Danish waters (Denmark) (Marinho *et al.*, 2019).

Conclusions

The algal biomass harvested from April to May (spring) and October to November (autumn) showed seasonal differences in *S. illicifolium* biomass, except for IC50. However, it exhibited a blend of characteristics from the Tang, Chababar, and Gowatr sites in terms of FRAP content, total flavonoids, total anthocyanin, and β-carotene, highlighting the importance of considering these factors when evaluating its antioxidant potential. The FRAP content positively affected the radical scavenging activity, especially IC 50 in spring (pre-monsoon). In autumn (post-monsoon), β-carotene had a negative effect on the radical scavenging activity, especially TFC and anthocyanin content.

Conflicts of interest

The author does not have any conflicts of interest to declare.

References

- Aghajanoor, N., Babakhani, A. and Tabarsa, M., 2020.** Optimization of the extraction of antioxidant compounds from brown algae *Sargassum angustifolium* in the Persian Gulf by Response Surface Methodology (RSM). *Aquaculture Sciences*, 8(1), 43-58.
- Aleem, A.A., 1993.** The Marine algae of Alexandri, Faculty of Science, University of Alexandria, Egypt. 139 P.
- Butler, T.O., McDougall, G.J., Campbell, R., Stanley, M.S. and Day, J.G., 2018.** Media screening for obtaining *Haematococcus pluvialis* red motile macrozooids rich in astaxanthin and fatty acids. *Biology*, 7(1), 2. DOI:10.3390/biology7010002
- Chidambarajan, P., Keerthana, V., Priyadharshini, K. and Sakthivel, B., 2019.** *In vitro* antioxidant and anticancer activity of *Ulva lactuca* L. using molt-3 cell line. *Asian Journal of Pharmaceutical and Clinical Research*, 12(5), 75-78. DOI:10.22159/ajpcr.2019.v12i5.29825
- Connan, S., Goulard, F., Stiger, V., Deslandes, E. and Ar Gall, E., 2004.** Interspecific and temporal variation in phlorotannin levels in an assemblage of brown algae. *Botanica Marina*, 47(5), 410-416. DOI:10.1515/BOT.2004.057
- Duan, X.J., Zhang, W.W., Li, X.M. and Wang, B.G., 2006.** Evaluation of antioxidant property of extract and fractions obtained from a red alga, *Polysiphonia urceolata*. *Food Chemistry*, 95(1), 37-43. DOI:10.1016/j.foodchem.2004.12.015
- Eijkelhoff, C. and Dekker, J.P., 1995.** Determination of the pigment stoichiometry of the photochemical reaction center of photosystem. *Biochimica et Biophysica Acta (BBA)*, 1231(1), 21 - 28.
- Farvin, K.H.S. and Jacobsen, C., 2015.** Antioxidant activity of seaweed extracts: *in vitro* assays, evaluation in 5% fish oil-in-water emulsions and characterization. *Journal of the American Oil Chemists' Society*, 92(4), 571-587. DOI:10.1007/s11746-015-2624-5
- Fellah, F., Louaileche, H., Dehbi-Zebboudj, A. and Touati, N., 2017.** Seasonal variation in the phenolic compound content and antioxidant activities of three selected species of seaweeds from Tiskerte islet, Bejaia, Algeria. *Journal of Materials and Environmental Sciences*, 8(12), 4451 - 4456.
- Fortes, M.D. and Lüning, K., 1980.** Growth rates of North Sea macroalgae in relation to temperature, irradiance and photoperiod. *Helgol Meeresunters*, 34, 15-29. DOI:10.1007/BF01983538
- Gharanjik, B. and Mahdi Abkenar, A., 1999.** Identification of seaweed in Sistan and Baluchestan province coastal zone. *Iranian Scientific Fisheries Journal*, 9(1), 37-48. (In Persian)
- Gobbo-Neto, L. and Lopes, N.P., 2007.** Plantas medicinais: fatores de influência no conteúdo de metabólitos secundários. *Química nova*, 30(2), 374-381. DOI:10.1590/S0100-40422007000200026
- Goiris, K., Muylaert, K., Voorspoels, S., Noten, B., De Paepe, D., E Baart, G.J. and De Cooman, L., 2014.** Detection of flavonoids in microalgae from different

- evolutionary lineages. *Journal of Phycology*, 50(3), 483–492. DOI:10.1111/jpy.12180
- González del Val, A., Platas, G., Basilio, A., Cabello, A., Gorrochategui, J., Suay, I., Vicente, F., Portillo, E., Jiménez del Río, M., Reina, G.G. and Peláez F., 2001.** Screening of antimicrobial activities in red, green and brown macroalgae from Gran Canaria (Canary Islands, Spain). *International Microbiology*, 4(1), 35–40. DOI:10.1007/s101230100006
- Heavisides, E., Rouger, C., Reichel, A.F., Ulrich, C., Wenzel-Storjohann, A., Sebens, S. and Tasdemir D., 2018.** Seasonal variations in the metabolome and bioactivity profile of *Fucus vesiculosus* extracted by an optimised, pressurised liquid extraction protocol. *Marine Drugs*, 16(12), 503. DOI:10.3390/md16120503
- Jensen, A., 1966.** Carotenoids of Norwegian brown seaweeds and of seaweed meals. *Norwegian Institute of Seaweed Research*. Report No. 31, 1–138.
- Klejduš, B., Lojková, L., Plaza, M., Šnóbllová, M. and Štěřbová, D., 2010.** Hyphenated technique for the extraction and determination of isoflavones in algae: Ultrasound-assisted supercritical fluid extraction followed by fast chromatography with tandem mass spectrometry. *Journal of Chromatography A*, 1217(51), 7956–7965. DOI:10.1016/j.chroma.2010.07.020
- Mahdi Abkenar, A., Yahyavi, M., Esmaeili, M. and Rombenso, A., 2021.** High bioaccumulation factors and ecological risk index of Cd, and Hg in Indian white shrimp, hooded oyster, brown algae and sediment in northern coasts of the Gulf of Oman before and after a monsoon. *Regional studies in Marine Science*, 41, 101552. DOI:10.1016/j.rsma.2020.101552
- Marinho, G., Holdt, S., Jacobsen, C. and Angelidaki, I., 2015a.** Lipids and composition of fatty acids of *Saccharina latissima* cultivated year-round in integrated multi-trophic aquaculture. *Marine Drugs*, 13(7), 4357–4374. DOI:10.3390/md13074357
- Marinho, G.S., Holdt, S.L. and Angelidaki, I., 2015b.** Seasonal variations in the amino acid profile and protein nutritional value of *Saccharina latissima* cultivated in a commercial IMTA system. *Journal of Applied Phycology*, 27, 1991–2000. DOI:10.1007/s10811-015-0546-0
- Marinho, G.S., Sørensen, A.D.M., Safafar, H., Pedersen, A.H. and Løvstad Holdt, S. 2019.** Antioxidant content and activity of the seaweed *Saccharina latissima*: a seasonal perspective. *Journal of Applied Phycology*, 31(2), 1343–1354. DOI:10.1007/s10811-018-1650-8
- Nosrati, A., Taheri, A. and Khajeh-Amiri, C., 2021.** Evaluation of antioxidant properties of brown algae (*Sargassum glaucescens*) extract and optimization of extraction of its antioxidant compounds with artificial neural network. *Journal of Marine Medicine*, 3(3), 153–161. (In Persian)
- Palanisamy, S., Vinosha, M., Marudhupandi, T., Rajasekar, P. and Prabhu, N.M., 2017.** Isolation of

- fucoidan from *Sargassum polycystum* brown algae: Structural characterization, in vitro antioxidant and anticancer activity. *International Journal of Biological Macromolecules*, 102, 405-412.
DOI:10.1016/j.ijbiomac.2017.03.182
- Petrushkina, M., Gusev, E., Sorokin, B., Zotko, N., Mamaeva, A., Filimonova, A., et al., 2017.** Fucoxanthin production by heterokont microalgae. *Algal Research*, 24, 387–393.
DOI:10.1016/j.algal.2017.03.016
- Ramage, C., 1971.** Monsoon Meteorology. International Geophysics Series, Academic Press, USA. 296 P.
- Ramus, J., Lemons, F. and Zimmerman, C., 1977.** Adaptation of light-harvesting pigments to downwelling light and the consequent photosynthetic performance of the eulittoral rockweeds *Ascophyllum nodosum* and *Fucus vesiculosus*. *Marine Biology*, 42(4), 293-303.
DOI:10.1007/BF00402191
- Rodríguez-Bernaldo de Quirós, A., Lage-Yusty, M.A. and López-Hernández, J., 2010.** Determination of phenolic compounds in macroalgae for human consumption. *Food Chemistry*, 121(2), 634–638.
DOI:10.1016/j.foodchem.2009.12.078
- Rohani-Ghadikolaei, K., Abdulalian, E. and Ng, W.K., 2012.** Evaluation of the proximate, fatty acid and mineral composition of representative green, brown and red seaweeds from the Persian Gulf of Iran as potential food and feed resources. *Journal of Food Science Technology*, 49(6), 774-780.
DOI:10.1007/s13197-010-0220-0
- Rozema, J., Björn, L.O., Bornman, J.F., Gaberščik, A., Häder, D.P., Trošt, T., Germ, M., Klisch, M., Gröniger, A., Sinha, R.P. and Lebert, M., 2002.** The role of UV-B radiation in aquatic and terrestrial ecosystems-an experimental and functional analysis of the evolution of UV-absorbing compounds. *Journal of Photochemistry and Photobiology B: Biology*, 66(1), 2–12.
DOI:10.1016/s1011-1344(01)00269-x
- Sahragard, K., Tabarsa, M. and Ahmadi Gavlighi, H., 2021.** Isolation, purification, anti-diabetic and antioxidant properties of fucoidan from brown seaweed *Sargassum ilicifolium*. *Journal of Fisheries*, 74 (2), 223-234.
DOI:10.22059/JFISHERIES.2021.319673.1233
- Schmid, M., Guihéneuf, F. and Stengel, D.B., 2017.** Ecological and commercial implications of temporal and spatial variability in the composition of pigments and fatty acids in five Irish macroalgae. *Marine Biology*, 164(8), 158. DOI:10.1007/s00227-017-3188-8
- Sharifian, S., Shahbanpour, B., Taheri, A. and Kordjazi M., 2019.** Effects of different solvents on the phenolic compounds and antioxidant properties of brown seaweeds, *Nizimuddinia zanardinii* (Schiffner) P.C. Silva and *Padina australis* Hauck. *Journal of Aquatic Ecology*, 8(4), 76-86. (In Persian)
- Stengel, D.B. and Dring, M.J., 1998.** Seasonal variation in the pigment content and photosynthesis of different thallus regions of *Ascophyllum nodosum* (Fucales, Phaeophyta) in relation to position in the canopy. *Phycologia*, 37

- (4), 259-268. DOI:10.2216/i0031-8884-37-4-259.1
- Taheri, A., Ghaffari, M., Bavi, Z. and Sohili, F., 2018.** Cytotoxic effect of the extract of seaweed *Sargassum glaucescens* against breast (MCF-7) and colorectal (HT-29) cancer cell lines. *Feyz Medical Sciences Journal*, 22(3), 292-301. (In Persian)
- Wang, P., Clemens, S.C., Tada, R. and Murray, R.W., 2019.** Blowing in the monsoon wind. *Oceanography*, 32(1), 48-59. DOI:10.5670/oceanog.2019.119.
- Wang, B., Biasutti, M., Byrne, M.P., Castro, C., Chang, C.P., Cook, K., Fu, R., Grimm, A.M., Ha, K., Hendon, H., Kitoh, A., Krishna, R., Lee, J., Li, J., Liu, J., Moise, A., Pascal, S., Roxy, M.K., Seth, A., Sui, C., Turner, A., Yang, S., Yun, K., Zhang, L. and Zhou, T., 2021.** Monsoons climate change assessment. *Bulletin of the American Meteorological Society*, 102(1), E1-E19. DOI:10.1175/BAMS-D-19-0335.1
- Yende, S.R., Arora, S.K. and Ittadwar, A.M., 2021.** Antioxidant and Cognitive Enhancing Activities of *Sargassum ilicifolium* and *Padina tetrastromatica* in Scopolamine Treated Mice. *Journal of Biologically Active Products from Nature*, 11(1), 11-21. DOI:10.1080/22311866.2020.1871071