

## Microsatellite DNA diversity of *Coilia mystus* (Clupeiformes:Engraulidae) in three Chinese estuaries

Fang Chen<sup>1, 2</sup>; Qiqun Cheng<sup>1,\*</sup>

Received: July 2011

Accepted: November 2011

### Abstract

An analysis of eight microsatellite loci in 90 individuals was performed to define the genetic structure and variability of three estuarine populations of *Coilia mystus*: 30 individuals each collected from ChangJiang River (Yangtze River) estuary-CJ, MinJiang River estuary-MJ, and ZhuJiang River (Pearl River) estuary-ZJ. The results showed polymorphic information contents (*PIC*) of 0.78, 0.77, and 0.64 in CJ, MJ, and ZJ, respectively. The mean observed number of alleles ( $N_a$ ) ranged from 7.38 to 11.88. The mean observed and expected heterozygosities ranged from 0.09 to 0.21 and from 0.68 to 0.81, respectively. The ZJ population was least polymorphic. Highly significant deviations from Hardy-Weinberg equilibrium, mostly due to deficits of heterozygote, were found in these three populations. Pairwise  $F_{ST}$  and Reynolds' distance indicated that the three estuarine populations were genetically distinct, in accordance with the principal component analysis (PCA) and the Bayesian model-based clustering algorithm analysis. The results showed that pairwise genetic differentiation among these three estuarine populations were relatively high, with possible divergence to subspecies level. This study provides microsatellite DNA evidence for assessing the genetic distinctness of *C. mystus* populations and will benefit fishery resource management and sustainable utilization of *C. mystus*.

**Keywords:** *Coilia mystus*, Microsatellite, Genetic diversity, Population structure, Genetic divergence

1- Key Laboratory of East China Sea and Oceanic Fishery Resources Exploitation, Ministry of Agriculture, East China Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences, Shanghai 200090, China.

2- College of Marine Sciences, Shanghai Ocean University, Shanghai 201306, China.

\*Corresponding author's email: qiquncheng@gmail.com

## Introduction

*Coilia mystus* (Linnaeus 1758) is a small to moderate-sized anchovy in the genus *Coilia*, family Engraulidae and order Clupeiformes. It is distributed widely in Chinese estuaries, and is one of the most important species to China's estuarine fishery economy. *C. mystus* is a short-distance anadromous fish, which migrates from waters near the ocean to brackish water areas every year during the spawning season. Mature *C. mystus* migrates upriver, spawns in the brackish waters, and never enters pure freshwater areas (East China Sea Fisheries Research Institute, Chinese Academy of Fishery Sciences and Shanghai Fisheries Research Institute, 1990). The ChangJiang River (Yangtze River), MinJiang River, and ZhuJiang River (Pearl River) are three important rivers, which are separated from each other by long distances. Therefore, local adaptation, overfishing, and environmental conditions may contribute to genetic divergence of the three populations.

Previously, we have studied the divergence of these three populations by using morphological and mitochondrial indices. We measured morphological differences between these three populations of *C. mystus*, which were difficult to differentiate using the meristic index, and found great morphological differences between them which can be discriminated with synthetical indices. The Minjiang population was not closely clustered with either the Changjiang or the Zhujiang

populations (Cheng et al., 2005; Cheng, 2010). We also compared the partial cytochrome *b* and 16S rRNA gene sequences of these three populations, and hypothesized that they were diverged at the subspecies level (Cheng et al., 2006, 2008). The significant morphological differences between populations of *C. mystus* were in concordance with genetic divergence of mitochondrial DNA data.

Microsatellites markers, which are neutral genetic markers, abundant in fish genomes, with high levels of polymorphism (Tautz, 1989), are simple tandemly repeated sequence motifs consisting of repeat units of 1-6 bp. Microsatellite markers are very useful for molecular ecology and population genetic studies of fishery resources. They have been widely used in many species for evaluating population genetic diversity, constructing genetic maps, and determining pedigrees (McDonald et al., 2004; Guyomar et al., 2006; Pettay and Lajeunesse, 2009). We have developed several microsatellite markers in *C. ectenes*, a closely related species of *C. mystus*, and found these markers are suitable for cross-species amplification in *C. mystus* (Ma et al., 2011).

The research of genetic diversity of species can help us understand the genetic structure of species and population history and then comprehend their evolution for a conservation strategy. In the present study, we applied eight microsatellite primer pairs to study genetic diversity and variability of

three estuarine populations of *C. mystus*. The present study provides microsatellite DNA evidence for assessing the genetic distinctness of *C. mystus* populations and will benefit fishery resource management and sustainable utilization of *C. mystus*.

## Materials and methods

### *Sample collection and DNA extraction*

A total of 90 *C. mystus* were obtained from Changjiang (CJ), Minjiang (MJ), and Zhujiang (ZJ) river estuaries, with 30 individuals of each (Table 1 and Fig. 1). All individuals were transferred to the laboratory in dry ice and then stored at -80°C before DNA extraction. Genomic DNA was extracted from muscle tissues. Muscle tissues were dissected and then digested by proteinase K, followed by phenol-chloroform extraction and 100% ethanol precipitation (Sambrook et al., 1989).

### *Microsatellite analysis*

We genotyped all 90 individuals at 8 microsatellite loci (D13, D55, D58, D63, D81, D95, D104, D114) following our established methods (Ma et al., 2011). PCR reactions were carried out in a 25 µL volume comprising 100 ng of genomic DNA, 10×PCR buffer (Mg<sup>2+</sup>), 10 uM of each primer, 2.5 mM of dNTP mix, and 2.5 U of *Taq* polymerase (Takara). The thermocycling conditions were as follows: initial denaturation at 94°C for 5 min, followed by 35 cycles of 94°C for 30 s,

annealing temperature (see Table 1 in Ma et al., 2011) for 30 s, and 72°C for 1 min, and then a final extension of 72°C for 7 min. PCR products were separated in 3% agarose gels. The range of allele sizes was determined by referring to the *MseI* marker (TianGen Biotech) and molecular tools in Smartview Analysis Software (FuRi Tech).

### *Statistical analysis*

For each marker and population, POPGENE 1.31 software (Yeh et al., 1999) was used to estimate allele frequencies ( $P$ ), observed number of alleles ( $N_a$ ), effective number of alleles ( $N_e$ ), observed heterozygosity ( $H_o$ ), expected heterozygosity ( $H_e$ ). Polymorphism information content ( $PIC$ ) was computed according to the following formula (Botstein et al., 1980):

$$PIC = 1 - \sum_{i=1}^n P_i^2 - \sum_{i=1}^{n-1} \sum_{j=i+1}^n 2P_i^2 P_j^2, \quad \text{where } P_i \text{ and } P_j \text{ are the frequencies of the } i\text{th and } j\text{th alleles at one locus, and } n \text{ is the number of alleles at one locus.}$$

GENEPOP 1.2 software (Raymond and Rousset, 1995) was used to estimate departures from Hardy-Weinberg equilibrium at each locus, which implements Fisher's exact tests for multiple alleles (Guo and Thompson, 1992). The null allele frequency was estimated by MICRO-CHECKER version 2.2.3 (Van Oosterhout et al., 2004).

Genetic differences between populations were evaluated by calculating pairwise  $F_{ST}$  values and Reynold's genetic

Genetic differences between populations were evaluated by calculating pairwise  $F_{ST}$  values and Reynold's genetic

distance using ARLEQUIN 3.0 (Excoffier et al., 2005). As an alternative approach to represent the genetic relationship among the *C. mystus* populations, a Principal Component Analysis (PCA) was also performed based on gene frequencies of all the variable loci, using the GENETIX software package (Belkhir et al., 2000). A Bayesian model-based clustering algorithm implemented in the computer program STRUCTURE 2.0 (Pritchard et al., 2000) was also applied. This model-based algorithm places the given individuals into  $K$  clusters, where  $K$  is chosen in advance but can be varied across independent runs of the algorithm. All runs were carried out under “admixture model”, with a burn-in period of 10,000 iterations and a period of data collection of 10,000 iterations.

## Results

### *Within population variation*

All the eight microsatellite markers were successfully amplified in all 90 individuals of these three populations and were found to have high polymorphic levels. Alleles ranged in size from 100 to 310 bp over the eight loci. In these three populations, the observed number of alleles across loci in the populations ranged from 2 alleles in ZJ to 20 alleles in CJ. The mean observed number of alleles ( $N_a$ ) ranged from 7.38 to 11.88 across populations. The mean effective alleles ( $N_e$ ) per population ranged from 3.97 to 7.68. Polymorphism information contents (PIC) were between 0.22 and 0.92, and the

averages were 0.78, 0.77, and 0.64 in CJ, MJ, and ZJ, respectively. The mean observed and expected heterozygosities for the three populations ranged from 0.09 to 0.21 and from 0.68 to 0.81, respectively (Table 1). The average observed heterozygosity was lower than the expected heterozygosity at all the loci. These parameters of genetic diversity indicated that there were moderate-to-high levels of polymorphism within these three populations. An exact  $P$ -value test indicated that all loci significantly deviated from HWE and was associated with heterozygote deficiencies (Table 1). Based on average  $F_{IS}$  values, it can be seen that the pattern of heterozygote deficiencies was pronounced in all populations.

### *Genetic differentiation and relationships between populations*

The pairwise  $F_{ST}$  and Reynolds' distance values for the three populations are shown in Table 2. The three populations were well differentiated from each other ( $F_{ST} = 0.116$ -0.161). MJ and ZJ was the most divergent ( $F_{ST} = 0.161$ ) followed by CJ and ZJ ( $F_{ST} = 0.150$ ). The divergence between the CJ and MJ was lowest ( $F_{ST} = 0.116$ ). The pairwise  $F_{ST}$  comparisons revealed a relatively high degree of genetic differentiation between these three populations. The Reynolds' distances ranged from 0.156 to 0.231. The genetic distance tended to be the least (0.156) between CJ and MJ and the widest (0.231) between CJ

and ZJ. PCA based on the correlation matrix obtained from the allele frequencies was also performed (Fig. 2). The first axis explained 29.5% of the inertia and distinguished ZJ from the rest of the populations. The second axis contributed 17.7% of the inertia and separated CJ from MJ. The result indicated that these three populations could be separated clearly by

these eight microsatellite markers. The STRUCTURE 2.0 software was used to evaluate the assignments of individuals to 1-3 assumed populations with three simulations (Fig. 3). The three populations were separated at  $K = 3$ , which represented the membership of each individual to the three clusters. It matches the result of the Principal Component Analysis.




Figure 1: Locations of sampling sites (numbers). 1. ChangJiang River (Yangtze River) estuary, Shanghai; 2 .MinJiang River estuary, Fuzhou; 3. ZhuJiang River (Pearl River) estuary, Guangzhou




Figure 2: Scattergram showing relative positions of three *Coilia mystus* populations defined by PC1 vs PC2 of eight microsatellite loci




Figure 3: Clustering assignment of the three populations provided by  
STRUCTURE analyses

Table 1: Estimation of genetic variability of three *Coilia mystus* populations at 8 microsatellite loci

Pop\Locus		D13	D55	D58	D63	D81	D95	D104	D114	Mean
CJ	$N_a$	15	16	10	14	12	4	20	3	11.75
	$N_e$	10.58	8.85	4.84	8.84	4.59	2.94	10.45	2.65	6.72
	$H_o$	0.10*	0.07*	0.00*	0.26*	0.93*	0.00*	0.32*	0.00*	0.21
	$H_e$	0.91	0.89	0.79	0.89	0.78	0.66	0.9	0.62	0.81
	$PIC$	0.9	0.88	0.78	0.88	0.76	0.59	0.9	0.55	0.78
MJ	$N_a$	18	17	12	12	6	3	17	10	11.88
	$N_e$	13.24	11.2	9.89	7.5	3.91	1.77	11.11	2.82	7.68
	$H_o$	0.24*	0.25*	0.00*	0.11*	0.00*	0.00*	0.13*	0.00*	0.09
	$H_e$	0.92	0.91	0.9	0.87	0.74	0.44	0.91	0.64	0.79
	$PIC$	0.92	0.9	0.89	0.85	0.7	0.37	0.9	0.63	0.77
ZJ	$N_a$	9	10	9	8	5	2	9	7	7.38
	$N_e$	6.48	6.28	3.58	2.41	2.78	1.34	4.89	4.04	3.97
	$H_o$	0.22*	0.24*	0.14*	0.21*	0.00*	0.00*	0.13*	0.00*	0.12
	$H_e$	0.85	0.84	0.72	0.59	0.64	0.25	0.8	0.75	0.68
	$PIC$	0.83	0.82	0.67	0.55	0.58	0.22	0.77	0.72	0.64
$F_{IS}$		0.79	0.79	0.94	0.75	0.57	1.00	0.77	1.00	0.83

\*: Significant deviation from Hardy–Weinberg equilibrium

Table 2: Genetic distances (Reynold's distance, above diagonal) and differentiation ( $F_{ST}$ , below diagonal)  
between *Coilia mystus* populations

	CJ	MJ	ZJ
CJ	—	0.156	0.231
MJ	0.116	—	0.217
ZJ	0.150	0.161	—

## Discussion

### *Genetic diversity within populations*

Biological population diversity is an important foundation for evaluating species resources. It is true that the species to adapt to different environments. It is a precondition to use the species resource persistently, to keep the highest level of genetic variation.  $N_a$ ,  $H_o$ ,  $H_e$  and  $PIC$  are all parameters for measuring population genetic variations.

On the basis of the present study, analyses of eight microsatellite loci revealed relatively high levels of genetic variation. The mean effective alleles and mean  $PIC$  per population was from 3.97 to 7.68 and from 0.64 to 0.78, respectively. Bostein et al. (1980) indicated that a locus was considered highly polymorphic when  $PIC \geq 0.50$ , low polymorphic when  $PIC \leq 0.25$ , and moderate polymorphic when the  $PIC$  value was between 0.25 and 0.50. The  $PIC$  here ranged from 0.64 to 0.78, which means there is a high level of polymorphism in these three estuarine populations and it is suitable for estimating genetic diversity of *C. mystus* populations. The observed allele numbers and allele ranges here were relatively higher than that observed in a previous study of *C. mystus* based on 12 microsatellite loci (Ma et al., 2011). The mean observed and expected heterozygosities were from 0.09 to 0.21 and from 0.68 to 0.81, respectively. It can be speculated that these discrepancies may be due to the sharp decline in number of *C.*

*mystus* in the recent years, which may result in low observed genetic heterozygosities in populations. This may indicate loss of genetic diversity and the simpler genetic structure in the three estuarine populations, which would reduce the environmental adaptability.

Cheng et al. (2008) studied the partial mitochondrial cytochrome *b* and 16S rRNA gene sequences of these three populations, and indicated that genetic diversity was lowest within the ZJ population, which is in concordance with the present microsatellite analysis. Human activity may be influencing genetic diversity within the three populations. Guangzhou is highly developed economically with human activities, which may be lowering the genetic diversity of *C. mystus* populations through overharvesting.

This study showed that all loci significantly deviated from the Hardy-Weinberg equilibrium. Biased samples and small sample size may lead to this phenomenon. However, these two reasons can be excluded in this study. *C. mystus* is one of the main economic fish in the Changjiang, Minjiang, and Zhujiang river estuaries and the samples were collected randomly. Barker et al. (1993) proposed that at least 25 individuals should be analyzed for each species/population for an acceptable statistical resolution. In this study, 30 individuals were analysed for each population, and they were therefore representative of the population genetic

information. Generally, heterozygote deficit may be linked to multiple causes: population substructure, inbreeding, null alleles, the Wahlund effect, and natural selection (Ardren et al., 1999; Lade et al., 1996; Castric et al., 2002; Morand et al., 2002). *C. mystus* is inhabited in Chinese estuary and no obvious geographical barriers separate migration and gene flow between individuals, so population substructure may not exist. Fixed factor  $F_{IS}$  is an important parameter to evaluate the occurrence of inbreeding in populations.  $F_{IS}$  values for each population are relatively high, suggesting that inbreeding exists in each population. Over-fishing and environmental pollution could provide a reasonable explanation for inbreeding within population. Null alleles within population may be another factor to account for the increased homozygosity (Pemberton et al., 1995) as has been observed in related species due to the sequence mutation of microsatellite flanking (Harper et al., 2003). However, in all samples, though each locus deviated from the Hardy-Weinberg equilibrium, at least one allele can be stably amplified, so the frequency of null alleles was not high enough to affect the results of this study significantly (Goodman et al., 2001; Abbott and Double, 2003; Harper et al., 2003; Lucchini et al., 2004). The Wahlund effect may also be excluded, because no significant hidden substructure was detected in the *C. mystus* population. Natural selection, which usually acts on

long-term evolutionary processes, was also considered as a possible cause of heterozygote deficiency. The different environmental factors such as water temperature, water current, salinity, food availability, and fishing intensity in these three rivers may influence populations throughout the sampling period.

#### *Relationships and genetic differentiations between populations*

The values of  $F_{ST}$  and Reynolds' distance suggested that the three populations have evolved independently and have developed significant genetic structure with possible divergence to subspecies level. This result was in concordance with morphological differences (Cheng, 2010) and genetic divergence of mitochondrial DNA data (Cheng et al., 2006, 2008).

According to this research, both  $F_{ST}$  and Reynolds' distance values showed that MJ and CJ were closely related while CJ and ZJ populations were farthest. The molecular results matched the results of the previous morphological study. Cheng et al. (2005) also found that there were great morphological differences between the Changjiang and the Zhujiang populations. The most likely reason for this finding could be due to geographic proximity. The long distance separating Guangzhou and Shanghai (1,689 km), coupled with the short-distance migrating habit of *C. mystus*, affords little chance for gene flow between the ZJ and CJ populations.



PCA and Structure analyses clearly revealed the genetic structuring of the investigated population into three isolated clusters with restricted gene flow among them. The potential capacity of populations to adapt and evolve as independent biological entities in different environmental conditions is restricted by the exchange of individuals between populations. A sufficient degree of isolation may result in notable phenotypic and genetic differentiation among fish populations within species, which may be recognizable as a basis for separation and management of distinct populations (Turan, 2004). Geographic isolation, living environment, population genetic bottleneck, gene flow, and selection have large effects on the genetic construction of populations. Heredity and variation are necessary to adapt to the changing environment. Population genetic research on a wide distributing species can elucidate population genetic variability and structuring.

In summary, the three estuarine populations of *C. mystus* maintain a moderate-to-high level of genetic diversity within populations and higher genetic differentiation among populations. The status of genetic diversity in these three estuary populations may affect the management and utilization of this species. Some necessary measures should be implemented to decrease harmful human activities, such as overfishing, environmental pollution, and irrigation

projects. Only when the pressures on these populations are reduced, can the genetic diversity of *C. mystus* in these waters be maintained and restored.

### Acknowledgements

The author would like to thank Mrs. Jindi Han and Mr. Qizhe Liu for assistance in sample collection. This work was supported by the Special Research Fund for national non-profit institutes (East China Sea Fisheries Research Institute, Project No.2008Z02) and the National Natural Science Foundation of China (No.30600456).

### References

- Abbott, C. L. and Double, M. C., 2003.** Genetic structure, conservation genetics and evidence of speciation by range expansion in shy and white, capped albatrosses. *Molecular Ecology*, 12, 2953-2962.
- Ardren, W.R., Borer, S., Thrower, F., Joyce, J. E. and Kapuscinski, A. R., 1999.** Inheritance of 12 microsatellite loci in *Oncorhynchus mykiss*. *Journal of Heredity*, 90, 529-536.
- Barker, J. S. F., Bradley, D. G., Fries, R., Hill, W. G., Nei, M. and Wayne, R. K., 1993.** An Integrated Global Programme to Establish the Genetic Relationships Among the Breeds of Each Domestic Animals Species. Food and Agriculture Organization of the United Nations, Rome, 24 p.

- Belkhir, K., Borsa, P., Chikhi, L., Raufaste, N. and Bonhomme, F., 2000.** GENETIX 4.01, logiciel sous Windows™ pour la géne'tique des populations. Laboratoire Ge'nome, Populations, CNRS UPR 9060.Interactions, Universite' de Montpellier II, Montpellier, France.
- Botstein, D., White, R. L., Skolnick, M. and Davis, R. W., 1980.** Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *American Journal of Human Genetics*, 32, 314-331.
- Castric, V., Bernatchez, L., Belkhir, K. and Bonhomme, F., 2002.** Heterozygote deficiencies in small lacustrine populations of brook charr *Salvelinus fontinalis* Mitchill (Pisces, Salmonidae): a test of alternative hypothesis. *Heredity*, 89, 27-35.
- Cheng, Q. Q., Lu, D. R. and Ma, L., 2005.** Morphological differences between close populations discernible by multivariate analysis: a case study of genus *Coilia* (Teleostei: Clupeiformes). *Aquatic Living Resources*, 18(2), 187-192.
- Cheng, Q. Q., Han, J. D., Wang, Y. L. and Lu, D. R., 2006.** Genetic diversity and evolutionary characters in two populations of *Coilia mystus* inferred from cytochrome *b* gene segment sequence of mitochondrial DNA. *Journal of Fishery Sciences of China*, 13(3), 337-343.
- Cheng, Q. Q., Ma, C. Y., Cheng, H. P. and Zhang, Q. Y., 2008.** Mitochondrial DNA diversity of *Coilia mystus* (Clupeiformes: Engraulidae) in three Chinese estuaries. *Environmental Biology of Fishes*, 83(3), 277-282.
- Cheng, Q. Q., 2010.** Morphological variation of *Coilia mystus* (Clupeiformes:Engraulidae) in three Chinese estuaries. *Journal of Life Sciences*, 4(6), 29-34.
- East China Sea Fisheries Research Institute, Chinese Academy of Fisheries Science and Shanghai Fisheries Research Institute, 1990.** The fishes of Shanghai Area. Shanghai Scientific and Technical Publishers, Shanghai.
- Excoffier, L., Laval, G. and Schneider, S., 2005.** ARLEQUIN. Ver 3.0. An integrated software package for population genetic data analysis. *Evolutionary Bioinformatics Online*, 1, 47-50.
- Goodman, S. J., Tamate, H. B., Wilson, R., Nagata, J., Tatsuzawa, S., Swanson, G. S., Pemberton, J. M. and McCullough, D. R., 2001.** Bottlenecks, drift and differentiation : the population structure and demographic history of sika deer ( *Cervus nippon* ) in the Japanese archipelago. *Molecular Ecology*, 11, 1357-137.
- Guo, S. and Thompson, E., 1992.** Performing the exact test of Hardy-Weinberg proportion for

- multiple alleles. *Biometrics*, 48, 361-372.
- Guyomar, R. S., Mauger, K., Tabet-Canale, S., Martieau, C., Genet, F. K. and Quillet, E., 2006.** A type I and II microsatellite linkage map of rainbow trout (*Oncorhynchus mykiss*) with presumptive coverage of all chromosome arms. *BMC Genomics*, 7, 302-314.
- Harper, G. L., Maclean, N. and Goulson, D., 2003.** Microsatellite markers to assess the influence of population size, isolation and demographic change on the genetic structure of the UK butterfly *Polyommatus bellargus*. *Molecular Ecology*, 12, 3349-335.
- Lade, J. A., Murray, N. D., Marks, C. A. and Robinson, N. A., 1996.** Microsatellite differentiation between Philip Island and mainland Australian populations of the red fox (*Vulpes vulpes*). *Molecular Ecology*, 5, 81-87.
- Lucchini, V., Galov, A. and Randi, E., 2004.** Evidence of genetic distinction and long-term population decline in wolves (*Canis lupus*) in the Italian Apennines. *Molecular Ecology*, 13, 523-536.
- Ma, C. Y., Cheng, Q. Q. and Zhang, Q. Y., 2011.** Development of 12 polymorphic microsatellite markers in *Coilia ectenes* Jordan and Seale, 1905 (Clupeiformes: Engraulidae) and cross-species amplification in *Coilia mystus* Linnaeus, 1758. *Environmental Biology of Fishes*, 91, 243-249.
- McDonald, G. J., Danzmann, R. G. and Ferguson, M. M., 2004.** Relatedness determination in the absence of pedigree information in three cultured strains of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 233, 65-78.
- Morand, M. E., Brachet, S., Rossignol, P., Dufour, J. and Frascaria-Lacoste, N., 2002.** A generalized heterozygote deficiency assessed with microsatellites in French common ash populations. *Molecular Ecology*, 11, 377-385.
- Pemberton, J. M., Slate, J., Bancroft, D. R. and Barrett, J. A., 1995.** Nonamplifying alleles at microsatellite loci: a caution for parentage and population studies. *Molecular Ecology*, 4, 249-252.
- Pettay, D. T. and Lajeunesse, T. C., 2009.** Microsatellite loci for assessing genetic diversity, dispersal and clonality of coral symbionts in 'stress-tolerant' clade D Symbiodinium. *Molecular Ecology Resources*, 9, 1022-1025.
- Pritchard, J. K., Stephens, M. and Donnelly, P., 2000.** Inference of population structure using multilocus genotype data. *Genetics*, 155, 945-959.
- Raymond, M. and Rousset, F., 1995.** GENEPOP (Version 1.2): population genetics software for exact tests and ecumenicism. *Journal of Heredity*, 86, 248-249.
- Sambrook, J., Fritsch, E. F. and Maniatis, T., 1989.** Molecular cloning: a

- laboratory manual, 2nd edn. Cold Spring Harbor Laboratory Press, New York.
- Tautz, D., 1989.** Hypervariability of simple sequences as a general source for polymorphic DNA markers. *Nucleic Acids Research*, 17(16), 6463-6471.
- Turan, C., 2004.** Stock identification of Mediterranean horse mackerel (*Trachurus mediterraneus*) using morphometric and meristic characters. *ICES Journal of Marine Sciences*, 61, 774-781.
- Van Oosterhout, C., Hutchinson, W. F., Wills, D. P. M. and Shipley, P., 2004.** MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, 4, 535-538.
- Yeh, F. C., Yang, R. C. and Boyle, T., 1999.** POPGENE version 1.31. Microsoft window-bases freeware for population genetic analysis. University of Alberta and the Centre for International Forestry Research.  
<http://www.ualberta.ca/~fyeh/>.