2023

### Research Article Culture strategy for production of Indian white prawn, *Fenneropenaeus indicus* in semi-arid conditions using biofloc technology

Sambhu C.<sup>1\*</sup>; Al Harbi M.<sup>1</sup>; Broom M.<sup>2</sup>; Nasser A.K.V.<sup>2</sup>; Sherly D.<sup>3</sup>

Received: March 2018

Accepted: March 2019

#### Abstract

An experiment was conducted to test efficiency of biofloc technique for production of Indian white prawn, *Fenneropenaeus indicus* in HDPE liner ponds (300 m<sup>2</sup>) for 90 days. There were control (water exchange to maintain transparency at 40-50 cm) and treatment (zero water exchange) ponds and both were triplicated. Soya hull and molasses were added to treatment ponds as carbon sources to induce biofloc formation. Post larvae (PL<sub>20</sub>) were stocked at the rate of  $50/m^2$  and fed with a standard fishmeal based supplementary pellet feed. Physico-chemical parameters of water, microbial and plankton population, immune response, physical quality and shrimp growth were monitored during the period. High growth and survival was observed in treatment ponds compared to control. Heterotrophic bacteria, phytoplankton population and total haemocyte count (THC) were found to be enhanced in treatment ponds. A strong linear relationship (R<sup>2</sup>=0.8758) was found between growth rate and biofloc content. Shrimp raised through biofloc culture strategy showed better colour and quality compared to control. Biofloc technology is an ideal culture method for biosecure production of white shrimp in semi arid lands.

**Key words**: *Fenneropenaeus indicus*, Biofloc technology, Water quality, Plankton, Immune response, Growth

<sup>1-</sup> Department of Marine Biology, Faculty of Marine Sciences, King Abdulaziz University, Jeddah 21589, Saudi Arabia

<sup>2-</sup>National Aquaculture Group, Al Lith 21961, Saudi Arabia

<sup>3-</sup>Department of Zoology, All Saints' College, University of Kerala, Kerala, India

<sup>\*</sup>Corresponding author's Email: sambhu@kau.edu.sa

#### Introduction

**Biofloc** systems were developed primarily to improve environmental over production and control for enhanced biosecurity. Biofloc systems need limited water exchange and prevent introduction of disease to a farm from incoming water (Ogello et al., 2014). It uses a counter-intuitive approach which allows or encourages associated solids and microbial community to accumulate in water. Biofloc technology was developed by Avnimelech (2000, 2005 and 2012) and initially implemented commercially in Belize by Belize Aquaculture (McIntosh, 2000). It also has been applied with success in shrimp farming in Indonesia, Malaysia and Australia (Nyan Taw, 2010). Bioflocs are aggregates (flocs) of algae, bacteria, protozoans, and other kinds of particulate organic matter such as faeces and uneaten feed (Baloi et al., 2013). Each floc is held together in a loose matrix of mucus that is secreted by bacteria, bound by filamentous microorganisms, or held by electrostatic attraction (Supono et al., 2014). The biofloc community also includes animals that are grazers of flocs, such as some zooplankton and nematodes (Nyan Taw, 2014). Large bioflocs can be seen with naked eye, but most are microscopic (Avnimelech, 2012). Flocs in a typical green water biofloc system are rather large, around 50 to 200 microns, and will settle easily in calm water (Hargreaves, 2006).

Other than less water exchange, advantage with biofloc system is

capacity to recycle waste nutrients through microbial protein into fish or shrimp. About 20 to 30% of nitrogen in feed is assimilated by fish, implying that 70 to 80 percent of nitrogen (85% according to Crab et al. (2012) is released to culture environment. In biofloc systems some of this nitrogen is incorporated into bacterial cells that are a main component of biofloc. Biofloc are nutritious with dry-weight protein content of 30 to 45%, fat content of 1 to 5% and good sources of vitamins and minerals, especially phosphorus (De Schryver et al., 2008). Bioflocs may also have probiotic effects (Rivera et 2014). Research with shrimp al., suggests that for every unit of growth derived from feed, an additional 0.25 to 0.50 units of growth are derived from microbial protein in biofloc systems (Crab et al., 2012). This benefit is reflected in improved feed conversion. Considering improved biosecurity, high production capacity, low FCR and cost of production per kg, a study was conducted to evaluate the effect of biofloc culture strategy for production of Indian white prawn, F.indicus an ideal candidate species for coastal aquaculture practice.

#### Materials and methods

### Experimental set up and study management

The study was conducted in HDPE liner ponds (300m<sup>2</sup>) for a period of 90 days at University Fish Farm at Obhur, Jeddah. Culture ponds were limed and sundried for one week prior to water culture. In order to develop algal bloom

(40-50cm Sechi disc transparency), water culture was done by the modified methods of Boyd (1990). On day 1, culture ponds were filled with seawater (30%) and manured by applying Urea (400 g), Molasses (1.5 liter) and Diammonium phosphate (200g) and the dose repeated on 4<sup>th</sup> and 8<sup>th</sup> day of culture. Pond water level increased to 60 and 100% before applying second and third dose. Two unit of aspirator (1 hp) (Force-7, Acquaeco, aerator Italy) were installed at 40 cm below water level with 35 cm angle downward in each pond. Ponds which were maintained transparency at 40-50 cm by water exchange at10cm/day were considered as control. Whereas the ponds in which zero water exchange was done (topping up was done to maintain the loss of water due to evaporation) were designated as treatment and both of them were triplicated. To boost up heterotrophic bacterial growth in treatment ponds, soya hull and molasses (2 kg) were applied as carbon sources once in three days. On 12<sup>th</sup> day of water culture, healthy and uniform size juvenile (1.84±0.2g: 5.7±1.8cm) produced at Farm Hatchery were stocked at the rate of 50 pieces  $/m^2$  in each pond (hapa survival >95%). A standard fish meal based pellet feed having 35% protein (NAQUA, Jeddah) was supplemented to shrimp based on a standard feed table at 7:00am, 1:00 and 6:00pm daily.

Water quality control, heterotrophic bacteria, plankton and immune test

Water quality parameters such as temperature, dissolved oxygen, pH, salinity (YSI Incorporated, Yellow Springs, Ohio, USA) and biofloc were recorded daily. Ammonia (unionized), nitrates  $(NO_3)$ . nitrites  $(NO_2)$ . orthophosphates  $(PO_4)$  and alkalinity (as CaCO<sub>3</sub>) were recorded (JBL Test kit, GmbH & Co., Germany) weekly. volume (mL/liter) was The floc determined by using Imhoff cones, which recorded the suspended organic solids for 20 minutes of settling (Taw, 2014). Heterotrophic bacterial population (Total plate count for colony forming units (CFU/mL) and plankton community (phyto and zooplankton) in the pond water were examined every month (APHA, 1995; Smith and Johnson, 1996). Immune response of shrimp was tested a week prior to harvest. Shrimp from control and treatment ponds were brought to laboratory and subjected to cold challenge test at 20°C for 24 hours. After the cold challenge, haemolymph sample was drawn from the heart of shrimp and subjected to haemocyte count using a compound microscope (Krupesha et al., 2009).

## Sampling, Growth analysis and physical quality test

Shrimp sampling (200pcs/sampling) was done biweekly to assess the growth in weight and the feed quantity was re adjusted after every sampling. Upon harvest, all shrimp from control and

ponds collected. treatment were survival and biomass were recorded. A panel of experts evaluated physical quality of shrimp such as colour, loose shell, soft shell and taste. Specific growth rate was calculated as Log<sub>e</sub> W2-Log<sub>e</sub>W1/T2-T1 (where W2 is the weight of shrimp at time T2 and W1 is the weight of shrimp at time T1). Oneway analysis of variance (ANOVA) was employed to find out the statistical difference between growth. water quality parameters, biofloc content, bacterial population and haemocyte count of control and treatment. Linear regression analysis was done to find out relationship between biofloc content and growth of shrimp.

#### Results

#### *Growth performance*

Growth performance of control and treatment shrimp recorded during the culture period is shown in Figure 1 and production details are presented in Table 1. Shrimp grown in treatment ponds showed better growth in weight when compared to control and the observed difference was found to be non significant (*p*>0.01). Survival. average weekly growth, specific growth rate and biomass were high in treatment ponds; whereas, feed conversion rate was found to be low in treatment ponds. Significant relationship  $(R^2=0.8758)$ observed between biofloc was concentration and average body weight (g) in treatment ponds (Fig. 2).



Figure 1: Growth of F.indicus during culture period.

# Water quality parameters and biofloc content

Water quality parameters recorded during the culture period were found to be conducive for shrimp growth (Table 2). Significant difference was not observed in water temperature, dissolved oxygen, pH, salinity, nitrite, phosphate and ammonia between control and treatment ponds whereas, alkalinity and nitrate contents showed significant difference between control and treatment ponds.

biomass of shrimp					
	Control	Treatment			
Parameters	Mean ±	Mean ±			
	SD	SD			
Initial weight (g)	1.84 + 0.2	1.84±0.2			
Final weight (g) <sup>NS</sup>	$14.8 \pm 2.3^{a}$	$15.6 \pm 3.3^{a}$			
Net weight gain (g)	$13.0{\pm}1.2$	13.8±1.9			
Average weekly	1.0±0.1	1.2±0.1			
growth (g)					
Specific Growth	$1.03 \pm 0.01$	$1.06\pm0.02$			
Rate (%)					
Survival (%)	$77\pm8$	83±12			
Feed Conversion	2 6+0 10	2 3+0 20			
Rate	2.0±0.10	2.5±0.20			
Biomass/pond (kg)	171±12	194±23			
Biomass (kg)/ha	$5700 \pm 121^{a}$	6466±309 <sup>b</sup>			
NC 0.01, $(100)$ , $** <0.01$					

Table 1: Growth perfromance, survival and

NS-*p*>0.01; (*n* =100); \*\* *p*<0.01

a, b. Means with the same superscript do not differ from each other.

#### Biofloc and Plankton community

Biofloc concentration during the culture period is depicted in Figure 3. An increase in biofloc concentration was noticed in treatment ponds compared to attained control and it a high concentration during the last week of culture. Plankton community in the ponds is presented in Figure 4. Phytoplankton community was found to be decreased in both control and treatment ponds in increasing culture days. However, it was high in treatment ponds during every sampling.



Figure 2: Linear relationship between Biofloc and ABW in treatment pond.

Table 2: Water quality parameters during the culture period

the culture period				
Paramatars	Control	Treatment		
1 ai aincui s	Mean ±SD	Mean ± SD		
Temperature (°C) <sup>NS</sup>	$27.22 \pm 2.72$	27.30±2.76		
Dissolved Oxygen $(mg L^{-1})^{NS}$	5.90±0.51	6.03±0.47		
pH <sup>NS</sup>	$8.03 \pm 0.10$	8.22±0.11		
Salinity (g L <sup>-1</sup> ) <sup>NS</sup>	$39.20{\pm}1.05$	41.20±0.52		
Alkalinity $((mg L^{-1})^*)$	136.60±7.3	$144.00 \pm 8.0$		
Nitrate (mg $L^{-1}$ ) <sup>*</sup>	$1.68 \pm 0.66$	$1.48 \pm 0.40$		
Nitrite (mg L <sup>-1</sup> ) <sup>NS</sup>	$0.02 \pm 0.01$	$0.03 \pm 0.01$		
Orthophosphate (mg $L^{-1}$ ) <sup>*</sup>	$0.66 \pm 0.11$	$1.50\pm0.12$		
Unionized Ammonia (mg $L^{-1}$ ) <sup>NS</sup>	$0.08 \pm 0.01$	0.13±0.07		

Species like Tetraselmis, Peridinium, Trichodesmium, Ntzchia, Malassiosira, Cylotella, Ceratium, Amphora, Navicula, Psuedonizchium and Blue green algae were the dominant groups in the community. Zooplankton community was found to be high in treatment and low in control ponds. Ciliates, protozoans, rotifers, crustacean larvae and copepods were observed in the group. Treatment ponds had low levels of micro flora, organic substances and other organisms compared to control.



Figure 3: Biofloc (mL/L) in control and treatment ponds.



Figure 4: Plankton community in control and treatment ponds.

Heterotrophic bacteria and THC count Total heterotrophic bacterial population in pond water during the culture period is presented in Table 3. Significant difference (p<0.01) in Total plate count (TPC) for colony forming units (CFU) was found between control treatment ponds. Total Yellow *Vibrio* (TYV) colonies were found to be very low in treatment ponds when compared to control. Total Green *Vibrio* (TGV) colonies were found to be decreased in treatment ponds. It is noted that harmful *Vibrio* colonies were found to be considerably reduced in treatment pond water compared to control ponds. Details on total haemocyte count/mL of haemolymph are shown in Figure 5. Shrimp grown in Treatment ponds showed high haemocyte count compared to control and it did not

Table 3: Heterotrophic bacterial population in control and treatment ponds					
Days Ponds	Donda	TPC	TYV	TGV	
	ronus	(CFU/mL)	(CFU/mL)	(CFU/mL)	
(	Control	10000±232 <sup>a</sup>	10000±432 <sup>a</sup>	130±0.001 <sup>a</sup>	
30*	Treatment	8500±134 <sup>b</sup>	8200±267 <sup>b</sup>	13±2 <sup>b</sup>	
60* Control Treatment	Control	12333±366 <sup>a</sup>	3833±384 <sup>a</sup>	170±21 <sup>a</sup>	
	10833±376 <sup>b</sup>	1333±98 <sup>b</sup>	313±13 <sup>b</sup>		
90* Control Treatment	Control	63000±298 <sup>a</sup>	11093±399 <sup>a</sup>	$690 \pm 28^{b}$	
	Treatment	47333±243 <sup>b</sup>	1066±201 <sup>a</sup>	$23\pm9^{a}$	

change even after cold challenge test done at  $20^{\circ}$ C for 24 hours.

TPC-Total plate count; TYV- Total Yello Vibrio; TGV- Total Green Vibrio

CFU- Colony forming unit

\* p < 0.01; n =3; ANOVA showing significant difference between control and treatment in each sampling. a, b. Means with the same superscript do not differ from each other.



Figure 5: Total haemocyte count (THC) before and after cold challenge test.

Post harvest status and shrimp quality

Pond bottom status after harvest showed that an accumulation of sludge  $at102\pm12kg$  at the centre of treatment ponds; whereas in control ponds, it was  $29.4\pm5.8kg$ . Physical quality test of shrimp reveals that treatment shrimp had better colour than that of the control. The percentage of hard shell, loose shell and soft shell was 89.1, 5.5 and 5.4 respectively in treatment shrimps and 78.5, 8.5 and 13 was in control. No significant difference in taste observed between control and treatment shrimp.

#### Discussion

Biofloc technology is a technique of enhancing water quality in aquaculture through balancing carbon and nitrogen in the system (Crab *et al.*, 2012, Ray, 2014, Taw, 2014). Results of the present study show that biofloc strategy influences shrimp growth and production. Even though high growth was observed in biofloc ponds, the difference on growth between control and treatment was found to be non significant. This indicates that zero water exchange could bring growth and production as equal as traditional semi extensive culture and is an effective method to control the frequent discharge of to pond water the environment. The positive linear relationship found between biofloc content and growth rate in treatment ponds shows that biofloc formation influences to create favorable а environment for shrimp growth. Similar enhanced growth coupled with biofloc were observed in pacific white shrimp, Litopenaeus vannamei grown in zero water exchange system (Avnimelech, 2012; Manecas et al., 2013; Rivera et al., 2014; Ray, 2014; Taw, 2014). Therefore, it is suggested that biofloc culture strategy can be considered as is an ideal method to improve growth and production of Indian white shrimp, F.indicus.

All the water quality parameters remained within the ranges reported as conducive for the culture of white shrimp (Wickins, 1976; Van Wyk and 1999). The high salinity Scarpa, recorded in biofloc ponds may be the result of evaporation of water due to zero water exchange (Emerenciano, 2012). Studies evaluating water quality in zero-exchange systems report low concentrations of nitrate (Ray et al., 2010a,b; Vinatea et al., 2010). Low concentrations of nitrate observed during culture period suggest oxidation of ammonia (Cohen et al., 2005). According to Avnimelech and Ritvo (2003), only about 25% of the feed nutrients are converted into harvestable products hence contributing to high nitrogen residues in pond water, total ammonia nitrogen especially (TAN), which is the sum of both ammonia and ammonium which will adversely affect shrimp growth. This may be a reason for accumulation of ammonia in treatment ponds. Orthophosphate level in pond water regulates proliferation of phytoplankton and diatoms (Chaignon et al., 2002). The increased phosphate level found in treatment pond may lead the phytoplankton growth and pursued the formation of biofloc in treatment ponds.

Biofloc is explained as a medium rich in organic matter made of particulate biomass, friendly bacteria and phytoplankton (Rivera et al., 2014). From a nutritional point of view, it helps shrimp to gain weight owing to an abundance of native protein sources from protozoa, filamentous bacteria, nematodes, ciliates, flagellates, and rotifers (Decamp et al., 2002; Ray et al., 2010a,b). This is true in the case of present study, which shows that treatment pond has high phyto and zooplankton community compared to control. Presence of these organism in biofloc may served as source of natural food and this probably a reason for the high survival recorded in biofloc ponds as stated by Burford et al. (2003).

It has been reported that natural productivity in zero-exchange shrimp production systems provide supplemental food resources, reducing feed costs and improving shrimp growth rate (Otoshi et al., 2011, Manecas et al., 2013). Divakaran and Moss (2004) correlated higher shrimp growth with higher concentration of Chlorophyll a in phytoplankton. Becker (1994) and Olvera-Novoa et al. (1998) reported that all type of microalgae biomass are rich in polyunsaturated fatty acids and can be an important source of essential fatty acids for shrimp growth. The better colour of shrimp grown in biofloc pond may be due to the presence of high plankton (due to carotenoids) in the ponds. This assumption is in agreement the findings of Ju et al. (2009) who that microalgae reported in the microbial floc play a key role in improving shrimp growth rates and quality. Zooplankton consumes algae and bacteria and they can play an important role in transfer of nutrients from primary producers to secondary consumers (Moss al.. 2001). et Zooplankton such as rotifers can contribute significantly to protein and energy requirements of shrimp (Focken et al., 1998). This may be another reason for the enhanced survival and quality of shrimp recorded in biofloc ponds.

Natural production of some substances by bacteria in biofloc has been reported to inhibit growth of cohabiting pathogenic species such as *Vibrio harveyi* (Hsieh *et al.*, 2007; Iyapparaj *et al.*, 2013). Results of the present study show that harmful *Vibrio* colonies were found to be decreased in biofloc water and this can be attributed to the inhibitory effect of substance in bioflocs as stated above. Bianchi (1979) observed that bacteria in biofloc fluctuate and can have some antibiotic activity. The low yellow and green *Vibrio* colonies noticed in biofloc water may be due to the antibiotic activity of bacteria present in bioflocs. It is suggested that biofloc can control the growth and proliferation of pathogenic bacteria and harmful *Vibrio* colonies in pond water and thereby creating a favorable environment for shrimp growth.

Shrimp health is influenced by a range of factors, one of the most important being environmental stress (Xu and Pan, 2013). Under culture conditions, wide range of stresses by various caused adverse environmental factors damage the host defense system resulting in an increased susceptibility to infections (Perazzolo et al., 2002; Vazquez et al, 2009). Recently, scientists have hypothesized possibilities of immunostimulatory features of the bioflocs leading to enhancement of the immunity to provide broad-based resistance towards many infections (Crab et al., 2012). According to Wang et al. (2008), existing immunostimulants are group of live and synthetic compounds including bacteria and bacterial products, complex carbohydrates, nutritional animal extracts, factors, cytokines, lectins and plant extracts. Therefore, bioflocs might also contain immunostimulatory compounds since biofloc technology deals with bacteria and bacterial products. Total haemocyte count (THC) in haemolymph is

246

considered as index of shrimp immunological functions and a higher THC is responsible for high immune status (Rodriguez and Moullac, 2000; Krupesha et al., 2009). Results of the present study show that shrimp grown under stressed environment (treatment ponds had high ammonia, pH and salinity) had high THC. It is interesting to note that the treatment shrimp had a capacity to maintain the high THC even after a cold challenge was given at 20°C for 24 hours. This suggests that shrimp grown in biofloc environment may have strong immune capacity/response than those grown in clean water.

High accumulation of organic sludge was observed in all treatment ponds after harvest. This sludge formation was due to the deposition of dead algae and other organic compounds formed as a result of zero water exchange (Mikkelsen et al., 1996; Gonzalez-Felix et al., 2007). Studies show that this formed sludge would spoil pond bottom environment and act as medium for the proliferation and growth of harmful bacteria (Ju et al., 2008: Crab et al., 2012). Pathogenic bacteria act as agents to convert nitrogenous compounds into toxic ammonia and foul gases at pond bottom, elevating ammonia and pH level of water (Ebeling et al., 2006). This can be correlated as a factor for the enhanced ammonia, and pH observed in the biofloc ponds.

Overall performance of biofloc culture technique in the present study shows that it enhances shrimp growth similar to clean water culture. Heterotrophic bacterial population and phytoplankton community in biofloc ponds enhanced the colour and quality of shrimp. If better management methods are implemented to control ammonia and pH, biofloc technology would certainly be an ideal strategy for farmers for the biosecured production of shrimp and also to reduce environmental pollution due to coastal aquaculture.

#### Acknowledgements

The authors express sincere thanks to the Dean, Faculty of Marine Sciences and Managing Director, National Aquaculture Group for the facilities and support extended for the successful completion of the study.

### References

- APHA (American Public Health Association), 1995. Standard methods for the examination of water and wastewater, 19th Ed. American Public Health Association, Washington, DC.
- Avnimelech, Y., 2000. Nitrogen control and protein recycle. Activated suspension pond. *Global Aquaculture Advocate*, April, 23-24.
- Avnimelech, Y., 2005. Tilapia harvest microbial flocs in active suspension research pond. *Global Aquaculture Advocate*, October, pp.57-58.
- Avnimelech, Y., 2012. Biofloc technology A practical guide book.
  2nd edition, The World Aquaculture Society, Baton Rouge, Louisiana, USA, 272 P.
- Avnimelech, Y. and Ritvo, G., 2003. Shrimp and fish pond soils:

Processes and management. *Aquaculture*, 220, 549-567.

- Baloi, M., Arantes, R., Schveitzer, R., Magnotti, C. and Vinatea, L., 2013. Performance of Pacific white shrimp, *Litopenaeus vannamei* raised in biofloc systems with varying levels of light exposure. *Aquacultural Engineering*, 52, 39-44.
- Becker, E.W., 1994. Microalgae: Biotechnology and Microbiology. Cambridge University Press, Cambridge, UK, pp. 250-260.
- Bianchi, M.A.G., 1979. Polyphasic study of the microbial ecology of bacteria–phytoplankton interactions.
  Presented at: Aquatic Microbial Ecology: Proceedings of the Conference / Sponsored by the American Society of Microbiology, 7-10 Feb 1979, Clearwater Beach, FL (USA).
- **Boyd, C.E., 1990.** Water quality in ponds for aquaculture. Alabama Agriculture Experiment Station, Auburn University, Alabama, pp. 379-380.
- Burford, M.A., Thompson, P.J., McIntosh, R.P., Bauman, R.H. and Pearson, D.C., 2003. Nutrient and microbial dynamics in highintensity, zero-exchange shrimp ponds in Belize. Aquaculture, 219, 393-411.
- Chaignon, V., Lartiges, B.S., El Samrani, A. and Mustin, C., 2002. Evolution of size distribution and transfer of mineral particles between flocs in activated sludges: an insight

into floc exchange dynamics. *Water Research*, 36, 676-684.

- Cohen, J,M., Samocha, T.M., Fox, J.M., Gandy, R.L. and Lawrence, A.L., 2005. Characterization of water quality factors during intensive raceway production of juvenile Litopenaeus vannamei using limited discharge and biosecure management tools. *Aquacultural Engineering*, 32, (3-4), 425-442.
- Crab, R., Defoirdt, T., Bossier, P. and Verstraete, W., 2012. Biofloc technology in aquaculture: Beneficial effects and future challenges. *Aquaculture*, 356-357, 351-356.
- De Schryver, P., Crab, R., Defroit, T., Boon, N. and Verstraete W., 2008. The basics of biofloc technology: the added value for aquaculture. *Aquaculture*, 277, 125-137.
- Decamp, O., Conquest, L., Forster, I. and Tacon, A.G.J., 2002. The nutrition and feeding of marine shrimp within zero-water exchange aquaculture production system: role of Eukaryotic microorganisms. In: Lee, C.S., O'Bryen, P. (Eds.), Microbial Approaches to Aquatic Nutrition within Environmentally Sound Aquaculture Production World Aquaculture Systems. Society, Baton Rouge, FL, USA, pp. 79-86.
- **Divakaran, S. and Moss, S.M., 2004.** In vitro evidence of laminarinase activity in the digestive gland of juvenile Pacific white shrimp *Litopenaeus vannamei. Journal of*

the World Aquaculture Society, 35, 546-550.

- Ebeling, J.M., Timmons, M.B. and Bisogni, J.J., 2006. Engineering analysis of the stoichiometry of photoautotrophic, autotrophic and heterotrophic removal of ammonianitrogen in aquaculture systems. *Aquaculture*, 257, 346-358.
- Emerenciano, M., Ballester, E., Cavalli, R.O. and Wasielesky, W., 2012. Biofloc technology application as a food source in a limited water exchange nursery system for pink shrimp, *Farfantepenaeus* brasiliensis (Latreille, 1817). Aquaculture Research, 43, 447-457.
- Focken, U., Groth, A., Coloso, R.M. and Becker, K., 1998. Contribution of natural food and supplemental feed to the gut content of *Penaeus monodon* (Fabricius) in a semi intensive pond system in the Philippines. *Aquaculture*, 164, 105-116.
- Gonzalez-Felix, M.L., Ponce-Palafox, J.T., Valenzuela-Quinonez, W., Arredondo Figueroa, J.L. and Garcia-Ulloa, G.M., 2007. Nitrogen budget for a low salinity, zero-water exchange culture system. I. Effect of protein level dietary on the performance of Litopenaeus vannamei (Boone). Aquaculture and Research, 38, 798-808.
- Hargreaves, J.A., 2006. Photosynthetic suspended-growth systems in aquaculture. *Aquacultural Engineering*, 34, 344-363.
- Hsieh, S.L., Ruan, Y.H., Li, Y.C., Hsieh, P.S., Hu, C.H. and Kuo,

C.M., 2007. Immune and physiological responses in Pacific white shrimp (*Penaeus vannamei*) to *Vibrio alginolyticus. Aquaculture*, 275, 335-341.

- Iyapparaj, P., Maruthiah, T., Ramasubburayan, R., Prakash, S., Kumar, C., Immanuel, G. and Palavesam, A., 2013. Optimization of bacteriocin production by *Lactobacillus sp.* MSU3IR against shrimp bacterial pathogens. *Aquatic Biosystems*, 9 (1), 12-15.
- Ju, Z.Y., Forster, I., Conquest, L., Dominy, W.G, Kuo, W.C. and Horgen, F.D., 2008. Determination of microbial community structures of shrimp floc cultures by biomarkers and analysis of floc amino acid profiles. *Aquaculture and Research*, 39, 118-133.
- Ju, Z.Y., Forster, I.P. and Dominy, W.G., 2009. Effects of supplementing two species of marine or their fractions algae to a formulated diet on growth, survival and composition of shrimp (Litopenaeus vannamei). Aquaculture, 292, 237–243.
- Krupesha, S., Seema, J., Philipose, K.
  K. and Radhakrishnan, E.V., 2009.
  Effect of salinity and pH on selected immune functions of the Indian white shrimp, *Fenneropenaeus indicus* (H. Milne Edwards, 1837). *Indian Journal of Fisheries*, 56,183-187.
- Manecas B., Rafael, A., Rodrigo, S., Caio, M. and Luis, V., 2013. Performance of Pacific white shrimp, *Litopenaeus vannamei* raised

in biofloc systems with varying levels of light exposure. *Aquaculture Engineering*, 52, 39-44.

- McIntosh, **R.P.**, 2000. Changing paradigms in shrimp farming. heterotrophic Establishment of communities. bacterial Global Aquaculture Advocate. December. 52-54.
- Mikkelsen, L.H., Gotfredsen, A.K., Agerbxk, M.L., Nielsen, P.H. and Keiding, K. 1996. Effects of colloidal stability on clarification and dewatering of activated sludge. *Water Science and Technology*, 34, (3-4), 449-457.
- Moss, S.M., Divakaran, S. and Kim, B.G., 2001. Stimulating effects of pond water on digestive enzyme activity in the Pacific white shrimp, *Litopenaeus vannamei* (Boone). *Aquaculture Research*, 32, 125-131.
- Ogello, E.O., Musa, S.M., Aura, C.M., Abwao, J.O. and Munguti, J.M., 2014. An appraisal of the feasibility of tilapia production in ponds using biofloc technology: A review. *International Journal of Aquatic Science*, 5, 21-39.
- Olvera-Novoa, M.A., Doiminguez-Cen, L.J., Olivera-Castillo, L. and Martinez-Palacios, C.A., 1998. Effect of the use of the micro alga *Spirulina maxima* as fish meal replacement in diets for tilapia, *Oreochromis mossambicus* (Peters) fry. *Aquaculture Research*, 29, 709-715.
- Otoshi, C.A., Moss, D.R. and Moss, S.M., 2011. Growth-enhancing effect of pond water on four size classes of pacific white shrimp,

Litopenaeus vannamei. Journal of the World Aquaculture Society, 42, 417-422.

- Perazzolo, L. M., Gargioni, R., Ogliari, P. and Barracco, M.A., 2002. Evaluation of some hematoimmunological parameters in the shrimp *Farfantepenaeus paulensis* submitted to environmental and physiological stress. *Aquaculture*, 214, 19-33.
- Ray, A.J., 2014. Biofloc trial results in fast shrimp growth, low FCR, high survival. *Global Aquaculture Advocate*, January/February, 22-23.
- Ray, A.J., Lewis, B.L., Browdy, C.L. and Leffler, J.W., 2010b. Suspended solids removal to improve shrimp (Litopenaeus vannamei) production and an evaluation of a plant based feed in minimal-exchange, super intensive culture systems. Aquaculture, 299, 89-98.
- Ray, A.J., Seaborn, G., Leffler, J.W., Wilde, S.B., Lawson, A. and Browdy, CL., 2010a. Characterization of microbial communities in minimal-exchange, intensive aquaculture systems and the effects of suspended solids management. *Aquaculture*, 310, 130-138.
- Rivera, D.A., Davo, A.P., Escalante, K., Chavez, C., Cuzon, G. and Gaxiola, G., 2014. Probiotic effect of FLOC on Vibrios in the pacific white shrimp, *Litopenaeus vannamei. Aquaculture*, 424-425, 215-219.

- Rodriguez, J. and Le Moullac, G., 2000. State of the art of immunological tools and health control of penaeid shrimp. *Aquaculture*, 191, 109-119.
- Smith, D. and Johnson, K. B., 1996. A guide to marine coastal plankton and marine invertebrate larvae. Dubuque, Iowa: Kendall/Hunt Pub. Co., 221p.
- Supono, U., Hutabarat, J., Prayitno, B.S. and Darmanto, Y.S., 2014. White shrimp (*Litopenaeus vannamei*) culture using heterotrophic aquaculture system on nursery phase. *International Journal* of Waste Resources, 4 (2), 1-4.
- Taw, N., 2010. Biofloc technology expanding at white shrimp farms. *Global Aquaculture Advocate*, May/June, 20-22.
- Taw, N., 2014. Intensive farm in Bali, Indonesia, produces shrimp in biofloc system. *Global Aquaculture Advocate*, January/February, 24-25.
- Van Wyk, P. and Scarpa, J., 1999. Water quality requirements and management. In: Van Wyk, P., Davis-Hodgkins, M., Laramore, R., Main, K.L., Scarpa, J. (Eds.), Farming Marine Shrimp in Recirculating Freshwater Systems. Florida Department of Agriculture

and Consumer Services, Tallahassee, FL, pp. 128-138.

- Vazquez, L., Alpuche, J. and Maldonado, G., 2009. Immunity mechanisms in crustaceans. *Innate Immunity*, 15, 179-188.
- Vinatea, L., Galvez, A.O., Browdy, C.L., Stokes, A., Venero, J., Haveman, J., Lewis, B.L., Lawson, A., Shuler, A. and Leffler, J.W., 2010. Photosynthesis, water respiration and growth performance of *Litopenaeus vannamei* in a superintensive raceway culture with zero water exchange: interaction of water quality variables. *Aquacultural Engineering*, 42, 17-24.
- Wang, J.C., Chang, P.S. and Chen, H.Y., 2008. Differential time-series expression of immune-related genes of pacific white shrimp, *Litopenaeus vannamei* in response to dietary inclusion of  $\beta$ -1, 3-glucan. *Fish* & *Shellfish Immunology*, 24, 113-121.
- Wickins, J.F., 1976. The tolerance of warm-water prawns to re-circulated water. *Aquaculture*, 9, 19-37.
- Xu, W.J. and Pan, L.Q., 2013. Enhancement of immune response and antioxidant status of *Litopenaeus vannamei* juvenile in biofloc based culture tanks manipulating high C/N ratio of feed input. *Aquaculture*, 412-413, 117-124.