

Catalog of ophiuroidea from southeastern coast of Iran (Northern parts of Oman sea)

Beygmoradi A.¹; Attaran-Fariman G.^{1*}

Received: July 2016

Accepted: August 2019

1-Chabahar Maritime University, Faculty of Marine Sciences, Department of Marine Biology, Chabahar, Iran.

*Corresponding author's Email: gilan.attaran@gmail.com

Keywords: Ophiuroidea, Morphology, Chabahar Bay, Iran

Introduction

Chabahar Bay is located in southeastern coast of Iran and the northern part of the Oman Sea in the geographic area of 25°22' 47"N and 60°39' 90"E. This bay with a variety of coverings, such as, muddy, stone, sandy and rocky, has provided possibility of life for different taxa, such as, shellfish, mollusks, echinoderm (Nikouyan and Savari, 1999; Ghanbarifardi and Malek, 2009; Yazdani *et al.*, 2009; Sadeghi and Loghmani, 2010; Khaleghi and Owfi, 2011; Fazeli *et al.*, 2013; Attaran *et al.*, 2014). Echinoderms with around 7,000 species can be found at every ocean depth of the marine ecosystems (Mirzaei *et al.*, 2019). Ophiuroids are relatively small echinoderms that all live in marine habitats, such as, sandy, rocky, stony, muddy, on sponge and corals (Barnes *et al.*, 2001). Ophiuroids, with more than 2,000 species, are one of the largest group of echinodermata that are distributed in all marine environments (from polar to equatorial

regions and from the intertidal to subtidal zones) (Pawson, 2007).

Materials and methods

Ophiuroidea specimens have been collected from four stations of intertidal and subtidal areas located in eastern part of Chabahar Bay by grab, hand and scuba diving from October 2013 to July 2014 (Fig. 1). Sampling stations were selected according to the accessibility to the beach, geomorphological varieties and ecological diversities. The samples were placed into plastic jars contain seawater and transferred to the zoology laboratory of Chabahar Maritime and Marine Sciences University. All ophiuroids specimens were examined under microscope and photographed by Cannon camera model C-DS and also stereo-microscope equipped with a Cannon camera model T4AL250 V. Species identification to the species level was carried out by using the following sources: Clark (1953); Clark and Rowe (1971); Clark and Courtman-Stock (1976);

Based on this research, seven brittle stars is recorded for the first time; *O. venusta* 'O. tigris' 'Ophiothela sp.' 'Ophiothrix sp.' 'Macrophiothrix sp₁,

Macrophiothrix sp₂. and *M. longipedata* from study locations. Morphology of identified species from Chabahar bay is shown in Figs. 2-17.




Figure 2: *Ophiothela venusta*; Chabahar Bay (North of Oman Sea, Iran). A: The full view B: dorsal view C: ventral view D: dorsal view of an arm E: ventral view of an arm with ventral arm plates. Scale bars = A 5 mm; B and E 400 μ m.




Figure 3: *Ophiothela venusta*, (showing color pattern), A: The full view with 6 arms B: dorsal view, detail of the radial shields C: ventral view D: dorsal view of an arm E: ventral view of an arm with ventral arm plates. Scale bars = A 5 mm; B 400 μ m; C and E 200 μ m.



Figure 4: *Ophiothela venusta*; (showing color pattern), A: The full view B: dorsal view, detail of the radial shields C: ventral view D: dorsal view of an arm E: ventral view of an arm with ventral arm plates. Scale bars = A 2 mm; B and E 200 μ m




Figure 5: *Ophiothela venusta*; (showing color pattern), A: The full view B: ventral view C: dorsal view of an arm D: ventral view of an arm with ventral arm plates. Scale bars = A 5 mm; B and D 200 μ m.




Figure 6: Other color patterns in *Ophiothela venusta* in Chabahar bay Scale Bars= A and E 5mm; F200 μ m




Figure 7: *Ophiothela tigris*; A: The full view B: dorsal view C: ventral view D: dorsal view of an arm E: ventral view of an arm with ventral arm plates. Scale bars = A 9 mm; B 600 μ m; C and E 400 μ m.



Figure 8: *Ophiothela tigris* with dark color pattern. Scale bar= 9mm




Figure 9: *Ophiothela* sp.; A: dorsal view, showing radial shields B: ventral view C: dorsal view of an arm D: ventral view of an arm with ventral arm plates. Scale bars = A 2mm; B 300 μ m; C and D 200 μ m.




Figure10: *Macrophiothrix* sp1.; on *Gorgonian* A: dorsal view B: ventral view C: dorsal view of an arm with dorsal arm plates D: ventral view of an arm with ventral arm plates. Scale bar = A 20 mm; B and C 400 μ m ; D 200 μ m.




Figure 11: *Macrophiothrix* sp2.; A: dorsal view B: ventral view C: dorsal view of an arm with dorsal arm plates D: ventral view of an arm with ventral arm plates. Scale bar = A 5 mm; B and D 400 μ m.




Figure 12: *Macrophiothrix longipeda*; A: dorsal view B: ventral view C: dorsal view of an arm with dorsal arm plates D: ventral view of an arm with ventral arm plates. Scale bar = A 2 cm; B 5 mm ; C and D 200 μ m.




Figure 13: *Macrophiothrix elongate*; A: dorsal view, detail of the radial shield B: ventral view C: dorsal view of an arm with dorsal arm plates D: ventral view of an arm with ventral arm plates. Scale bar = A 20 mm; B 7 mm; C and D 400 μ m.




Figure 14: *Ophiothrix* sp.; A: dorsal view B: ventral view C: dorsal view of an arm with dorsal arm plates D: ventral view of an arm with ventral arm plates. Scale bar = A and B 5 mm; C and D 200 μ m.




Figure 15: *Ophiothrix* *savigny*; A: dorsal view B: ventral view C: dorsal view of an arm with dorsal arm plates D: ventral view of an arm with ventral arm plates. Scale bar = A and B; 5 mm; C and D; 400 μ m.




Figure 16: *Ophionereis* *dubia*; A: dorsal view B: ventral view C: dorsal view of an arm with dorsal arm plates D: ventral view of an arm with ventral arm plates, tentacle scales and banded spines visible. Scale bars = A and B 1 mm; C 400 μ m; D 250 μ m




Figure 17: *Amphipholis squamata*; A: dorsal view, detail of the radial shields B: ventral view C: month frame D: dorsal view of an arm with dorsal arm plate E: ventral view of an arm with ventral arm plates. Scale bar = A-B 2 mm; C and E 100 μ m.

Ophiuroids echinoderms are well distributed in the tropical regions. Price and Rowe (1996), reported 44 species of brittle stars from the Indian Ocean.

The identification of this species is not, generally, easy due to their similar morphological characteristics and their accurate identification based on morphological features need to be assessed in detail and carefully (Stöhr *et al.*, 2012). In the present study 11 species of brittle star recorded from Chabahar Bay located in northern part of Oman Sea, which has been presented in this catalog. Of which *M. elongata* and *O. savignyi* reported for the first time by Khaleghi (2010), and subsequently by Khaleghi and Owfi (2011) from Chabahar Bay. Attaran-Fariman *et al.* (2014) recorded *O. dubia* from that area for the first time. *A. squamata* has also been documented from Chabahar Bay (Attaran-Fariman and Beygmoradi, 2016). *Ophiothrix* and *Macrophiothrix* genera occupy dominantly the shallow habitats of the tropical zones (Stöhr *et al.*, 2012). Price

(1981) encountered *O. venusta* between Gorgonians of Persian Gulf. in the present study this species also found on the gorgonians (coral; order: Alcyonacea) bed in sutida zones of hotel Lipar and Shahid Behashti stations. This is the first record of *O. venusta* for this location. In the Chabahar Bay *Ophiothela* species were found with multiple color patterns and morphological variety. Generally *O. venusta* has been reported with 5 arms, however, is rarely observed with 6 arms (Price, 1981). In this study, a specimen was found with 6-arms (Fig. 3A) from the subtidal zone of hotel Lipar (site 2). Since all the species belonging to the genus *Ophiothela* recorded from Chabahar Bay (during the present study) were collected from gorgonians substratum, it seems that gorgonians could be an appropriate and sustainable habitat for them. Due to the position of Chabahar Bay along the Oman Sea and open waters of the Indian Ocean and also presence of the variety of biological substrates, such as sandy,

muddy, stone, rocky, coral and spongy for distribution of Ophiuroids, findings a variety of ophiuroids in Chabahar Bay would not be so surprising and unexpected.

Acknowledgements

The present work is part of the MSc work of A.B. at the Chabahar Maritime University. Hereby, we would like to express our deep appreciations to laboratory members.

References

Attaran-Fariman, G., Beygmoradi, A. and Boos, K., 2014. First record of *Ophionereis dubia* (Echinodermata: Ophiuroidea) from Chabahar Bay (Oman Sea, Iran). *Marine Biodiversity Records*, 7(2), 1-5.

Attaran-Fariman, G., Beygmoradi, A., 2016. The first report of *Amphipholis squamata* (Delle Chiaje, 1829) (Echinodermata: Ophiuroidea) from Chabahar Bay – northern Oman Sea. *Iranian Journal of Fisheries Sciences*, 15(3), 1254-1261.

Barnes, R.S.K., Calow, P., Olive, P., Golding, D. and Spicer, J., 2001. The Invertebrates: A Synthesis. *Blackwell Publishing*, 497 P.

Cherbonnier, G. and Guille, A., 1978. Echinoderms: Ophiurides. Faune de Madagascar. *Le Centre national de la recherche scientifique, Le Journal*, 48(1), 272.

Clark, H.L., 1938. Echinoderms from Australia: An account of collections made in 1929 and 1932. *Memoirs of the Museum of Comparative Zoology*, 55, 1-596.

Clark, A.M., 1953. A revision of the genus *Ophionereis*. *Proceedings of the Zoological Society of London*, 123, 65-94.

Clark, A.M. and Rowe, F.W., 1971. Monograph of Shallow-water Indo-west Pacific Echinoderms. Trustees of the British Museum (Natural History), London. 234 P.

Clark, A.M. and Courtman-Stock, J., 1976. The echinoderms of South Africa. London: British Museum (Natural History). 277 P.

De Loriol, P.de., 1900. Notes Pour Servir a Letude des Echinoderms. *Revue Suisse Zool*, 8, 55-96.

Fazeli, N., Savari, A., Nabavi, S.M.B. and Zare, R., 2013. Seasonal variation of zooplankton abundance, composition and biomass in the Chabahar Bay, Oman Sea. *International Journal of Aquatic Biology*, 1, 294-305.

Ghanbarifardi, M. and Malek, M., 2009. Distribution, diversity, and abundance of rocky intertidal fishes in the Persian Gulf and Gulf of Oman, Iran. *Marine Biology Research*, 5, 496-502.

Khaleghi, M., 2010. Echinoderms species of Chabahar Bay. MSc thesis. University of Khoramshahr, Iran, 80 P.

Khaleghi, M. and Owfi, F., 2011. Identification of Echinoidea species in the intertidal zones of Chabahar Bay. *Journal of Animal Environment*, 4, 31-36.

Lamarck, J.P.B.A., 1816. Histoire naturelle des animaux sans

veretèbres. First Edition, Paris. 2, 522-568.

Ljungman, A., 1867. Om Nagra nya arter of Ophiuroider. *Ofversigt af Kongliga Vetenskaps-Akademiens Forhandlingar*, 23, 163-166.

Lütken, C.F., 1859. Additamenta ad historiam Ophiuidarum. Beskrivelser af nye eller hidtil kun ufuldstændigt kendte Arter af Slangestjerner. *Anden Afdeling—Kongelige Danske Videnskabernes Selskabs Skrifter*, 5, 177-271.

Lyman, T., 1871. Supplement to the Ophiuridae and Astrophytidae. *Illustrated Catalogue of the Museum of Comparative Zoölogy at Harvard College*, 6, 1-17.

Mirzaei, M.R., Vahabnezhad, A. and Owfi, F., 2019. Short communication: The first report of *Asterodiscides fourmanoiri* (Echinodermata: Asteroidea) from the North Coast of Oman Sea, Iran. *Iranian Journal of Fisheries Sciences*, 18(3), 583-587.

Müller, J. and Troschel, F.H., 1840. Über die Gattungen der Asterien. Berlin: Wiegemanns Archiv fur Naturgeschichte, VI(1), 318.

Müller, J. and Troschel, F.H., 1842. System der Asteriden. Braunschweig: Verlag Friedrich Vieweg und Sohn. 134 P.

Nikouyan, A. and Savari, A., 1999. Distribution and biomass of macrobenthic fauna in the Chabahar Bay (North Eastern Sea of Oman). *Iranian Journal of Fisheries Sciences*, 1(2), 23-39.

Pawson, D.L., 2007. Phylum Echinodermata, *Zootaxa*, 1668, 749-764.

Pomory, M.C., 2007. Key to the common shallow-water brittle stars (Echinodermata: Ophiuroidea) of the Gulf of Mexico and Caribbean Sea. *Caribbean Journal of Science*, Special Publication, 10, 1-42.

Price, A.R.G., 1981. Studies on the echinoderm fauna of the western Arabian Gulf. *Journal of Natural History*, 15:, 1-15.

Price, A.R.G., 1983. Echinoderms of Saudi Arabia. Echinoderms of the Arabian Gulf coast of Saudi Arabia. *Fauna of Saudi Arabia*, 5, 28-108.

Price, A.R.G. and Rowe, F.W.E., 1996. Indian Ocean echinoderms collected during The Sindbad Voyage (1980-81): 3. Ophiuroidea and Echinoidea. *Bulletin Natural History of Museum London (Zoology)*, 62(2), 71-82.

Sadeghi, P. and Loghmani, M., 2010. First record of *Acanthopleura vaillantii* (Mollusca: Polyplacophora) from Iran-Chabahar Bay in the Oman Sea. *Marine Biodiversity Records*, 3(7), 1-2.

Stöhr, S., O'Hara, T.D. and Thuy, B., 2012. Global diversity of brittle stars (Echinodermata: Ophiuroidea), *PLoS One*, 7, 1-14.

Yazdani, M., Taheri, M., Sanjani, M.S. and Seyfabadi, J., 2009. New observation of two species, *Thalamita admete* and *Thalamita prymna* (Decapoda: Portunidae), from Chabahar Bay (Gulf of Oman). *Marine Biodiversity Records*, 2(56), 1-3.