The effects of feeding rates on growth performance, feed conversion efficiency and body composition of juvenile snow trout, *Schizothorax zarudnyi*

Khandan Barani H.^{1*}; Dahmardeh H.²; Miri M.³; Rigi M.¹

Received: May 2016

Accepted: January 2017

Abstract

This study was performed to investigate the effects of feeding rates on growth performance, feed conversion and body composition of juvenile snow trout (*Schizothorax zarudnyi*). For this purpose, fish with an initial body weight of 2.17 ± 0.1 g were fed a commercial diet (40% protein, 3.8 Kcal g⁻¹ diet) at five ration sizes of 2%, 4%, 6%, 8% and 10% of their body weight per day, for 60 days. The feeding trial was conducted in triplicate. Maximum weight gain, best feed conversion ratio (FCR), best specific growth rate (SGR %), highest energy retention (ERE %) and protein efficiency ratio (PER %) were evident for rations of 4–6% body weight. Second-degree polynomial regression analysis for PER, FCR and ERE indicated the break-points occurred at 4.9, 5.1 and 5.2% BW day⁻¹, respectively. Maximum body protein content was obtained for 4% and 6% rations. Body moisture, fat and ash content remained unchanged. Based on the above results, it may be concluded that the 4.9-5.2% body weight per day ration is optimal for the growth, conversion efficiencies and body composition of juvenile *S. zarudnyi* and it entails favorable economic benefits.

Keywords: Growth, Body composition, Ration, Schizothorax zarudnyi

¹⁻Department of Fisheries, Hamoun International Wetland Research Institute, University of Zabol, Zabol, Iran

²⁻Department of Fisheries, Faculty of Natural Resources, University of Zabol, Zabol, Iran.

³⁻Department of Natural ecosystems management, Hamoun International Wetland Research Institute, University of Zabol, Zabol, Iran.

^{*}Corresponding author's Email: Hashem.Barani@uoz.ac.ir

Introduction

А crucial factor in commercial aquaculture is feeding, especially when it affects growth, health of fish and production costs (Cho et al., 2003; Khan and Abidi, 2010; Tian et al., 2015). Feed costs make up a large fraction of the total fish production costs (40-50 % of production costs) (Anderson et al., 1997). Moreover, in practical fish farming, growth optimization is important to ensure profitability (Wang et al., 2009). The growth of fish at all stages in its life history is largely regulated by a number of factors, such as food type, food intake, feeding rate, feeding frequency and its ability to absorb nutrients (Gelineau et al., 1998; Akbulut et al., 2013). Among these, feeding rate is an important factor for the survival and growth of fish at the early life stages (Goddard, 1996). The influence of feeding rate on fish growth received performance has much attention (Goddard, 1996; Oberg et al., 2014; Luo et al., 2015; Li et al., 2016). Most studies of feeding rate have confirmed that there should be optimal feeding rates in fish culture. Determination of optimal values for this factor helps to enhance growth (Saether and Jobling, 1999) or feed utilization (Fiogbé and Kestemont, 2003); to reduce feeding costs (Silva et al., 2007; Luo et al., 2015) or to improve water quality (Allan et al., 1995; Zakes et al., 2006) and reduce fish size variations (Johnston et al., 2003). Consequently, for the commercial success of any aquaculture venture, it is important to

have adequate knowledge of the optimal ration level.

Several studies have been conducted to evaluate the effect of feeding rate on the growth, survival, feed intake, body composition, among other parameters, in various culturable finfish species, such as common carp (Desai and Singh, 2009), mangrove red snapper (Abbas and Siddiqui, 2009), pigfish (Oberg et al., 2014), blunt snout bream (Tian et al., 2015), hybrid sturgeon (Luo et al., 2015). It has been shown in these studies that optimum rate might vary based on fish species, size, age, feed quality and culture system. Thus, to achieve optimum production, it is feeding important that rate be standardized for the target species in aquaculture.

It is important to choose new fish good species with potential for aquaculture to develop this growing industry. Schizothorax zarudnyi, also known as snow trout, is a native fish of southeast Iran and Afghanistan that belongs to the cyprinid family. In general, cyprinid species have high commercial value and their meat is appreciated by human consumers. Also, snow trout has high market value and is a promising candidate species for Iranian aquaculture diversification (Aghili, 2013; Rahdari et al., 2014). However, there is no information on the feeding application rates for juvenile snow trout. Therefore, the aim of this study was to assess the effects of feeding rate on the growth performance, feed conversion and body composition of juvenile snow trout, and determine the best feeding rate for this species

Downloaded from jifro.ir on 2025-06-16

under commercial production conditions. Fish nutritionists and farmers will benefit from the results of this study.

Materials and methods

This study was performed at Hamoun International Wetlands Research Institute of Zabol University. Healthy juvenile of snow trout were obtained from a local commercial fish farm in Zehak city, Iran. Fish were acclimated for 14 days before the start of the experiment. During the acclimation and experimental period, the water quality parameters were maintained as follows: temperature (22.6±1.5 °C). pН (7.6 ± 0.3) , dissolved oxygen (6.9 ± 0.5) mg L^{-1}) and ammonia concentration was kept around zero mg L^{-1} . After the acclimation period, fish with an average initial body weight of 2.17±0.3 g (mean±SD) were distributed randomly among 15, 80 L glass tanks at the rate of 20 fish per tank. There were no significant differences in the initial weight between groups. The study design included five feeding rates (2%, 4%, 6%, 8% and 10%) with three replicates for each treatment. The daily rations were supplied each day in two equal halves, at 08.00 and 17.00 hours. Fish were weighed every 2 weeks to regulate the feeding rates. The feeding trial lasted for 60 days. The fish were offered a commercial diet (Esfahan Mokammel Company, Isfahan, Iran), containing 40% protein, 12% lipid, 9-10% moisture and 10% ash.

Body composition analysis

Before the start of the experiment, 20 stocked fish were killed and analyzed for initial body composition. At the end of the experiment, ten fishes from each replicate were sacrificed for the analysis of proximate composition of whole body. The ash, moisture, crude protein and lipid contents of fish were estimated by AOAC (2000).

Growth parameters

The growth parameters were evaluated follows: Survival (%)as (Survival%=100×(final fish number/ initial fish number), Weight gain rate (WGR=100×(average final body weight-average initial body weight)/average initial body weight), Feed conversion ratio (FCR=total feed intake/total weight gain), Specific growth rate (SGR=100×(ln final mean weight-ln initial mean weight)/days), retention efficiency Energy (ERE=100×(energy gain/total energy in the diet), Protein efficiency ratio (PER=100×(weight gain/protein fed) (Tacon, 1990; De Silva and Anderson, 1995; Khan and Abidi, 2010).

Data analysis

Data were analyzed by one-way (ANOVA) using SPSS, version 16 for Windows, and Duncan's test was used to compare differences between the means at 5% probability. All experimental values are expressed as mean \pm SD. The estimation of optimum feeding rate for growth and feed conversion efficiency of snow trout was carried out using second-degree polynomial regression analysis ($Y=aX^2$ +bX+c) (Zeitoun *et al.*, 1976).

Results

Growth performance and feed conversion efficiencies of juvenile S. zarudnyi subjected to different feeding rates are presented in Table 1. All fish survived throughout the trial. Maximum weight gain rate (%) was recorded with a diet of 6% BW day⁻¹. Best FCR, and highest specific growth, Energy retention efficiency (ERE) and Protein efficiency ratio (PER) where obtained for fish fed by 4-6% ration. Feed conversion ratio (FCR) decreased as the ration rose up to 6%. No significant (p>0.05) improvement in FCR was seen for fish fed with the 8% ration and increasing the ration further indicated in no improvement or even poor FCR. In order to obtain more exact information on the ration level, second-degree polynomial regression analysis was conducted for the growth parameters. Using the second-order polynomial regression analysis model, the optimal feeding rate for juvenile snow trout based on PER, FCR and RER was estimated to be 4.9, 5.1 and 5.2 % BW day⁻¹, respectively (Figs. 1, 2 and 3).

 Table 1: Growth, conversion efficiencies and nutrient retention efficiency of juvenile Schizothorax zarudnyi fed at different feeding rates.

Parameters	Feeding rate (%weight day ⁻¹)				
	2	4	6	8	10
Initial body weight (g)	2.21±0.07	2.13±0.10	2.14±0.17	2.20±0.06	2.09±0.06
Final body weight (g)	$2.75{\pm}0.07^{a}$	3.73±0.24 ^b	4.36±0.14 ^c	$4.09 \pm 0.19^{\circ}$	3.48 ± 0.15^{b}
Weight gain (%)	54±3.46 ^a	152±15.39 ^b	222.33±4.50 ^d	188.6±22.2 3 ^c	139±19.15 ^b
Feed conversion ration	4.82 ± 0.49^{b}	3.48 ± 0.24^{a}	3.47 ± 0.35^{a}	$5.67 \pm 0.83^{\circ}$	9.13±1.36 ^d
Specific growth rate (%)	0.36±0.02 ^a	0.86 ± 0.05^{b}	1.19±0.08 ^c	1.02 ± 0.10^{bc}	$0.84{\pm}0.10^{b}$
PER (%)	52.49±0.05 ^b	$71.27 \pm 0.05^{\circ}$	73.79±0.04°	42.63±0.09 ^b	$27.87{\pm}0.04^{a}$
ERE (%)	16.19±0.01 ^b	21.46±0.02 ^c	22.31±0.005 ^c	14.51 ± 0.01^{b}	8.76±0.01 ^a

Mean value \pm SEM from three replicate analyses; mean values with the same superscript are not significantly different (*p*>0.05)



Figure 1: Second-order polynomial fitting of protein efficiency ratio to feeding rate of *Schizothorax zarudnyi*.



Figure 2: Second-order polynomial fitting of feed conversion efficiency to feeding rate of *Schizothorax zarudnyi*.



Figure 3: Second-order polynomial fitting of energy retention to feeding rate of *Schizothorax zarudnyi*.

Whole-body composition of juvenile snow trout is shown in Table 2. Wholebody protein content was found to be significantly higher (p<0.05) for 4% and 6% rations compared with fish fed with other rations. Body ash, fat and Moisture did not differ among fish fed with different rations.

 Table 2: Whole body composition of juvenile Schizothorax zarudnyi fed different feeding rates

 Parameters
 Fooding rate (% waight(day))

Parameters	Feeding rate (%weight/day)							
	2	4	6	8	10			
Moisture (%)	76.41±0.52	76.85±0.28	76.12±0.20	77.41±0.63	77.90±0.46			
Protein (%)	66.79 ± 1.25^{a}	69.58 ± 1.09^{bc}	$71.24 \pm 1.38^{\circ}$	68.14 ± 1.2^{ab}	68.38 ± 1.03^{b}			
Fat (%)	9.78±0.33	9.74 ± 0.42	9.49±0.11	9.16±0.82	9.38±0.16			
Ash (%)	3.01±0.19	2.86±0.26	2.80±0.19	2.63±0.27	2.79±0.12			

Mean values \pm SEM from three replicate analyses; mean values with the same superscript are not significantly different (p>0.05)

Discussion

Feeding rate is an important factor affecting growth in fish. Determination of optimal values for this factor is necessary to the success of aquaculture production. In the present study, ration levels had significant effects on the growth, SGR, FCR, ERE and PER of S. zarudnyi. Similarly, studies by Ahmed (2007) on Labeo rohita, Khan and Abidi (2010) on Heteropneustes fossilis and Oberg et al. (2014) on pigfish reported that ration levels very often affect the growth performances. survival, feed intake and conversion efficiencies of cultured fish.

Our study revealed that growth in terms of final weight and other weight related indices SGR and WGR in juveniles of snow trout could be improved when they were reared with increasing feeding ration up to a certain level as was observed in Cirrhinus mrigala (Khan and Abidi, 2004) and hybrid sturgeon (Luo et al., 2015). growth However, significant improvement was recorded with increasing rations up to 6%, and was not significantly different from that achieved by fish fed at 4% BW day⁻¹. This indicates that feeding fish in the range of 4-6% BW day⁻¹ results in maximum utilization of food for growth. On subjecting PER, FCR and RER data to second degree polynomial regression analysis, however, breakpoints occurred for rations of 4.9, 5.1 and 5.2%, respectively. These breakpoints show that rations in the range 4.9–5.2% BW day⁻¹ are optimal for growth of snow trout. This enables producers to grow healthy fish as

quickly and/or efficiently as desired and using optimal ration prevents overfeeding, which is costly and reduces water quality. The above range is higher than the ration allocation reported for grass carp, Ctenopharyngodon idella (1.97%) (Du et al., 2006), Atlantic salmon, Salmo salar (2%) (Austreng et al., 1987), Nile tilapia, O. niloticus (2%) (El-Saidy and Gaber, 2005), European sea bass, Dicentrarchus labrax (3 - 3.5%)(Eroldogan al., 2004), et hybrid sturgeon (3.7%) (Lue et al., 2015), in the range determined for big head carp. Aristichthyes nobilis (2-5.7%)(Opuszynski and Shireman, 1991), and lower than the values for H. fossilis (5.9-6.8%) (Khan and Abidi, 2010), Common carp, Cyprinus carpio (6%) (Desai and singh, 2009), Indian major carp, Labeo rohita (6.5-7%) (Ahmed, 2007). African catfish, Clarias gariepinus (6%) (Al-Hafedh and Ali, white sturgeon, 2004), Acipenser transmontanus (6%) (Deng et al., Tambaqui, 2003). Colossoma *macropomum* (10%) (Silva et al., 2007), clownfish, Amphiprion percula (10.26%) (Johnston et al., 2003) and brown trout, Salmo trutta (11.3%) (Elliot, 1975). Thus, the difference in nutrient requirement could also be related to species and age of fish, composition of the diet, stock density and water quality.

In this study, poor growth and FCR for fish fed lower rations of 4% BW day⁻¹, lower rations than the optimum, suggests these rations approximate to maintenance requirements only and that most of the ingested nutrients are used

to maintain life and a small portion probably remained for growth. The present findings on snow trout also seem to be in agreement with the observations of Khan et al. (2004) on C. mrigala. Moreover, poor FCR for higher rations can be due to loss of nutrients and waste of food, because it takes longer for fish to consume food to reach satiation. Similar results were also reported in Heteropneustes fossilis (Khan and Abidi, 2010), Labeo rohita (Ahmed. 2007) and Acipenser transmontanus (Hung and Lutes, 1987). The whole-body composition of fish is known to indicate the fish quality. Many studies have confirmed that body composition is significantly affected by feeding rate (Ahmed, 2007; Desai and Singh, 2009; Khan and Abidi, 2010). The findings of the present study showed that ration significantly affects whole-body protein content. The highest protein contents were observed at 4%-6% feeding ration. Similar observations were made by Khan et al. (2004) in C. mrigala. Also in the present study, ash, fat and moisture were not significantly influenced by ration level. This is in agreement with results of Desai and Singh (2009) who reported that ash and moisture content remained insignificantly different among the ration levels in common carp. Generally, lack of difference between fish fed with 4% and 6% ration indicates that all fishes in these treatment groups were adequately fed. In conclusion, results of the present study show that a feeding rate of 4.9-5.2% BW day^{-1} is the most efficient feeding strategy for juvenile snow trout, because it results in greater growth and production. Therefore the application of this feeding rate for juvenile Schizothorax zarudnyi would insure feed consumption optimum with minimum waste, improve nutritional efficiency food conversion and efficiency, decrease production cost and reduce water pollution.

Acknowledgements

The authors would like to thank Narjes Sanchooli, Alireza Noorafar and Mahdi Keikha for their help during the analyses.

References

- Abbas, G. and Siddiqui, P.J.A., 2009. Effects of different feeding level on the growth, feed efficiency and body composition of juvenile mangrove red snapper, *Lutjanus argentimaculatus* (Forsskal 1775). *Aquaculture Research*, 40, 781-789.
- Aghili, K., 2013. Replacement of snow trout, *Schizothorax zarudnyi*, by common carp in poly culture. *Animal Research*, 26, 435-442 (In Persian).
- Ahmed, I., 2007. Effect of ration size on growth, body composition, and energy and protein maintenance requirement of fingerling Indian major carp, *Labeo rohita* (Hamilton). *Fish Physiology and Biochemistry*, 33, 203–212.
- Akbulut, B., Feledi, T., Lengyel, S. and Ronyai, A., 2013. Effect of feeding rate on growth performance, food utilization and meat yield of starlet, *Acipenser ruthenus* (Linne, 1758). *Journal of Fisher Scientific*, 7, 216-224.

- Al-Hafedh, Y.S. and Ali, S.A., 2004. Effects of feeding on survival, cannibalism, growth and fed conversion of African catfish, *Clarias gariepinus* (Burchell) in concrete tanks. *Journal of Applied Ichthyology*, 20, 225–227.
- Allan, G.L., Moriarty, D.J.W. and Maguire, G.B., 1995. Effects of pond preparation and feeding rate on production of *Penaeus monodon* fabricius, water quality, bacteria and benthos in model farming ponds. *Aquaculture*, 130, 329-349.
- Anderson, J.S., Higgs, D.A., Beams, R.M. and Rowshandeli, M. 1997. Fish meal quality assessment for Atlantic salmon, *Salmo salar*, reared in seawater. *Aquaculture Nutrition*, 3, 25-38.
- AOAC, 2000. Official Methods of Analysis. 17th Edition, The Association of Official Analytical Chemists, Gaithersburg, MD, USA.
- Austreng, E., Storebakken, T. and Asgard, T., 1987. Growth rate estimates for cultured Atlantic salmon and rainbow trout. *Aquaculture*, 60, 157–160.
- Cho, S.H., Lim, Y.S., Lee, J.H. and Park, S., 2003. Effect of feeding rate and feeding frequency on survival, growth, and body composition of ayu post-larvae *Plecoglossus altivelis. Journal of the World Aquaculture Society*, 34, 85–91.
- Deng, F., Koshio, S., Yokoyama, S., Bai, S.C., Shao, Q., Cui, Y. and Hung, S.S.O., 2003. Effects of feeding rate on growth performance of white sturgeon *Acipensor*

transmontanus larvae. *Aquaculture*, 217, 589–598.

- Desai, A.S. and Singh, R.K., 2009. The effects of water temperature and ration size on growth and body composition of fry of common carp, *Cyprinus carpio. Journal of Thermal Biology*, 34, 276–280.
- De Silva, S.S. and Anderson, T.A., 1995. In: fish nutrition in aquaculture. Chapman and Hall, London. 319 P.
- Du, Z.Y., Liu, Y.J., Tian, L.X., He, J.G., Cao, J.M. and Liang, G.Y., 2006. The influence of feeding rate on growth, feed efficiency and boy composition of juvenile grass carp *Ctenopharyngodon idella*. *Aquaculture International*, 14, 247–257.
- Elliot, J.M., 1975. The growth rate of brown trout *Salmo trutta* L. fed on reduced rations. *Journal of Animal Ecology*, 44, 823–842.
- El-Saidy, D.M.S.D. and Gaber, M.A., 2005. Effect of dietary protein levels and feeding rates on growth performance, production traits and body composition of Nile tilapia, *Oreochromis niloticus* cultured in concrete tanks. *Aquaculture Research*, 36, 163–171.
- Eroldogan, O.T., Kumlu, M. and Aktas, M., 2004. Optimum feeding rates for European sea bass *Dicentrarchus labrax* L. reared in seawater and freshwater. *Aquaculture*, 231, 501–515.
- Fiogbé, E.D. and Kestemont, P., 2003. Optimum daily ration for Eurasian perch *Perca fluviatilis* L. reared at its

optimum growing temperature. *Aquaculture*, 216, 243-252.

- Gelineau, Corraze, A., G. and 1998. Boujar, Т., Effects of restricted ration, time-restricted access and reward level on voluntary food intake, growth and growth heterogeneity of rainbow trout Oncorhynchus mykiss fed on demand with self-feeders. Aquaculture, 167, 247-258.
- Goddard, S., 1996. Feed Management in Intensive Aquaculture. Chapman and Hall, New York. 194 P.
- Hung, S.S.O. and Lutes, P.B., 1987. Optimum feeding rate of hatchery produced juvenile white sturgeon *Acipenser transmontanus* at 20 °C. *Aquaculture*, 65, 307–317.
- Johnston, G., Kaiser, H., Hecht, T., and Oellermann, L., 2003. Effect of ration size and feeding frequency on growth, size distribution and survival of juvenile clownfish, *Amphiprion percula*. *Journal of Applied Ichthyology*, 19, 40–43.
- Khan, M.A. Ahmed, I. and Abidi, S.F., 2004. Effect of ration size on growth, conversion efficiency, and body composition of fingerling mrigala, *Cirrhinus mrigala* (Hamilton). *Aquaculture Nutrition*, 10, 47–53.
- Khan, M.A. and Abidi, S.F., 2010. Optimum ration level for better growth, conversion efficiencies and body composition of fingerling *Heteropneustes fossilis* (Bloch). *Aquaculture International*, 18, 175– 188.
- Li, X.F., Xu, C., Tian, H.Y., Jiang, G.Z., Zhang, D.D. and Liu, W.B.,

2016. Feeding rates affect stress and non-specific immune responses of juvenile blunt snout bream *Megalobrama amblycephala* subjected to hypoxia. *Fish and Shellfish Immunology*, 49, 298-305.

- Luo, L., Li, T., Xing, W., Xue, M., Ma, Z., Jiang, N. and Li, W., 2015. Effects of feeding rates and feeding frequency on the growth performances of juvenile hybrid sturgeon, Acipenser schrenckii Brandt^Q × Acipenser baeri Brandt^J. Aquaculture, 448, 229–233.
- Oberg, E.W., Faulk, C.K. and Fuiman, L.A., 2014. Optimal dietary ration for juvenile pigfish *Orthopristis chrysoptera* grow-out. *Aquaculture*, 433, 335–339.
- **Opuszynski, K. and Shireman, J.V., 1991.** Food passage time and daily ration of bighead carp, *Aristichthyes nobilis*, kept in cages. *Environmental Biology of Fishes*, 30, 387–393.
- Rahdari, A., Gharaei, A. and Ghaffari, M., 2014. Spawning latency period in hormonal induced reproduction of snow trout, *Schizothorax* zarudnvi Nikolskii, Iranian 1897. Journal of Biotechnology, 12, 61-65.
- Saether, B.S. and Jobling, M., 1999. The effects of ration level on feed intake and growth, and compensatory growth after restricted feeding in turbot *Scophthalmus maximus* L. *Aquaculture Research*, 30, 647-53.
- Silva, C.R., Gomes, L.C. and Brandao, F.R., 2007. Effect of feeding rate and frequency on tambaqui *Colossoma macropomum*

growth, production and feeding costs during the first growth phase in cages. *Aquaculture*, 264, 133–139.

- Tacon, A.G.J., 1990. Standard method for nutritional and feeding of farmed fish and shrimp. Argent liberations press. Redmond, Wash, 1, 117.
- Tian, H.Y., Zhang, D.D., Li, X.F.,
 Zhang, C.N., Qian, Y. and Liu,
 W.B., 2015. Optimum feeding frequency of juvenile blunt snout bream *Megalobrama amblycephala*.
 Aquaculture, 437, 60–66.
- Wang, N., Xu, X. and Kestemont, P., 2009. Effect of temperature and feeding frequency ongrowth performances, feed efficiency and body composition of pikeperch juveniles *Sander lucioperca*. *Aquaculture*, 289, 70–73.
- Zakes, Z., Demska-Zakes, K., Jarocki, P. and Stawecki, K., 2006. The effect of feeding on oxygen consumption and ammonia excretion of juvenile tench *Tinca tinca* (L.) reared in a water recirculating system. *Aquaculture International*, 14, 127-140.
- Zeitoun, I.H., Ullrey, D.E., Magee,
 W.T., Gill, J.L. and Bergen, W.G.,
 1976. Quantifying requirements of
 fish. Journal of the Fisheries
 Research Board of Canada, 33, 167–172