Metal bioaccumulation in representative organisms from different trophic levels of the Caspian Sea

Mirzajani A.R.^{1,2}; Hamidian A.H.^{1*}; Karami M.¹

Received: May 2015 Accepted: October 2015

Abstract

The bioaccumulations of metals Cu, Cd, Ni, Cr, Co, Mn, Zn and Fe were measured in bivalves, Cerastoderma glucaum, and four species of fishes including Alburnus chalcoides, Liza aurata, Rutilus frisii and Sander lucioperca from various trophic levels of the Caspian food web. The concentrations of Cd, Cr, Co and Ni in most samples of fish were below the detection limits; while the concentrations were detected in most samples of bivalve C. glucaum. The stable nitrogen isotope ratios varied among the samples from C. glucaum (δ^{15} N=3.5 %) to S. lucioperca (δ^{15} N=13.1%). Among the four fish species, while the highest concentrations of Mn, Ni and Fe were observed in L. aurata, the lowest concentrations of Mn and Fe were observed in S. lucioperca. These species also had the lowest and highest trophic levels with an average of 3.3 and 4.2, respectively. No accumulation of metals with increase in body size was observed in muscles of species from different trophic levels. The comparison of metal concentrations with the health guidelines for human consumption showed that those intakes were lower than the legislated limits. While there was a strong relationship between trophic levels and body size of A. chalcoides and R. frisii, no significant slopes were observed between the total lengths (TLs) and the Ln concentrations of metals. It is necessary to determine metal concentrations in food resources of fish species, particularly in R. frisii that has significantly different $\delta^{15}N$ in relation to body size.

Keywords: Metal pollution, Bioaccumulation, Stable isotope, Caspian Sea

[DOR: 20.1001.1.15622916.2016.15.3.9.7]

¹⁻Department of Environment, Faculty of Natural Resources, University of Tehran, P.O. Box 4314, 31587-77878 Karaj, Iran

²⁻ Present address; Inland waters Aquaculture Research Center, Iranian Fisheries Science Research Institute, Agricultural Research Education and Extension Organization (AREEO), Badar Anzali, P.O. Box 66, Iran

^{*}Corresponding author's Email: a.hamidian@ut.ac.ir

Introduction

The Caspian Sea, the largest landenclosed drainage area in the world, supports substantial fisheries, while it has been endangered by massive loads of contaminants. The various environmental pollutants produced by natural and anthropogenic sources are discharged from coastal catchment (Glantz, 2000; UNEP/UNDP, 2011). The hazardous effects of heavy metals on human health are reviewed by international organizations such as the WHO (Järup, 2003).

Caspian fishes, an important source of protein, are consumed by millions of people living around the Caspian Sea; therefore monitoring their metal pollution is essential (Fallah *et al.*, 2011).

Bioaccumulation is defined as an increase in the concentration pollutants by an increase in age, which sometimes is reflected in size of the organism (Fernandes et al., 2007). Metal bioaccumulation is influenced by various environmental and biological factors particularly the feeding source (Agusa et al., 2004; Wang and Rainbow, 2008). Although referring to scientific literature, very few data have been published on metal accumulation related to size and age (Anan et al., 2005; Pourang et al., 2005; Foroughi Fard et al., 2008; Pazooki et al., 2009; Hoseini and Tahami, 2012). There have been some research (Sadeghirad, 2002; Agusa et al., 2004; Sadeghirad, 2007; Shahryari et al., 2010; Eslami et al., 2011; Fallah et al., 2011; Monsefrad et al., 2012) presenting metals

concentrations in the Caspian fishes. While nearly all of these reports were focused on the Acipenseridae species (particularly *Acipenser persicus*) and *Rutilus frisii*, other commercial fishes were less concerned. Furthermore, no research has been done according to trophic levels of Caspian fishes.

Four species, *A. chalcoides, Liza aurata, R. frisii* and *Sander lucioperca,* have a high commercial value and good market demands in the southern Caspian regions. These species are traditionally known as zooplankton feeders, detritivrous, benthivorous and piscivorous, respectively (Ghadirnejad, 1996; Zarinkamar, 1996; Abbasi *et al.*, 2004; Abdolmaleki and Ghaninezhad, 2007; Neamatparast, 2007).

Stable isotope analysis has been widely used, as a powerful tool, for source identification of changes in aquatic and estuarine ecosystems, and also for assessing pollutant flux in marine ecosystems (Broman *et al.*, 1992; Cabana and Rasmussen, 1996; Vander Zanden and Rasmussen, 1996; Lake *et al.*, 2001; Jardine *et al.*, 2006).

However the metal concentrations in muscle tissues are reported to be usually lower than in other organs (Pourang *et al.*, 2005; Kojadinovic *et al.*, 2007; Mashroofeh *et al.*, 2013), the turnover rate of δ^{15} N in muscle protein is much longer and represents the integrated diet and contamination of fishes for a long time (Cabana and Rasmussen, 1996; Vander Zanden and Rasmussen, 1996).

Typically the ¹³C/¹²C and ¹⁵N/¹⁴N ratios of a consumer are slightly higher

than those in its diet. On average, δ^{13} C and $\delta^{15}N$ values of consumers are expected to increase by around 1% and 3.4% compared to those of their food, respectively (Minagawa and Wada, 1984). The nitrogen isotope compositions are particularly wellsuited for examining trophic dynamics and provide an estimate of their trophic position (Minagawa and Wada, 1984; Post, 2002) while carbon isotope elucidates the origin and pathways of organic matter in food webs (Le Loc'h et al., 2008).

In this study, we determined metal concentrations in *A. chalcoides, L. aurata, R. frisii* and *S. lucioperca.* Furthermore, we studied the metal bioaccumulation based on the trophic levels in the Caspian food web.

Material and methods

Sample collection

The collections included Pike-perch (S. lucioperca) (n=16), Kutum (R. frisii) (n=25), Danube bleak (A. chalcoides) (n=20) and Golden grey mullet (L. aurata) (n=16). These species were freshly collected from Anzali region (~37 28 N 49 33 E) during winterspring of 2013. Two fishing methods; beach seine and gill net (Gerami and Dastbaz, 2013) were used for sampling. Furthermore about 360 individuals of the benthic invertebrates bivalvia C. glucaum were freshly collected along the beach down to 20 meter depths by hand and by using a bottom sampler (Van-Veen grab: 400 respectively. The total length of each sample was measured by Vernier calipers; and then they were weighed using a balance at the nearest 0.01g precision.

The dorsal muscles of fish samples were individually dissected using a high quality stainless steel knife; rinsed with distilled water and dried at 60 °C for 48 h. The dried samples were ground and homogenized with a pestle and mortar, stored in distilled water washed in plastic bags and kept at -20°C before the laboratory experiments. Gut organs and shells of mollusks were carefully removed under stereomicroscope and their soft tissues were pooled (20-30 as one sample). individuals samples were oven dried at 60 °C for 24 h. The preparation of samples followed the ROPME instructions (1989).

Chemical analyses

For metal (Cu, Cd, Ni, Cr, Co, Mn, Zn and Fe) analyses, a mass of around 0.5 g of each sample was digested in4 ml high purity nitric acid (Merck, 65%) and 1 mL perchloric acid (Merck, 70%) temperature. Digestion at room continued on a sand bath with a maximum temperature of 115 °C, for about 4 h and until the solutions were cleared. After cooling, the solutions were filtered through Whatman filter membranes and were diluted up to 10 ml in a volumetric flask with deionized distilled water. The digestion procedure followed the ROPME instructions (1989) and Pourang (1995, 1996).

A HP 4500 ICP-MS, equipped with Asx-520 Autosampler (England) was

used for the determination of metal concentrations in the samples. In order to control the quality of the measurement for each 25 samples, 2 blanks and 2 spiked samples were used. Instrumental detection limits for most metals was 0.02 µg/g, except for Cd $(0.01 \mu g/g)$ and Fe $(0.1 \mu g/g)$. The recoveries for Cd, Cu, Co, Cr, Ni, Mn, Zn and Fe were 84%, 107%, 79%, 94%, 101%. 86%, 72% 72%, and respectively.

Stable isotope analysis and trophic level calculation

The stable isotope signatures were used to calculate the trophic levels. Around 1 mg of homogenized powder of each sample was weighed in tin capsules. No further steps were taken for the elimination of carbonates and lipids because this helps us to avoid dispersion of $\delta^{13}C$ and $\delta^{15}N$.

All stable isotope measurements were performed at the University of California, Davis, USA. Isotope values were expressed in standard d-notation relative to the international V-PDB (Vienna PeeDee Belemnite) for carbon and atmospheric N₂ for nitrogen, using the standard equation;

 δ^{13} C or δ^{15} N = [(R _{Sample} /R _{Standard})-1]×1000

where R is ${}^{13}C/{}^{12}C$ or ${}^{15}N/{}^{14}N$.

The trophic level (TL) was calculated using the following equation: $TL_i = [(\delta^{15}N_i - \delta^{15}N_{pc})/3.4] + 2$

where TLi is the trophic level of fish species i, $\delta^{15}N_i$ represents the $\delta^{15}N$ of fish species i, $\delta^{15}N_{pc}$ represents the $\delta^{15}N$

of primary consumers (PC; i.e. bivalve C. glucaum), 3.4 and 2 are the mean δ¹⁵N trophic enrichment occurring per trophic level and the trophic position of the baseline organism, respectively. Commonly, the primary consumers preferred (TL₂)are to primary producers (TL₁) for evaluation of trophic levels (Cabana and Rasmussen, 1996; Vander Zanden et al., 1999; Vander Zanden and Rasmussen, 2001). We did not calculate the trophic levels of C. glucaum individuals because the stable isotopes of particle organic matter (POM) as its baseline had high variations in our sampling.

Statistical analyses

The Kolmogorov-Smirnov and Levene's tests were used to control the normality of distribution and homogeneity of variances, respectively. Simple linear regressions performed using ln-transformed metal concentrations in muscle vs. trophic levels. The slopes of regressions were used to determine the rate bioaccumulation through the body size of the Caspian fishes and bivalves. One way ANOVA and Tukey test were used to assess differences among the δ^{13} C, δ^{15} N values and size groups of fishes. statistical analyses The were not conducted when >50% of the observations were below the detection statistical limit. The tests were performed at the significance level of p<0.05 and the software SPSS version 13 for Windows was used.

Results

According to our results, the content of C and N in samples was the lowest in bivalves, C. glucaum while those amounts were the highest in fish A. chalcoides (Table 2). The nitrogen isotope ratios varied among the samples from C. glucaum (δ^{15} N=3.5 ‰) to *S. lucioperca* (δ^{15} N=13.1‰). The stable carbon isotope ratios varied from A. chalcoides $(\delta^{13}C = -25.2\%)$ to S. lucioperca (δ^{13} C = -17.9‰) (Fig. 1). The average $\delta^{15}N$ was nearly similar for three species R. frisii, L. aurata and A. chalcoides (Table 2). Among fish species, L. aurata and S. lucioperca had the lowest and highest TLs; with TLs average of 3.3 and 4.2. respectively.

The statistical test showed that only δ^{13} C had significant difference (F=6.4; p<0.05; ANOVA) with small size (total length<60 mm) and big size groups (total length>200 mm) of *A. chalcoides*.

Furthermore, the δ^{15} N and TL values in the small size of *R. frisii* (total length<140mm) was more than medium (total length 141-500 mm) and big size (total length>500 mm) (F=7.3 and F=5.5, respectively; p<0.05).

Cd, Cr, Co and Ni concentrations in most of fish samples were below the detection limits; while the concentrations were detected in most samples of bivalves, *C. glucaum* (Table 1). Furthermore, the *C. glucaum* were shown to have the highest metal concentrations. Among the fish samples, however the highest mean concentration of Cu was found in *S.*

lucioperca (average 9.3 μg g⁻¹), and the highest concentrations of most metals were observed in *L. aurata* (Table 1). The lowest concentrations of Mn and Fe were observed in *S. lucioperca* (average 0.7, 7.1 and 9.4 μg g⁻¹ dw), respectively.

There was strong correlation between body length and body weight (Spearman's correlation coefficient = 0.99, p<0.05). Therefore, body size was used instead of body length and weight (Table 3).

An increase in trophic levels of A. chalcoides and R. frisii vs. their body size increase was observed. Generally accumulation of metals increase in body size was observed in muscles of all species from different trophic levels (Figs 2 and 3). Sizedependent concentrations of metals were only observed for Cr and Fe in C. glucaum (Fig. 2). While no significant correlation were observed between metal concentrations and fish size (also TLs) of L. aurata, negative significant relationships were observed between Mn and Zn concentrations and fish size for A. chalcoides (Fig. 3). These relationships were also observed for Cu and Zn in S. lucioperca and for Mn in R. frisii. In our study, no significant decrease in concentrations of metals was observed with increasing trophic levels in fishes (Table 3, Fig 3).

Table 1: The biological characteristics and metal concentrations in different fishes and bivalve species collected from Anzali region ($C_{\rm m}$; number of samples for chemical analysis, F: female, M: male, U: undefined sex, D; percentage of detected sample numbers; the average metals concentrations have been calculated from detected data, minimum values (min) have been recorded from the minimum detected data).

		Weight (g)	Cd	Со	Cr	Cu	Mn	Ni	Zn	Fe
Species	С ш	Length (mm) Age (y)	Mean ± Sd (Min-Max) D	Mean ± Sd (Min-Max) D	Mean ± Sd (Min-Max) D	Mean ± Sd (Min-Max) D	Mean ± Sd (Min-Max) D	Mean± Sd (Min-Max) D	Mean ± Sd (Min-Max) D	Mean ± Sd (Min-Max) D
Percidae Sander lucioperca	16 (8F, 8M)	121- 549 250-365 2-3			0.77 ± 0.46 (0.36-1.47) 38%	9.3 ± 7.76 (0.22-24.7) 100%	0.77± 0.67 (0.01-0.59) 88%	1.16	23.3 ± 7 (11.2-38.9) 100%	7.12 ± 5.04 (2.1-14.6) 60%
Cyprinidae Rutilus frisii	25 (10F, 10M, 5U)	2-1248 65-500 1-6	0.54 3%		1.4 ± 0.34 (1.1-1.7) 9%	0.22 ± 0.67 (0.12-2.55) 30%	2.6 ± 3.8 (0.26-15.7) 82%	0.81 3%	26.8 ± 13.6 (11.2-69.7) 100%	140.1 ± 198.2 (2.9-548.1) 90%
Alburnus chalcoides	20 (6F, 8M, 6U)	2.1-159 70-264 1-3	0.48 5%			1.46 ± 1.97 (0.45-5.75) 70%	3.2 ± 4.1 (0.25-14.3) 85%	0.42 5%	38.5 ± 30.4 (8.9-98.9) 80%	80.9 ± 66.3 (21-178) 94%
Mugilidae Liza aurata	16 (3F, M, 8U)	2.1-545 68-372 1-5		1.34		3.6 ± 11.0 (0.1-40.1) 50%	3.2 ± 4.4 (0.4-12.7) 75%	1.9 ± 0.6 (1.4-2.3) 15%	27.5 ± 12.7 (11.7-51.2) 94%	217 ± 171 (61-510) 94%
Cardidae Cerastoderma glucaum	13	Height= 14.8-22.4 - 2-11	0.54 8%	1.44 ±0.38 (0.9-2.1) 85%	1.08±0.42 (0.44-1.62) 54%	8.94±5.52 (0.2-19.8) 100%	14.8±8.5 (5.4-35.8) 100%	15.5± 4.7 (10.4-24.9) 92%	60.4±14.2 (42.6-95.3) 100%	546.3±430.1 (96.2-1558) 100%

Table 2: The results of isotope analysis (C $_i$: number of samples for isotope analysis, TL_{PC} : trophic level based on primary consumer, TL_{Pom} : trophic level based on particulate organic matter, Averages with dissimilar letters (a, b, c) specify significant difference, F: statistic value, P: significant levels.

	C _i	Length classes (mm)	δ ¹³ C	$\delta^{15}N$	C amount (ug)	N amount (ug)	TL PC
Percidae		<300	-19.97 ± 0.1	12.9 ± 0.2			
Sander	9	300>	-19.96 ± 1.6	12.2 ± 0.8	463.9 ± 183.1	130.5 ± 48.1	4.2 ± 0.24
lucioperca			(F=0.01,P=0.99)	(F=2.6, P=0.15)			
		< 260	-21.7 ± 0.7	8.9 ± 0.4^{a}			
Cyprinidae	13	261-370	61-370 -20.4 ± 1.4 10.1 ± 0.9^{b}		326.7 + 220.7	007.550	3.4 ± 0.27
Rutilus frisii		370>	-21.6 ± 10.5	10.5 ± 0.2^{b}	320.7 ± 220.7	88.7 ± 55.0	3.4 ± 0.27
			(F=2.6, P=0.12)	(F=7.4, P=0.01)			
		<100	-24.2 ± 1.4^{a}	8.6 ± 1.9			
Alburnus	8	101-200	-21.9 ± 1.1^{ab}	10.6 ± 0.8	550.1 × 220.6	128.5 ± 68.9	3.7 ± 0.50
chalcoides		201>	-21.2 ± 0.2^{b}	11.7 ± 1.0	559.1 ± 330.6		
			(F=6.4, P=0.04)	(F=4.3, P=0.08)			
M:1: J		<100	-20.1 ± 0.2	8.2 ± 0.1			
Mugilidae	5	100>	-18.5 ± 0.6	10.1 ± 1.3	404.3 ± 229.6	117.9 ± 71.2	3.3 ± 0.42
Liza aurata			(F=6.3, P=0.14)	(F=1.4, P=0.42)			
Cardidae		Height <17	-20.3 ± 0.4	4.5 ± 0.43			
	9	(mm) 17-20	-21.3 ± 1.1	4.6 ± 1.6	267.9 : 121	63.9 ± 32.5	$\begin{array}{c} TL_{Pom} \\ 1.41 \pm 0.21 \end{array}$
Cerastoderma		20>	-21.7 ± 1.0	5.0 ± 0.1	267.8 ± 121		
glucaum			(F=2.1.P=0.2)	(F=0.4. <i>P</i> =0.69)			




Figure 1: Distribution of carbon and nitrogen stable isotope ratios among fishes and bivalves from southwest of the Caspian Sea.

Table 3: The Spearman correlation test for dependence of Ln metal concentrations on body size (length or weight) and trophic level (* significant level <0.05).

Species		Body size	Co	Cr	Cu	Mn	Ni	Zn	Fe
Sander lucioperca	Body size TL	-0.18			-0.65* -0.03	-0.05 0.12		-0.84* -0.03	0.97
Rutilus frisii	Body size TL	0.72*			0.86	-0.65* -0.57		-0.39 -0.48	0 -0.5
Alburnus chalcoides	Body size TL	0.83*			-0.29 -0.8	-0.54* -0.25		-0.61* -0.64	-0.6 -0.2
Liza aurata	Body size TL	0.67			0.14 0.6	-0.5 -0.86		-0.44 0.46	0.1 0.32
Cerastoderma glucaum	Body height TL	0.49	0.11 -0.11	-0.93* 0.4	-0.19 0.5	-0.55 -0.11	0.22 0.05	0.08 0.03	-0.63 -038




Figure 2: Regression between Ln metal concentration on height and trophic levels of bivalves (significant differences have been showed by lines).




Figure 3: Regression between Ln metal concentration on body length and trophic levels of fishes (significant differences slopes have been showed by lines; vertical lines in horizontal axis indicated the body size classes).

Table 4: Variations of metals concentrations in *Liza aurata* and *Rutilus frisii* in the Southern Caspian Sea as reported in previous investigations (*based on wet weight).

							`		0 /
Species	Cd	Co	Cr	Cu	Mn	Ni	Zn	Fe	References
Liza aurata	ND		0.43						Pazooki et al., 2009
	0.04					0.12			Fallah et al., 2011*
	0.02								Shahryari et al., 2010
	0.02			1.3			3.4		Ardalan et al., 2012
	0.03		0.08						Tabari et al., 2010
	0.63	ND		10.85		`0.41	217.8	125.6	Varedi, 2011
Rutilus frisii	0.001	0.009	0.33	1.01	0.45		17.2		Anan et al., 2005
				2.7			22.3		Foroughi Fard et al., 2008
	ND			2.76			15.4		Monsefrad et al., 2012
	0.17					0.52			Fallah et al., 2011*
	0.11								Hoseini and Tahami, 2012
	0.02								Shahryari et al., 2010
	0.325		ND	1.68	0.78	2.65	0.63	5.6	Eslami et al., 2011*
	0.04		0.06						Tabari et al., 2010
	ND	ND		0.013		ND	0.4	0.08	Varedi, 2011

ND: not detected

Discussion

The study showed that the concentrations of Mn, Zn and Fe in Rutilus frisii and Liza aurata were higher than those reported in previous studies (except Cu level; Tables 1 and 4). It might be related to temporal rise of the Caspian pollutants. According to Karpinsky (1992) the concentrations of metals in sediments of the Caspian Sea have shown high temporal increases. There are a few investigations showing variations in pollutant concentrations in the southern Caspian Sea. Among them, several authors have reported high variations in metals concentrations of water and substrate (Laloei, 2001; Parizanganeh et al., 2007; Hashemian kafshgari, 2008; Sohrabi et al., 2010).

While only Cu concentration had the highest concentrations in piscivorous *S. lucioperca*, no significant difference in metals was observed in deposit feeder *L. aurata* and benthivorous (*R. frisii* and *A. chalcoides*) (Table 1). According to Goodyear and McNeill (1999) the animals took up Zn in direct proportion to levels found in sediments while Cu had similar uptake rates from the water and sediment. Therefore the high concentration of Cu in *S. lucioperca* might be related to its different feeding habits.

The low concentration of Cr and Cd, similar to what found in previous studies (Tables 1 and 4), might be to their concentrations related sediments. The local sources might be responsible for metal pollution in the Caspian Sea. Similarly, Anan et al. (2005) found higher concentrations of Co, Cd in fishes from our study area than that found in other parts of the Caspian. Also higher concentrations of metals such as Cd were observed in the central and southern part of the Caspian Sea than those in the northern sector (de Mora et al., 2004) because of their rich natural background resource.

In our investigated species, the depleted and enriched $\delta^{15}N$ values can be due to their diets; i.e. deposit feeders and piscivorous, respectively (Table 1, Fig. 1). A. chalcoides has been known as a zooplankton feeder (Abbasi et al., 2004) but its mean TL value is relatively higher than benthivorous species R. frisii (Table Comparatively, R. frisii seems to have longer-distance movements and depleted δ^{15} N probably feeds on sources; while A. chalcoides exists in littoral areas where inshore eutrophication and subsequently the trophic base line is higher than offshore regions. The $\delta^{15}N$ values of primary consumers and other organisms in coastal areas would be increased after eutrophication and upward N loading phenomena (Cabana and Rasmussen, 1996; Martinetto et al., 2006). Asante et al. (2008) also found the higher $\delta^{15}N$ in fishes from shallow water rather than deep water. Furthermore, a broad range of δ^{13} C values were observed in *A. chalcoides* (Fig. 1). Hobson *et al.* (2002) reported that the high variation in δ^{13} C of benthivorous species might be related to the broad range of δ^{13} C in benthic compared to pelagic organisms. A higher δ^{13} C value observed in *L. aurata* (Fig. 1) also might be due to their diet that mostly contains the substrate invertebrates and the meiofauna (Ghadirnejad, 1996).

The significant slopes of metals concentrations, the fish size and the TLs showed that the metals do not bioaccumulate, but possibly biodilute by fish size and the TLs (Table 3, Fig. 2). Similarly, correlation between $\delta^{15}N$ and metals concentrations in fishes was not observed in other investigations (Campbell *et al.*, 2005; Asante *et al.*, 2008; Ikemoto *et al.*, 2008; Zhang and Wang, 2012).

A significant positive correlation between TLs and fish size was only observed in Cyprinid species (Table 3). It is probably related to change or variety of feeding habits for different sizes; while the Percidae and Mugilidae samples are known as piscivorous and deposit feeders, respectively. Atherin and kilka followed by Mugilidae were found to be the main food items of S. lucioperca (Abbasi et al., 2004). According to Vinagre et al. (2011), there was no correlation between the Carbon and Nitrogen stable isotopes of muscles and body size. It seems, there is a large range between $\delta^{13}C$ and $\delta^{15}N$ values in fish particularly A. chalcoides (Fig. 1) due to different food items size. The findings showed, the δ^{13} C value in the small size of A. chalcoides was significantly lower than in big sizes of the fish (F=6.4; p<0.05; Table 1). The small size of A. chalcoides might feed on the micro-zooplankton while the big size of this fish feed on the crustacean and macro-planktons (Abbasi *et al.*, 2004).

The low significant $\delta^{15}N$ (F=7.3; p<0.05; Table 1) in the small size groups of R. frisii as compared with the big size groups could be related to variations in their diets. Additionally, Zarinkamar (1996), Abdolmaleki et al. (2009) and Afraei Bandepei et al. (2009) noted that the big size groups of R. frissi feed on bivalves, while the diet of small size groups was crustacean in the Caspian Sea.

Although there were a strong relationship between TLs and body size in Cyprinid species (Table 3), there were no significant slopes between TLs and the Ln concentrations of metals (Fig. 2).

A significant negative relationship was found between the concentrations of Zn vs. body size for most fish species (Table 3). However, in the biomagnification of Zn documented in a few investigations on food web (Quinn *et al.*, 2003; Campbell *et al.*, 2005), there was also a significant negative correlation observed between $\delta^{15}N$ and Zn concentration (Yoshinaga *et al.*, 1992).

The negative relationship between body size and metal concentration in muscle (Table 3) has been previously reported in many Caspian fishes (Anan et al., 2005; Pourang et al., 2005; Pazooki et al., 2009). The relationships between metals concentrations in the muscle and length have been widely documented by several investigators. Some contradictory results have been discussed for fishes in the world by Pourang et al. (2005). Growth (body weight, length and age) dependent decreases in metals concentrations have been reported for many fish species particularly for Zn, Cu, Cd and Mn in muscle tissues (Anan et al., 2005). Significant negative correlations were found for Co, Zn, Cu in Rutilus caspicus (Anan et al., 2005). While Hoseini and Tahami (2012) found a significant relationship between Cd concentrations in muscle and R. frisii size, no significant relationship was observed for Cu and Zn concentrations by Foroughi Fard et al. (2008). Pourang et al. (2005) did not observe any age differences related in metal concentrations in different species; and only the concentration of cadmium had a negative relationship with weight of the samples (p=0.038). A significant positive relationship was found between bioaccumulation factors and fish age in Liza saliens (Fernandes et al., 2007).

Various environmental and biological factors, nourishment source and seasonal changes affect differences in metal bioaccumulation (Agusa et al., 2004; Wang Rainbow, 2008). The findings showed, there was no bioaccumulation of metals in S. lucioperca, A. chalcoides and L. aurata, so no significant differences

were observed between $\delta^{15}N$, TLs and the fish sizes.

The metabolic rate and dilution of metal during growth period can be responsible for this relationship; as smaller fish have higher metabolic rates and are able to accumulate metals, via food and water, more rapidly than larger fish (Anan *et al.*, 2005). The *R. frisii* had a significant difference in δ^{15} N among their different body sizes which could be related to difference in metal concentration in their food diet.

The mean metals concentrations in muscles the were lower than provisional tolerable daily/weekly/monthly intake of the metals, as established by the United States Environmental Protection European Food Agency, Safety Ministry Authority, of Agriculture **Fisheries** and Food. and FAO/WHO (the values presented by Fallah et al.(2011), Zhang and Wang (2012), Mashroofeh et al. (2013)).

The metals concentrations in the muscles of fishes were lower than those reported in the previous studies. These concentrations could not pose any threat to human health. In addition, no bioaccumulation was observed for metals in different trophic levels of *R. frisii* in the Caspian Sea. It is suggested that monitoring of metals in aquatic organisms and the Caspian environment should be performed to further discover those trends.

Acknowledgments

We wish to thank J. Hassan, N. Pourang, H. Babaei, J. Matthews and

M. Farhadinia for their help to do the laboratory work.

References

Abbasi, K., Sabkara, J., Rahimi, R. and Mahisefat, F., 2004. Feeding studies of Caspain fishes in southern Caspian Sea. Inland Water Auaculture Research Center, Bandar Anzali, 4P.

Abdolmalaki, S., Kor, D. and G., 2009. Bandani, Stock assessment of the bony fishes in Iranian coastal waters of the Caspian Sea (2005-2007). Iranian Fisheries Research Organisation, Inland Waters Aquaculture Research Center, 158P.

Abdolmaleki, S. and Ghaninezhad, D., 2007. Stock assessment of the Caspian Kutum *Rutilus frisii Kutum* in the Iranian coastal waters of Caspian Sea. *Iranian Journal of Fisheries Science*, 16(1), 103-114.

Afraei Bandepei, M., Mashhor, M., Abdolmalaki, S. and El-Sayed, M.A.F., 2009. Food and feeding habits of the Caspian kutum, *Rutilus frisii* kutum (Cyprinidae) in Iranian waters of the Caspian Sea. *Cybium*, 33, 193-198.

Agusa, T., Kunito, T., Tanabe, S., Pourkazemi, M. and Aubrey, D.G., 2004. Concentrations of trace elements in muscle of sturgeons in the Caspian Sea. *Marine Pollution Bulletin*, 49, 789-800.

Anan, Y., Kunito, T., Tanabe, S., Mitrofanov, I.and Aubrey, D.G., 2005. Trace element accumulation in

- fishes collected from coastal waters of the Caspian sea. *Marine Pollution Bulletin*, 51, 882-888.
- Ardalan, A., Sharif, A.A. and Farahani, Z., 2012. Trace elements accumulation and nutritive value in muscle tissue of golden mullet from the Kiashahr port (Caspian Sea). *Advances in Environmental Biology*, 6, 232-235.
- Asante, K.A., Agusa, T., Mochizuki, K., H., Ramu, Inoue, S., Kubodera. T., Takahashi, **S..** Subramanian, A.and Tanabe, S., 2008. Trace elements and stable isotopes (δ^{13} C and δ^{15} N) in shallow and deep-water organisms from the China Sea. Environmental Pollution, 156, 862-873.
- Broman, D., Rolff, C., Näf, C., Zebühr, Y., Fry, B. and Hobbie, J., 1992. Using ratios of stable nitrogen isotopes to estimate bioaccumulation and flux of polychlorinated dibenzo-p-dioxins (PCDDs) dibenzofurans (PCDFs) in two food chains from the northern Baltic. Environmental *Toxicology* and Chemistry, 11, 331-345.
- Cabana, G. and Rasmussen, J.B., 1996. Comparison of aquatic food chains using nitrogen isotopes. *Proceedings of the National Academy of Sciences*, 93, 10844-10847.
- Campbell, L.M., Norstrom, R.J., Hobson, K.A., Muir, D.C., Backus, S. and Fisk, A.T., 2005. Mercury and other trace elements in a pelagic Arctic marine food web (Northwater

- Polynya, Baffin Bay). Science of the Total Environment, 351, 247-263.
- De Mora, S., Sheikholeslami, M.R., Wyse, E., Azemard, S.and Cassi, R., 2004. An assessment of metal contamination in coastal sediments of the Caspian Sea. *Marine Pollution Bulletin*, 48, 61-77.
- Eslami, S., Moghaddam, A.H., Jafari, N., Nabavi, S.F., Nabavi, S.M. and Ebrahimzadeh, M.A., 2011. Trace element level in different tissues of *Rutilus frisii* kutum collected from Tajan River, Iran. *Biological Trace Element Research*, 143, 965-973.
- Fallah, A.A., Zeynali, F., Saei-Dehkordi, S.S., Rahnama, M. and Jafari. T.. 2011. Seasonal bioaccumulation of toxic trace elements in economically important fish species from the Caspian Sea using GFAAS. Journal für Verbraucherschutz. und Lebensmittelsicherheit, 6, 367-374.
- Fernandes, C., Fontainhas-Fernandes, A., Peixoto, F., and Salgado, M. A., 2007. Bioaccumulation of heavy metals in Liza saliens from the Esmoriz— Paramos coastal lagoon, Portugal. Ecotoxicology and Environmental Safety, 66(3), 426-431.
- Foroughi Fard, R., Esmaeli Sari, A. and Ghasempouri, S.M., 2008. The correlation of length and weight of Kutum (*Rutilus frisii kutum*) in the central south of Caspian Sea with copper and zinc concentration in muscle and liver tissues. *Iranian*

- Journal of Fisheries Sciences, 16 (4), 121-128.
- Gerami, M.H. and Dastbaz, M., 2013. Commercial fishing methods in Iran. World Journal of Fish and Marine Sciences, 5(1), 63-70.
- **Ghadirnejad, H., 1996**. Population dynamics of grey mullet species (*Liza aurata* and *L. saliens*) in the southern Caspian Sea. University of Wales Swansea, 207P.
- Glantz, M., 2000. Global environmental problems in the Caspian region. available on the Internet:

http://www.caspianstudies.com, 32P.

- Goodyear, K. and McNeill, S., 1999.

 Bioaccumulation of heavy metals by aquatic macro-invertebrates of different feeding guilds: a review. Science of the Total Environment, 229, 1-19.
- Hashemian kafshgari, A., 2008.

 Hydrology and hydrobiology and environmental pollutions in lower than 10 meters depths of Caspian Sea. Iranian Fisheries research organization, Caspian Sea Ecology Research Center, Sari, 103P.
- **Hobson, K.A., Fisk, A., Karnovsky, N., Holst, M., Gagnon, J.M. and Fortier, M., 2002.** A stable isotope (δ¹³C,δ¹⁵N) model for the north water food web: implications for evaluating trophodynamics and the flow of energy and contaminants. *Deep Sea Research Part II: Topical Studies in Oceanography*, 49, 5131-5150.

- Hoseini, H. and Tahami, M.S., 2012. Study of heavy metals (Pb and Cd) concentration in liver and muscle tissues of *Rutilus frisii Kutum*, Kamenskii, 1901 in Mazandaran Province. *Global Veterinaria*, 9(3) 329-333.
- Ikemoto, T., Tu, N.P.C., Okuda, N., Iwata, A., Omori, K., Tanabe, S., Tuyen, B.C. and Takeuchi, I., 2008. Biomagnification of trace elements in the aquatic food web in the Mekong Delta, South Vietnam using stable carbon and nitrogen isotope analysis. Archives of Environmental Contamination and Toxicology, 54, 504-515.
- Jardine, T.D., Kidd, K.A. and Fisk, **A.T..** 2006. Applications, considerations. and sources of uncertainty when using stable isotope analysis in ecotoxicology. Environmental Science and Technology, 40, 7501-7511.
- **Järup, L., 2003.** Hazards of heavy metal contamination. *British Medical Bulletin*, 68, 167-82.
- Karpinsky, M., 1992. Aspects of the Caspian Sea benthic ecosystem. *Marine Pollution Bulletin*, 24, 384-389.
- Kojadinovic, J., Potier, M., Le Corre, M., Cosson, R.P. and Bustamante, P., 2007. Bioaccumulation of trace elements in pelagic fish from the Western Indian Ocean. *Environmental Pollution*, 146, 548-566.
- Lake, J.L., McKinney, R.A., Osterman, F.A., Pruell, R.J.,

- **Kiddon, J., Ryba, S.A. and Libby, A.D., 2001.** Stable nitrogen isotopes as indicators of anthropogenic activities in small freshwater systems. *Canadian Journal of Fisheries and Aquatic Sciences,* 58, 870-878.
- **Laloei, F., 2001.** Investigation of hydrology and hydrobiology in the south of Caspian Sea. Iranian Fisheries Research Organization, Sari, Iran (In Persian).
- Le Loc'h, F., Hily, C. and Grall, J., 2008. Benthic community and food web structure on the continental shelf of the Bay of Biscay (North Eastern Atlantic) revealed by stable isotopes analysis. *Journal of Marine Systems*, 72, 17-34.
- Martinetto, P., Teichberg, M. and Valiela, I., 2006. Coupling of estuarine benthic and pelagic food webs to land-derived nitrogen sources in Waquoit Bay, Massachusetts, USA.
- Mashroofeh, A., Bakhtiari, A.R., Pourkazemi, M. and Rasouli, S., 2013. Bioaccumulation of Cd, Pb and Zn in the edible and inedible tissues of three sturgeon species in the Iranian coastline of the Caspian Sea. *Chemosphere*, 90, 573-580.
- Minagawa, M. and Wada, E., 1984. Stepwise enrichment of 15 N along food chains: Further evidence and the relation between δ^{15} N and animal age. *Geochimica et Cosmochimica Acta*, 48, 1135-1140.
- Monsefrad, F., Imanpour Namin, J. and Heidary, S., 2012.

- Concentration of heavy and toxic metals Cu, Zn, Cd, Pb and Hg in liver and muscles of *Rutilus frisii kutum* during spawning season with respect to growth parameters. *Iranian Journal of Fisheries Sciences*, 11, 825-839.
- Neamatparast, A., 2007. Studying of population structure, morphology, diet and growth of *Liza saliens* in Anzali sea shore. Sciences and research branch Islamic Azad University, Tehran, 147P. (In Persian).
- Parizanganeh, A., Lakhan, V. and Jalalian, H., 2007. A geochemical and statistical approach for assessing heavy metal pollution in sediments from the southern Caspian coast. International Journal of Environmental Science and Technology, 4, 351-358.
- Pazooki, J., Abtahi, B. and Rezaei, F., 2009. Determination of heavy metals (Cd, Cr) in the muscle and skin of *Liza aurata* from the Caspian Sea (Bandar Anzali). *Environoment Sciences*, 7, 21-32 (In persian).
- **Post, D.M., 2002.** Using stable isotopes to estimate trophic position: models, methods, and assumptions. *Ecology*, 83, 703-718.
- **Pourang, N., 1996.** Heavy metal concentrations in surficial sediments and benthic macroinvertebrates from Anzali Wetland, Iran. *Hydrobiologia*, 331, 53-61.
- Pourang, N., Tanabe, S., Rezvani, S.and Dennis, J., 2005. Trace elements accumulation in edible

- tissues of five sturgeon species from the Caspian Sea. *Environmental Monitoring and Assessment*, 100, 89-108.
- Quinn, M.R., Feng, X., Folt, C.L. and Chamberlain, C.P., 2003.

 Analyzing trophic transfer of metals in stream food webs using nitrogen isotopes. *The Science of the Total Environment*, 317, 73-89.
- ROPME, 1989. Manual of oceanographic and pollutant analysis. Regional Organization for the Protection of the Marine Environment, Kuwait, 290P.
- Sadeghirad, M., 2002. Determination of heavy metals (Zn, Cu, Cd, Pb, Hg) caviar and muscle tissue in Persian sturgeon (Acipencer and Stellate persicus) sturgeon (Acipenser stellatus) in southern Caspian sea. Iranian **Fisheries** Research organization, International sturgeon research institute, Rasht, 82P.
- **Sadeghirad, M., 2007.** Accumulation of trace elements in *Acipenser persicus* tissues in relation to feeding habits and mode of absorption of these pollutants. Iranian fisheries research organisation, Internatinoal sturgeon research institute, 42P.
- Shahryari, A., Golfirozy, K. and Noshin, S., 2010. Muscular concentration of cadmium and lead in carp, mullet and kutum of the Gorgan Bay, Caspian Sea. *Iranian Journal of Fisheries Sciences*, 19(2), 95-100.

- Sohrabi, T., Ismail, A. and Nabavi, M., 2010. Distribution and normalization of some metals in surface sediments from south Caspian Sea. Bulletin of Environmental Contamination and Toxicology, 85, 502-508.
- Tabari, S., Saravi, S.S.S., Bandany, G., Dehghan, A. and Shokrzadeh, M., 2010. Heavy metals (Zn, Pb, Cd and Cr) in fish, water and sediments sampled form Southern Caspian Sea, Iran. *Toxicology and Industrial Health*, 26, 649–656
- **UNEP/UNDP, 2011.** Caspian Sea State of Environment. 102P.
- Vander Zanden, M.J. and Rasmussen, J.B., 1996. A trophic position model of pelagic food webs: impact on contaminant bioaccumulation in lake trout. *Ecological Monographs*, 451-477.
- Vander Zanden, M.J. and Rasmussen, J.B., 2001. Variation in $\delta 15N$ and $\delta 13C$ trophic fractionation: implications for aquatic food web studies. *Limnology and Oceanography*, 2061-2066.
- Vander Zanden, M.J., Shuter, B.J., Lester, N. and Rasmussen, J.B., 1999. Patterns of food chain length in lakes: a stable isotope study. *The American Naturalist*, 154, 406-416.
- Varedi, S.E., 2011. Survey on envirnomental pollutants (heavy metals, hydrocarbons, surfactant and chlorinated pesticides) in southern part of the caspian sea. Iranian Fisheries research organization,

- Caspian Sea Ecology Research Center, Sari, 200P.
- Vinagre, C., Máguas, C., Cabral, H. and Costa, M., 2011. Effect of body size and body mass on δ^{13} C and δ^{15} N in coastal fishes and cephalopods. *Estuarine*, *Coastal and Shelf Science*, 95, 264–267.
- Wang, W.-X.and Rainbow, P.S., 2008. Comparative approaches to understand metal bioaccumulation in aquatic animals. Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology, 148, 315-323.
- Yoshinaga, J., Suzuki, T., Hongo, T., Minagawa, M., Ohtsuka, R., Kawabe, T., Inaoka, T. and

- Akimichi, T., 1992. Mercury concentration correlates with the nitrogen stable isotope ratio in the animal food of Papuans. *Ecotoxicology and Environmental Safety*, 24, 37-45.
- **Zarinkamar, H., 1996.** Survey feeding habits of Kutum in coastal of Bandar Anzali. Islamic Azad University, Tehran, 150P. (In Persian).
- Zhang, W.and Wang, W. X., 2012. Large-scale spatial and interspecies differences in trace elements and stable isotopes in marine wild fish from Chinese waters. *Journal of Hazardous Materials*, 215, 65-74.