Age, growth and mortality of Caspian Spirlin, *Alburnoides* eichwaldii (De Filippi, 1863), from Aras River Basin in Turkey

Cicek E. 1*; Sigirci U.1; Birecikligil S.1; Saylar Ö2.

Received: February 2015 Accepted: October 2015

- 1- Department of Biology, Faculty of Art and Sciences, Nevşehir Hacı Bektaş Veli University, Nevşehir, TURKEY
- 2-Division of ScienceTeaching, Faculty of Education, Gazi University, Beşevler, 06500 Ankara, TURKEY

Keywords: Caspian spirlin, *Alburnoides eichwaldii*, Aras River, Çıldır Lake, Balık Lake

Introduction

The genus Alburnoides belongs to the Leuciscine cyprinids and is distributed in Europe and eastern parts of Asia. two subspecies have consistently reported for Alburnoides bipunctatus (Heckel, 1858) Turkey (Kuru, 2004), A. bipunctatus eichwaldii and A. bipunctatus fasciatus. Just a few years ago, some taxonomic on work focusing the Alburnoides reported 4 new species from Turkey (Turan et al., 2013; 2014).

Alburnus eichwaldii (De Filippi, 1863), described from the Kurpresso Tiflis (Kura River near Tbilisi, Georgia), is usually regarded as a Caspian Sea basin subspecies of *A. bipunctatus* (Bogustkaya, 1997; Nazari et al., 2009). Recently, the species was considered to be a full species (Fricke et al., 2007).

A. eichwaldii inhabits both lentic and lotic water bodies, with welloxygenated clear water (Stepanyan et al., 2013). Because of low tolerance to water pollution it is considered as a biological indicator good environmental quality (Abbasi et al., 2013a). A. eichwaldii does not have economic value however it plays an important role because it is important prey item of many predator fish species while its main food items are plankton and some invertebrates, mostly diptera and ephemeroptera (Abbasi *et al.*, 2013a).

There are several papers dealing with the Karyotype (Stepanyan *et. al.*, 2013) in Armenia and diet (Abbasi *et al.*, 2013a), hematological parameters (Kohanestani *et al.*, 2013), body shape (Eagderi *et al.*, 2013), some population parameters (Seifali *et al.*, 2012; Abbasi

^{*}Correspondingauthor's Email: erdogancicek@yahoo.com

et al., 2013b; Monajjemi et al., 2014; Nowferesti et al., 2014) in Iran. However, there has not been any study conducted on population dynamic parameters of the species in Turkey. Therefore this study aimed investigate population parameters including age, growth, mortality and exploitation rates of A. eichwaldii from Aras River basin in Turkey.

Material and methods

This investigation was carried out on August 2-6, 2014 and September 25-29, 2014 from Aras River Basin at stations in both rivers and lakes (Çıldır, Balik and Aygir). A total of 162 specimens were caught using a backpack electrofisher (SAMUS 725MP) in the river and 90 specimens were caught using multi-mesh gillnets (5, 6.25, 8, 10, 12.5, 15.5, 19.5, 24, 29, 35, 43, 55 mm, knot to knot) in the lakes. Collected specimens were fixed in 10% formalin, transferred to the laboratory and stored in 70% ethanol for further processing. Five of the specimens were cataloged (NHVIC 201408080) in the Ichthyology Collections of Nevsehir **BektasVeli** University, Department of Biology, Nevsehir. Turkey.

Morphometric measurements and meristic counts based on Bogutskaya and Coad (2009) were made for comparison with other studies. The taxonomic key given by Berg (1949a, b), Bogutskaya and Coad (2009), Coad and Bogustkaya (2009) were used to identify the samples to the lowest

taxonomic level. Morphometric characters were taken with digital calipers to the nearest 0.1 mm and meristicones were counted under magnifier or stereo microscope. All measurements and counts were made on the left side of the fish.

In order to determine the population parameters, the total length and weight were determined to the nearest 1 mm and 0.01 g respectively. The scale samples were removed from the left side of the fish body, from ventral to the dorsal fin for the age determination. Scales were soaked in water and examined under the stereo binocular microscope, independently twice with no reference to the previous readings and without any knowledge of the length or weight of the fish. The precision was measured by the percentage of agreement between the two readings (Chang, 1982). The assessment of age was based on the determination of the number of annuli on each scale.

The length-frequency data were plotted with 1 cm length intervals. The length-weight relationships were determined according to the power equation given by Sparre and Venema (1998): $W=a*L^b$. In this equation, W is total weight, a and b are regression constants and L is total length. Growth in length and weight were expressed in terms of the von Bertalanffy equation $L_t=L_{\infty}[1-e^{-k(t-to)}]$. The growth parameters L_{∞} , k and t_o were estimated using the Least Squares Method recommended by Sparre and Venema (1998).

Correspondence between empirical data and an expected distribution was tested by Khi^2 test. The b value was tested by t-test to verify whether it was significantly different from the isometric growth (b: 3).

The growth performance index (Φ') was calculated using the formula (Pauly and Munro, 1984): $\Phi = \log k + 2\log L_{\infty}$. Fulton's condition factor (K) was calculated by the following equation: $K = 100 \frac{W}{L^{b}}$ where; W: total weight, L: total length, b: regression constant (Sparre and Venema, 1998).

The instantaneous rate of total mortality coefficient Z was estimated using Beverton and Holt's Z Equation (1956): $Z = k \frac{(L_{\infty} - L)}{(L - L')}$, where; \bar{L} is the mean length of the entire catch, L' is the lower limit of corresponding length intervals (Sparre and Venema, 1998). The natural mortality coefficient (M)was estimated following Pauly's empirical formula (Pauly, 1980), linking the natural mortality with the von Bertalanffy parameters, L_{∞} (cm), kand mean annual temperature (T,°C) of water in habitat (in this case 9.3°C): Log10M = -0.0152-

 $0.279\log 10L_{\infty}+0.6543 \log 10k+0.463 \text{Lo}$ g10*T*. Fishing mortality rates (*F*) was calculated as the difference between *Z* and *M* (*F*=*Z*-*M*). The value of the average annual exploitation rate (*E*) was obtained by E=F/Z (Sparre and Venema, 1998).

Wilcoxon signed-rank test was performed to compare river and lakes samples. Chi-Square test (χ^2) was used

to compare observed and estimated parameters.

Results and discussion

Age, frequency distributions in length and weight of *A. eichwaldii* are presented in Table 1. There were significant differences in the parameters between the specimens of river and lake (p<0.05). Therefore, all analyses and evaluations were performed separately for river samples and lake samples.

Age of *A. eichwaldii* varied from 0 to III year age groups and most frequent groups for river and lakes were 0 (42.0%) and I (33.3%) year age groups, respectively. The total length ranged between 3.2 and 13.0 cm (8.98±2.11 cm) for river and 7.3 and 13.0 cm (9.63±2.08 cm) for lake samples. It was evident that *A. eichwaldii* grew rapidly in their first year after which growth rate declined (Table 1).

Monajjemi et al. (2014) reported that age of A. eichwaldii varied from 0 to III year age groups and the most frequent groups was the age group 0 in Shirud River (Iran). The maximum total length and weight observed was 111.4 mm and 15.5 g with the mean of 68.33±16.92 mm and 4.9 ± 3.2 respectively in this study. distribution has been revealed as 0-II, and the length distribution as 2.53-11.02 cm according to Abbasi (2013b) in Tilaban Creek. In another study the length range was reported as 2.5-11.02 cm (n: 47) in Tilaban Creek, 4.06-11.68 cm (n: 66) in Shirabad Creek and 5.47-11.15cm in Kaboodval Creek (Abbasi et al., 2013a).

Table	1:	Age,	length	and	weight-frequency	$\ distribution$	of	Caspian	spirlin,
	F	Alburn	oides ei	chwa	<i>ldii</i> , in Aras River I	Basin.			

				River San	ıples		
A	N	0/		Total Length	(cm)	Total W	eight (g)
Age	IN	%n	Mean	Range	Growth rate (%)	Mean	Range
0	68	42.0	6.97±1.33	3.2-9.8		3.85±2.43	0.21-9.20
I	44	27.2	9.48 ± 0.67	8.0-11.6	36.00	9.13 ± 2.05	5.15-15.52
II	30	18.5	10.90 ± 0.62	9.9-12.3	14.96	14.33 ± 2.65	9.77-19.94
III	20	12.3	11.77±0.85	10.0-13.0	7.90	18.16±3.39	11.12-23.17
Σ	162		8.98 ± 2.11	3.2-13.0		8.99 ± 5.68	0.21-23.17
				Lake Sam	ples		
0	9	16.7	5.54±0.91	7.3-9.0		2.50±1.28	4.01-7.39
I	30	33.3	9.45 ± 0.82	9.0-10.5	17.28	12.12±3.26	7.47-13.58
II	24	26.7	11.25±0.81	9.5-12.0	13.43	20.05 ± 4.73	8.12-24.28
III	21	23.3	14.75 ± 2.33	11.6-13.0	11.22	39.31±15.27	19.51-27.67
Σ	90		9.63 ± 2.08	7.3-13.0		13.96±7.91	4.01-27.67

Growth performance, condition factor, mortality and exploitation rate of the Caspian spirlin have been reported in a few studies.

Growth performance index (Φ') was estimated as 4.11 by Seifali *et al.* (2012) and 4.24 Abbasi *et al.* (2013b).The index was not estimated for this species previously.

There is a striking similarity among all river populations whereas the parameters estimated for lake were quite different from others. The results showed a significant difference in *A. eichwaldii* population inhabiting the river and the dam in terms of body shape.

The observed differences were related to head, snout, caudal peduncle regions and eye diameter that could be an adaptation to altered habitat and hydrological conditions of upstream and downstream due to damming (Eagderi *et al.*, 2014).

For a long time, the influence of environmental factors on fish has been studied in respect to their effects on fish growth. Because fish are ectotherms, growth of fish is highly dependent on temperature. But other factors are also involved in the control of physiological functions (Boeuf and Le Bail, 1999). Consequently, all of these factors have impact on population parameters. The length-weight relationship and von Bertalanffy growth parameters for A. eichwaldii are presented in Fig. 1. The relationship determined as $W=0.00644L^{3.2221}$ (95% CI of b: 3.151-3.293) for rives and $W=0.00651L^{3.3714}$ (95% CI of b: 3.225-3.518) for lakes. The b value was not significantly different than 3.0 (p<0.001), which indicates isometric growth of A. eichwaldii.




Figure 1: Length-weight relationship of Caspian spirlin, *Alburnoides eichwaldii*, sampled from the stations of river (a) and lakes (b) in Aras River Basin.

Table 2: Length-weight relationship and von Bertalanffy growth parameters for Caspian spirlin, *Alburnoides eichwaldii* (*females, **males, ***combined sexes)

В	а	L_{∞} (cm)	k (year ⁻¹)	t_{θ} (year)	Ф′	K	Habitat	Country	References
3.122***	0.000006	10.45	1.19		4.11		River	Iran	Seifali et al. 2012
2.919*	0.005	12.45	0.248	-0.39	4.24		River	Iran	Abbasi et al., 2013b
2.876**	0.005	11.66	0.281	-0.39			River	Iran	Abbasi et al., 2013b
2.944***	0.0145	12.08	0.55	-0.47			River	Iran	Monajjemi et al., 2014
3.290***	0.0046						River	Iran	Nowferesti et al., 2014
3.221***	0.0064	12.91	0.548	-1.41	1.96	1.08	River	Turkey	This study
3.371***	0.0065	16.36	0.237	-2.89	1.80	1.19	Lake	Turkey	This study

The length-weight relationship reported from Iran is given in Table 2. As can be seen in the table, b value was close to 3 or above 3. The highest value (3.371) was estimated to be lake population in The length-weight this study. relationship may be influenced by sex, maturity, geographical location and environmental conditions in the given year (Weatherley and Gill, 1987). Additionally, sampling bias due to the collection method could influence the size frequency distribution and finally the estimation of parameters.

Considering the values determined in the previous studies, all values were higher than that of this study (Table 2). When reporting as the annual average water surface temperature of 12.46 °C in Iran, it has been measured at approximately 5-6°C in this studied area. Certain environmental factors such as water temperature, food supply, poor water quality, physical disturbance and biology of fish such as maturity and hormones have an obvious and major influence on growth rate (Kapoor and

Khanna, 2004). Therefore the difference among studies linked with these factors.

Instantaneous total (Z), natural (M) and fishing (F) mortalities were estimated as 0.88, 0.75 and 0.13 year⁻¹, respectively for river samples and 0.39, 0.37 and 0.02 year⁻¹, respectively for lake samples. The exploitation rate (E) was calculated as 0.15 and 0.05 for river and lake samples, respectively.

Mortality and exploitation rates were estimated by Seifali *et al.* (2012) and Monajjemi *et al.*, (2014) given in Table 3. There was over fishing pressure on the population of *A. eichwaldii in* both of these studies. However over fishing pressure was not observed neither in the river nor in the lake station in this study. Indeed, *A. eichwaldii* population inhabiting in rivers or lakes is not commercially fished in Aras River Basin in Turkey.

Z	M	F	E	Habitat	Country	References
3.40	0.97	2.43	0.71	River	Iran	Seifali et al. 2012
2.41	1.19	1.22	0.51	River	Iran	Monajjemi et al., 2014
0.88	0.75	0.13	0.15	River	Turkey	This study
0.39	0.37	0.02	0.05	Lake	Turkey	This study

Table 3: Mortality rates (year-1) of Caspian spirlin, Alburnoides eichwaldii.

Acknowledgements

A special thanks to my students Selda Öztürk, Burak Seçer, Yasemin Celepoğlu, Muhammed Kelleci, Batuhan Keskin and Elçin Keşir for their assistance in the laboratory. The specimens were collected during the survey of "Establishment of Country Specific Our Project Ecological Water Quality Evaluation System" conducted by the Turkish Republic, Ministry of Water Affairs and Forestry.

References

Abbasi, F., Ghorbani, R. and Yelghi, S., 2013a. Comparative analysis of the diet of the spirlin (*Alburnoides eichwaldii*) in the Tilabad, Shirabad and Kaboodval Streams Golestan Province, Iran. *World Journal of Fish and Marine Sciences*, 5, 79-83.

Abbasi, F., Ghorbani, R., Yelghi, S., Hajimoradloo, A. andFazel, A., 2013b. Study of some growth parameters of spirlin (Alburnoides eichwaldii), in Tilabad Stream of the Golestan Province. Journal of Fisheries, 6, 59-70.

Berg, L.S., 1949a. Freshwater Fishes of the U.S.S.R. and Adjacent Countries. 2, Israel Program for Scientific

Translations 1964 (Translated from Russian), Jerusalem.

Berg, L.S., 1949b. Freshwater Fishes of the U.S.S.R. and AdjacentCountries. 3, Israel Program for Scientific Translations 1964 (Translated from Russian), Jerusalem.

Boeuf, G. and Le Bail, P.Y., 1999. Does light have an influence on fish growth? *Aquaculture*, 177, 129-152.

Bogustkaya, N.G. and Coad B.W., 2009. A review of vertebral and fin-ray the counts in genus Alburnoides (Teleostei: Cyprinidae) with species. description of six new Zoosystematica Rossica, 18, 126-173.

Chang, W.Y.B., 1982. A statistical method for evaluating the reproducibility of age determination. Canadian Journal of Fisheries and Aquatic Sciences, 39, 1208-1210.

Coad B.W. and Bogustkaya, N.G., 2009. *Alburnoides qanati*, a new species of cyprinid fish from southern Iran (Actinopterygii, Cyprinidae). *Zookeys*, 13, 67-77.

Eagderi, S., Esmaeilzadegan, E. and Maddah, A., 2013. Body shape variation in riffle minnows (Alburnoides eichwaldii De Filippii, 1863) populations of Caspian Sea

- basin. *Taxonomy and Biosystematics*, 5, 1-9.
- Eagderi, S., Esmaeilzadegan, E. and Pirbeigi, **2014.** Morphological A., Capoetagracilis responses of Alburnoides eichwaldii populations (Cyprinidae) fragmented due to Tarik Dam (Sefidrud River, Caspian Sea Iranian basin. Iran). Journal Ichthyology, 1, 114-120.
- Fricke, R., Bilecenoğlu, M. and Sarı, H.M., 2007. Annotated checklist of fish and lamprey species (Gnathostomata and Petromyzontomorphi) of Turkey, including a Red List of threatened and declining species. Stuttgarter Beiträgezur Naturkunde, 706, 1-169.
- **Kapoor, B.G. and Khanna, B., 2004.** Ichthyology handbook. Springer-Verlag Berlin, 1063P.
- Kohanestani, Z.M., Hajimoradloo, A., Ghorbani, R., Yulghi, S., Hoseini, A. and Molaee, M., 2013. Seasonal variations in hematological parameters of *Alburnoides eichwaldii* in Zaringol Stream-Golestan Province, Iran. *World Journal of Fish and Marine Sciences*, 5, 121-126.
- **Kuru, M., 2004.** Recent Systematic Status of Inland Water Fishes of Turkey. *GÜ Gazi Eğitim Fakültesi Dergisi*, 24, 1-21.
- Monajjemi, M., Ghorbani, R., Vesaghi, M.J., Rajabi, A.R.N., 2014. Age structure, growth and mortality index of spirlin (*Alburnoides eichwaldii* De Filippii 1863) in Shirud River, Mazandaran Province. *Journal of*

- Fisheries and Sciences and Technology, 2, 67-72.
- Nazari, S:, Pourkazemi, M. and Porto Rebelo, J.I., 2009. Comparative karyotype analysis of two Iranian cyprinids, *Alburnoides bipunctatus* and *Alburnus filippii* (Cypriniformes, Cyprinidae). *Iranian Journal of Animal Biosystematics*, 5, 23-32.
- Nowferesti, H., Asgardun, S., Zare, P., 2014. Length-weight relationships of six freshwater cyprinid fishes of Iran. *Journal of Applied*, 30, 1069-1070.
- **Pauly, D. and Munro, J.L., 1984.** Once more on the comparison of growth in fish and invertebrates. *Fishbyte*, 2, 21, 1984.
- Pauly, D., 1980. On the inter relationships between natural mortality, growth parameters, and mean environmental temperature in 175 fishstocks. *Conseil International pourl'Exploration de la Mer*, 39, 175-192.
- Seifali, M., Arshad, A., Amin, S.M.N., Kiabi, B.H., Esmaeili, H.R. and Moghaddam, F.Y., 2012. Population dynamics of Caspian spirlin (Actinopterygii: Cyprinidae) in the Kesselian Stream, Iran. *African Journal of Biotechnology*, 11, 9214-9222.
- Sparre, P. and Venema, S.C., 1998.
 Introduction to tropical fish stock assessment. Part 1. Manual. FAO Fisheries Technical Paper. No. 306.1 Rev.2, Rome, FAO. 407P.
- Stepanyan, I., Vasilyan, E., Pipoyan, D. and Kh., S., 2013. Karyotype of the riffle minnow *Alburnoides bipunctatus* eichwaldii (Filippi, 1863)

(Cypriniformes, Leuciscinae) from the Hrazdan River. *Biological Journal of Armenia*, 65, 79-84.

- Turan, D., Ekmekçi, F.G., Kaya, C. and Güçlü, S.S., 2013. Alburnoides manyasensis (Actinopterygii, Cyprinidae), a new species of cyprinid fish from Manyas Lake basin, Turkey. ZooKeys, 276, 85-102.
- Turan, D., Kaya, C., Ekmekçi, F.G. and Doğan, E., 2014. Three new species of *Alburnoides* (Teleostei: Cyprinidae) from Euphrates River, Eastern Anatolia, Turkey. *Zootaxa*, 2 (3754), 101-116.
- Weatherley, A.H. and GillH. S., 1987. The biology of fish growth. Academic Pres Inc. London, 443P.