Shape variation of the Indian Ocean slender mudskipper, Scartelaos tenuis (Day, 1876) from the Persian Gulf and Oman Sea (Gobioidei: Gobiidae)

Ghanbarifardi M.^{1*}; Aliabadian M.^{2,3}; Esmaeili H.R.⁴

Received: January 2017

Accepted: September 2017

Abstract

The shape diversity among populations of *Scartelaos tenuis* (Day 1876) from five stations (194 specimens) in the Persian Gulf, Strait of Hormuz (Qeshm Island), and the Oman Sea were investigated. The shape variation was assessed by geometric morphometrics, using 14 morphological landmarks. The two sexes were not distinguishable by the shape. Fish samples were classified into two distinct groups, the fish from the Persian Gulf (Heleh and Asaluyeh), and the specimens of Qeshm Island and the Oman Sea (Chahoo, Dargahan and Jask). The Persian Gulf and the Oman Sea are characterized by specific environmental parameters of surface waters, including temperature and salinity. The shape discrepancies of *Scartelaos tenuis* specimens from different stations are probably because of ecological parameters of the Persian Gulf and Oman Sea. Our results suggest that the observed pattern of morphological variation is almost some kind of polyphenism induced by distinct environmental factors.

Keywords: Geometric morphometrics, Environmental factors, Polyphenism, *Scartelaos tenuis*, Persian Gulf, Oman Sea

1-Department of Biology, Faculty of Sciences, University of Sistan and Baluchestan, Zahedan, Iran

2-Department of Biology, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran

3-Research Department of Zoological Innovations (RDZI), Institute of Applied Zoology, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran.

4-Ichthyology and Molecular Systematics Research Lab., Department of Biology, College of Sciences, Shiraz University, Shiraz 71454, Iran

*Corresponding author's Email: Meghfardi@gmail.com

Introduction

Three species of mudskippers are distributed in the Persian Gulf and Oman including Sea. **Boleophthalmus** dussumieri, Periophthalmus waltoni, and Scartelaos tenuis (Ghanbarifardi and Malek, 2007). Based on recent molecular phylogenetic trees for mudskippers, the Indian Ocean slender mudskipper, Scartelaos tenuis (Day, 1876) is included the "Periophthalmus lineage" in of gobionelline-like gobies (Gobiidae) and sister taxa for Boleophthalmus genus; moreover, ecomorphological adaptations of three Iranian mudskippers have been examined by geometric morphometric method (Ghanbarifardi et al., 2016; Polgar et al., 2017). S. tenuis specimens from Chabahar and Qeshm Island have been compared using meristic and morphometric characteristics; this study proves some differences between fish from stations regarding length-weight two relationship and condition factor (Hashemi et al., 2015). Geometric morphometric and traditional morphometric methods have employed investigate been to the morphologic diversity among various populations of P. waltoni (Ghanbarifardi et al., 2014a,b). These two studies prove that P. waltoni populations from the Persian Gulf differ from those of the Oman Sea the morphological regarding characteristics. Age, growth (Abdoli et al., 2010) and feeding habits (Abdoli et al., 2012) of Scartelaos tenuis have been surveyed. However, there is no study to compare different populations of this species in a wide geographical range.

Geometric morphometrics (GM) is a method to compare the shape of organisms (Rohlf and Marcus, 1993). Two- or three-

dimensional coordinates of biologically landmarks definable are used for Landmark-based geometric morphometric Superimposition methods. approaches remove non-shape variation (variation in position, orientation, and scale of the specimens) in configurations of landmarks by overlaying them and employing some optimization criteria. Generalized Procrustes Analysis (GPA) is usually used to superimpose landmark configurations and remove the non-shape variation. After the non-shape diversity has been eliminated, the shape variables could be compared via multivariate statistics, and graphical representations of shape (thinplate spline transformation grids) are generated for comparison. GM presents shape diversification, and it is employed as a substitute to the linear measurements, counts, and ratios used in conventional morphometrics (Adams et al., 2004). GM has been used to study fish ecology, population biology and stock discrimination (Silva, 2003; Costa and Cataudella, 2007; Mostafa et al., 2010). It has also been established that several environmental elements have transiently or permanently affected the morphological traits of fish. Temperature (Loy et al., 1996a), fish body condition (Fitzgerald et al., 2002), adaptations to specific habitat types (Langerhans et al., 2003), predation risk (Eklov and Svanback, 2006), and salinity 1996b) (Lov al., et are among the most important ecological f actors.

The Persian Gulf is a limited waterbody, linked to the Oman Sea via the narrow Strait of Hormuz (Fig. 1); therefore, its water transmission with the Oman Sea is restricted. The average depth of the Persian Gulf is only ~35 m. The surface water of the Persian Gulf experiences broad seasonal temperature alterations and superior salinity readings because of the low rainfall and high evaporation values (Yao, 2008). On the other part, the Oman Sea is widely connected to the Indian Ocean and much deeper than the Persian Gulf (~1000 m). Enough circulation, direct connection with the Indian Ocean and confined extension of shallow regions are three characters of the Oman Sea that make ecological conditions of this waterbody steadier than those of the Persian Gulf (Reynolds, 1993).

This survey explores shape variation among populations of *S. tenuis* from five stations in the Persian Gulf, Strait of Hormuz (Qeshm Island) and Oman Sea, utilizing GM.

Materials and methods

One hundred and ninety four specimens of *S. tenuis* (standard length > 70 mm) were collected from five stations (Heleh 39, Asaluyeh 34, Chahoo 38, Dargahan 39,

Jask 44) along the Iranian coasts of the Persian Gulf and Oman Sea, in February 2013 (Fig. 1, Table 1). The specimens are transferred to the Zoological Museum of Ferdowsi University of Mashhad (ZMFUM, Table 1). Specimens were caught by hand net during low tide and preserved in 5% formalin. The urogenital papilla is pointed in males and spatulate in females, and this character is used to sex determination of all fish (Miller, 1986; Ghanbarifardi et al., 2014a). To evaluate the trustworthiness of this method, 20 accidentally chosen specimens were also sex determined by dissection. The left side of each sample was photographed using a Canon SX150 digital camera mounted on a bench-top camera stand. Fish images were converted into TPS files with tpsUtil v. 1.38 (Rohlf, 2006), and 14 definable homologous landmarks (Fig. 2) were digitized from each image using tpsDig v. 2.17 (Rohlf, 2013).

 Table 1: Sampling coordinates, area, number and call numbers of Scartelaos tenuis specimens collected at five stations.

Station	Coordinates	Area	Specimens	ZMFUM call numbers
Heleh	29° 14.93' N 50° 40.72' E	Persian Gulf	13 females, 26 males	104 hs-142hs
Asaluyeh	27°26'N 52°40'E	Persian Gulf	15 females, 19 males	3as-36as
Chahoo	26° 41.47' N 55° 28.75' E	Qeshm Island	18 females, 20 males	143qws-149qws
		(Strait of		18qws-49qws
		Hormuz)		
Dargahan	26° 58.39' N 56° 4.85' E	Qeshm Island	19 females, 20 males	50qes-88qes
		(Strait of		
		Hormuz)		
Jask	25° 33.85' N 58° 10.61'E	Oman Sea	21 females, 23 males	100js-118js
				74js-99js



Figure 1: Sampling stations. 1: Heleh, 2: Asaluyeh, 3: Chahoo, 4: Dargahan, 5: Jask.

Landmark data from each specimen was aligned using Generalized Procrustes (GPA) Analysis for superimposition (Rohlf, 1999) with MorphoJ v. 1.04a (Klingenberg, 2011). The GPA method translates, rotates, and scales all landmarks to eliminate the non-shape variation and size factor (Zelditch et al., 2004). After Procrustes analysis, discriminant function analysis (DFA) was carried out with MorphoJ v. 1.04a to assess any shape difference between sexes. Canonical variate analysis (CVA) of landmark data was performed to evaluate any significant dissimilarity in shape among fish samples. CVA was conducted using SPSS 20.0 (© IBM); for CVA, 34 specimens of each station were used. To illustrate body shape variation, thin-plate spline transformation grids and lollipop plot (vectors showing transpositions from mean landmark positions) are generated with MorphoJ v. 1.04a. In this method, the shape changes from mean landmark positions are depicted as functions of their position along CV axes (e.g. CV1).

Results

DFA method could not distinguish significant difference in shape between females and males of S. tenuis (Table 2). specimens Therefore, all of each population were considered as a whole and CVA was conducted. The thin-plate spline (transformation grids) and lollipop plot along CV1 were used to depict the deformation in shape. The shape of the snout (Landmark 1), the eye situation (landmarks 3 and 4) and the position of the first and second dorsal fins and anal fin (landmarks 5, 6, 7, 8, 10 and 11) are disfiguring for the most (Fig. 3). The initial three canonical variates (CVs) achieved 66.0%, 19.1% and 10.6% of total variance, correspondingly. CV1 obviously detaches Chahoo, Dargahan (Oeshm Island) and Jask (Oman Sea) from Heleh and Asaluyeh (Persian Gulf) (Fig. 4). Totally, 91.8% of the grouped specimens of S. tenuis from different stations were properly categorized by CVA (Table 3).

	~~~~~~,			
Heleh P-value: 0.1628		Predicted group	Total	
		female male		
Count	female	8	5	13
Count	male	8	18	26
Asaluyeh P-value: 0.2801		female	male	
		Termate	niaic	
Count	female	7	8	15
	male	9	10	19
Chahoo				
P-value:	0.0913	female	male	
Counts	female	9	9	18
Counts	male	10	10	20
Dargahan				
P-value:	0.8694	female	male	
Counts	female	7	12	19
Counts	male	11	9	20
Jask				
P-value:	0.6968	female	male	
Counts	female	8 13		21
Counts	male	12	11	23

#### Table 2: DFA of male and female specimens of Scartelaos tenuis from different sampling stations, based on GM data.

Table 3: CVA of Scartelaos tenuis from different populations. The percentage of specimens correctly classified to each station is written in bold. Overall, 94.3% of the specimens from different stations were accurately classified.

	Heleh	Asaluyeh	Chahoo	Dargahan	Jask
Heleh	85.3	11.8	2.9	0.0	0.0
Asaluyeh	14.7	85.3	0.0	0.0	0.0
Chahoo	0.0	0.0	97.1	2.9	0.0
Dargahan	0.0	0.0	11.8	79.4	8.8
Jask	0.0	0.0	0.0	11.8	85.3



Figure 2: 14 morphological landmarks shown on *Scartelaos tenuis*. 1: anterior tip of the snout on the upper jaw. 2: rictus. 3 and 4: eye's anterior and posterior edges, correspondingly. 5 and 6: anterior and posterior points of insertion of the first dorsal fin, respectively. 7 and 8: anterior and posterior points of insertion of the second dorsal fin, sequentially. 9: ventral point of insertion of the caudal fin. 10 and 11: posterior and anterior points of insertion of the anal fin, respectively. 12: Ventral point of attachment of the left branchiostegal membrane. 13 and 14: anterior and posterior points of insertion of the left pelvic fin (landmarks are named after Ghanbarifardi *et al.*, 2014a).



Figure 3: Transformation grids and lollipop plot of all specimens along CV1 axis, accounting for 66.0% of total morphometric variance.



Figure 4: Scatter plot of canonical variate analysis for the group centroids of *Scartelaos tenuis* from the five stations.

#### Iranian Journal of Fisheries Sciences 19(2) 2020

## Discussion

The shape diversities between S. tenuis specimens separate western Persian Gulf populations (Heleh and Asaluyeh) from the other three populations (Chahoo, Dargahan and Jask) (Fig. 4). S. tenuis specimens from Chabahar and Qeshm Island are distinguishable regarding morphometric and meristic characteristics (Hashemi et al., 2015). These results are also coherent with recent morphological studies conducted on the other sympatric mudskipper **Periophthalmus** waltoni Koumans, 1941 (Ghanbarifardi et al., 2014a,b). Moreover, investigations on Lutjanus johnii (Bloch, 1792) (otolith shape) and Leptodius exaratus (H. Milne Edwards, 1834) (morphometric characteristics) exhibit discrepancies between populations from the Persian Gulf and the Oman Sea (Sadighzadeh et al., 2014; Afkhami et al., 2016). Aphanius dispar populations from the Persian Gulf and the Oman Sea have been compared using morphometric and otolith characteristics (Teimori et al., 2012a,b). The results from these studies proved that both coastal and landlocked populations of the Oman Sea are distinctive from those of the Persian Gulf. The mentioned studies assume otolith characters to be under genetic control; therefore, the phenotype diversities among populations of this specie s would be a result of geographical isolation and not related to environmental variations.

Such morphological variation (Fig. 3) is almost some kinds of polyphenism induced by distinct environmental factors (e.g. temperature and salinity); notwithstanding populations of these intertidal species (*S. tenuis* and *P. waltoni*) are genetically connected (West-Eberhard, 1989). Surface waters of the Persian Gulf (Heleh and Asaluyeh) have different physical conditions compare to the Oman Sea (Jask) (Thoppil and Hogan, 2010; Yao, 2008; Pous et al., 2004; Swift and Bower, 2003; Reynolds, 1993). Chahoo and Dargahan are located in Qeshm Island (Strait of Hormuz); however, mudskipper populations collected from these two stations are probably influenced by the currents entering the Persian Gulf from the Oman Sea (Fig. 1). Therefore, the specimens from Qeshm Island are similar to those of Jask (Oman Sea) (Fig. 4). Such different conditions might induce intraspecific phenotypic plastic responses. In fact, temperature and salinity can both affect growth and shape (Meador and Kelso, 1990; Loy et al., 1996a,b; Turan et al., 2005).

Mudskippers can tolerate wide ranges of temperature; salinity and however, changing these environmental factors could alter some behavioral and physiological characteristics of these fish. Temperature and salinity affect oxygen consumption and  $Na^+$ ,  $K^+$ , ATPase activity in mudskippers, respectively (Clayton, 1993). In specific, a study on P. waltoni in Iran showed that increasing salinity significantly induces elevation of RBC, Hct and Hb. (Soltanian et al., 2016). These mentioned physiological responses could influence the shape of fish in different stations with distinct temperature and salinity readings. Future phylogeographic molecular studies on S. tenuis could determine if these shape differences bear genetic variation.

# Acknowledgments

We are thankful to the Ferdowsi University of Mashhad for financial support to M. Gh (3/24103) and Shiraz University. Thanks also to Abdolwahed Pehpouri, Ahmad Mahmoudi and Mojtaba Masoudi for their help during sampling trips and lab work.

# References

- Abdoli, L., Kamrani, E., Abdoli, A. and Kiabi, B., 2010. Age and growth of the Mudskipper, *Scartelaos tenuis* (Day, 1876) (Gobiidae: Oxudercinae), in the coastal areas of the Persian Gulf at Bushehr province, Iran. *Zoology in the Middle East*, 51(1), 113-115.
- Abdoli, L., Kiabi, B., Kamrani, E., Abdoli, A., Rezazadeh katehsari, E. and Keshavarz, M., 2012. Feeding habits of Scartelaos tenuis in bushehr province, Iran. Journal of Fisheries (Iranian Journal of Natural Resources). 309-318.
- Adams, D. C., Rohlf, F. J. and Slice, D.
  E., 2004. Geometric morphometrics: ten years of progress following the "revolution". *Italian Journal of Zoology*, 16, 5–16.
- Afkhami, M., Schubart, C.D. and Naderloo, R., 2016. Morphometric differentiation among populations of *Leptodius exaratus* (H. Milne Edwards, 1834) (Brachyura, Xanthidae) from the Persian Gulf and the Oman Sea. *Crustaceana*, 89(3), 259-271.
- Annabi. A., Said, K. and Reichenbacher, **B..** 2013. Interpopulation differences in otolith morphology are genetically encoded in the killifish *Aphanius* fasciatus

(Cyprinodontiformes). *Scientia Marina*, 77(2), 269-279.

- Bookstein, F. L., 1991. Morphometric tools for landmark data: geometry and biology. Cambridge: Cambridge University Press.
- Cadrin, S. X. and Friedland, K. D., 1999. The utility of image processing techniques for morphometric analysis and stock identification. *Fisheries Research*, 43, 129–139.
- Clayton, D. A., 1993. Mudskippers. Oceanography and Marine Biology: an annual review.
- Costa, C. and Cataudella, S., 2007. Relationship between shape and trophic ecology of selected species of sparids of the Caprolace coastal lagoon (Central Tyrrhenian Sea). *Environmental Biology of Fishes*, 78, 115–123.
- Eklov, P. and Svanback, R., 2006. Predation risk influences adaptive morphological variation in fish populations. *The American Naturalist*, 167, 440–452.
- Fitzgerald, D. G., Nanson, J. W., Todd, T. N. and Davis, B. M., 2002. Application of truss analysis for the quantification of changes in fish condition. *Journal of Aquatic Ecosystem Stress and Recovery*, 9, 115– 125.
- **Ghanbarifardi, M. and Malek, M., 2007.** Permanent intertidal fish from the Persian Gulf and Oman Sea, *Iran. Iranian Journal of Animal Biosystematics*, 3, 1–14.
- Ghanbarifardi, M., Aliabadian, M.,
  Esmaeili, H. R. and Polgar, G.,
  2014a. Morphological divergence in the
  Walton's mudskipper, *Periophthalmus*

*waltoni* Koumans, 1941, from the Persian Gulf and Oman Sea (Gobioidei: Gobiidae). *Zoology in the Middle East*, 60(2), 133-143.

- Ghanbarifardi, M., Aliabadian, M., and Esmaeili, H. R., 2014b. Morphometric variation of *Periophthalmus waltoni* Koumans, 1941(Teleostei: Gobiidae) in the Persian Gulf and Oman Sea. *Iranian Journal of Animal Biosystematics*, 10(2), 137-144.
- Ghanbarifardi, M., Esmaeili, H. R., Gholami, Z., Aliabadian, M. and Reichenbacher, B., 2016. Molecular phylogeny of three mudskippers (Gobiidae) from the Persian Gulf and Oman Sea. Journal of Applied Ichthyology, 1-5.
- Hashemi, M., Moodi, S. and abdollahi , A.D., 2015. A study of morphological analysis of mudskipper fish (*Scartelaos tenuis* Day, 1876) in the two regions of Pozm gulf (Chabahar) and Dargahan port (Qeshm island).
- Klingenberg, C. P., 2011. MorphoJ: an integrated software package for geometric morphometrics. *Molecular Ecology Resources*, 11, 353–357.
- Kon, T. and Yoshino, T., 2002. Diversity and evolution of life histories of gobioid fishes from the viewpoint of heterochrony. *Marine and Freshwater Research* 53, 377–402.
- Langerhans, R. B., Layman, C. A., Langerhans, A. K. and Dewitt, T. J., 2003. Habitat-associated morphological divergence in two Neotropical fish species. *Biological Journal of the Linnean Society*, 80, 689–698.
- Lee, J. A., Kim J. W., Oh, S. Y., Yi, S. K., Noh, I., Ishimatsu, A. and Kim W.S., 2012. Effect of low dissolved

oxygen on the oxygen consumption rate and rhythm of the mudskipper *Scartelaos gigas* (Pisces, Gobiidae). *Fisheries Science* 78(5), 1013-1022.

- Loy, A., Ciccotti, E., Ferrucci, L. and Cataudella, S., 1996a. An application of automated feature extraction and geometric morphometries: temperaturerelated changes in body form of *Cyprinus carpio* juveniles. *Aquacultural Engineering*, 15, 301– 311.
- Loy, A., Cataudella, S., and Conti, M, 1996b. Shape changes during the growth of the sea bass, *Dicentrarchus labrax* (Teleostea: Perciformes), in relation to different rearing conditions. An application of thin-plate spline regression analysis. In: *Advances in morphometrics*, Marcus L.F. et al. eds, Plenum Press, New York, p. 399-405.
- Meador, M. R. and Kelso, W. E., 1990. Growth of largemouth bass in lowsalinity environments. *Transactions of the American Fisheries Society*, 119, 545-552.
- Miller, P. J., 1986. Reproductive biology and systematic problems in gobioid fishes. In T. Uyeno, R. Arai, T. Taniuchi, & K. Matsuura (Eds.), *Pacific Fish Biology: Proceedings of the 2nd International Conference on Indo-Pacific Fishes* (pp. 640–647). Tokyo: Ichthyological Society of Japan.
- Mostafa, A. R. H., Nahiduzzaman, M. D., Debasish, S., Habiba Khanam, M. U. and Alam, Md. S., 2010. Landmark-based morphometric and meristic variations of the endangered carp, kalibaus *Labeo calbasu*, from stocks of two isolated rivers, the

Jamuna and Halda, and a hatchery. *Zoological Studies*, 49, 556–563.

- Murdy, E. O., 1989. A taxonomic revision and cladistic analysis of the oxudercine gobies (Gobiidae: Oxudercinae). *Records of the Australian Museum*, (Supplement 11), 1–93.
- Polgar, G., Ghanbarifardi, M., Milli S., Agorreta, A., Aliabadian, М., Esmaeili, H.R. and Khang, **T.F.** 2017. Ecomorphological adaptation in three mudskippers (Teleostei: Gobioidei: Gobiidae) from the Persian Gulf and the Oman Sea. Hydrobiologia, 795(1), 91-111.
- Pous, S. P., Carton, X. and Lazure, P., 2004. Hydrology and circulation in the Strait of Hormuz and the Oman Sea. Results from the GOGP99 Experiment:
  1. Strait of Hormuz. *Journal of Geophysical Research* 109, PART 12, SECT 3, pp. C12037.
- Reynolds, R. M., 1993. Physical oceanography of the Gulf, Strait of Hormuz, and the Oman Sea. Results from the Mt. Mitchell expedition. *Marine Pollution Bulletin*, 27, 35–59.
- Rohlf, F. J., 1990. Morphometrics. Annual Review of Ecology and Systematics, 21, 299-316.
- Rohlf, F. J., 1999. Shape statistics: Procrustes superimpositions and tangent spaces. *Journal of Classification*, 16, 197–223.
- Rohlf, F. J., 2006. tpsUtil v1.38. Retrieved from http://life.bio.sunysb.edu/morph.
- Rohlf, F. J., 2013. tpsDig v2.17. Retrieved from http://life.bio.sunysb.edu/morph.

- Rohlf, F. J. and Marcus, L. F., 1993. A revolution in morphometrics. *Trends in Ecology and Evolution*, 8 (4), 129-132.
- Sadighzadeh, Z., Valinassab, T., Vosugi,
  G., Motallebi, A.A., Fatemi, M. R.,
  Lombarte, A. and Tuset, V.M., 2014.
  Use of otolith shape for stock identification of John's snapper,
  Lutjanus johnii (Pisces: Lutjanidae),
  from the Persian Sea and the Oman Sea.
  Fisheries Research, 155, 59-63.
- Silva, A., 2003. Morphometric variation among sardine (*Sardina pilchardus*) populations from the northeastern Atlantic and the western Mediterranean. *Journal of Marine Science*, 3139(03), 1352–1360.
- Soltanian, S., Vazirzadeh, A. and fallahi, R., 2016. Effects of sudden salinity changes on short-term hematological and biochemical responses in Mudskipper Periophthalmus waltoni Koumans 1941 (Gobiidae: Perciformes). Iranian Journal of Ichthyology, 3(1), 31-42.
- Swift, S. A. and Bower, A. S., 2003. Formation and circulation of dense water in the Persian Gulf. *Journal of Geophysics Research*, 108(C1), 3004.
- Teimori, A., Jawad, L. A. J., Al-Kharusi, L. H., Al-Mamry, J. M. and Reichenbacher, **B.**, 2012a. Late Pleistocene to Holocene diversification and historical zoogeography of the Arabian killifish (Aphanius dispar) inferred from otolith morphology. 637-645. Scientia Marina. 76. doi:10.3989/scimar.03635.26C.
- Teimori, A., Schulz-Mirbach, T.,Esmaeli, H. R. and Reichenbacher,B., 2012b. Geographical differentiation

of *Aphanius dispar* (Teleostei: Cyprinodontidae) from Southern Iran. *Journal of Zoological Systematics and Evolutionary Research*, 1-16. doi: 10.1111/j.1439-0469.2012.00667.x.

- Thoppil, P. G. and Hogan, P. J., 2010. A modelling study of circulation and eddies in the Persian Gulf. *Journal of Physical Oceanography*, 40, 2122– 2134.
- Turan, C., Yalçin, S., Turan, F., Okur, E. and Akyurt, I., 2005. Morphometric comparisons of African catfish, *Clarias* gariepinus, populations in Turkey. *Folia Zoologica*, 54(1/2), 165.

- West-Eberhard, M. J., 1989. Phenotypic plasticity and the origins of diversity. Annual review of Ecology and Systematics, 20(1), 249-278.
- Yao, F., 2008. Water mass formation and circulation in the Persian Gulf and water exchange with the Indian Ocean. Dissertation thesis, Miami (USA): University of Miami.
- Zelditch, M. L., Swiderski, D. L., Sheets,
  H. D. and Fink, W. L., 2004.
  Geometric morphometrics for
  biologists: a primier. San Diego:
  Elsevier Academic.