Downloaded from jifro.ir on 2025-10-29]

Determination of components and *in vitro* antistreptococcal properties of *Mentha piperita* L., *Satureja khuzistanica* Jamzad, *Matricaria recutica* L., *Zataria multiflora* Boiss. and *Rosmarinus officinalis* L. hydroethanolic extracts

Tafi A.A.1*; Meshkini S.2; Tukmechi A.3; Alishahi M.4; Noori F.5

Received: June 2017 Accepted: October 2017

Abstract

Streptococcosis is an infectious disease with significant economic and sanitary repercussions for trout farms in the world. The objective of this study was characterize the chemical constituents and in vitro antibacterial properties of Mentha piperita L., Satureja khuzistanica Jamzad, Matricaria recutica L., Zataria multiflora Boiss and Rosmarinus officinalis L. extracts against streptococcus iniae. The chemical components of the selected hydroalcoholic extracts were also analyzed by GC/MS. The most constituents were menthol (24%) in the M. piperita L., carvacrol (85.88%) in the S. khuzistanica Jamzad, guaiazulene (25.6%) in the M. recutica L. carvacrol (51.12%) in the Z. multiflora Boiss and α-pinene (12.5%) in the R. officinalis L. Among plant extracts under study Z. multiflora had the lowest MIC (4.896 mg ml⁻¹) and MBC (9.792 mg ml⁻¹) and the maximum MIC and MBC were belonged to M. piperita (18.55 mg ml⁻¹) and R. officinalis (33.645 mg ml⁻¹) respectively. Also, the inhibition zone diameter of bacteria had determined by disc diffusion method and compared to erythromycin. The highest and the lowest Inhibition zone diameter were belonged to Z. multiflora (14.43±0.55 mm) and M. recutita (13.23±0.35 mm) respectively. Results showed that Z. multiflora extract exhibited highest antibacterial effect against S. iniae among all plant extracts in this study.

Keywords: Herbal extracts, Antistreptococcal property, Minimum inhibitory concentration, Minimum bactericidal concentration

¹⁻Department of Fisheries, Faculty of Natural Resources, Urmia University, Urmia, Iran.

²⁻Department of Food Hygiene and Quality Control, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

³⁻Department of Microbiology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

⁴⁻Department of Clinical Science, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Ahvaz, Iran.

⁵⁻Department of Aquaculture, Artemia and Aquaculture Institute, Urmia University, Urmia, Iran

^{*}Corresponding author's Email: aliakbartafi@gmail.com

Introduction

Research directed towards the development of new antimicrobial agents is necessary for several reasons like the emergence and rapid spread of drug resistant against bacterial pathogens and raising the spectra of terrestrial and aquatic animals' untreatable disease (Payne and Tomasz, 2004). Moreover, experts rightly warn that if a new antimicrobial agent is not continued we will not able to improve sensitive veterinary and improving aquaculture industry (Charles Grayson, 2004). It is therefore imperative to find/develop safer and more potent compounds for the future that are not toxic to the host (Amin et al., 2010).

There is an increasing interest in medical plants as new sources of natural antibacterial agents in aquaculture (Jeevan et al., 2004). There are many reports about components antibacterial some medicinal plants against aquatic animals disease agents (Issabeagloo et al., 2012; Roomiani et al., 2013). According to researchers reports wide spectra of infectious bacteria sensitive are to medicinal plant extracts (Zlotkin et al., 1998; Agnew and Barnes, 2007; Bairwa et al., 2012; Mesbah et al., 2015).

The well-known and widely used peppermint (*Mentha piperita* L.) (Lamiaceae) is a cultivated natural hybrid of *M. aquatica* L. (water mint) and *M. spicata* L. (Spearmint). Although a native genus of the Mediterranean region, it is cultivated all over the world for flavor, fragrance, medicinal, and pharmaceutical applications. Mint essential oils are generally used externally for antipruritic, astringent, rubefacient, antiseptic, and antimicrobial purposes (Zargari, 1990).

Satureja species are native to warm temperate regions and may be annual or perennial (Tepe, 2015). Significant proportions of Satureja species are plants that have an important role in the pharmaceutical industry. Satureja khuzistanica Jamzad (Marzeh-e-Khuzestani in Persian) is one of the endemic Satureja species that grows in foothills, limestone gaps, and the southwestern parts of Iran. The aerial parts of antiparasitic, this species have carminative, digestive and diuretic properties (Ghodrati et al., 2015). The essential oil and extract of S. khuzistanica have biological compounds including antibacterial, antifungal, antioxidant and anti-inflammatory properties (Sahin et al., 2003; Tampieri et al., 2005).

Chamomile (Matricaria recutica L.), a member of the daisy family, is one of the most widely used and well-documented medicinal plants in the world. It is widely distributed in Asian tropics and subtropics areas and grows freely everywhere. The chamomile volatile oils are shown to have antimicrobial activity against certain species of bacteria, fungi and viruses in vitro; however, Gram-positive bacteria were found more susceptible than Gramnegative ones (Gholipour Kanani et al., 2012). The beneficial effects are related to different classes of therapeutically interesting ingredients such as essential oil components, flavonoids, and coumarin derivatives (Moricz et al., 2012).

Zataria multiflora Boiss is a plant belongs to the Labiatae family that is distributed only in Iran, Pakistan and Afghanistan. It is greatly used for medicinal and condimental purposes in these countries. This plant with the vernacular name of Avishan Shirazi in Iran several traditional uses such as antiseptic (Zargari, 1990), anesthetic Rohani et al., (Sharif 2007) and antispasmodic (Zargari, 1990; Nakhai et al., 2007). Most aspects of their medicinal use are related to the essential oil which contains various levels of thymol and/or carvacrol, phenolic derivatives with strong and wide-spectrum antimicrobial activity (Dorman and Deans, 2000; Mohagheghzadeh etal..2000). Z. multiflora Boiss is a perennial herbaceous plant, distributed in central and southern of Iran (Amin et al., 2010).

Rosemary (Rosmarinus officinalis L.) originally grows in southern Europe. Its herb and oil are commonly used as spice and flavoring agents in food processing for its desirable flavor, high antioxidant activity and lately as antimicrobial agent. Rosemary plants are rich sources of phenolic compounds with high antimicrobial activity against both Gram positive and Gram-negative bacteria (Rozman and Jersek, 2009).

Streptococcus is a genus of coccus (spherical) Gram-positive bacteria belonging to the phylum Firmicutes and the order Lactobacillales (lactic acid bacteria), which normally live on the body of humans or animals and may cause disease (sometimes severe) in both humans and animals especially in fish (Weinstein et al., 1997; Agnew and Barnes, 2007). Streptococcus iniae has become one of the most serious aquatic pathogens in the last decade causing high losses in farmed marine and freshwater finfish in warmer regions. This species is capable of causing invasive disease and outbreaks in

aquaculture farms (Agnew and Barnes, 2007).

This study aims to determine the composition and antimicrobial activities of *M. piperita* L., *S. khuzistanica* Jamzad, *M. recutica* L., *Z. multiflora* Boiss , and *R. officinalis* L. hydro alcoholic extracts against *S. iniae*.

Materials and methods

Plant material

Herbal samples (aerial organs of M. piperita L., S. khuzistanica Jamzad, M. recutica L. Z. multiflora Boiss. and R. officinalis L.) were collected from local areas in Khuzestan Province of Iran and identified in the Department of Botany, Faculty of Agriculture, Urmia University, Iran. After identification the plants were transferred to central laboratory of Urmia University and thoroughly washed in running water to remove debris and dust particles and then rinsed in distilled water. Plant samples were air-dried and ground. Twenty grams of grinded powders from each plant was soaked in 100 ml solvent (ethanol (95 %) + distilled water) (50:50) for 15 min with occasional shaking at 60°C. After they were dissolved, the materials were filtered through Buchner funnel and Whatman No.1 filter paper. Then, the filtrates were evaporated using concentrated rotary evaporator and (Khakzadie *et al.*, 2015).

Identification of extracts' components by GC/MS

GC/MS analyses were performed on a Thermo Finnegan capillary gas chromatograph directly coupled to the mass spectrometer system (model GC TRACE; TRACE MS plus). HP-5MS non-

polar fused silica capillary column (30m×0.250 mm, 0.25µm film thickness) was used. Temperature profile was as follows: at first, the temperature of the oven was fixed on 40°C for 2 min. and then increased to 160°C with the temperature rate of 3°C min⁻¹, and finally increased to 280°C at 5°C min⁻¹ for 2 min. The carrier gas was helium at a flow rate of 1 ml min⁻¹, and ionization energy was 70 eV (Van den and Kratz, 1963).

Bacteria preparation and morphological, physiological and biochemical determination tests

The bacterial strain streptococcus iniae was provided by bacteria collection of Laboratory for Microbiology, Faculty of Sciences, Ghent University (LMG 3740). After preparation, bacteria strain was subjected the some standard morphological, physiological and biochemical tests (Chang et al., 2002). Gram stain and cellular morphologies were examined at scale 1000x. Catalase activity was determined by flowing drops of 3% hydrogen peroxide down the colonies of bacteria on a clean glass slide. Growth features were determined by culturing bacteria in nutrient broth provided at temperatures (10 and 37 °C), salinities (1.5 and 5 % NaCl) and pH (7.5 and 9). Motility was determined from a hanging drop preparation of bacteria using phase contrast microscopy and inoculation of bacteria into brain heart infusion agar. Acid from carbohydrates was performed using inoculation of the bacteria into nutrient broth containing the respected Dglucose and Lactose. Hydrolysis of arginine and lysine was examined (Soltani et al., 2005).

Antibacterial susceptibility measured by disc diffusion

The antibacterial activity of the plant extracts was determined by disc diffusion according to NCCLS (National Committee for Clinical Laboratory Standard) 1999. Briefly, $100\mu L$ of guidelines suspension of Streptococcus iniae (half McFarland 1.5×10⁸ cfu mL⁻¹, calculated by McFarland tubes) was spread out on Mueller-Hinton Agar (CONDALAB, Spain) plates and sterile 6mm discs, each containing 10µL of plant extract placed on the microbial lawns. Antibiotic discs including Erythromycin (PADTANTEB, Tehran, Iran) were also included. The tests were carried out in triplicate and plates were incubated at 37°C for 24h. Inhibition zone diameters of bacteria were measured after incubation period and reported in mm (NCCLS, 1999).

Determination of minimum inhibitory concentrations and minimum bactericidal concentrations

Minimum inhibitory concentration (MIC) determined values were by broth microdilution as recommended by NCCLS (NCCLS, 1999). Serial two-fold were obtained from the hydroalcoholic extracts of M. piperita, S. khuzistanica, M. recutica, Z. multiflora and R. officinalis in heart infusion broth brain [OUELAB, Canada] in 96-well micro titer plates. Fresh bacterial suspensions (S. iniae), prepared from overnight grown cultures in BHI, were added to give a final concentration of 1.5×10⁸ cfu ml⁻¹. For each plant serial dilution, a control group of bacteria, a medium control group, and an extract control group were also included.

The microplates were incubated at 37°C for 24h and the first dilution with no turbidity (no growth) was considered as MIC. Minimal the bactericidal concentrations (MBC) were determined by spreading 10µl of the contents of the MIC wells (all wells without turbidity) that showed no bacterial growth on Mueller Hinton Agar (MHA) [CONDA LAB, Spain] plates followed by incubation at 37°C for 24h. The first well with colony counts of <5 was considered to be negative for growth and reported as the MBC.

Statistical analysis

Inhibition zone diameters data of the plant extracts and erythromycin antibiotic against *Streptococcus iniae* were analyzed in SPSS statistical software (v. 20) using one-way ANOVA followed by least significant differences Dunnett's test (p<0.05).

Results

According to gas chromatography-mass spectrometry (GC-MS) analysis, plant extracts compositions of M. piperita, S. khuzistanica, M. recutica, Z. multiflora and R. officinalis are presented in Table 1. The main components were Menthol (24.6%) followed by Menthone (18.4%) for M. piperita, Carvacrol (85.88%) for S. Guaiazulene khuzestanica, (25.6%)followed by (E)- β -Faranesens (20.1%) and Chamazulene (12.4%) for *M. recutita*, Carvacrol (51.12%) followed by Gammaterpinene (10.35%) for Z. multiflora and α pinene (17.5%) followed by 1,8-cineole (10.2%) for *R. officinalis*.

Results of some standard morphological, physiological and biochemical tests to identify bacterial strain are shown in Table 2.

Table 1: Chemical composition of hydroalcoholic extracts of plants.

Mentha piperita		Satureja khuzestanica		Matricaria recutita		Zataria multiflor	Rosmarinus officinalis		
Compound	(%)	Compound	(%)	Compound	(%)	Compound	(%)	Compound	(%)
Menthofuran	5.2	α- pinene	0.26	Limonene	0.5	Thujene	0.19	α- pinene	17.5
Isomenthone	5.9	β-Myrcene	0.29	-Terpineney	0.5	Alpha-pinene	4.26	Camphene	6.3
Menthone	18.4	α-Terpinene	1.49	(E)-β-Faranesens	20.1	Beta-pinene	0.43	β- pinene	3.5
Menthol	24.6	p-Cymene	3.11	Germacrene-D	3.1	Beta-myrcene	0.85	p-cymene	3.0
1,8-cineole	6.0	β-Phellandrene	3.19	α-Muurolene	0.8	Eucaliptol	3.37	1,8-cineole	10.2
1-limonene	4.1	□-Terpinene	1.14	Germacrene-A	0.5	Gamma-terpinene	10.35	1-limonene	2.4
Linalool	4.8	Linalool	0.91	Z-γ-Bisabolene	2.6	Linalool	5.68	Linalool	2.6
β -caryophyllene	2.5	Borneol	0.36	Caryophyllene oxide	1.2	Thymyl methyl ether	0.47	Camphor	4.2
Borneol	0.7	Terpinene-4-o1	0.65	Spathulenol	1.7	Carvacrol methyl ether	0.46	Borneol	9.6
α-terpineol	0.7	Thymol	0.19	α -Bisabolol oxide B	7.3	Carvacrol	51.12	4-terpineol	2.6
Pulegone	14.4	Carvacrol	85.88	α -Bisabolol	7.3	Trans-caryophyllene	0.41	Verbenone	8.6
Terpinen-4-ol	4.1	(Z)-Caryophyllene	0.15	Chamazulene	12.4	Globulol	2.32	Borneol acetate	3.1
		β-Bisabolene	0.21	α- Bisabolol oxide A	1.9			Tricyclene	0.5
		Caryophyllene oxide	0.18	Guaiazulene	25.6			3-octanone	0.8
				Hexadecanole	5.6			β-myrcene	1.6
				n-nonadecane	1.4			Crysanthenone	1.0
				Sclarene	0.4			Geraniol	1.8
				n-pentacosane	0.5			Caryphyllene oxide	1.7

Table 2: Morphological, physiological and biochemical characteristics of *Streptococcus iniae* (LMG 3740) + = positive reaction/growth, - = negative reaction/growth.

	Morphology	•		Motility	Growth									
Gram			Catalase activity		Temperature °C		NaCl % (after 72 h)		pH (after 72 h)		Acid from		Hydrolysis of	
					10	37	1.5	5	7.5	9	D-glucose	Lactose	Arginine	Lysine
+	cocci	<0.5mm	-	-	-	+	+	+	+	+	+	-	+	-

Antibacterial effects of the plant extracts are presented in Table 2. The MIC and MBC values of *M. piperita*, *S. khuzistanica*, *M. recutica*, *Z. multiflora*, *R.*

officinalis against *S. iniae* and inhibition zone diameters are presented in Table 3. Table 3 indicates MIC and MBC of erythromycin against *S. iniae*.

Table 3: MIC and MBC of plant extracts against *Streptococcus iniae* and comparison of inhibition zone diameters.

	MIC	MBC	Inhibition zone diameter (mm) $(Mean\pm SD)^*$				
Plant extracts	(mg ml ⁻¹)	(mg ml ⁻¹)	Plant extracts	Erythromycin			
Mentha piperita	18.550	18.550	13.97±0.50 ^a	16.07±0.58 ^b			
Satureja khuzestanica	10.870	10.870	14.20±0.30 ^a	16.07±0.58 ^b			
Matricaria recutita	8.285	16.570	13.23±0.35 ^a	16.07±0.58 ^b			
Zataria multiflora	4.896	9.792	14.43±0.55 ^b	16.07±0.58 ^b			
Rosmarinus officinalis	16.822	33.645	14.20±0.61 ^a	16.07±0.58 ^b			

^{*} Inhibition zones diameters were subjected to analysis of variance (SPSS, One-Way ANOVA) followed by least significant differences Dunnett's test. The different superscript alphabets in the same row are significantly different at *p*<0.05.

Table 4: MIC and MBC of Erythromycin against Streptococcus iniae.

	Concentration (µg ml ⁻¹)										
500	250	125	62.5	31.25	15.62	7.81	3.90				
					MBC	MIC					

Discussion

In recent years, plant species have been used in the focus of scientific scrutiny due to their unique phytochemical contents. As it is well-known, these compounds have excellent pharmacological properties (Tepe, 2015).

Many authors have suggested that the differentiated effects of *M. piperita* L. (Talpur, 2014; Trevisan *et al.*, 2017), *S.*

khuzistanica Jamzad (Tepe, 2015), M. recutica L. (Owlia et al., 2007) Z. multiflora Boiss. (Khakzadie et al., 2015) and R. officinalis L. (Issabeagloo et al., preparations selected 2012) on microorganisms depend on their composition, the characteristics of the plant material, and concentration used. Ndakalimwe et al., (2015) reported that 4% Aloe vera powder supplemented diet enhanced immune responses and haematobiochemical indices of Oreochromis niloticus against aquatic pathogenic Streptococcus iniae. Antimicrobial activity of R. officinalis (Rosemary), Z. multiflora (Oregano), Anethum graveolens (Dill) and Eucalyptus globulus (Eucalyptus) essential oils and extracts against Lactococcus garvieae was reported by Mahmoodi et al., (2012). They demonstrated that among four plant extracts they have been studied. Multiflora extract with carvacrol (71.1 %) was the most effective plant extract with the highest antibacterial effect against Lactococcus garvieae. Thus, in the present study, three hydroalcoholic extracts obtained from these various medicinal plants (M. piperita, S. khuzistanica, M. recutica, Z. multiflora and R. officinalis) were analyzed by GC/MS (Table 1) to determine their main constituents and evaluated for their antimicrobial effects against gram positive zoonosis pathogenic bacteria S. iniae.

All results of bacterial identification tests (Table 2) are compatible with standard characterizations of S. iniae reported by Soltani et al. (2005). S. iniae may colonize in the surface of fish or causes invasive disease associated with 30 to 50 percent mortality in affected fish ponds (Eldar et al., 1994). The results of the present study indicated that S. iniae was inhibited or eliminated after exposure the extracts of M. piperita, S. khuzistanica, M. recutica, Z. multiflora and R. officinalis in culture broth (Table 3). Such delay in inhibition of microbial growth is particularly useful in terms of public health and safety. The difference in microbial susceptibility is attributable to the chemical composition of extracts. The

lower effectiveness of some extracts against S. iniae in this study (Table 3) might reflect the lower amount of antibacterial compounds in the plants. A possible explanation for could be due to some of the plant extracts may have contained antibacterial constituents but sufficient were present in concentrations to be effective. Moreover, microorganism's features are reasons for miscarriage of extracts to eliminate or inhibit microorganisms (Owlia et al., 2007).

The results obtained in the present study, referring to the activity of M. piperita, S. khuzistanica, M. recutica, Z. multiflora and R. officinalis extracts against S. iniae are consistent with findings of some authors (Iscan et al., 2002; Owlia et al., 2007; Issabeagloo et al., 2012; Esmaeili et al., 2015; Khakzadie et al., 2015). This effect is attributable to the considerable concentration of Menthol (24.6%) and Menthone (18.4%) in M. piperita, Carvacrol (85.88%) in Satureja khuzistanica, Guaiazulene (25.6%), (E)-β-Faranesens (20.1%) and Chamazulene (12%) in *M. recutita*, Carvacrol (51.12%) and Gamma-terpinene (10.35%) in Z. multiflora Boiss and α- pinene (17.5%) and 1.8-cineole (10.2%) in R. officinalis L. (Table 1). However, in some cases, the results of the present study are a little bit different from findings of some authors (Iscan et al., 2002). These differences in extracts' compositions the can attributed to the several environmental factors such as climatic, seasonal and geographical or ontogenesis variations, plant harvest time, error in the analysis of extracts, time between extraction and analysis of compounds' constituents.

Antibacterial and antiparasitical effects of *M. piperita*, *S. khuzistanica*, *M. recutica*, *Z. multiflora* and *R. officinalis* extracts in fishes were reported in *in vivo* research of many authors (Abutbul *et al.*, 2004; Soltani *et al.*, 2010; Gholipour Kanani *et al.*, 2012; Khansari *et al.*, 2013; Adel *et al.*, 2015; Adel *et al.*, 2016). These researchers reported promotes growth performance and increases the main humoral immune parameters (both at mucosal and systemic levels) of dietary administration of these herbal extracts to farmed fish.

Mahmoodi et al. (2012) reported that Z. multiflora extract had the most active antibacterial plant against L. garvieae extract among plants in his study, showing both MIC and MBC of 15.6 µg mL⁻¹ and inhibition zone diameter 28±0.7 mm. Inhibition zone diameter of herbal extracts under study have lower current antistreptococcal effect than erythromycin, but Z. multiflora extract with MIC (4.896 mg mL⁻¹) and MBC (9.792 mg mL⁻¹) has good potentials as antistreptococcal agent in combating S. iniae and its inhibition zone diameter (14.43±0.55 mm) has no significant difference beside erythromycin $(16.07\pm0.58 \text{ mm})$ statistically (p<0.05)(Table 3). So Z. multiflora may be more acceptable to consumers and the regulatory agencies than chemical compounds. Antibacterial effects of Z. multiflora extract on the inactivation or elimination of S. iniae will help to develop or modify a new factor used for safety of aquatic animals.

References

Abutbul, S., Golan-Goldhirsh, A., Barazani, O. and Zilberg, D., 2004. Use of *Rosmarinus officinalis* as a

treatment against *Streptococcus iniae* in tilapia (*Oreochromis* sp.). *Aquaculture*, 1(4), 97-105.

Adel, M., Safari, R., Pourgholam, R., Zorriehzahra, J. and Esteban, M.A., 2015. Dietary peppermint extracts (Mentha piperita) promote growth performance and increase the main humoral immune parameters (both at mucosal and systemic level) of Caspian brown trout (Salmo trutta caspius Kessler, 1877). Fish and Shellfish Immunology, 47(1), 623-629.

Adel, M., Pourgholam, R., Zorriehzahra, J. and Ghiasi, Μ. 2016. Hemato-Immunological and biochemical parameters, skin antibacterial activity, and survival in rainbow trout (Oncorhynchus mykiss) following the diet supplemented with Mentha piperita against Yersinia r uckeri. Fish and Shellfish Immunology, 55, 267-273.

W. A.C., Agnew, and Barnes, 2007. Streptococcus iniae: An aquatic pathogen of global veterinary significance and challenging a candidate reliable for vaccination. Veterinary Microbiology, 122, 1-15.

Amin, M., Kalantar, E., Mohammad Saeid, N. and Ahsan, B. 2010. Antibacterial effect and physicochemical properties of essential oil of *Zataria multiflora* Boiss. *Asian Pacific Journal of Tropical Medicine*, 3(6), 439-442.

Bairwa, M.K., Jakhar, J.K., Satyanarayana, Y. and Reddy, A.D. 2012. Animal and plant originated immunostimulants used in aquaculture.

- Journal of Natural Product and Plant Resources, 2 (3), 397-400.
- Chang, P.H., Lin, C.W. and Lee, Y.C., **2002.** Lactococcus garvieae infection of cultured rainbow trout, Oncorhynchus mykiss, in Taiwan and associated biophysical characteristics and histopathology. Bulletin the of European Association of Fish Pathologists, 22, 319-327.
- Charles, G.P.P. and Grayson, M.L., 2004. The dearth of new antibiotic development: why we should be worried and what we can do about it. *The Medical Journal of Australia*, 181(10), 549-53.
- **Dorman, H.J. and Deans, S.G., 2000.**Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *Journal of Applied Microbiology*, 88, 308-316.
- Eldar, A., Bejerano, Y. and Bercovier, H., 1994. Streptococcus shiloi and Streptococcus difficile: two new streptococcal species causing a meningoencephalitis in fish. Current Microbiology, 28, 139-43.
- Esmaeili, D., Moradi, F., Imani Foladi, A.A. and Bahador, A., 2015. The study of inhibitory effects of Satureja khuzestanica essence against Exos gene of MDR Pseudomonas aeruginosa by RT-PCR technique. Pharmaceutical Regulatory Affairs, 4(3), 1-4.
- Ghodrati, L., Alizadeh, A. and Ketabchi, S., 2015. Essential oil constituents and antimicrobial activities of Iranian Satureja khuzistanica Jamzad. International Journal of Biosciences, 6(3), 249-257.
- Gholipour Kanani, H., Sahandi, J. and Taheri, A., 2012. Influence of garlic

- (Allium sativum) and mother worth (Matricariachamomilla) extract on Ichthyophtirius multifilus parasite treatment in sail fin molly (Poecilia latipinna) ornamental fish. APCBEE Procedia, 4, 6–11.
- Iscan, G., Kirimer, N., Kurkcuogly, M., Husnu Can Baser, K. and Demirci, F., 2002. Antimicrobial screening of *Menthapiperita* essential oils. *Journal of Agricultural and Food Chemistry*, 50, 3943-3946.
- Issabeagloo, E., Kermanizadeh, P., Taghizadieh, M. and Forughi, R., 2012. Antimicrobial effects of rosemary (Rosmarinus officinalis L.) essential oils against Staphyloccus spp. African Journal of Microbiology Research, 6(23), 5039-5042.
- Jeevan, R., Bhakshu, M. and Venkata, R., 2004. *In vitro* antimicrobial activity of certain medicinal plants from Eastern Ghats, India, used for skin diseases. *Journal of Enthnophamacology*, 90(2-3), 353-357.
- Khakzadie, M., Eslami, H., Vazifehdust, S., Behboudi, M.R. and Shahabzadeh, S.H., 2015. Investigating the antibacterial effects of aqueous and alcoholic extracts of thyme on *Bacillus subtilis* bacterias. *International Journal of Biology*, *Pharmacy and Allied Sciences*, 4(8), 5531-5538.
- Khansari, A., Yavari, V., Alishahi, M., Mousavi, S.M., Ghorbanpoor, M., Darvish Bastami, K. and Azizi, S.H., 2013. Effects of *Oliviera decumbens* and *Satureja khuzestanica* extract on some immunological and haematological parameters of *Cyprinus carpio. Comparative Clinical Pathology*, 22, 339–342

- Mahmoodi, A., Soltani, M., Akhondzadeh Basti, A., Roomiani, L., Kamali, A. and Taheri, S., 2012. Chemical composition and antibacterial activity of essential oils and extracts from Rosmarinus officinalis, Zataria multiflora, Anethum graveolens and Eucalyptus globulus. Global Veterinaria, 9(1), 73-79.
- Mesbah, M., Mohammadian, T., Alishahi, M. and Jangaran Nejad, A., 2015. Effects of dietary *Aloe vera* and Echinacea on some nonspecific immunity in Shirbot (*Barbus grypus*). *Iranian Journal of Aquatic Animal Health*, 2(1), 24-36.
- Mohagheghzadeh, A., Shams-Ardakani, M. and Ghannadi, A., 2000. Volatile constitutes of callus and flower-bearing tops of *Zataria multiflora* Boiss (Lamiaceae). *Flavour and Fragrance Journal*, 15, 373-6.
- Moricz, A.M., Szarka, S., Ott, P.G., Hethelyi, E.B., Szoke, E. and Tyihak, E., 2012. Separation and identification of antibacterial chamomile components using OPLC, bioautography and GC-MS. *Medicinal Chemistry*, 8(1), 85-94.
- Nakhai, L.A., Mohammadirad, A., Yasa, N., Minaie, B., Nikfar, S. Ghazanfari, G., Zamani, M.J., Dehghan, G., Jamshidi, H., Boushehri, V.S., Khorasani, R. and Abdollahi, M., 2007. Benefits of *Zataria multiflora* Boiss in experimental model of mouse inflammatory bowel disease. *Evidence-Based Complementary and Alternative Medicine*, 4(1), 43-50.
- National Committee for Clinical Laboratory Standard (NCCLS).

 1999. Performance standards for antimicrobial susceptibility testing.

- 9th International Supplement; M100-S9. Wayne, Pennsylvania, USA.
- Ndakalimwe, N.G., Qiang, J., He, J., Ma, X.Y., Kpundeh, M.D. and Xu, P., 2015. Dietary Aloevera supplementation on growth performance, some haematobiochemical parameters and disease resistance against Streptococcus iniae in tilapia (GIFT). Fish and Shellfish Immunology, 44, 504-514.
- Owlia, P., Rasooli, I. and Saderi, H., 2007. Antistreptococcal and antioxidant of essential oil from *Matricaria chamomilla*. Research Journal of Biological Sciences, 2(2), 155-160.
- Payne, D. and Tomasz, A., 2004. Antimicrobials: The challenge of antibiotic resistant bacterial pathogens: The medical need the market and prospects for new antimicrobial agents. *Current Opinion in Microbiology*, 7(5), 435-438.
- Roomiani, L., Soltani, M., Akhondzadeh Basti, A., Mahmoodi, A., Taheri Mirghaed, A. and Yadollahi, F., 2013. Evaluation of the chemical composition and in vitro antimicrobial activity of Rosmarinus officinalis, Zataria multiflora, Anethum graveolens and **Eucalyptus** globulus against Streptococcus iniae; the cause of zoonotic disease in farmed fish. Iranian Journal of Fisheries Sciences, 12(3), 702-716.
- Rozman, T. and Jersek, B., 2009. Antimicrobial activity of rosemary extracts (*Rosmarinus officinalis* L.) against different species of *Listeria*. *Acta Agriculturae Slovenic*, 93(1), 51–58.

- Sahin, F., Karaman, I., Gulluce, M., Ogutcu, H. and Sengul, M., 2003. Evaluation of antimicrobial activities of Satureja hortensis L. Journal of Ethnopharmacology, 87, 61-65.
- Sharif Rohani, M., Haghighi, M., Assaeian, H. and Lashtoo Aghaee, G.H.R., 2007. A study of the anesthetic effect of *Zataria multiflora* Boiss (Labbiatae) essence on *Oncorhynchhus mykiss* and cultured *Salmo trutta caspius*. *Iranian Scientific Fisheries Journal*, 16(4), 99-106.
- **Soltani, M., Jamshidi, and Sharifpour, I., 2005.** Streptococcosis caused by *Streptococcus iniae* in farmed rainbow trout (*Oncorhynchys mykiss*) in Iran: Biophysical characteristics and pathogenesis. *Bulletin of the European Association of Fish Pathologists*, 25(3), 95-106.
- M., Soltani, Sheikhzadeh, N., E brahimzadeh-Mousavi, H.A. and Zargar, A., 2010. Effects of Zataria multiflora oil on innate immune of responses Common carp (Cyprinus carpio). Journal of Fisheries and Aquatic Science, 5(3), 191-199.
- Talpur, A.D., 2014. *Mentha piperita* (Peppermint) as feed additive enhanced growth performance, survival, immune response and disease resistance of Asian seabass, *Lates calcarifer* (Bloch) against *Vibrio harveyi* infection. *Aquaculture*, pp. 420–421, 71–78.
- Tampieri, M.P., Galuppi, R., Macchioni,
 F., Carelle, M.S. and Falcioni, L.,
 2005. The inhibition of *Candida albicans* by selected essential oils and their major

- components. *Mycopathologia*, 159, 339-345.
- **Tepe, B., 2015.** Inhibitory effect of *Satureja* on certain types of organisms. *Records of Natural Products*, 9(1), 1-18.
- Trevisan, S.C.C., Menezes, A.P.P., Barbalho, S.M. and Guiguer, E.L., 2017. Properties of Mentha piperita: a Brief review. World Journal of Pharmaceutical and Medical Research, 3(1), 309-313.
- Van den, D.H. and Kratz, P.D.A., 1963.

 Generalization of the retention index system including linear temperature programmed gas-liquid partition chromatography. *Journal of Chromatography*, 11, 463-471.
- Weinstein, M.R., Litt, M., Kertesz, D.A., Wyper, P., Rose, D., Coulter, M., McGeer, A., Facklam, R., Ostach, C., Willey, B.M., Borczyk, A. and Low, D.E., 1997. Invasive infections due to a fish pathogen, *Streptococcus iniae*. The New England Journal of Medicine, 337(9), 589-594.
- **Zargari, A., 1990. Medicinal Plants.** 4th edition. Tehran University Publications, Tehran. 269 P.
- **Zlotkin, A., Hershko, H. and Eldar, A., 1998.** Possible transmission of *Streptococcus iniae* from wild Fish to cultured marine fish. *Applied and Environmental Microbiology*, 64(**10**), 4065–4067.