# Dietary ginger improve growth performance, blood parameters, antioxidant capacity and gene expression in Cyprinus carpio

19(3) 1237-1252

Jafarinejad R.<sup>1</sup>; Gharaei A.<sup>2</sup>\*; Mirdar Har<u>ij</u>ani J.<sup>1</sup>

Received: November 2017 Accepted: February 2018

#### **Abstract**

The present study evaluates the effect of dietary ginger (Zingiber officinale) on growth performance, haematological parameters, antioxidant status of common carp (Cyprinus carpio). Fish with initial weight of 14±1.95 g were allocated into twelve tanks at density of 18 fish per tank. Fish were fed four experimental diets containing 0 (as control diet), 0.5, 2 and 5% ginger powder for 56 days. After 56 days of culture, fish fed with 2 and 5% ginger diets showed significant increase in final weight, weight gain, specific growth rate (SGR) and feed conversion ratio (FCR) (p<0.05) compared to the control diet. Significant higher number of leucocytes (WBC), erythrocytes (RBC), haematocrit, lymphocyte, monocyte and neutrophil were found in the fish fed with 2 and 5% ginger diets compared to the control diet. Superoxide dismutase (SOD), catalase (CAT) and Glutathione peroxidase (GPX) enzyme activities and Malondialdehyde (MDA) levels were significantly higher in the groups fed 2 and 5% ginger diets (p<0.05). SOD, CAT and GPX mRNA expression levels of fish fed with 2 and 5% ginger diets were significantly down-regulated compared to the control diet (p<0.05). The results suggested that dietary ginger can improve the growth performance, health status and antioxidant capacity of common carp.

Keywords: Ginger, Antioxidant enzyme activity, Haematological indices, Reactive oxygen species, Growth performance.

<sup>1-</sup>Department of Fisheries, Faculty of Natural Resources, University of Zabol, Zabol,

<sup>2-</sup>Department of Fisheries, Faculty of Natural Resources and Hamoon International Wetland Research Institute, University of Zabol, Zabol, Iran.

<sup>\*</sup>Corresponding author's Email: agharaei551@uoz.ac.ir

#### Introduction

Plants are natural sources of bioactive substances, which have useful effects in animal nutrition including appetite and feed intake induction, improvement of endogenous digestive enzyme secretion, activation of immune responses and antibacterial, antiviral and antioxidant actions (Lee *et al.*, 2014).

Improvement and protection of fish health in commercial production practices is an important factor in aquaculture industry. Medicinal plants are very useful in aquaculture because they have natural organic components that help reducing the mortality of fishes because of various pharmacological actions. In addition, they do not have any adverse effect to the environment and organism's health al.. (Vahedi 2015). Today, aquaculture products have gained a special place among other human food sources. One of the basic practices in order to supply part of human need to protein is fish culture. Therefore, due to increasing human population in recent decades, the researchers try to present the new ways for increasing the production in the shortest possible time, along with the least side effects to the cultured organisms. This subject has led the researchers to focus more on using, additives and biological diets, such as enzymes, probiotics, proteins, medicinal plants.

In many studies, the effects of different parts of plant species (flowers, leaves, seeds and roots) had been reported to influence in different ways such as growth improvement, antioxidant status, none specific

immune response and increasing the survival rate in some commercial fish species such as African catfish, *Clarias gariepinus* (Dada and Ikuerowo, 2009; Soosean *et al.*, 2010), tilapia *Orechromis mossambicus* (Immanuel *et al.*, 2009) and common carp, *Cyprinus carpio* (Alishahi *et al.*, 2010; Pakravan *et al.*, 2012; Fallahpour *et al.*, 2014), and Major carp, *Catla catla* (Arulvasu *et al.*, 2013).

Ginger (Zingiber officinale) is a medicinal plant that is widely used in traditional medicine and available as dietary supplement in medical sciences. From ancient times due to diversity of the active compounds in ginger it had been used in treatment of various ginger generally contains diseases alkaloids. flavonoids, polyphenols, saponin, steroids. tannin, fiber. carbohydrate, vitamins, carotenoids and minerals (Otunola, 2010; Prakash, 2010); and various pharmacological properties of ginger such as antioxidant, anti-apoptosis, anti-tumor, antiinflammation, anti-hyperglycemia, anticough and anti-colds have been known and reported (Dadfar et al., 2014).

Common carp that belongs to the family Cyprinidae, is one of the most important species due to its rapid growth rate, high economic profit and ease of cultivation that can be seen in most natural environments all over the world (Tokur *et al.*, 2006).

Some studies have reported that dietary ginger could enhance the growth performance, survival rate, nonspecific immune response, physiological status, resistance to bacterial diseases and pH stress (Chang

et al., 2012; Arulvasu et al., 2013; Chan et al., 2014; Levy et al., 2015; Nan et al., 2015).

Previous studies have shown that the antioxidant defense system, including antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPX) and some functional molecules (such as: albumin, ceruloplasmin, ascorbic acid, etc), is developed in animals in order to stabilize the minimum levels of ROS in the cell (García-Triana *et al.*, 2010; Sheikhzadeh *et al.*, 2012; Yuan *et al.*, 2015; Zhang *et al.*, 2013).

Generally, environmental changes (such as seasonal and daily temperature fluctuations) or anthropogenic factors can make hypoxic status (Zhang et al., 2013). The hypoxia subjection would cause to increase the production and the release of reactive oxygen species (ROS) and oxidative stress (Jusman et al., 2010). ROS are well known to be responsible for lipid peroxidation, protein degradation, DNA damage, and apoptosis in vertebrates. In addition, these effects can permanently lead to the break of free radicals in the destructive process of lipid peroxidase (Ahmad et al., 2000). The oxidative stress happen when a misbalance occurs between the production of ROS and antioxidant systems in the body of organism that can be quantified by measurement of the antioxidant enzyme activity and antioxidant gene expression procedures (Trenzado et al., 2006).

The aim of this study was to further improve our knowledge of the antioxidant mechanism of dietary ginger on the growth parameters,

survival and antioxidant enzymes activities, CAT, SOD, GPX mRNA levels and some hematological parameters of cultured Common carp (*Cyprinus carpio*) Juveniles.

## Materials and methods

Fish, experimental design and preparation of ginger feed

Common carp juveniles with mean weight of 14±1.95 g (mean±SD) were collected from the Zahak Fish Reproduction and Rearing Co. Ltd, Zabol, Iran. The fishes were fed with the basal diet for 2 weeks for adaptation to the experimental conditions and then were divided into four treatment groups and three replication (18 fishes in tanks with 75 L capacity). Fishes were fed with various concentrations of ginger (supplied by Golha corporation, Iran) powder (0 "control", 0.5, 2 and 5 % of diet) for 56 days (Table 1). The daily feeding rate was 3% of body weight in three times (08:00, 14:00 and 18:00). During the experimental period, water temperature, dissolved oxygen and pH were measured daily and maintained at  $19.8\pm1.5^{\circ}$ C,  $6.2\pm0.6$  mg L<sup>-1</sup>, and  $8.1\pm0.2$ . respectively. Continuous aeration was provided in each tank through an air stone connected to a central air compressor. In order to prepare the experimental diets, all ingredients were thoroughly mixed with various levels of ginger powder, and water was added to produce pellets of approximate 2 mm in diameter (Cerezuela, 2008). The pelleted diets were dried, sieved and stored at -20 °C until used.

In order to evaluate the bioenergetics parameters at the end of experimental period, Weight Gain (WG%), Specific Growth Rate (SGR %/day), Feed Conversion Ratio (FCR) and Survival Rate (%) were calculated according to the following equations (Mahghani *et al.*, 2014): WG(%)=(Wt-W<sub>0</sub>)×100/W<sub>0</sub>, SGR=(Ln Wt-Ln W<sub>0</sub>)×100/t, FCR=dry

feed fed/ Wet weight gain, Survival rate  $=(Nt/N_0)\times100$ . Here Wt and W<sub>0</sub> are final and initial body weights (g), respectively, t is duration of experimental days, N<sub>0</sub> is the initial number of fishes and Nt is the final number of fishes.

Table 1: Formulation and proximate composition of experimental diets (%).

Inquadianta	Experimental diet				
Ingredients	0	0.5%	2%	5%	
Fish meal	37.5	37.5	37.5	37.5	
Soybean meal	16.5	16.5	16.5	16.5	
Wheat	20	20	20	20	
Canola oil	6.5	6.5	6.5	6.5	
Corn	5	5	5	5	
DCP	2.05	2.05	2.05	2.05	
Carboxy methyl mellulose	2	2	2	2	
CaCO <sub>3</sub>	1.89	1.89	1.89	1.89	
Vitamin permix <sup>1</sup>	0.4	0.4	0.4	0.4	
Mineral premix <sup>2</sup>	0.25	0.25	0.25	0.25	
Vitamin C	0.08	0.08	0.08	0.08	
Colin	0.15	0.15	0.15	0.15	
Filler	5	4.5	3	0	
ginger powder	0	0.5	2	5	
Proximate composition (% DM	<sup>3</sup> )				
Protein	32	32.5	31.9	32.3	
Fat	13.4	12.5	12.8	13.1	
Moisture	4.8	5	4.9	5.1	
Ash	14.5	14.6	14.4	14.2	

<sup>&</sup>lt;sup>1</sup>Provided per kilogram of feed: vitamin D, 1.28 mg; vitamin E, 136 mg; thiamine, 8 mg; riboflavin, 16 mg; 3 pyrodoxine-Cl,16 mg; vitamin C, 80 mg; calcium pantothenate, 17.4 mg; biotin, 0.16 mg; folic acid, 4.8 mg; niacin, 40 mg; vitamin B,0.016 mg; menadion bisulphite, 20 mg

# Haematological and biochemical parameters

At the end of trial, the hematological parameters were measured by taking 2 mL blood samples from the caudal vein of three fishes from each tank. The blood samples were then suspended in heparinized tubes, and then values of red blood cell (RBC), white blood cell (WBC), hematocrit (PCV), Neutrophil,

Lymphocyte and Monocytes were measured (Faggio *et al.*, 2014).

For biochemical analysis, blood samples of fishes were taken from the caudal vessels by using heparinized syringes. Prior to sampling, the fish were anesthetized with 200 mg L<sup>-1</sup> MS 222 (Faggio *et al.*, 2013). Blood sera were separated by centrifuging the sample at 3,000 rpm for 10 min

<sup>&</sup>lt;sup>2</sup>Provided per kilogram of feed: magnesium, 625 mg; potassium, 509 mg; calcium, 465 mg; zinc, 100 mg; iron, 62 mg; manganese, 13 mg; copper, 6 mg

<sup>&</sup>lt;sup>3</sup>DM=dry matter

(Heraeus Labofuge 400) and the sera were removed with a disposable transfer pipette (Akrami et al., 2015). Then, the SOD (Superoxide dismutase) activity was determined based on the method previously described by Beyer and Fridovich (1987). The SOD activity was reported in units of enzyme/gram hemoglobin. One unit of SOD corresponds to the enzyme concentration required to inhibit the produced chromogen nitro tetrazolium by 50% in 1 min under standard conditions. GPX (Glutathione peroxidase) activity was measured according to the method explained by the previously described procedure (Paglia and Valentine, 1967). GPX activity was expressed in milliunits/mg hemoglobin. One specific unit is defined as 1 mmol of nicotinamide dinucleotide adenine phosphate (NADPH) needed with hemoglobin to be converted to NADPH in 1 min of reaction. CAT (Catalase) activity was measured according to the method proposed previously (Aebi, 1984). The CAT activity is defined in specific units/mg hemoglobin. One unit of CAT corresponds to the amount of enzyme needed to decompose H<sub>2</sub>O<sub>2</sub> in phosphate buffer, at pH 7.0, in 1 sec of reaction. Malondialdehyde (MDA) was measured by using a competitive enzyme-linked immunosorbent assay kit (Zellbio Co., GmbH, Germany) according to the manufacturer's instructions as previously described by Zhang et al. (2013).

Expression of CAT, SOD and GPX mRNAs

In order to compare the expression of mRNA levels, individual liver tissue from each treatment group (n=6) were randomly collected at the end of trial and frozen and kept at -80°C until use. Total RNA extractions from liver samples were carried out by using Takapou Kit (Tehran, Zist Iran) following the manufacturer's instruction. RNA integrity was verified by ethidium bromide staining of 28S and 18S ribosomal RNA bands on 1.2% gel. agarose To remove **DNA** contaminants, the extracted RNA was treated with RNA-Free DNase (Takara, Japan) and reverse transcribed to cDNA by a Superscript cDNA synthesis kit (AccuPawer® CycleScript RT PreMix, Germany) following the manufacturer's instructions.

The mRNA expression levels of CAT, SOD and GPX genes in the liver of common carp were evaluated by fluorescent real-time quantitative PCR. The specific primers for CAT, SOD, GPX and β-actin (housekeeping gene) were designed according to the cDNA sequences of common carp in Gen Bank (Table 2) (Jiang et al., 2016), and thermocycling conditions as indicated Table All primers synthesized by Takapou Zist Co., Ltd. and amplified fragment length of 75-295bp. Real-time quantitative PCR was conducted in a quantitative thermal cycler (Mastercycler® eprealplex; Eppendorf, Germany). Each reaction was performed in a total volume of 20 μL, containing 0.5 μL of each primer (15 mM), 2 µL of the diluted first strand cDNA product, 10 µL of Green qPCR Master Mix (Yekata Tajhiz Azma Co., Iran) and 7 µL of sterilized doubledistilled water. Three technical each replicates for sample were performed and the threshold cycle (CT) was determined manually for each run. PCR efficiencies for each set of primers were determined using serial 10-fold dilutions of cDNA and resulting plots of CT vs. the logarithmic cDNA dilution, using the efficiency equation (E):

$$E = 10^{(-1/\text{slope})}$$

Gene expression data were analyzed using the  $2^{-\Delta\Delta CT}$  method after verification that the primers amplified with an efficiency of from 97 to 99% (Gharaei *et al.*, 2011), and data for all treatment groups were compared to the control group.

Table 2: Real-time PCR primer sequences and thermocycling condition.

Target genes	Primer	Primer sequence (5 — 3 )	Thermocycling condition	Accession no.	Product length (bp)
SOD	F	TGGCGAAGAAGGCTGTTTGT	95 °C 30 s, 35 cycles of 95 °C 5 s, 60.4 °C 30 s and 72 °C 30 s	TE2/12255	85
SOD	R	TTCACTGGACCCGTCT	93 -0 30 s, 33 cycles 01 93 -0 3 s, 00.4-0 30 s and 72 -0 30 s	JF342333	85
CAT	F	CTGGAAGTGGAATCCGTTTG	95 °C 30 s, 35 cycles of 95 °C 5 s, 54.5 °C 30 s and 72 °C 30 s	TE/1160/	75
CAI	R	CGACCTCAGCGAAATAGTTG	23 030 s, 33 cycles 0223 0 2 s, 34.3 0 30 s and 72 0 30 s	31411004	75
GPX	F	CCTTCCCATCCCACCAGTTT	95 °C 30 s, 35 cycles of 95 °C 5 s, 60 °C 30 s and 72 °C 30 s	EI656211	118
GFA	R	TGCGGAGTCACCGTTCACAT	95 °C 50 8, 55 Cycles 01 95 °C 5 8, 00 °C 50 8 and 72 °C 50 8	F3030211	110
β-Actin	F	CGTGATGGACTCTGGTGATG	95 °C 30 s, 35 cycles of 95 °C 5 s, 60 °C 30 s and 72 °C 30 s	M24113	295
p-ricini	R	TCGGCTGTGGTGAAG		1412-4115	

# Statistical analysis

Data were tested for normality and homogeneity of variance before application of parametric tests. The data related to growth performance indices, Hematological indices, the liver mRNA gene expression levels were subjected to one-way analysis of variance (ANOVA) and if significant differences (p<0.05) were found, Tukey's post hoc test was used for ranking the groups by using of the SPSS (version 18).

#### **Results**

## Growth performance

The effects of ginger powder in common carp diet on the growth performance and survival rate are represented in Table 3. The results of statistical analysis of growth performance indices showed significant improvement (p<0.05) in final weight, weight gain, SGR and FCR in fishes fed 2 and 5% ginger diets compared with the control diet at the end of experiment. No mortality was recorded during feeding trial and survival rate was 100% in all treatment groups.

Table 3: Growth parameters and survival rate of common carp juveniles fed with different ginger level diets for 56 days feeding trial (data expressed as Mean $\pm$ SE). Means marked by different letters are significantly different (p<0.05).

		9	<u> </u>		
Performance	Dietary groups				
indices	control	0.5 %	2%	5%	
Initial weight (mg)	$14.51 \pm 0.30$	$14.87 \pm 0.49$	$13.88 \pm 0.63$	$13.90 \pm 0.66$	
Final weight (mg)	$42.51\pm3.46^{a}$	$45.67 \pm 2.56^{a}$	$60.08 \pm 3.10^{b}$	$71.20 \pm 4.31^{c}$	
Weight gain (%)	$19.3\pm2.9^{a}$	$20.7\pm3.3^{a}$	$33.3 \pm 4.12^{b}$	$41.2 \pm 5.74^{\circ}$	
SGR (% day <sup>-1</sup> )	$1.92\pm0.13^{a}$	$2.00\pm0.17^{a}$	$2.6 \pm 0.25^{b}$	$2.9 \pm 0.22^{c}$	
FCR	$1.79\pm0.05^{a}$	$1.75\pm0.07^{a}$	$1.55 \pm 0.09^{b}$	$1.42 \pm 0.04^{b}$	
Survival rate (%)	100	100	100	100	

Haematological and biochemical parameters

The effects of different levels of dietary ginger on the haematological parameters are presented in Table 4.

WBC, RBC, haematocrit, lymphocyte, monocyte and neutrophil levels were significantly higher in the fish fed with 2 and 5% ginger powder diets, compared with the other groups (p<0.05).

Table 4: Hematological parameters of common carp juveniles fed with different ginger level diets for 56 days feeding trial (data expressed as M  $\pm$  SE). Means marked by different letters are significantly different (p<0.05).

Danamatana	Dietary groups				
Parameters	Control	0.5 %	2 %	5 %	
WBC (10 <sup>4</sup> cell mm <sup>-3</sup> )	24.83±3.1 <sup>a</sup>	$32.41 \pm 2.2^{b}$	47.43±3.1°	49.33±3.3°	
RBC $(10^6 \text{cell mm}^{-3})$	$1.6\pm0.3^{a}$	$1.5\pm0.2^{a}$	$1.8\pm0.15^{b}$	$1.9 \pm 0.25^{b}$	
Haematocrit(%)	$30.12\pm2.23^{a}$	$31.4\pm3.1^{a}$	$38.2 \pm 2.5^{b}$	$41.4\pm3.8^{b}$	
Lymphocytes(%)	32.00±1.7 a	35.00±1.6 a	$42.30\pm2.3^{\text{ b}}$	$47.50\pm2.60^{\text{ b}}$	
Monocytes(%)	19.66±0.2 a	20.5±0.17 a	$26.7\pm0.2^{\mathrm{b}}$	$28.5\pm0.3^{\text{ b}}$	
Neutrophils (%)	28.33±0.71 a	30.53±0.55 a	$35.60\pm0.66^{b}$	36.36±0.95 b	

The Antioxidant enzyme activity and malondialdehyde levels were presented in Table 5. SOD, CAT and GPX enzyme activities and MDA level of the

fish fed 2 and 5% ginger-supplemented diets were significantly higher than that of fish fed with the control and 0.5% ginger-supplemented diets (p<0.05).

Table 5: Antioxidant capacity parameters of common carp juveniles fed with different ginger level diets for 56 days feeding trial (data expressed as Mean $\pm$  SE). Means marked by different letters are significantly different (p<0.05).

Parameter	Diet group				
r ar ameter	control	0.5%	2%	5%	
SOD(unit ml <sup>-1</sup> )	46.28±0.62 <sup>a</sup>	$42.42\pm1.29^{ab}$	34.92±1.33 <sup>b</sup>	31.65±3.98 <sup>b</sup>	
CAT(unit ml <sup>-1</sup> )	$3.98\pm0.70^{\rm s}$	$3.71\pm0.13^{s}$	$2.20\pm0.39^{b}$	$2.32\pm0.33^{b}$	
GPX (unit ml <sup>-1</sup> )	$30.68\pm7.20^{a}$	$28.60\pm4.50^{ab}$	$23.50\pm3.25^{b}$	$20.91\pm3.1^{b}$	
MDA (nmol ng <sup>-1</sup> )	5.8±0.52 <sup>a</sup>	5.5±0.48 <sup>a</sup>	4.3±0.26 <sup>b</sup>	3.9±0.3 <sup>b</sup>	

Expression of CAT, SOD and GPX mRNAs

The mRNA levels of common carp liver genes expression relative to  $\beta$ -

Actin are showed in Table 6. The expression of SOD, CAT and GPX genes in fish liver was affected by the ginger supplementation. SOD and CAT

mRNAs expression levels had significant differences (p<0.05) between fishes fed with 2 and 5% ginger-supplemented diets and fish fed in other treatment groups. GPX gene

expression levels showed significant increase (p<0.05) in fishes received 5% ginger-supplemented diet in comparison to the other treatment groups.

Table 6: Relative mRNA levels of common carp juvenile's liver exposed to four dietary ginger powder levels for 56 days feeding trial (data expressed as Mean±SE and n=5). Means marked by different letters are significantly different (p<0.05).

Comos		Diet group			
Genes	control	0.5%	2%	5%	
SOD	6.90±0.51 <sup>a</sup>	5.79±0.11 <sup>a</sup>	$3.53\pm0.24^{b}$	$3.06\pm0.20^{b}$	
CAT	$9.79\pm0.68^{a}$	$8.12\pm0.22^{a}$	$5.18\pm0.32^{b}$	$3.20\pm0.49^{c}$	
GPX	$8.24\pm0.31^{a}$	8.17±0.41 <sup>a</sup>	$6.76\pm0.42^{ab}$	$4.15\pm0.18^{b}$	

#### **Discussion**

Among phytochemical the agents application in aquaculture, ginger as a dietary supplementation has widely used in recent decade. The present study indicated that dietary ginger levels of 2 and 5% could improve the final weight, weight gain, SGR and FCR of common carp. The previous studies demonstrated that administration of ethanolic extract and powdered ginger increases the growth performance in Oncorhynchus mykiss (Shaluei et al., 2016), Lates calcarifer (Talpur et al., 2013), Macrobrachium rosenbergii (El-Desouky et al., 2012) and O. mykiss (Nya and Austin, 2009). However, in our study the higher growth was dependent to the ginger level, suggesting that 2 and 5% ginger diets were most favorable for growth performance of common carp. So, improvement of FCR could be due to the role of ginger as an appetizer. Thus, the increase of the growth rate could be due to enhancement of digestibility and metabolic process. Common carp as an omnivorous fish has a long intestine, in order to have a

better absorption capacity related to antiflammatory and antiproliferation effects (Pérez-Sánchez *et al.*, 2015). In addition, the functional mechanisms of dietary ginger effect on the growth of fishes are still not well understood and need to be further studied.

Generally, haematological and biochemical parameters are indicators of fish health condition as well as the physiological status of an organism (Gharaei *et al.*, 2016). White blood cells are identified as indicators of health status in fish and play an important role in regulation of immunologic function (Ballarin *et al.*, 2004; Van Hai, 2015), because their numbers enhance immediately when infections arise.

In this study, the WBC and RBC counts, haematocrite, lymphocyte and other blood cells increased in the 2 and 5% ginger diets which demonstrate the immunostimulatory and anti-infection properties of ginger. Improvement of blood cell counts were also reported by Talpur *et al.* (2013) and Nya and Austin (2009) who obtained enhanced blood cells in *L. calcarifer* and *O.* 

mykiss after feeding them with ginger. It could also be related to bioactive compounds of ginger including polyphenols, flavonoids, tannins and saponins that play anti-infection role in fish. Moreover, the increase of RBC count in fish fed with 2 and 5% ginger diets implies the suitable health effect in them due to bioactive components in ginger. Similar results have been reported about the ginger effect on RBC count in L. calcarifer (Talpur et al., 2013) and O. mykiss (Nya and Austin 2009). So, many studies demonstrated the effect of immunostimulant herbal plants on blood cells because they induce immune system in aquatic animals (Sahu et al., 2007; Harikrishnan et al., 2011; Talpur and Ikhwanuddin., 2012; Hwang et al., 2013; Van Hai, 2015).

Similar to other blood cells, lymphocytes, monocytes and neutrophils levels were improved by 2 and 5% ginger diets. These results could be implied to the task of these cells, because they increase when the organism is exposed to infections caused by pathogens and other foreign toxic material. These finding are comparable with Talpur *et al.* (2013), Talpur and Ikhwanuddin (2012) and Nya and Austin (2009).

The haematocrite (Hct) % increase in 2 and 5% ginger fed groups which could be explained by immunostimulatory effect of ginger. Typically, a higher Hct level signifies the blood sample's ability to transport oxygen and also serve as an indicator of health conditions (Birchard, 1997). Similarly, increased Hct % was

observed in *L. calcarifer* and *O. mykiss* fed by ginger diet in Talpur *et al.*, (2013) and Nya and Austin (2009) studies.

The earlier studies stated that more than 60 different medicinal plant species were used for the improvement of fish health and disease management in aquaculture (Van Hai, 2015) and some of them could enhance the antioxidant enzyme activities in adaptation organisms for environmental stress which may lead excessive ROS production (Livingston, 2001; Lukaszewicz-Hussain, 2010). SOD, CAT and GPX enzymes involving in antioxidant system are identified as the first defense mechanism against oxidative by stress radical process phagocytosis within damaged tissue (Abhijith et al., 2016). However, the major role of these enzymes is to neutralize the ROS produced different organs (Zhang et al., 2011). The significant decrease which was observed in SOD, CAT and GPX enzyme activities in liver of fishes fed with 2 and 5% ginger diets suggested that the ginger stronger O<sub>2</sub> quenching activity than these enzymes, dietary ginger could relieve oxidative stress and maintain healthy condition of individuals. Our finding corroborates the study by Zhang et al., (2011) who reported significant decreases in SOD and CAT expression levels following administration of astaxanthin in L. vannamei.

Generally, the production of reactive aldehyde is used as a biomarker to

measure the level of oxidative stress in an organism (Del Rio et al., 2005). Besides, ROS degrade polyunsaturated lipids by forming malondialdehyde (MDA) (Zheng et al., 2012). The decline of MDA level in fish fed with 2 and 5% ginger diets indicated that ginger could prevent the lipid damage caused by free radicals (Talas and Duran, 2012). The results may be explained that some of the existing compounds in ginger such flavonoids could reduce the oxidative stress by ROS scavenging (Fuat Gulhan et al., 2012).

Many studies reported that the ingredients or nutritional factors in feed could affect the expression of antioxidant enzymes of fish (Lin *et al.*, 2007; Tovar-Ramírez *et al.*, 2010; Zheng *et al.*, 2012).

But the effects of dietary ginger on antioxidant enzymes mRNA levels in fish are not consistent in previous studies. In present study, the expression levels of CAT, SOD and GPX mRNAs were examined among fishes fed with all treatment diets. The lower expression levels of these genes recorded in fishes fed with 2 and 5% ginger diets and their expression profiles were similar together and parallel with antioxidant enzymes activity levels (Table 5). Also, it was demonstrated that increase in the expression level of these genes was attributed to increase in expression level of Nfr2 gene and decrease in expression level of Keap1 (Sukumaran et al., 2016). Because the Nfr2 gene plays as a key moderator in primarily regulating the expression antioxidant enzymes and have been shown to increase the expression levels of SOD and GPX associated with increasing of the Nfr2 gene expression (Chen et al., 2015). The results of this study suggested that dietary ginger could partially alleviate oxidative stress by involvement in the expression of Nfr2 gene in common carp, which could help to enhance the scavenging ability of ROS in tissue cells.

In conclusion, the present study revealed that dietary ginger improves performance and feed growth utilization efficiency of common carp. In addition, the supplementation of ginger in fish feed can enhance the antioxidant capacity in response to oxidative stress in enzymatic and molecular functional levels in the tissue cells, which results in the increase of stress resistance in fish.

# Acknowledgments

We thank Dr. Ghaffari for manuscript review, Mr. Khandan Barani and Mr. Rahdari for their time and energy, Mrs. Miri for technical assistance in experimental analysis, and all staff of Hamoon International Wetland Research Institute for financial support and cooperation. The research project was funded by University of Zabol (Grant cod: UOZ-GR-9517-16).

#### References

- Abhijith, B.D., Ramesh, M. and Poopal, R.K., 2016. Responses of metabolic and antioxidant enzymatic activities in gill, liver and plasma of *Catla catla* during methyl parathion exposure. *The Journal of Basic and Applied Zoology*, 77, 31-40. DOI:10.1016/j.jobaz.2015.11.002
- **Aebi, H., 1984.** Catalase in vitro. *Methods in Enzymology,* 105, 121-126. DOI:10.1016/S0076-6879(84)05016-3
- Ahmad, I., Hamid, T., Fatima, M., Chand, H.S., Jain, S.K., Athar, M. and Raisuddin, S., 2000. Induction of hepatic antioxidants in freshwater catfish (*Channa punctatus* Bloch) is a biomarker of paper mill effluent exposure, *Biochim Biophys Acta*, 1523, 37-48.
- Akrami, R., Nasri Tajan, M., Jahedi, A., Jahedi, M., Razeghi Mansour, M. and Jafarpour, S.A., 2015. Effects of dietary synbiotic on growth, survival, lactobacillus bacterial count, blood indices and immunity of beluga (*Huso huso* Linnaeus, 1754) juvenile. *Aquaculture Nutrition*, 21(6), 952-959. DOI: 10.1111/anu.12219
- Alishahi, M., Ranjbar, M., Ghorbanpoor, M., Peyghan, R. and Mesbah, M., 2010. Effects of dietary Aloe vera on some specific and nonspecific immunity in the common carp (Cyprinus carpio). International Journal of Veterinary Research, 4, 189-95.
- Arulvasu, C., Mani, K., Chandhirasekar, D., Prabhu, D. and Sivagnanam, S., 2013. Effect

- of dietary administration of Zingiber officinale on growth, survival and immune response of Indian major carp, Catla catla (Ham.). International Journal of Pharmacy and Pharmaceutical Sciences, 5, 108-115.
- Ballarin, L., Dall'Oro, M., Bertotto, D., Libertini, A., Francescon, A. and Barbaro, A., 2004. Haematological in Umbrina cirrosa parameters (Teleostei, Sciaenidae): a comparison between diploid and triploid specimens. Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology, 138, 51.DOI:org/10.1016/j.cbpb.2004.02.01 9.
- Beyer, W.F. and Fridovich, I., 1987.
  Assaying for superoxide dismutase activity: some large consequences of minor changes in conditions.

  Analytical Biochemistry, 161(2), 559-566.
- **Birchard, G.F., 1997.** Optimal hematocrit: theory, regulation and implications. *American Zoologist,* 37(1), 65-72.
- Cerezuela, R., Cuesta, A., Meseguer, J. and Esteban, M.Á., 2008. Effects of inulin on gilthead sea bream (*Sparus aurata* L.) innate immune parameters, *Fish & Shellfish Immunol*ogy, 24, 663-668. DOI: 10.1016/j.fsi.2007.10.002.
- Chan, T.I., Roy, S.D., Sharma, A., Biswas, P., Siddhya, G.M. and Das, A., 2014. Anti stress potential of acetone extract of *Zingiber officinale* Roscoe on biochemical and oxidative stress parameters in *Labeo calbasu* (Hamilton 1822) fingerlings subjected to acid stress. *Indian Journal Fisheries*, 61, 69-77.

- Chang, Y.P., Liu, C.H., Wu, C.C., Chiang, C.M., Lian, J.L. and Hsieh, S.L., 2012. Dietary administration of zingerone to enhance growth, nonspecific immune response, and resistance to *Vibrio alginolyticus* in Pacific white shrimp (*Litopenaeus vannamei*) juveniles, Fish & Shellfish Immunology, 32, 284-290. DOI: 10.1016/j.fsi.2011.11.017.
- Chen, B., Lu, Y., Chen, Y. and Cheng, J., 2015. The role of Nrf2 in oxidative stress-induced endothelial injuries. *Endocrinology*, 225(3), 83-99. DOI:10.1530/JOE-14-0662
- Dada, A.A. and Ikuerowo, M., 2009.

  Effects of ethanolic extracts of
  Garcinia kola seeds on growth and
  haematology of catfish (*Clarias*gariepinus) brood stock. *African*Journal of Agricultural Research, 4,
  344-347.
- Dadfar, F., Hosseini, S.E. and Bahaoddini, A., 2014. A review of phytochemical, pharmacological and physiological properties of ginger (*Zingiber officinale*). *Clinical Excellence*, 3(1), 72-86.
- Pellegrini, N., 2005. A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutrition Metabolism Cardiovascular Diseases*, 15(4), 316-328. DOI:10.1016/j.numecd.2005.05.003
- El-Desouky, H., El-Asely, A., Shaheen, A.A. and Abbass, A., 2012. Effects of *Zingiber officinalis* and *Cyanodon dactylon* on the growth performance and immune parameters of *Macrobrachium*

- rosenbergii. World Journal of Fish and Marine Sciences, 4(3), 301-307.
- DOI:10.5829/idosi.wjfms.2012.04.0 3.62120
- Faggio, C., Casella, S., Arfuso, F., Marafioti, S., Piccione, G. and Fazio, F., 2013. Effect of storage time on haematological parameters in mullet, *Mugil cephalus. Cell Biochemistry and Function*, 31(5), 412-416. DOI:10.1002/cbf.2915
- Faggio, C., Piccione, G., Marafioti, S., Arfuso, F., Trischitta, F., Fortino, G. and Fazio, F., 2014. Monthly variations of haematological parameters of *Sparus aurata* and *Dicentrarchus labrax* reared in Mediterranean land off-shore tanks. *Cahiers de Biologie Marine*, 55, 437-443.
- Fallahpour, F., Banaee, M. and Javadzade, N., 2014. Effects of dietary marshmallow (Althaea officinalis L.) extract on growth performance and body composition of common carp (Cyprinus carpio), International Journal of Advanced Biological and Biomedical Research, 2, 2453-60
- Fuat Gulhan, M., Duran, A., Selamoglu Talas, Z., Kakoolaki, S. and Mansouri, S.M., 2012. Effects of propolis on microbiologic and biochemical parameters of rainbow trout (*Oncorhynchus mykiss*) after exposure to the pesticide. Iran. *Iranian Journal of Fisheries Sciences*, 11(3), 490-503.
- García-triana, A., Gómez-Jiménez, S., Peregrino-Uriarte, A.B., López-Zavala, A., González-

- Aguilar, G., Sotelo-Mundo, R.R., Valenzuela-Soto, E.M. and Yepiz-Plascencia, G., 2010. Expression and silencing of Selenoprotein M (SelM) from the white shrimp *Litopenaeus* vannamei: Effect on peroxidase activity and hydrogen peroxide concentration in gills and hepatopancreas. Comparative Biochemistry and Physiology - Part A: Molecular and Integrative Physiology, 155(2), 200-204. DOI: 10.1016/j.cbpa.2009.10.037.
- Gharaei, A., Ghaffari, M., Keyvanshokooh, S. and Akrami, R., 2011. Changes in metabolic enzymes, cortisol and glucose concentrations of beluga (Huso huso) exposed to dietary methylmercury. Fish Physiology and Biochemistry, 37(3), 485-493. DOI:10.1007/s10695-010-9450-3
- Gharaei, A., Rayeni, M.F., Ghaffari, M., Akrami, R. and Ahmadifar, E., 2016. Influence of dietary prebiotic mixture a-mune) on growth performance, haematology and innate immunity of Beluga sturgeon (Huso huso) juvenile. International Journal of Aquatic Biology, 4(4), 277.
- Harikrishnan, R., Balasundaram, C. and Heo, M.S., 2011. Impact of plant products on innate and adaptive immune system of cultured finfish and shellfish. *Aquaculture*, 317(1), 1-15. DOI:10.1016/j.aquaculture.2011.03. 039
- Hwang, J.H., Lee, S.W., Rha, S.J., Yoon, H.S., Park, E.S., Han,

- **K.H.,and Kim, S.J., 2013.** Dietary green tea extract improves growth performance, body composition, and stress recovery in the juvenile black rockfish, *Sebastes schlegelii*, *Aquaculture International.*, 21, 525-538.
- R.P., Immanuel, G., Uma, P., Iyapparaj, Citarasu, T., Punitha Peter, S. M., Michael Babu, M. and Palavesam, A., 2009. Dietary medicinal plant extracts improve growth, immune activity and survival of tilapia Oreochromis mossambicus. Journal of Fish Biology, 74(7), 1462-1475. DOI:10.1111/j.1095-8649.2009.02212.x
- Jiang, W., Zhu, J., Yang, Q., Zhao, H., Chen, M., He, H. and Yu, W., 2016. Complete mitochondrial DNA genome of *Polytremis nascens* (Lepidoptera: Hesperiidae). *Journal of Mitochondrial DNA*, 27(5), 3131-3132. DOI: 10.3109/19401736.2015.1007298
- Jusman, S. W., Halim, A., Wanandi, S. I. and Sadikin, M., 2010.

  Expression of hypoxia-inducible factor-1alpha (HIF-1alpha) related to oxidative stress in liver of ratinduced by systemic chronic normobaric hypoxia. *Acta Medical Indonesiana*, 42(1), 17.
- Lee, D.H., Lim, S.R., Han, J.J., Lee, S.W., Ra, C.S. and Kim, J.D., 2014. Effects of dietary garlic powder on growth, feed utilization and whole body composition changes in fingerling sterlet sturgeon, *Acipenser ruthenus*.

- Asian-Australasian Journal of Animal Science, 27(9), 1303-1310.
- Levy, G., Zilberg, D., Paladini, G., and Fridman, S., 2015. Efficacy of ginger-based treatments against infection with *Gyrodactylus turnbulli* in the guppy (*Poecilia reticulata* (Peters)). *VeterinaryParasitology*,209, 235-241. DOI: org/10.1016/j.vetpar.2015.03.002.
- Lin, R., Wang, X., Luo, Y., Du, W., Guo, H. and Yin, D., 2007. Effects of soil cadmium on growth, oxidative stress and antioxidant system in wheat seedlings (*Triticum aestivum* L.). Chemosphere, 69, 89-98. DOI: 10.1016/j.chemosphere.2007.04.041.
- **Livingston, J.C., 2001.** Anatomy of the Sacred: An introduction to Religion, Prentice Hall.
- **Lukaszewicz-Hussain, A., 2010.** Role of oxidative stress in organophosphate insecticide toxicity—Short review. *Pesticide Biochemistry and Physiology*, 98, 145-50. DOI: org/10.1016/j.pestbp.2010.07.006.
- Mahghani, F., Gharaei, A., Ghaffari, M. and Akrami, R., 2014. Dietary synbiotic improves the growth performance, survival and innate immune response of Gibel carp (Carassius auratus gibelio) juveniles, International Journal of Aquatic Biology, 2, 99-104. DOI: org/10.22034/ijab.v2i2.37.
- Nan, F.H., Agus Putra, A.S., Margie, B. and Lee, M.C., 2015. The effects of *Curcuma zedoaria* and *Zingiber zerumbet* on non-specific immune responses of grouper *Epinephelus coioides*. *Journal of Fisheries Sciences*, 14(3), 598-611.
- **Nya, E.J. and Austin, B., 2009.** Use of dietary ginger, *Zingiber*

- officinale Roscoe, as an immunostimulant to control Aeromonas hydrophila infections in rainbow Oncorhynchus trout, mykiss (Walbaum). Journal of Fish 971-977. Diseases, 32(11), DOI:10.1111/j.1365-2761.2009.01101.x
- Otunola, G.A., Oloyede, O.B. and Oladiji, **A.T.**, 2010. Afolayan, Comparative analysis of the chemical composition of three spices-Allium sativum L. Zingiber officinale Rosc. And Capsicum frutescens commonly consumed in Nigeria, African Journal. Biotechnology, 9, 6927-6931.
- **Paglia, D.E. and Valentine, W.N., 1967.** Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. *Translational Research*, 70(1), 158-169.
- Pakravan, S., Hajimoradloo, A. and Ghorbani, R., 2012. Effect of dietary willow herb, Epilobium hirsutum extract on growth performance, body composition, haematological parameters Aeromonas hydrophila challenge on common carp, Cyprinus carpio. Aquaculture Research, 43(6), 861-869. DOI:10.1111/j.1365-2109.2011.02901.x
- Pérez-Sánchez, J., Benedito-Palos, L., Estensoro, I., Petropoulos, Y., Calduch-Giner, J.A., Browdy, C.L. and Sitjà-Bobadilla, A., 2015. Effects of dietary NEXT ENHANCE® 150 on growth performance and expression of immune and intestinal integrity related genes in gilthead sea bream

- (Sparus aurata L.). Fish and Shellfish Immunology, 44(1), 117-128. DOI:10.1016/j.fsi.2015.01.039
- **Prakash, J., 2010.** Chemical composition and antioxidant properties of ginger root (*Zingiber officinale*). *Journal of Medicinal Plants Research*, 4(24), 2674-2679. DOI: 10.5897/JMPR09.464
- Sahu, S., Das, B.K., Mishra, B.K., Pradhan, J. and Sarangi, N., 2007. Effect of *Allium sativum* on the immunity and survival of *Labeo rohita* infected with *Aeromonas hydrophila*. *Journal of Applied Ichthyology*, 23, 80-86. DOI: org/10.1111/j.1439-0426.2006.00785.x
- Shaluei, F., Nematollahi, A., Naderi Farsani, H.R., Rahimi, R. and Kaboutari Katadj, J., 2016. Effect of ethanolic extract of *Zingiber officinale* on growth performance and mucosal immune responses in rainbow trout (*Oncorhynchus mykiss*). Aquaculture Nutrition, 23(4), 814–821. DOI:10.1111/anu.12448.
- Sheikhzadeh, S., Kade, C., Keyser, Stuhrmann, M., B., Arslan Kirchner, M., Rybczynski, M., Bernhardt, A.M., Habermann, C. R., Hillebrand, M., Mir, T. and Robinson, P.N., 2012. Analysis of phenotype and genotype information for the diagnosis of Marfan syndrome. Clinical Genetic, 240-247. DOI: 82(3), 10.1111/j.1399-0004.2011.01771.x
- Soosean, C., Marimuthu, K., Sudhakaran, S. and Xavier, R., 2010. Effects of mangosteen (Garcinia mangostana L.) extracts

- as a feed additive on growth and haematological parameters of African catfish (*Clarias gariepinus*) fingerlings. *European Review for Medical and Pharmacological Sciences*, 14, 605-611.
- Sukumaran, V., Park, S.C. and Giri, S.S., 2016. Role of dietary ginger Zingiber officinale in improving growth performances and immune functions of Labeo rohita fingerlings. Fish and Shellfish Immunology, 57, 362-370. DOI:10.1016/j.fsi.2016.08.056
- **Talas, Z.S. and Duran, A., 2012.** The effects of slaughtering methods on physical and biochemical changes in fish. *Energy Education Science and Technology*, 29(2), 741-748.
- Talpur, A.D. and Ikhwanuddin, M., 2012. Dietary effects of garlic (Allium sativum) on haematoimmunological parameters, survival, growth, and disease resistance against Vibrio harveyi infection in Asian sea bass, Lates calcarifer (Bloch). Aquaculture, 364, 6-12.DOI:10.1016/j.aquaculture.2012.09. 020
- Talpur, A.D., Ikhwanuddin, M. and Bolong, A.M.A., 2013. Nutritional effects ginger (Zingiber officinale Roscoe) on immune response of Asian sea bass, Lates calcarifer (Bloch) and disease resistance against Vibrio harveyi. 400, 46-52. Aquaculture, DOI:10.1016/j.aquaculture.2013.02. 043
- Tokur, B., Ozkütük, S., Atici, E., Ozyurt, G. and Ozyurt, C.E.,

- **2006.** Chemical and sensory quality changes of fish fingers, made from mirror carp (*Cyprinus carpio* L., 1758), during frozen storage (-18 °C). *Food Chemistry*, 99(2), 335-341.
- DOI:10.1016/j.foodchem.2005.07.0
- Tovar-Ramírez, D., Mazurais, D., Gatesoupe, J.F., Quazuguel, P., Cahu, C.L. and Zambonino-Infante, J.L., 2010. Dietary probiotic live yeast modulates antioxidant enzyme activities and expression of sea bass (Dicentrarchus *labrax*) larvae. Aquaculture, 300(1),142-147. DOI:10.1016/j.aquaculture.2009.12. 015
- Trenzado, C., Hidalgo, M. C., García-Gallego, M., Morales, A. E., Furné, M., Domezain, A., Domezain, J. and Sanz, A., 2006. Antioxidant enzymes and lipid peroxidation in sturgeon Acipenser naccarii and trout Oncorhynchus comparative mykiss. A study. 758-767. Aquaculture, 254(1), DOI:10.1016/j.aquaculture.2005.11. 020
- Vahedi, G., Kanno, Y., Furumoto, Y., Jiang, K., Parker, S. C., Erdos, M. R., Davis, S. R., Roychoudhuri, R., Restifo, N.P., Gadina, M. and Tang, Z., 2015. Super-enhancers delineate disease-associated regulatory nodes in T cells. *Nature*, 520(7548), 558-562. DOI:10.1038/nature14154
- Van Hai, N., 2015. The use of medicinal plants as immunostimulants in aquaculture:

- A review. Aquaculture, 446, 88-96. DOI:10.1016/j.aquaculture.2015.03. 014Zhang, J., Khvorostov, I., Hong, J. S., Oktay, Y., Vergnes, L., Nuebel, E., Wahjudi, P.N., Setoguchi, K., Wang, G. and Do, A. Jung, H.J., McCaffery, J.M., Kurland, I.J., Reue, K., Lee, W.N., Koehler, C.M. and Teitell, M.A., 2011. UCP2 regulates energy metabolism and differentiation potential of human pluripotent stem cells. EMBO, 30, 4860-4873. DOI:10.1038/emboj.2011.401
- Yuan, J., Wang, X., Gu, Z., Zhang, Y. and Wang, Z, 2015. Activity and Transcriptional Responses of Hepatopancreatic Biotransformation and Antioxidant Enzymes in the Oriental River Prawn *Macrobrachium nipponense* Exposed to Microcystin-LR. Toxins, 7, 4006-4022. DOI: 10.3390/toxins7104006.
- Zhang, Y., Pan, Y., Yang, W., Liu, W., Zou, H. and Zhao, Z.K., 2013. Protein arginine allylation and subsequent fluorophore targeting. *ChemBioChem*, 14(12), 1438-1443. DOI: 10.1002/cbic.201300176
- Zheng, Q., Wen, X., Han, C., Li, H. and Xie, X., 2012. Effect of soybean meal replacing with cottonseed meal growth. on hematology, antioxidant enzymes activity and expression for juvenile grass carp, Ctenopharyngodon Fish Physiology idellus. 38(4), 1059-1069. Biochemistry, DOI:10.1007/s10695-011-9590-0