# Effects of 2-phenoxyethanol (2-PE) anesthesia on some haematological and biochemical indices of silver carp (Hypophthalmichthys molitrix)

# Hedayati A.A.\*

Received: November 2015 Accepted: July 2016

### **Abstract**

In this study, the anesthetic effects of 2-phenoxyethanol (2-PE) on possible primary (cortisol level) and secondary (haematological indices and glucose level) stress responses were studied in silver carp (Hypophthalmichthys molitrix). Fish were first exposed to 0.1, 0.3, 0.5, 0.7 and 0.9 mL L<sup>-1</sup> 2-PE, and the time of induction (deep anesthesia) and recovery were measured. At concentrations of 0.5, 0.7 and 0.9 mL L<sup>-1</sup> 2-PE, all fish were anaesthetized within 3 min of exposure. For assessing possible stress effects caused by effective concentrations of 2-PE, the haematological indices, serum cortisol and glucose were determined in the deeply anaesthetized fish as stress indicators. 2-PE exposure resulted in a significant increase in red blood cell (RBC) amounts at 0.1 and 0.3 mL L<sup>-1</sup>. A parallel increase in hemoglobin and Haematocrite amounts at 0.1 concentrations (p<0.05) was observed. The serum cortisol levels had the highest amount in 0.1 mL L<sup>-1</sup> of 2-PE. Moreover 2-PE exposure resulted in a significant increase in glucose amounts first at the 0.1 and later in the 0.3 mL L<sup>-1</sup> concentrations. This study shows that rapid induction of deep anesthesia with a relatively high concentration of 2-PE (0.5, 0.7 and 0.9 mL L<sup>-1</sup>) was associated with the lowest effects on the haematological and serum biochemical indices in silver carp and so 0.7 mL L<sup>-1</sup> could be suggested as a suitable dose for haematological studies in this species.

**Keywords**: Blood parameters, Anesthesia, Stress response, Silver carp

Department of Aquatics Production and Exploitation, Faculty of Fisheries and Environmental Sciences, Gorgan University of Agricultural Sciences and Natural Resources, Gorgan, Iran. \*Corresponding author's Email: Hedayati@gau.ac.ir

# Introduction

Anesthetics are physical or chemical operatives that make animals insensitive to pain by initially inducing a relaxing effect and afterwards inducing loss of balance, mobility, ability to perceive and reflex action (Summer and Smith, 1990). Anesthetics are greatly used in ordinary aquaculture performances to immobilize animals transportation, during spawning, vaccination and handling. Overdose of an anesthetic material may be utilized as a moral way to euthanize fish (Shaluei et al., 2012). The desirable characteristics of anesthetics used for finfish include: short induction and recovery time, non-toxicity to fish and humans, no long physiological effects, quick release from the body, steady under normal hatchery situations (light, heat) and non-foaming, readily attainable and cost efficient (Gholipour Kanani et al., 2011).

The efficacy and security of any anesthetic factor change among species, phases and environmental life conditions. A perfect anesthetic should cause anesthesia quickly with minimum hyperactivity or stress. It should be easy to supply and should preserve the animal in the selected state. When the animal is eliminated of the anesthetic, should be recoverv auick. anesthetic should be efficient at low doses and the toxic dose should greatly exceed the effective dose so that there is a vast border of security (Coyle et al., 2004). The energetic material of 2-PE is ethylene glycol monophenyl ether, with

the chemical formula C<sub>8</sub>H<sub>10</sub>O<sub>2</sub>, molar weight 138.17 g L<sup>-1</sup>, density 1.107 -1.108; peroxide content less than 0.005% and the boiling point 245 °C. The anesthetic is a little soluble in water (26.7 g L<sup>-1</sup>) but readily soluble in ethanol. 2-PE is a murky, greasy liquid. solution is bactericidal fungicidal, practical during surgery. It comparatively inexpensive remains active in the thinned state for at least 3days. 2-PE has a comparatively wide margin of security and has been announced to produce a range of effects light sedation surgical from anesthesia at concentrations of 100-600 mg L<sup>-1</sup>. 2-PE is not confirmed by FDA for use in fish in the U.S (Coyle et al., 2004). The anesthetic 2-PE is utilized not only for short-term immobilization of fish before reproduction, but also at any time fish are handled outside water, e.g. during veterinary interferences.

Estimation of blood cells, serum biochemistry and hormones are utilized as indicators of the physiological or sub lethal stress response in fish to endogenous or exogenous changes. In this regard, the rising of plasma cortisol and glucose levels is greatly used as a general indicator of stressful conditions (Hedayati *et al.*, 2015).

Silver carp is one of the prime cultured fresh water fish in polyculture and of great economic importance. It is vital to understand responses of silver carp to anesthetics. Therefore, this study was carried out to appraise the efficacy of 2-PE as an anesthetic on silver carp and its effect on

haematological and serum biochemical indices in this species.

### Material and methods

Preservation of fish

105 silver carp  $(71.6\pm5.8 \text{ g}, 20.3\pm3.1)$ cm) were supplied from the Bony Fish Propagation and Rearing Center of Sijeval (Bandar Torkaman, Gorgan, Iran). Fish were transferred to the aquaculture research center of Gorgan University of agricultural sciences and natural resources and acclimated to the laboratory conditions for 2 weeks. The average amounts for aerated and dechlorinated tap water were used during the acclimation period, water pH was 7.79±0.50, dissolved oxygen 7.93±0.25 mg L<sup>-1</sup>, temperature of 20.50±0.5°C and total hardness was 298±2.35 mg L<sup>-1</sup> as CaCO<sub>3</sub>. Water pH, temperature and dissolved oxygen were determined by Wagtech portable pH/temp meter and oxygen meter (Berkshire, UK). Water total hardness was determined using a portable photometer with commercial kits provided by the manufacturer (Wagtech Por-table Photometer 7100, Berkshire, UK). Throughout acclimation and experiment periods, fish were held under natural photoperiod conditions (11:13 lightdark).

### 2-PE treatment

The 2-PE stock solution was prepared by dissolving pure 2-PE (Sigma Chemicals, St. Louis, MO, USA, 99.5%, d=1.107–1.108 g L<sup>-1</sup>) in 95% ethanol (1: 10 ratio of 2-PE–ethanol). Concentrations of 2-PE tested were 0.1,

0.3, 0.5, 0.7 and 0.9 mL L<sup>-1</sup>. Each mentioned concentration of 2-PE was added to a 100-L tank and the water well ventilated for 2 min prior to the experiment. Ten fish were separately exposed to each concentration and time (surgical induction or deep anesthesia phase; loss of responsiveness, lying on the bottom of tank and not reacting to handling) registered with a stop monitor (Shaluei et al., 2012). As soon as the deep anesthesia stage was reached, they were transferred to fresh and strongly-aerated tanks and the time for total recovery was recorded. Recovery time was registered as the time required for fish to regain balance and begin active swimming.

Haematological and biochemical analysis

For the haematological and serum biochemical tests, 10 fish were used for each efficient concentration (0.1, 0.3, 0.5, 0.7 and 0.9 mL L-1 2-PE) and for the controls. Fish were separately exposed to each concentration. When fish arrived at the surgical or deep anesthesia phase, the midline posterior of the anal fin of the juveniles was cleaned with erasing paper to prevent possible pollution with water, and three milliliters of blood was drawn from the caudal vein. Blood collection was carried out by expert persons and lasted less than 20s immediately after netting the fish head with wet cloth to diminish stress (Chardeh Baladehi et al., 2017). Blood samples were rapidly collected from the caudal blood vessel by heparinized syringes and preserved without delay on ice (Abarghuei et al., 2015). Estimation of the blood indices was carried out on fresh blood. Leukocyte and erythrocyte count was measured by diluting heparinized blood with Giemsa stain at 1:30 dilution and cells were counted using a Neubauer under hemocytometer the light microscope (Hedavati and Niazi. 2015).

Haematocrit amounts (Ht%) were determined after sampling by placing fresh blood in glass capillary tubes and centrifuging for 5 min at 10000 rpm in micro Haematocrit centrifuge (Hettich, Germany). The packed cell volume was then calculated (Abarghuei et al., 2015); Haematocrit readings were performed with the aid of a micro Haematocrit reader. Hemoglobin levels (Hb mg  $L^{-1}$ ) were determined colorimetrically measuring by cyanomethemoglobin formation of following standard methods (Hedayati and Niazi, 2015).

Blood samples were placed into containing **EDTA** tubes as an anticoagulant for the hematology tests. These parameters were determined: total white blood cell (Leukocytes) count, red blood cell (Erythrocytes) hemoglobin count, content and Haematocrite level. For the biochemical experiments, blood was put in tubes and allowed to clot at room temperature (25°C), for 30 min. Serum was removed from the clotted sample centrifugation at 5,000 rpm for 5 min and frozen at -80°C until analysis. Glucose levels were calculated using a

spectrophotometric method (WPAS2000-UV/VIS, Cambridge, UK) with reagents supplied in standard (Pars Azmon, Iran). analyses kits Cortisol was determined with DRG kit commercial (ELISA, Diagnostics, Mountainside, NJ, USA) based on an enzyme immunoassay. Before using an ELISA kit for determining the serum cortisol level in silver carp, this diagnostic kit was validated for use by tests of linear response of the sample and cortisol overload (Weber al., et2011). Experimental kit, which was used for human samples, was functional to the measurement of cortisol content in fish. Prior studies approved that the kit could identify the cortisol molecule of the fish species (Hosseini et al., 2015).

### Statistical analyses

All results are expressed as mean±SD. Statistical analyses were carried out using the computerized package 18.0 for Windows. Normality of data was first estimated using a Kolmogorov-Smirnov's test, and homogeneity of variance was assessed with Levene's test. Based on these tests, all data were found to be normally distributed. To the effect of different evaluate concentrations of 2-PE on induction and recovery times and haematological and biochemical indices, all data were **ANOVA** subjected to one-way followed by Tukey's test at a 5 % significance level.

### Results

No mortality was monitored during the experiments. Effects of different concentrations of 2-PE on induction and recovery times are shown in Fig. 1. Induction times generally decreased significantly with increasing 2-PE concentrations (p<0.05). Contrary to

this trend, recovery time from anesthesia increased as expected (Fig. 1).

Effects of different concentrations of 2-PE on the haematological indices of silver carp are given in Table 1.




Figure1: Mean  $(\pm SD)$  induction time to anesthesia and recovery of silver carp (n = 10) exposed to different doses of 2-PE.

2-PE exposure resulted in significant increase in RBC amounts at 0.1 and 0.3 mL L<sup>-1</sup> and a parallel increase in hemoglobin and Haematocrite amounts were observed at 0.1 mL L<sup>-1</sup> 2-PE (Table 1).

Results indicated that deep anaesthetization of fish using different concentrations of 2-PE led to an alteration in cortisol and glucose levels (Figs. 2 and 3).

The serum cortisol levels reached the highest level with 0.1 mL L<sup>-1</sup> of 2-PE. Moreover, 2-PE exposure resulted in a significant increase in glucose amounts first at 0.1 and then at 0.3 mL L<sup>-1</sup> concentrations.

Table1: Haematological	indicae in	cilvar carn avnaca	d to differen	t doses of 2-PF
- Labiet: Haematological	-inaices in	i siiver card exdose	a to anteren	Looses of Z-PF.

Parameters	Control	2-phenoxyethanol (mL L <sup>-1</sup> )					
rarameters		0.9	0.7	0.5	0.3	0.1	
RBC (×106 cells mL-1)	$1.91 \pm 0.083^{c}$	$2.59\pm0.11^a$	$1.90 \pm 0.06$ c	$1.95\pm0.070~^{c}$	$2.00\pm0.115$ $^{\rm c}$	$2.28 \pm 0.11^{b}$	
WBC ( $\times 10^3$ cells mL <sup>-1</sup> )	$6.66\pm0.24^{ab}$	$6.570 \pm 0.27^b$	$6.77\pm0.17^{ab}$	$7.18\pm0.25^{ab}$	$7.04\pm0.50^{ab}$	$7.35\pm0.55^a$	
Hb (g $dL^{-1}$ )	$6.90\pm0.08^b$	$7.00\pm0.45^b$	$6.95\pm0.07^b$	$6.99\pm0.02^{\mathrm{b}}$	$7.26\pm0.24^{ab}$	$7.66\pm0.10^a$	
Hct%	$20.87 \pm 1.30^{bc}$	$20.30\pm1.78^{c}$	$21.34 \pm 1.56^{bc}$	$21.29 \pm 1.07^{bc}$	$23.40 \pm 0.50^b$	$27.63 \pm 1.45^{a}$	
Hct%	$20.87 \pm 1.30^{bc}$	$20.30\pm1.78^c$	$21.34 \pm 1.56^{bc}$	$21.29 \pm 1.07^{bc}$	$23.40\pm0.50$ $^{b}$	$27.63 \pm 1.45^a$	




Figure 2: Serum cortisol levels in silver carp exposed to different doses of 2-PE.




Figure 3: Serum glucose levels in silver carp exposed to different doses of 2-PE.

# **Discussion**

Anesthetics are essential for many actions in aquaculture, because species may be differ greatly in their response to the same anesthetic. Screening of dosage of different anesthetics is often

essential. The concentration of the anesthetic in the central nervous system determines the stage of anesthesia (Shaluei *et al.*, 2012). The fish takes up the anesthetic through a density gradient at the tissue/water interface,

and some time is needed for the concentration of the anesthetic in the blood to arrive at the threshold level at which the fish becomes anaesthetized. Present results indicate 2-PE is an efficient anesthetic that can induce quick induction and fast recovery time in juvenile silver carp. Gilderhus and Marking (1987)contrasted effectiveness of 16 anesthetic chemicals in rainbow trout, Oncorhynchus mykiss, and stated that 2-PE had all the criteria for efficacy. Weber et al. (2011) compared the effects of 2-PE. metomidate, clove oil and MS-222 in Senegalese sole, Solea senegalensis, and discovered that 2-PE was the most effective of the four anesthetic agents. **McCarter** (1992)announced negative effects of anesthesia with 0.20 mL L<sup>-1</sup> 2-PE on spermatozoa and their motility in grass carp (Ctenopharyngodon idella).

An appropriate anesthetic should cause anesthesia within 3 min or less, and permit recovery within 5 min or less. The maximum allowed time for the induction of deep anesthesia in fish is 10 min (Ross and Ross, 2008). There was no data in open literature on biochemical blood profile in silver carp anaesthetized with 2-PE. In a perusal of rainbow trout anaesthetized with 0.30 mL L<sup>-1</sup> 2-PE, no significant negative influences of anesthesia on parenchymatous tissues were discovered (Veliekl and Svobodovan, 2004). The lowest concentration of the 2-PE utilized here that was capable of inducing deep anesthesia in silver carp was 0.1 mL L<sup>-1</sup>. Anesthetics reactivate

stress by causing a depression in the nervous system; dosage and time can influence the physiology and blood profile of the fish particularly stress indices (Veliekl and Svobodovan, 2004; Bolasina, 2006; Akbary et al., 2016). Different stressors operate the pituitary-inter renal hypothalamus-(HPI) axis, resulting in a cortisol release that causes secondary stress answers. Stress hormones make active number of metabolic routes that result in alternation in blood chemistry and hematology indices. In the current study, RBC, hemoglobin concentration haematocrite and were raised significantly when fish were exposed to the 2-PE, first, at the concentration of 0.1 and then at 0.3. Adámek et al. (1993) discovered raised count of RBC and concentration of Hb in common carp (Cyprinus carpio) following 2phenoxyethanol (0.30)mL.  $L^{-1}$ anesthesia.

This raise may be because of the release of unripe red cells from the spleen, to increase the oxygen supply to the main organs in answer to the higher metabolic requirements. In addition, this phenomenon could be a prompt response to the acute stress mediated by catecholamines (Tort et al., 2002). It is well known that there is a reduction in the count of leukocytes in ill fish which are exposed to any type of infection and expansion related to protection in leukocyte cells (Coyle et al., 2004). While there were no changes in rainbow trout right after the anesthesia, an increase is discovered 24h later (Velisek et al., 2007).

In our study, the concentrations of glucose increased significantly in fish exposed to 2-PE, first at 0.1 and then at 0.3 mL L<sup>-1</sup>. Under stressful conditions, increases in glucose levels occurred because of catecholamines and glucocorticoids release from adrenal tissues of fish (Shaluei *et al.*, 2012). Ortuno *et al.* (2002) announced a raise in glucose and cortisol amounts in *Sparus aurata* anaesthetized with 2-phenoxyethanol.

The raised plasma levels of glucose are compatible with those reported in Senegalese sole (Soleasene galensis) after exposure to 2-PE (500 mg L<sup>-1</sup>) (Velisek et al., 2007). Ortuno et al. (2002) explained an increase of glucose in gilthead bream anaesthetized with 2-PE. Different findings have been reported in other fish anaesthetized using 2-PE (Velisek et al., 2007). In the present study, a significant increase in cortisol level was observed in fish exposed to 2-PE at 0.1 mL L-1. Molinero and Gonzalez (1995) showed that both the dose and exposure to 2-PE caused a cortisol reply in gilthead sea bream.

Weber *et al.* (2011) found a significant increase in plasma concentrations of cortisol in Senegalese sole after 20min exposure to 2-PE (500 mg L<sup>-1</sup>). King *et al.* (2005) found cortisol response by juvenile and adult black sea bass because of netting and capture, but no significant changes in cortisol levels beyond the 10 to 30 min exposure to 20-PE (300 mg L<sup>-1</sup>). Different responses were monitored in other fish after exposure to 2-PE and other

anesthetics (e.g., King et al., 2005; Bolasina, 2006; Weber et al., 2011).

Generally, this study shows that rapid induction of deep anesthesia with a relatively high concentration of 2-PE (0.5, 0.7 and 0.9 mL L<sup>-1</sup>) was related the lowest effects on haematological and serum biochemical indices in silver carp and therefore can be suggested as suitable doses for haematological studies in this species. Fast anesthesia was observed with a higher dose (0.9 mL L<sup>-1</sup>) but more recovery time was also recorded at the same dose, However in terms of blood and serum indices, by increasing 2-PE, the concentration of physiological parameters returned to levels at normal conditions (best in 0.9 mL L<sup>-1</sup>), eventually because of long recovery time at higher concentration. Based on these results, 0.7 mL L<sup>-1</sup> of 2-PE can be suggested as the best dose of anesthesia in silver carp.

# Acknowledgement

The author is thankful to Gorgan University of Agricultural Sciences and Natural Resources, Gorgan – Iran, for their technical and financial support.

# References

Abarghuei, S., Hedayati, A., Ghafari Farsani, H. and Gerami, M.H., 2015. Haematological responses of goldfish to different acute concentrations of silver sulfate as a toxicant. *Pollution*, 1(3), 247-256.

Adamek, Z., Faic, K. and Paul, K., 1993. The effect of 2-

- phenoxyethanol narcosis on blood of young carp (*Cyprinus carpio* L). *Veterinary Archive*, 63, 245-250.
- Pirbeigi, Akbary, P., A. and Jahanbakhshi, A. 2016. Analysis of primary and secondary stress responses in bighead carp (Hypophthalmichthys nobililis) by anesthetnization 2with phenoxyethanol. **International** Journal of Environmental Science and Technology, 13(4), 1009-1016.
- Bolasina, S.N., 2006. Cortisol and haematological response in Brazilian codling, *Urophycis brasiliensis* (Pisces, Phycidae) subjected to anesthetic treatment. *Aquaculture International*, 14, 569-575.
- Chardeh Baladehi, E., Hedayati, A. Miandareh. Kolangi H. Bagheri, T., 2017. The hematological improvement of rainbow trout during dietary supplementation with Vitamin C after exposure to zinc nano particles. **Fisheries** Iranian **Journal** of Sciences, 16(1), 162-169.
- Coyle, S.D., Durborow, R.M. and Tidwell, J.H., 2004. Anesthetics in aquaculture. SRAC Publication, Texas. 6P.
- Gholipour kanani, H., Mirzargar, S.S., Soltani, M., Ahmadi, M., Abrishamifar, A., Bahonar, A. and Yousefi, P., 2011. Anesthetic effect of tricaine methanesulfonate, clove oil and electroanesthesia on lysozyme activity of *Oncorhynchus mykiss. Iranian Journal of Fisheries Sciences*, 10(3), 393-402.

- **Gilderhus, P.A. and Marking, L.L., 1987.** Comparative efficacy of 16 anesthetic chemicals on rainbow trout. *North American Journal of Fish Management*, 7, 288-292.
- Hedayati, A. and Hassan Nataj Niazie, E., 2015. Haematological changes of silver carp in response to Diazinon pesticide. *Journal of Environmental Health Science and Engineering*. 13(52), 2-5
- Hedayati, A., Hoseini, S.M. and Hoseinifar, S.H., 2015. Response of plasma cooper, ceruloplasmin, iron and ions in carp to waterborne copper ion and nanoparticle exposure. *Comparative physiology and biochemistry* (part C), 179, 87-93.
- King, W.V.. Hooper, В. Hillsgrove, S., 2005. The use of oil. metomidate. tricainemethanesulphonate and 2phenoxyethanol for inducing anaesthesia and their effect on the cortisol stress response in black sea bass (Centropristis striata L.). Aquaculture Research, 36, 1442-1449.
- McCarter, N., 1992. Sedation of grass carp and silver carp with 2-phenoxyethanol during spawning. *Progressive Fish Culturist*, 54, 263-265.
- Molinero, A. and Gonzalez, J., 1995.

  Comparative effects of MS 222 and 2-phenoxyethanol on gilthead sea bream (*Sparus aurata* L.) during confinement. *Comparative Biochemistry and Physiology* (Part A), 3, 405-414

- Ortuno, J., Esteban, M.A. and Meseguer, J., **2002.** Effects of phenoxyethanol on the innate immune system of gilthead sea bream (Sparus aurata L.) exposed to crowding stress. Veterinary Immunopatholgy, 89, 29-36
- Ross, L.G. and Ross, B., 2008.

  Anesthetic and sedative techniques for aquatic animals. Blackwell Publishing, Oxford, 222P.
- Shaluei, F., Hedayati, A., Jahanbakhshi, A. and Baghfalaki, M., 2012. Physiological responses of great sturgeon to different concentrations of 2-phenoxyethanol as an anesthetic. Fish Physiology and Biochemistry, 38, 1627–1634.
- Summer, F.R.C. and Smith, L.S, 1990. Anesthesia, Surgery and Related Techniques. In: Methods for Fish Biology, Schreck, CB and PB Moyle (Eds.). American Fisheries Society, Bethesda, MD. 311P.
- **Tort, L., Puigcerver, M. and Crespo, S., 2002.** Cortisol and haaematological response in sea bream and trout subjected to the anesthetics clove oil and 2-phenoxyethanol. *Aquaculture Research*, 33, 907-910.
- Veliekl, J. and Svobodovan, Z., 2004. Anesthesia of rainbow trout (*Oncorhynchus mykiss*) with 2-phenoxyethanol: acute toxicity and biochemical blood profile. *Acta Vetrinaria Brno*, 73, 379-384.
- Velisek, J., Svobodova, Z. and Piackova, V., 2007. Effects of 2-phenoxyethanol anesthesia on haaematological profile on common

- carp (*Cyprinus carpio*) and rainbow trout (*Oncorhynchus mykiss*). *Acta Vetrinaria Brno*, 76, 487-492.
- Weber, R.A., Pérez-Maceira, J.J. and Peleteiro, J.B., 2011. Effects of acute exposure to 2-phenoxyethanol, clove oil, MS-222, and metomidate on primary and secondary stress responses in Senegalese sole (Soleasene galensis Kaup 1858). Aquaculture, 321, 108-112.