

## Hematite-biochemical and immune response of Caspian brown trout (*Salmo trutta caspius*, Kessler, 1877) juveniles fed different levels of spirulina (*Spirulina platensis*)

Meshkat Roohani A.<sup>1</sup>; Fallahi Kapoorchali M.<sup>1\*</sup>; Abedian Kenari A.<sup>2</sup>;  
Sayyad Borani M.<sup>1</sup>; Zorriezahra M.J.<sup>3</sup>

Received: July 2018

Accepted: August 2019

### Abstract

The effects of dietary *Spirulina platensis* on haemato-biochemical and immunity responses of Caspian brown trout (*Salmo trutta caspius*) juveniles was investigated. For this purpose, diets with five *S. platensis* inclusion levels (control, 2%FMR (13.2 g kg<sup>-1</sup> spirulina in diet), 4%FMR (26.4 g kg<sup>-1</sup> spirulina in diet), 6%FMR (39.6 g kg<sup>-1</sup> spirulina in diet), and 8%FMR (52.8 g kg<sup>-1</sup> spirulina in diet) were prepared. Six hundred juveniles with an average initial weight of 11±1.0 g were assigned to 15 experimental tanks. The experiment lasted for 10 weeks. At the end of the experiment, growth performance, haemato-biochemical parameters including white and red blood cell counts, neutrophils lymphocytes counts, hematocrit, hemoglobin, glucose, albumin, total protein, aspartate amino transferase (AST), alanine amino transferase (ALT), triglyceride, cholesterol, as well as immunity parameters including lysozyme, C3, C4, Immunoglobulin (IgM), ACH50 and respiratory burst activity were assessed. The results indicated that fish fed diets supplemented with 6%FMR and 8%FMR had a significantly higher weight gain (26.13 g and 25.88 g) and specific growth rate (1.74 %bw day<sup>-1</sup> and 1.71 %bw day<sup>-1</sup>) compared with control. Furthermore, 6%FMR and 8%FMR treatments had statistically higher protein efficiency (0.76 and 0.78), lipid efficiency (1.89 and 1.94) and statistically lower feed conversion ratio (2.91 and 2.84) compared to the other treatments respectively ( $p<0.05$ ). The physiological and immunological factors were improved when fish were fed a high level of *S. platensis* supplement. *S. platensis* inclusion also increased activity of Lysozyme C3, C4, IgM and ACH50 and respiratory burst activity and reduced AST and ALT formation. These results indicate that *S. platensis* supplement is promising for disease prevention in *S. trutta caspius* juveniles, at an optimum dietary level of 6% in diet.

**Keywords:** Spirulina, *Salmo trutta caspius*, Growth performances, Haemato-biochemical parameters, Immunity.

1-Inland Waters Aquaculture Research Center, Iranian Fisheries Science Research Institute, Agriculture Research, Education and Extension Organization (AREEO), Bandar Anzali, Gilan, Iran

2-Department of Aquaculture, Faculty of Natural Resources and Marine Sciences, Tarbiat Modares University, Noor, Mazandaran, Iran

3-Iranian Fisheries Science Research Institute, Agriculture Research, Education and Extension Organization (AREEO), Tehran, Iran

\*Corresponding author's Email: m\_fallahi2011@yahoo.com

## Introduction

Growth performance and disease control are the most important priorities in current aquaculture. Nowadays, use of antibiotics and chemotherapeutics has increased significantly, due to intensification of culture practices and followed by high prevalence of infectious diseases (Andrews *et al.*, 2011). Accumulation of antibiotic residues in fish tissues and environment causes human and animal health issues and led to an urgent need for alternative disease preventive substances and use of immune-stimulants in aquaculture (Andrews *et al.*, 2011; Krishnaveni *et al.*, 2013). Several types of stimulants with different mechanisms and functions such as bacterial products, complex carbohydrates, vaccines, immune enhancing drugs, nutritional factors, animal and plant extracts are considered, and their abilities to activate the innate immunity have been studied (Raa, 1996; Sakai, 1999; Shahbazi and Bolhassani, 2016). One supplement source, with ease of production, processing and distribution along with a wide range of macro- and micronutrients of health benefits is spirulina (Ravi *et al.*, 2010). Research has shown that this alga contains a wide variety of compounds such as phycobiliproteins, carotenoids, phycocyanin, polysaccharides, unsaturated fatty acids, superoxide dismutase, different vitamins, and other elements which by bring improvement in body coloration, growth and immunity of fish (Nakagawa *et al.*, 2007). Spirulina (*Spirulina platensis*), a blue-green filamentous, fresh water and

multi-cell microalgae is gaining more attention from medical scientists as a nutraceutical and source of potential pharmaceuticals. There are several new peer reviewed scientific studies about spirulina's ability to inhibit viral replication, strengthen and enhancing both the cellular and humoral arms of the immune system (Kozlenko *et al.*, 1998; Hirahashi *et al.*, 2002; Andrews *et al.*, 2011). It has been received the most research and public health attention due to its bioactive compounds including vitamins, essential amino acids, minerals, essential fatty acids (gamma linolenic acid), and antioxidant pigments such as phycocyanine (Belay *et al.*, 1996; Regunathan and Wesley, 2006; Ragap *et al.*, 2012); Ceballos *et al.*, 2006). Additionally, spirulina has high protein content (60–70% by dry weight) and is the richest natural source of vitamin B<sub>12</sub> (Estrada *et al.*, 2001). This alga contains a whole spectrum of natural mixed carotene and xanthophylls phytopigments (Belay *et al.*, 1996; Estrada *et al.*, 2001; Andrews *et al.*, 2011). It has been confirmed that the addition of small amounts of algae to fish feed can exert pronounced effects on growth, lipid metabolism, body composition and physiological response to stress and disease (Mustafa *et al.*, 1994; Mustafa and Nakagawa, 1995; Palmegiano *et al.*, 2008; Abdel-Tawwab and Ahmad, 2009; Teimouri *et al.*, 2013). Spirulina as an immune modulator not only stimulates the immune system, but also strengthens the body's ability to produce new blood cells (Andrews *et al.*, 2011). Important

parts of the immune system, such as bone marrow stem cells, macrophages cells (number of cells and phagocytosis), T cells (lymphocyte), NK (non-specific cytotoxic), spleen as well as thymus show tangible activities when treated with spirulina extract (Henrikson, 1998; Watanuki *et al.*, 2006; Tongsiri *et al.*, 2010; Yong-Chin *et al.*, 2010; Shahbazi and Bolhassani, 2016). Recently, spirulina has been speculated to be associated with modulation of the host immune system (Hironobu *et al.*, 2006).

Caspian brown trout (*Salmo trutta caspius*, Kessler 1877), is one of the world's nine subspecies of brown trout (Quillet *et al.*, 1992; Habibi *et al.*, 2013) and attains the greatest size and growth rate of all brown trout (Sedgwick, 1995; Rajabi *et al.*, 2016). This species is a critically endangered anadromous species distributed in southern region of the Caspian Sea. In 1999, this species was declared at risk according to IUCN conditions (Kiabi *et al.*, 1999; Coad, 2000; Kalbassi *et al.*, 2006). Artificial propagation and releasing fingerlings to the natural waters is an approach to prevent brown trout's extinction (Rajabi *et al.*, 2016). In addition, in the recent years, this species has attracted interest for aquaculture in cage and raceways in intensive culture systems (Kalbassi *et al.*, 2006). *S. trutta caspius*, like many other species is sensitive to stressors and pathogenic agents. Thus, strong defense mechanisms or immune system are needed against pathogens to improve the health of the fish. Despite the importance of Caspian salmon as an

endangered species, however, little is known about their nutritional requirements especially in the field of increasing immunity of species and types of immune-stimulants. Therefore, the main objective of this study was to evaluate the effects of *S. platensis* meal as feed additive on immune system, biochemistry and haematology in *S. trutta caspius* as a valuable species.

## Materials and methods

### Experimental diets

Five artificial diets were formulated using five levels of microalgae *S. platensis* (Sina microalgae Co., Qeshm, Iran) and in three replicates were examined in a randomized design. The experimental diets were formulated by partially replacing fishmeal in the basal diet (65.89% in control) with spirulina powder at inclusion levels of 0, 2%, 4%, 6% and 8%, respectively. Diet formulations were performed using Lindo software (Lindo copyright, release, 6.11998), and fishmeal and *S. platensis* powder were used as a protein source. Test diets provided, which were iso-nitrogenous (45-46% crude protein) and iso-caloric (20 MJ kg<sup>-1</sup>) were analyzed in this regard, only varied in terms of fish meal and *S. platensis* content. The proximate composition of experimental diets was measured according to the standard methods of Association of Official Analytical Chemists (AOAC, 1995) for moisture, protein, fat, and ash determination. Briefly, Moisture content was estimated by drying the samples to constant weight at 95 °C in a drying oven (GCA, model 18EM, Precision Scientific

Group, Chicago, IL, USA). N content was measured using the Labconco Micro Kjeldahl Apparatus (Labconco Corporation, Kansas, MO, USA) and crude protein was estimated by multiplying N content by 6.25.

Lipid content was determined by acetone extraction using the extraction Soxhlet apparatus (Lab-Line Instruments, Inc., Melrose Park, IL, USA) for 16 hours, and ash was determined by combusting dry samples in a muffle furnace (Thermolyne Corporation, Dubuque, IA, USA) at 550 °C for 6 hours. Carbohydrate (i.e., nitrogen-free extract plus fiber) was calculated by the difference:

$(100 - (\text{protein} + \text{fat} + \text{ash} + \text{moisture}))$

(Aksnes and Opstvedt, 1998).

Gross energy was calculated using the coefficient of 23.6, 39.5, and 17.2

(kJ g<sup>-1</sup>) for protein, fat, and carbohydrates, respectively (NRC, 1993). The experimental diet formulation and proximate composition are showed in Table1, respectively. Dietary feed ingredients were ground using a laboratory grinder (Philips HR7628, Finland) and then blended into a homogenous doughy matter by adding water and transferred to a meat grinder with a 2 mm-mesh (CGT Company, 2mec, Rome, Italy). After drying, diets were broken by hand to fit fish mouths and placed in a grinding machine for breakdown into small pieces. The resulting diets were then dried in food dryer and stored in plastic bags in a refrigerator at -2 °C until further use.

Table 1: Dietary formulation and proximate composition of experimental diets.

| Ingredient (g kg <sup>-1</sup> dry weight) | Experimental Diets |       |       |       |       |
|--|--------------------|-------|-------|-------|-------|
|  | 0%                 | 2%    | 4%    | 6%    | 8%    |
| Fish meal <sup>1</sup>                     | 658.9              | 645.7 | 632.4 | 619.4 | 606.2 |
| Spirulina <sup>2</sup>                     | 0                  | 13.2  | 26.5  | 39.5  | 52.7  |
| Wheat flour                                | 115.9              | 117.8 | 119.4 | 120.9 | 122.4 |
| Fish oil <sup>3</sup>                      | 45                 | 45    | 45    | 45    | 45    |
| Soybean oil <sup>4</sup>                   | 50.3               | 51.3  | 52.3  | 53.3  | 54.5  |
| Lethitin <sup>5</sup>                      | 5                  | 5     | 5     | 5     | 5     |
| Filler(sand)                               | 60                 | 57.1  | 54.6  | 52.1  | 49    |
| Mineral mixture <sup>6</sup>               | 25                 | 25    | 25    | 25    | 25    |
| Vitamin mixture <sup>7</sup>               | 15                 | 15    | 15    | 15    | 15    |
| Anti-oxidant <sup>8</sup>                  | 0.2                | 0.2   | 0.2   | 0.2   | 0.2   |
| Choline Chloride <sup>9</sup>              | 1                  | 1     | 1     | 1     | 1     |
| Mono calcium phosphate <sup>10</sup>       | 5                  | 5     | 5     | 5     | 5     |
| Anti fungal <sup>11</sup>                  | 2.5                | 2.5   | 2.5   | 2.5   | 2.5   |
| Lysine <sup>12</sup>                       | 5                  | 5     | 5     | 5     | 5     |
| Methionine <sup>13</sup>                   | 5                  | 5     | 5     | 5     | 5     |
| Vitamin C <sup>14</sup>                    | 1                  | 1     | 1     | 1     | 1     |
| <b>Chemical analysis (%)</b>               |                    |       |       |       |       |
| protein                                    | 45.1               | 45.2  | 44.89 | 45.07 | 45.12 |
| lipid                                      | 17.96              | 17.75 | 18.1  | 18.2  | 18.1  |
| moisture                                   | 7.9                | 8.02  | 8.00  | 7.9   | 8.1   |
| ash  | 8.9                | 8.8   | 8.7   | 8.9   | 9.1   |
| carbohydrate                               | 12.14              | 11.93 | 12.21 | 11.73 | 11.58 |
| Gross Energy(MJ kg <sup>-1</sup> )         | 19.83              | 19.73 | 19.84 | 19.84 | 19.8  |

<sup>1</sup> Herring meal, produced by Mirood, Mazandaran province, Iran.

<sup>2</sup> Produced in Sina micro algae, Qeshm. Hormozgan province, Iran.

<sup>3</sup> Produced by Mirood, Mazandaran province, Iran.

<sup>4,5</sup> Prepared of Mazandaran aquatic food factory, Sari, Mazandaran province, Iran.

<sup>6</sup> Mineral premix consisted of (mg kg<sup>-1</sup> premix): Man=2600mg, Cu=600mg, Fe=6000mg, Zn=4600mg, Se=100 mg, I=100 mg, Co=50 mg and carrier up to 1 kg.

<sup>7</sup> Vitamin premix consisted of (mg kg<sup>-1</sup> premix) : A=1200000 IU, D3=400000 IU, E= 3000IU, K3=1200mg, C= 5400mg, B1=200mg, H2=200 mg, B2=3360mg, B12=4mg, B9=600mg, B5=9000mg, B7=7200mg

<sup>8</sup> Preparation of the Mazandaran aquatic food factory and its combination including BHT, Etoxyquin, Synergist Propylgallate.

<sup>9</sup> Choline Chloride, 60%, Prepared of livestock and poultry pharmaceutical factory Aras Bazar.

<sup>10, 11, 12, 13</sup> Prepared of pharmaceutical factory Aras Taban, Amol, Mazandaran province, Iran.

<sup>14</sup> Prepared of pharmaceutical factory Aras Bazar, Amol, Mazandaran province, Iran.

<sup>15</sup> Aquatic vitamin C, production of pharmaceutical factory Aras Bazar, Amol, Mazandaran province, Iran.

### *Fish rearing*

Fish juveniles (n= 600; mean individual initial weight  $11 \pm 1.0$  g) were obtained from Cold Water Breeding and Restocking Hatchery, Shahid Bamonar Center in Kelardasht, Mazandaran, Iran, and then were transferred to NIAC multi-purpose co-operative farm fish in Baghbankola village, Amol, Mazandaran, Iran. Fish were allowed to acclimatize for two weeks in indoor fiberglass tanks prior to the experiment and during this period were fed a commercial diet twice daily. After the acclimation period, fish were randomly divided into five equal groups, each comprising three replicates (40 fish per replicate) in 15 separate 400 L fiberglass tanks. The fish were hand-fed three times daily (8:00, 12:00 and 16:00 h) to apparent satiation by visual observation for ten weeks (Sotoudeh *et al.*, 2015). Water temperature, dissolved oxygen and pH were monitored daily (during the experiment, temperature, dissolved oxygen and pH were  $15 \pm 2$  °C,  $8.3 \pm 1$  mg L<sup>-1</sup> and  $7.6 \pm 0.3$  respectively). Uneaten feed and fecal matter were siphoned off every day. Due to the sensitivity of these juveniles, and to prevent mortality, biometry was performed once at the beginning and once at the end of this period.

### *Growth parameters*

Growth performance was determined and feed utilization was calculated as following (Goytortua-Bores *et al.*, 2006):

Weight gain (WG %)= $100 \times (\text{final body weight} - \text{initial body weight}) / \text{initial body weight}$ .

Specific growth rate (SGR (%))= $100 (\ln W_2 - \ln W_1) / T$ ; where W1 and W2 are the initial and final weight, respectively, and T is the number of days in the feeding period;

Feed conversion ratio (FCR) = feed consumed (g dry weight)/weight gain (g);

Condition Factor (CF) =  $\text{Weight} / \text{Total length}^3 \times 100$ ;

Survival rate (%) =  $(\text{Number of fish at the beginning} / \text{Number of fish at the end}) \times 100$ .

### *Sampling*

At the end of our experiments, to evaluate haematological parameters, fish were fasted for 24 hours immediately prior to blood sampling and then six fish from each tank were sampled randomly. To prevent sampling stress, the fish were anesthetized with a stock solution (50 ppm) of clove oil and blood samples were collected quickly (approximately one min/fish) by puncturing the veins in the caudal peduncle with a sterile 5 ML syringe (Esmaeili *et al.*, 2017). The collected blood sample was divided into two portions. One portion was transferred into Eppendorf tubes containing heparin anti-coagulant (500 U L<sup>-1</sup>) for haematological tests and the second portion of blood sample was also transferred into Eppendorf tubes, left to clot at 4 °C and centrifuged at 5000 rpm for 5 min at room temperature for biochemical and Immunological tests.

*Haematology tests*

Red blood cells (RBC) and white blood cell (WBC) counts were determined using Neubauer haemocytometer following the methods of Blaxhall and Daisley (1973). Blood was diluted to 1:200 with Race solution. Five center cells of middle square in Neubauer chamber were used for counting RBC and multiplied by 10,000 (Esmaeili *et al.*, 2017). For counting WBC, four marginal squares in Neubauer chamber were used after blood dilution (1:50) with Race solution and the results were multiplied by 50. Then, blood samples were fixed in methanol and stained using Wright–Giemsa stain for determination of the differential WBC (lymphocyte and neutrophil portions of total WBC) count (Houston, 1990). At least 200 WBCs were counted for differential WBC determinations. Thin slices were used for counting WBC counting was done according to their shape and reported as percentages. To measure the Hemoglobin (Hb) levels were obtained by the Cyanmethemoglobin method (Houston, 1990). In this method, the RBC are haemolized and hemoglobin is converted to Cyanmethemoglobin. Twenty microliters of blood samples with 50- $\mu$ l Drabkin's reagent were mixed and placed in dark place for 10 min to form Cyanmethemoglobin. Then, absorbance was read at 540 nm in a UV with spectrophotometer (Jenway 6800, UK) and finally, hemoglobin value was calculated. Haematocrit (Ht) was measured using the standard micro-haematocrit method (Subhadra *et al.*, 2006) and reported as percentages.

Microhematocrit capillary tubes were centrifuged at 2500 rpm for 5 min, and then hematocrit value was calculated using graded plate. To calculate others haematological indices, MCV (mean corpuscular volume), MCH (mean corpuscular hemoglobin), MCHC (mean corpuscular hemoglobin concentration), the relevant formula (Bain *et al.*, 2011) was used as follows:

$$\text{MCV (fl)} = \text{hematocrit/RBC (million per mm}^3\text{)} \times 10$$

$$\text{MCH (pg)} = \text{hemoglobin/RBC (million per mm}^3\text{)} \times 10$$

$$\text{MCHC} = \text{hematocrit/hemoglobin} \times 100$$
*Blood biochemical parameters*

Total protein (modi-biuret method, Tietz, 1986) and Albumin (romocresol green binding method, Doumas and Peters, 1997) assays were performed with Zist Chemistry kits (Zist chemistry Company, Tehran, Iran). Total protein content (at wavelength of 560-520 nm) and Albumin (at wavelength of 630 nm) were determined using an auto analyzer system (Thecnicon, RA 1000, New York, NY, USA). Globulin levels of the samples were calculated from the difference in albumin from total protein (Kumar *et al.*, 2005). Total cholesterol, triglyceride and glucose measurements were performed using Pars Azmoon commercial kits (Pars Azmoon Company, Tehran, Iran) according to the company's protocol and using an auto-analyzer (Thecnicon, RA 1000, New York, NY, USA). Briefly, in this method, 10  $\mu$ l of plasma sample had mixed with 1000  $\mu$ l of reagent and incubated for 20 min in room temperature. Then, the absorbance of

the sample against the blank (10  $\mu$ l of distilled water mixed with 1000  $\mu$ l of reagents) had measured at a wavelength of 546 nm. Similar approach had done for other parameters, but with appropriate sample amount and wavelength according to each parameter's protocol (Esmaeili *et al.*, 2017). The Kinetic method was used to measure liver enzymes, aspartate aminotransferase (AST) and alanine aminotransferase (ALT), using Pars Azmoon commercial clinical investigation kits (Parsazmon Company, Tehran, Iran) and were analyzed with an auto analyzer (Technicon RA-1000; Technicon Instruments, New York, NY, USA).

#### *Immunological parameters*

The lysozyme activity in serum was determined turbidometrically using the method described by Ellis (1999). In this method, *Micrococcus luteus* bacteria was used (Sigma, M0508) with phosphate buffer (30 ml, 0.1 M, pH=7) as substrate. For this purpose, activated bacteria were used and cultivated at a concentration in 0.6–0.8 nm adjusted with spectrophotometer (Jenway 6800, UK). Then, 1990  $\mu$ l of substrate and 10  $\mu$ l of the sample were added to cuvette. The decrease of absorption at 450 nm each min for a period of 6 min had measured through spectrophotometer (Tukmechi *et al.*, 2011) with chicken egg white (Sigma, A5503) standard. Turbidometric method and Pars Azmoon commercial clinical investigation kits (Pars Azmoon Company, Tehran, Iran) were used to measure C3 and C4 serum complement

system (Tang *et al.*, 2008). The serum C3 and C4 reacted with the antibodies contained in the kit, and then OD had measured at a wavelength of 340 nm using an auto-analyzer (Thecnicon, RA 1000, New York, NY, USA), compared to the standard in kits and finally, the C3 and C4 amount was calculated based on  $\text{mg L}^{-1}$ . ACH50 (the activity of the complement system) was determined according to the indirect ELISA method using a commercial kit (Wielsa, comple 300 total complement functional screen kit, Sweden). The volume of the serum complement producing 50% hemolysis (ACH50) was determined, and the number of ACH50 units in ml was calculated for each experimental fish.

Total immunoglobulin levels were determined according to the method described by Siwicki and Anderson (1993). In this method, analysis of total Ig level in plasma is based on the biuret colorimetric method. First the immunoglobulins were separated from plasma by precipitation with polyethylenic glycol 10,000 (Sigma) and remaining supernatant was read. That number was subtracted from the total protein to give total immunoglobulin. Protein readings from supernatant gave the amount of protein taken out by absorption to polyethylene glycol. To calculate total immunoglobulin, subtracted these readings from total protein on individual samples and compared with standards for calculation of protein. The production of oxygen radicals by leukocytes was determined by the reduction of Nitro Blue Tetrazolium

(NBT, Sigma-Aldrich Chemical, St. Louis, MO, USA) according to Rook *et al.* (1985). Absorbance was converted to NBT units based on a standard curve of NBT diformazan per milliliter of blood.

### Statistical analysis

All the data have been analyzed using the SPSS (version 17, SPSS, Richmond, VA, USA) statistical package as described by Dytham (1999). Mean values of all the parameters were subjected to one-way analysis of variance (ANOVA) to study the treatment effect and comparison of any mean values had done by Duncan's Multiple Range Test (DMRT). Comparisons were made at the five percent probability level ( $p < 0.05$ ).

### Result

The growth performance of Caspian brown trout juvenile fed diets containing different levels of spirulina are presented in Table 2. Overall, the optimum growth performance has been

obtained at 6%FMR and 8%FMR treatments, whereas the fish fed control diet had the lowest growth performances. According to the results, fish fed diet supplemented with 6%FMR and 8%FMR had a significantly higher weight gain (26.13 g and 25.88 g) and specific growth rate ( $1.74 \%bw day^{-1}$  and  $1.71 \%bw day^{-1}$ ) compared with those fed the control diet (18.18 g and  $1.37 \%bw day^{-1}$ ). Furthermore, 6 %FMR and 8 %FMR treatments had statistically higher protein efficiency (0.76 and 0.78), lipid efficiency (1.89 and 1.94) and statistically lower feed conversion ratio (2.91 and 2.84) compared with other treatments ( $p < 0.05$ ). Moreover, there was no significant difference in the condition factor among experimental treatments. In addition, fish fed control, 2 %FMR and 4 %FMR diets exhibited equivalent FCR, protein efficiency and lipid efficiency. Survival rate in all treatments was 100%.

**Table 2: Mean growth performance of *Salmo trutta caspius* fed different levels of *Spirulina platensis*.**

| Growth performance   | Experimental Diets       |                           |                            |                           |                            |
|----------------------|--------------------------|---------------------------|----------------------------|---------------------------|----------------------------|
|                      | 0% control               | 2%                        | 4%                         | 6%                        | 8%                         |
| W <sub>1</sub> (g)   | 11.18±0.23               | 11.91±1.23                | 11.16±0.56                 | 10.91±1.61                | 11.19±0.33                 |
| WG (%)               | 62.73±12.3 <sup>a*</sup> | 93.93±11.31 <sup>ab</sup> | 101.41±14.29 <sup>ab</sup> | 163.35±67.22 <sup>c</sup> | 131.96±35.27 <sup>bc</sup> |
| SGR (% / day)        | 0.81±0.13 <sup>a</sup>   | 1.1±0.1 <sup>ab</sup>     | 1.16±0.12 <sup>ab</sup>    | 1.58±0.16 <sup>c</sup>    | 1.39±0.26 <sup>bc</sup>    |
| CF                   | 1.33±0.05 <sup>a</sup>   | 1.28±0.03 <sup>a</sup>    | 1.38±0.06 <sup>a</sup>     | 1.27±0.04 <sup>a</sup>    | 1.31±0.14 <sup>a</sup>     |
| FCR                  | 3.93±0.74 <sup>b</sup>   | 3.95±0.49 <sup>b</sup>    | 2.73±0.06 <sup>a</sup>     | 2.11±0.49 <sup>a</sup>    | 2.04±0.56 <sup>a</sup>     |
| Survival rate (100%) | 100                      | 100                       | 100                        | 100                       | 100                        |

W<sub>1</sub>: Primary weight

WG%= Weight gain

FCR: Feed conservation ratio

SGR: Specific growth ratio

Conditional Factor(CF)

\*Values are the least square means±standard errors of the means of triplicate samples. *P* values determined with one-way ANOVA are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different ( $p > 0.05$ ). Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.



The mean haematology parameters of *S. trutta caspius* juveniles fed with diets containing different levels of *S. platensis* are shown in Table 3. Based on obtained results, spirulina had a significant impact on haematological parameters (RBC, WBC, Hct, Hb, lymphocytes, neutrophils) in *S. trutta caspius* juveniles. RBC increased with increasing levels of dietary *S. platensis* to 6%, but in 8%, the rate decreased ( $p<0.05$ ). WBC, Hct, Hb, differential white blood cell percent, lymphocyte and neutrophil increased with

increasing levels of dietary *S. platensis*. So, the lowest and the highest levels for these parameters (with exception of neutrophils) were observed in the control and 8% treatments, respectively. Other haematological indices such as average volume of red blood cells (MCV), mean hemoglobin in red blood cells (MCH), average concentration of hemoglobin in a red blood cells (MCHC) also have increased with increasing levels of dietary *S. platensis* ( $p<0.05$ ).

**Table 3: Haematological parameters of *Salmo trutta caspius* fed with different levels of *Spirulina platensis*.**

| Haematological parameters               | Experimental Diets             |                                  |                                 |                                 |                                 |
|---|--------------------------------|----------------------------------|---------------------------------|---------------------------------|---------------------------------|
|   | 0% control                     | 2%                               | 4%                              | 6%                              | 8%                              |
| RBC ( $\times 10^6 \text{ n mm}^{-3}$ ) | 0.77 $\pm$ 0.17 <sup>a</sup>   | 1.22 $\pm$ 0.05 <sup>c</sup>     | 1.25 $\pm$ 0.16 <sup>cd</sup>   | 1.27 $\pm$ 0.09 <sup>d</sup>    | 1.12 $\pm$ 0.15 <sup>b</sup>    |
| WBC (cell $\text{ml}^{-1}$ )            | 9316 $\pm$ 3521.6 <sup>a</sup> | 10100 $\pm$ 1649.24 <sup>b</sup> | 11800 $\pm$ 2415.7 <sup>c</sup> | 13500 $\pm$ 2485.4 <sup>d</sup> | 15400 $\pm$ 2256.5 <sup>e</sup> |
| Hct (%)                                 | 24.66 $\pm$ 1.98 <sup>a</sup>  | 35.83 $\pm$ 1.94 <sup>b</sup>    | 37.17 $\pm$ 1.83 <sup>c</sup>   | 37.66 $\pm$ 0.81 <sup>c</sup>   | 37.83 $\pm$ 1.26 <sup>c</sup>   |
| Hb (g $\text{dl}^{-1}$ )                | 4.32 $\pm$ 0.61 <sup>a</sup>   | 7.95 $\pm$ 0.36 <sup>bc</sup>    | 7.59 $\pm$ 0.75 <sup>b</sup>    | 8.16 $\pm$ 0.31 <sup>c</sup>    | 8.9 $\pm$ 0.51 <sup>d</sup>     |
| LYM (% of WBC)                          | 85.5 $\pm$ 3.39 <sup>a</sup>   | 89.5 $\pm$ 1.76 <sup>b</sup>     | 92.66 $\pm$ 7.28 <sup>c</sup>   | 98.16 $\pm$ 1.16 <sup>d</sup>   | 98.66 $\pm$ 0.81 <sup>d</sup>   |
| Neut (% of WBC)                         | 1.16 $\pm$ 0.75 <sup>a</sup>   | 1.16 $\pm$ 0.75 <sup>a</sup>     | 1.33 $\pm$ 0.81 <sup>a</sup>    | 1.67 $\pm$ 0.81 <sup>c</sup>    | 1.5 $\pm$ 0.83 <sup>c</sup>     |
| MCV (fl)                                | 280.81 $\pm$ 6.96 <sup>a</sup> | 298.76 $\pm$ 25.03 <sup>ab</sup> | 300.98 $\pm$ 26.45 <sup>b</sup> | 332.15 $\pm$ 17.8 <sup>c</sup>  | 337.21 $\pm$ 12.2 <sup>c</sup>  |
| MCH (pg)                                | 52.37 $\pm$ 1.3 <sup>a</sup>   | 56.1 $\pm$ 1.72 <sup>a</sup>     | 61.23 $\pm$ 4.17 <sup>a</sup>   | 64.57 $\pm$ 2.3 <sup>ab</sup>   | 80.13 $\pm$ 6.7 <sup>b</sup>    |
| MCHC (%)                                | 17.63 $\pm$ 1.77 <sup>a</sup>  | 19.98 $\pm$ 0.71 <sup>b</sup>    | 20.4 $\pm$ 1.18 <sup>b</sup>    | 21.68 $\pm$ 1.04 <sup>b</sup>   | 23.78 $\pm$ 2.13 <sup>c</sup>   |

\*Values are least square means $\pm$ standard errors of the means of triplicate samples.  $p$  values determined with two-way ANOVA tests are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different ( $p>0.05$ ). Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

The mean blood biochemical parameters of *S. trutta caspius* juveniles fed with diets containing different levels of *S. platensis* are summarized in Table 4&5. Based on obtained results, with increasing levels of dietary *S. platensis* up to 6%, ALT and AST enzymes decreased ( $p<0.05$ ). Total

protein and globulin decreased and then increased in 8% treatment ( $p<0.05$ ). Other blood biochemical parameters such as glucose, cholesterol and triglycerides have significantly decreased with increasing levels of dietary *S. platensis* ( $p<0.05$ ).

**Table 4: Liver enzymes of *Salmo trutta caspius* fed different levels of *Spirulina platensis*.**

| Liver enzymes           | Experimental Diets     |                          |                        |                         |                         |
|-------------------------|------------------------|--------------------------|------------------------|-------------------------|-------------------------|
|                         | 0% control             | 2%                       | 4%                     | 6%                      | 8%                      |
| ALT(U L <sup>-1</sup> ) | 323±45.2 <sup>d*</sup> | 305.4±27.1 <sup>bc</sup> | 290±20.1 <sup>b</sup>  | 214.2±28.7 <sup>a</sup> | 212.3±16.2 <sup>a</sup> |
| AST(U L <sup>-1</sup> ) | 25.4±4.5 <sup>d</sup>  | 23.7±3.8 <sup>cd</sup>   | 22.9±6.6 <sup>bc</sup> | 21.3±4.3 <sup>b</sup>   | 12.2±2.5 <sup>a</sup>   |

\*Values are least square means ± standard errors of the means of triplicate samples. *p* values determined with two-way ANOVA tests are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different (*p*>0.05). Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

**Table 5: Blood biochemical parameters of *Salmo trutta caspius* fed different levels of *Spirulina platensis*.**

| Biochemical factors                 | Experimental Diets      |                         |                         |                         |                         |
|-------------------------------------|-------------------------|-------------------------|-------------------------|-------------------------|-------------------------|
|                                     | 0% control              | 2%                      | 4%                      | 6%                      | 8%                      |
| Total protein (g dl <sup>-1</sup> ) | 2.6±0.4 <sup>a*</sup>   | 2.6±0.4 <sup>a</sup>    | 2.8±0.7 <sup>b</sup>    | 3.1±0.3 <sup>c</sup>    | 4.3±0.5 <sup>d</sup>    |
| Albumin (g dl <sup>-1</sup> )       | 1.8±0.3 <sup>a</sup>    | 1.8±0.3 <sup>a</sup>    | 1.7±0.1 <sup>a</sup>    | 2±0.3 <sup>a</sup>      | 2±0.3 <sup>a</sup>      |
| Globulin (g dl <sup>-1</sup> )      | 0.8±0.1 <sup>a</sup>    | 0.8±0.1 <sup>a</sup>    | 1.1±0.2 <sup>b</sup>    | 1.1±0.3 <sup>b</sup>    | 2.3±0.3 <sup>c</sup>    |
| Glucose (mg dl <sup>-1</sup> )      | 172.5±16.3 <sup>e</sup> | 176±7.8 <sup>d</sup>    | 121.5±12.3 <sup>b</sup> | 133.1±15.9 <sup>c</sup> | 94.5±8.2 <sup>a</sup>   |
| Cholesterol (mg dl <sup>-1</sup> )  | 438.6±45.1 <sup>d</sup> | 412.5±47.3 <sup>c</sup> | 401.4±26.2 <sup>b</sup> | 346.8±42.2 <sup>a</sup> | 394.8±21.6 <sup>b</sup> |
| Triglyceride (mg dl <sup>-1</sup> ) | 369.9±5.3 <sup>e</sup>  | 356.4±40.5 <sup>d</sup> | 326.9±45.8 <sup>c</sup> | 322.7±58.9 <sup>b</sup> | 253.8±61.4 <sup>a</sup> |

\*Values are least square means ± standard errors of the means of triplicate samples. *p* values determined with two-way ANOVA tests are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different (*p*>0.05). Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

The mean immunity parameters of *S. trutta caspius* juveniles fed with diets containing different levels of *S. platensis* are shown in Table 6 and 7. Based on obtained results, lysozyme enzyme activity increased with increasing dietary *S. platensis* levels up to 6% and then decreased in 8%

treatment. Complement C3, C4, ACH50, free radical oxygen and immunoglobulin were significantly affected by dietary *S. platensis* levels. The lowest and the highest values for these parameters have been observed in the control and 8% treatments.

**Table 6: Lysozyme activity of *Salmo trutta caspius* fed different levels of *Spirulina platensis*.**

| Lysozyme activity                    | Experimental Diets      |                        |                        |                        |                        |
|--------------------------------------|-------------------------|------------------------|------------------------|------------------------|------------------------|
|                                      | 0% control              | 2%                     | 4%                     | 6%                     | 8%                     |
| After 15 sec (μg ml <sup>-1</sup> )  | 1.44±0.82 <sup>a*</sup> | 4.3±0.75 <sup>b</sup>  | 4.95±0.32 <sup>c</sup> | 7.33±0.22 <sup>d</sup> | 5.87±0.93 <sup>e</sup> |
| After 180 sec (μg ml <sup>-1</sup> ) | 6.22±0.35 <sup>a</sup>  | 6.32±0.38 <sup>a</sup> | 6.71±0.28 <sup>b</sup> | 8.59±1.06 <sup>c</sup> | 7.95±0.44 <sup>d</sup> |

\*Values are least square means ± standard errors of the means of triplicate samples. *p* values determined with two-way ANOVA tests are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different (*p*>0.05). Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

**Table 7: Immunological parameters of *Salmo trutta caspius* fed different levels of *Spirulina platensis*.**

| Immunological factor                  | Experimental Diets        |                          |                           |                           |                           |
|---------------------------------------|---------------------------|--------------------------|---------------------------|---------------------------|---------------------------|
|                                       | 0% control                | 2%                       | 4%                        | 6%                        | 8%                        |
| C <sub>3</sub> (mg ml <sup>-1</sup> ) | 21.1±3.3 <sup>a*</sup>    | 21.1±3.3 <sup>ab</sup>   | 23.4±2.5 <sup>b</sup>     | 25.7±3.2 <sup>c</sup>     | 35.7±6.9 <sup>d</sup>     |
| C <sub>4</sub> (mg ml <sup>-1</sup> ) | 13.9±3.6 <sup>a</sup>     | 14.7±2.7 <sup>a</sup>    | 15±2.4 <sup>a</sup>       | 19.2±1.8 <sup>b</sup>     | 20.8±1.5 <sup>b</sup>     |
| ACH50(unit ml <sup>-1</sup> )         | 220.1±3.87 <sup>a</sup>   | 225.16±4.9 <sup>b</sup>  | 228.23±7.1 <sup>c</sup>   | 234.33±8.73 <sup>d</sup>  | 247±4.77 <sup>e</sup>     |
| Igm(mg dl <sup>-1</sup> )             | 183.4±25.4 <sup>a</sup>   | 183.7±21.7 <sup>a</sup>  | 194.6±19.2 <sup>b</sup>   | 321.5±13.6 <sup>c</sup>   | 334.5±25.8 <sup>d</sup>   |
| O(RLU s <sup>-1</sup> )               | 384.96±21.05 <sup>a</sup> | 401.5±19.38 <sup>b</sup> | 415.51±29.73 <sup>c</sup> | 429.28±23.31 <sup>d</sup> | 471.03±23.77 <sup>e</sup> |

\*Values are least square means±standard errors of the means of triplicate samples. *p* values determined with two-way ANOVA tests are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different (*p*>0.05). Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

## Discussion

The present results indicated that spirulina significantly improved the growth performance of juvenile Caspian brown trout and the fish meal could be replaced with this microalga until 8% without impairments. During the experimental period, no mortality and disease in fish fed spirulina were observed compared to control. In agreement with our research, many studies have demonstrated a positive impact of spirulina on growth performance (Mustafa *et al.*, 1994; James *et al.*, 2006; Palmegiano *et al.*, 2008; Ramakrishnan *et al.*, 2008; Tongsiri *et al.*, 2010; Ungsethaphand *et al.*, 2010; Promya and Chitmanat, 2011; Teimouri *et al.*, 2013; Khanzadeh *et al.*, 2016; Cao *et al.*, 2018; Gogoi *et al.*, 2018). However, contradictory results have been reported with no beneficial effects of dietary spirulina on growth of common carp *Cyprinus carpio* (Nandeeshia *et al.*, 1998), Catla *Catla catla* (Nandeeshia *et al.*, 2001) and Nile tilapia *Oreochromis niloticus* (Lu and Takeuchi, 2002). It is likely that for carnivorous fish, this microalga has had better growth performance and better fish muscle quality and our study is in agreement with this hypothesis. Growth

improvement to dietary spirulina may be due to the improved feed intake and nutrient digestibility. Some researchers have been reported that spirulina improved intestinal microbial balance, leading to better growth by improving food absorption, digestive enzymes activities and fats transport system (James *et al.*, 2006; Teimouri *et al.*, 2013). In addition, the bio-compounds in spirulina, delayed absorption of dietary nutrients and improved carbohydrate and protein utilization in fish. Moreover, there are several nutrients especially vitamins, minerals, essential amino acids, fatty acids in spirulina, that may be beneficial to fish growth promotion and might activate metabolism and act as growth stimulants (Mustafa *et al.*, 1994). Generally, in the current study growth performance in the 6%FMR and 8%FMR treatments were higher than the other dietary groups. Differently, similar spirulina content to the 8%FMR (52.8 g kg<sup>-1</sup>) has been reported to decrease growth in rainbow trout (*Oncorhynchus mykiss*) (Teimouri *et al.*, 2013) and Nile tilapia (Olvera-Novoa *et al.*, 1998). There are no reports about anti-nutritional factors in spirulina (Vonshak *et al.*, 2014).

However, the slightly lower growth in 8%FMR compared to 6%FMR is due to the lower mineral content such as phosphorous in spirulina, compared to fish meal (Olvera-Novoa *et al.*, 1998). Perhaps the fish meal content in 6%FMR (620 g kg<sup>-1</sup> in diet) is optimum for Caspian brown trout growth and higher or lower contents is not enough to reach the highest growth.

There was a significant difference in haematology parameters among all tested treatments. Our results revealed that *S. platensis* has positive impact on the health status of *S. trutta caspius*. Previous studies such as those by Terry *et al.* (2000) on tilapia, Abdel Tawwab and Ahmad (2009) on tilapia, Yong-Chin *et al.* (2010) on vannamei shrimp (*Litopenaeus vannamei*), Andrews *et al.* (2011) on Nile tilapia, Promya and Chitmanat (2011) on African catfish, Ibrahim *et al.* (2012) on Nile tilapia, Ragap *et al.* (2012) on Nile tilapia, Krishnaveni *et al.* (2013) on catla, Zamini and Azimi (2015) on Carp koi; Salehi Farsani *et al.* (2014) on stellate sturgeon confirmed that *S. platensis* has positive effects on haematologic parameters. In addition, increasing haematology indices (e.g., RBC, Hb and Hct) of *S. trutta caspius* with increasing levels of *S. platensis* can result from a variety of reasons including iron compounds, vitamins (B12, A and E), and also the considerable antioxidant capacity contained in *S. platensis*. This high antioxidant capacity is due to the considerable amount of pigments, particularly Phycocyanobilin (e.g., Phycocyanin, Allophycocyanin and

Phycoerythrin). In fact, these pigments are responsible for the removal of peroxide radicals in the body and reduce the rate of haemolysis of RBC by oxidants as well as the ability to significantly increase iron absorption (Kop and Durmaz, 2008; Abdel Tawwab and Ahmad, 2009; Andrews *et al.*, 2011).

Reactions caused by WBC are mechanisms of cellular nonspecific or early defense in fish that occur in response to various conditions such as bacterial, viral, fungal, protozoan and parasitic infections, and it is the first indicator to determine the health of every living being (Andrews *et al.*, 2011). Increasing WBC and their differentiate percent with increasing levels of dietary *S. platensis* is due to stimulating the immune system via bioactive substances contained in the microalgae (Tort, 2003). Furthermore, phytocyanins contained in *S. platensis* activate types of WBC (e.g., macrophages and granulocytes) and are responsible for repairing damaged tissues by pathogens (Selmi *et al.*, 2011). This bioactive substance with induction of cytokine secretion (molecules involved in the immune system by activating macrophages and lymphocytes.) stimulates the production of new WBC marker molecules (Raa, 1996).

Based on these results, levels of liver enzymes (AST and ALT) decreased with increasing levels of *S. platensis*. Based on accepted hypothesis, the increase of these enzymes in animals (including fish), indicate damage to the liver cells (Jeney and Anderson, 1993).

Numerous studies have confirmed the effectiveness of *S. platensis* to reduce cholesterol and liver protection (Khan *et al.*, 2005; Pieretti and Meineri, 2011; Kim, 2013; Zeinab *et al.*, 2015). One of the most important approaches of antioxidants is liver cell protection. The antioxidant properties of *S. platensis* have attracted the attention of many researchers. One of the latest investigations Manoj *et al.* (1992) reported that alcoholic extract of *S. platensis* accelerates oxidation of fat significantly (65%) in comparison to the antioxidant chemicals such as alpha-tocopherol (35%), BHA or Butylated Hydroxy Anisole (45%), and beta-carotene (48%).

In this study, serum total protein, albumin and globulin increased with increasing levels of *S. platensis*. Several studies have been demonstrated the effects of *S. platensis* on blood biochemical parameters (Schaperclaus *et al.*, 1992; Duncan and Klesius, 1996; Abdel-Tawwab and Ahmad, 2009; Andrews *et al.*, 2011; Hernandez, 2005; Zeinab *et al.*, 2015). Blood serum protein show changes in health conditions affected by internal and external factors. Albumin is a transporter protein or public transporter of many organic and inorganic ligands, such as thyroxine, bilirubin, penicillin, cortisol, estrogen, free fatty acids, calcium, and magnesium during illness, malnutrition and stress, blood albumin is reduced (Misra *et al.*, 2006; Alexander *et al.*, 2011). Glucose, cholesterol and triglyceride levels fluctuate during protein catabolism and glycogenesis. Serum glucose levels are

often referred to as a non-specific marker of stress (Sheikhzadeh *et al.*, 2012). In this study, blood glucose, triglycerides and cholesterol decreased with increasing levels of *S. platensis*. Our results are in agreement with data reported by other researchers (Abdel-Tawwab and Ahmad, 2009; Andrews *et al.*, 2011; Hernandez, 2005; Salehi Farsani *et al.*, 2014).

Long-chain omega-3 fatty acids in the liver by reducing the secretion of lipoproteins (VLDL) can be reduced triacylglycerol levels (Harris *et al.*, 1984). *S. platensis* is low fat (5% fat) so, 10 gr. of *S. platensis* has only 36 gr. calorie and no cholesterol. *S. platensis* particularly in terms of alpha-linolenic acid (36% of the total PUFA) is very rich. The most essential fatty acids in *S. platensis* that in other food sources are scarce, is gamma linolenic acid. It has been grown under special lighting conditions in the cells, are also able to enrich and enhance it (Khan *et al.*, 2005). Gamma-linolenic acid is a precursor of prostaglandins (PGE) and has special effect on the control of blood cholesterol levels that leads to rapid cell growth. The role of PGE is blood pressure regulation, cholesterol synthesis control and cell survival (Harris *et al.*, 2002).

In this study, we evaluated the immunity status of *S. trutta caspius* fed with different levels of *S. platensis*, using lysozyme, complement C3 and C4, immunoglobulins, oxygen free radical and haemolytic activity of the complement factor. Based on our results, the activity of lysozyme after 15 and 180 seconds with increasing levels

of *S. platensis* significantly increased (up to 8%). Effect of *S. platensis* on lysozyme by various researchers have confirmed (Watanuki *et al.*, 2006; Abdel Tawwab and Ahmad, 2009; Andrews *et al.*, 2011; Promya and Chitmanat, 2011; Ragap *et al.*, 2012; Ibrahim *et al.*, 2012; Krishnaveni *et al.*, 2013; Kim, 2013; Shimaa, 2016; Salehi Farsani *et al.*, 2014). Lysozyme by WBCs (monocytes, macrophages, neutrophils) and using circulating the blood in different tissues of the body released and leads to non-specific immune organisms (Saurabh and Sahoo, 2008).

The complement system is a part of the immune system that enhances the ability of antibodies and phagocytic cells to clear microbes and damaged cells from an organism, promotes inflammation, and attacks the pathogen's plasma membrane (Claire *et al.*, 2002). Several factors have an impact on immunity, including divalent ions (especially magnesium and calcium), polysaccharides immunogenic, season, sexual maturity, prostaglandins, thromboxane and immunoglobulins (Tort *et al.*, 2003). The activity of the complement system and immunoglobulins is increased using microalgae (Ragap *et al.*, 2012; Krishnaveni *et al.*, 2013; Shimaa, 2016). C3 and C4 high levels are related to the health of the fish. High serum complement activity in fish fed *S. platensis* could be due to the presence of pigment carotenoids especially beta-carotene. In addition, lysozyme, antibodies and complement factors due to adhesion and

colonization with pathogenic microorganisms, prevent infection and pathogenesis. Free radicals of oxygen ( $O_2$ ,  $H_2O$  and  $O^{2-}$ ) together with nitric oxides, lysozyme, cytokines and other messenger molecules, (prostaglandins, leukotrienes and thromboxanes) increased phagocytic and Pynocytic capacity of macrophage cells (Taoka *et al.*, 2006). Measuring any of these factors can be effective in determining the activation of macrophages. Several studies have demonstrated a positive impact of immune-stimulants such as probiotics and micro-algae on amount of oxygen free radicals and phagocytic activity (Dunkan and Klesius, 1996; Rengpipat *et al.*, 2000; Li and Gatlin, 2004; Panigrahi *et al.*, 2005; Taoka *et al.*, 2006; Watanuki *et al.*, 2006; Abdel-Tawwab and Ahmad, 2009; Andrews *et al.*, 2011; Ragap *et al.*, 2012; Kim, 2013; Ibrahim *et al.*, 2012). Oxidative stress is the accumulation of ROS (Reactive Oxygen Species, ROS). It may be an important factor in the process of developing a variety of diseases. The immune system is vulnerable to oxidative damage. *S. platensis* also contains other pigments Phycocyanobilin including Phycocyanin, Allophycocyanin and Phycoerythrin that are mainly responsible for antioxidant activity and the removal of peroxide radicals in the body (Ragap *et al.*, 2012)

The results of different impacts levels of *S. platensis* on immune and haematology parameters in *S. trutta caspius* show that *S. platensis* may be beneficial in improve growth, immunity levels in Caspian Sea salmon and create

resistance to pathogens. These results indicate that *S. platensis* supplementation is promising for disease prevention in *S. trutta caspius* juveniles and can substitute for up to 8% of the fish meal in the diet, although the optimum performance is obtained with 6% substitution

## References

- Abdel-Tawwab, M. and Ahmad, M.H., 2009.** Live *Spirulina* (*Arthrospira platensis*) as a growth and immunity promoter for Nile tilapia, *Oreochromis niloticus* (L.), challenged with pathogenic *Aeromonas hydrophila*. *Aquaculture Research*, 40, 1037-1046.
- Aksnes, A. and Opstvedt, J., 1998.** Content of digestible energy in fish feed ingredients determined by the ingredient-substitution method. *Aquaculture*, 161, 45-53.
- Alexander, C., Sahu, N.P., Pal, A.K. and Akhtar, M.S., 2011.** Haematoimmunological and stress responses of *Labeo rohita* (Hamilton) fingerlings: Effect of rearing temperature and dietary gelatinized carbohydrate. *Journal of Animal physiology and Animal Nutrition*, 95, 653-663.
- Andrews, S.R., Sabu, N.P., Pal, A.K., Mukherjee, S.C. and Kumar, S., 2011.** Yeast extract, brewer's yeast and *Spirulina* in diets for *Labeo rohita* fingerlings affect haematoimmunological responses and survival following *Aeromonas hydrophila* challenge. *Research in Veterinary Science*, 91, 103-109.
- AOAC, 1995.** Official methods of analysis. 16th ed. Association of Official Analytical Chemists. Gaithersburg, MD, USA. 2200 P.
- Bain, B., Bates, I. and Laffan, M., 2011.** Dacie and Lewis Practical Haematology. 12th Edition, Published Edinburgh Churchill Livingstone.
- Belay, A., Kato, T. and Ota, Y., 1996.** *Spirulina* (*Arthrospira*): Potential application as an animal feed supplement. *Journal of Applied Phycology*, 8, 303-311.
- Blaxhall, P.C. and Daisley, K.W., 1973.** Routine haematological methods for use with fish blood. *Journal of Fish Biology*, 5(6), 771-781.
- Cao, S.P., Zou, T., Zhang, P.Y., Han, D., Jin, J.Y., Liu, H.K., Yang, Y.X., Zhu, X.M. and Xie, S.Q., 2018.** Effects of dietary fishmeal replacement with *Spirulina platensis* on the growth, feed utilization, digestion and physiological parameters in juvenile gibel carp (*Carassis auratus gibelio* var. CAS III). *Aquaculture Research*, 49, 1320-1328.
- Ceballos, J.B., Villareal, H., Garcia, T., Perez-Jar, L. and Alfonso, E., 2006.** Effect of *Spirulina platensis* meal as feed additive on growth, survival and development in *Litopenaeus schmitti* shrimp larvae. *Revista de Investigaciones Marinas*, 26(3), 235-241.
- Claire, H., Holland, M. and Lambris, J.D., 2002.** The complement system in teleosts. *Fish and Shellfish Immunology*, 12, 399-420.

- Coad, B.W., 2000.** Criteria for assessing the conservation status of taxa (as applied to Iranian freshwater fishes). *Biologia*, 55,539-557.
- Doumas, B.T. and Peters, J.R.T., 1997.** Serum and urine albumin: a progress report on their measurement and clinical significance. *Clinica Chimica Acta*, 258(1), 3-20.
- Dunkan, P.L. and Klesius, P.H., 1996.** Effects of feeding spirulina on specific and non-specific immune responses of channel catfish. *Journal of Aquaculture Animal Health*, 8, 308-313.
- Dytham, C., 1999.** Choosing and using statistics: A biologist's guide. *Journal of Ecology*, 87(4), 734-735.
- Ellis, A.E., 1999.** Lysozyme assays. Techniques in Fish Immunology. pp. 101-103.
- Esmaeili, M., Abedian Kenari, A. and Rombenso, A.N., 2017.** Immunohematological status under acute ammonia stress of juvenile rainbow trout (*Oncorhynchus mykiss* Walbaum, 1792) fed garlic (*Allium sativum*) powder-supplemented meat and bone meal-based feeds. *Comparative and Clinical Pathology*, DOI: 10.1007/s00580-017-2457-8.
- Estrada, J.P., Bescós, P.B. and Del Fresno, A.V. 2001.** Antioxidant activity of different fractions of *Spirulina platensis* protean extract. *Il farmaco*, 56, 497-500.
- Gogoi, S., Mandal, S. and Patel, A., 2018.** Effect of dietary *Wolffia arrhiza* and *Spirulina platensis* on growth performance and pigmentation of Queen loach *Botia dario* (Hamilton, 1822). *Aquaculture Nutrition*, 24, 285-291.
- Goytortua-Bores, E., Civera-Cerecedo, R., Rocha-Meza, S. and Green-Yee, A., 2006.** Partial replacement of red crab (*Pleuroncodes planipes*) meal for fish meal in practical diets for the white shrimp *Litopenaeus vannamei*, Effects on growth and in vivo digestibility. *Aquaculture*, 256, 414-422.
- Habibi, E., Kalbassi, M., Hosseini, S. and Qasemi, S., 2013.** Feasibility of Identification of Fall and Spring Migrating Caspian trout (*Salmo trutta caspius*) by Using AFLP Molecular Marker. *Turkish Journal of Fisheries and Aquatic Sciences*, 13, 241-248
- Harris, W.S., Connor, W.E., Inkeles, S.B. and Illinworth, D.R., 1984.** Dietary omega-3 fatty acids prevent carbohydrate-induced hypertriglyceridemia. *Metabolism* 33(11), 1016-1019.
- Harris, S.G., Josue, P., Koumas, L., Denise, R. and Phipps, R.P., 2002.** Prostaglandins as modulators of immunity. *TRENDS in Immunology*, 23, 3.
- Henrikson, R., 1998.** Earth food Spirulina. California/USA. Ronore Enterprises. 180 P.
- Hernandez, S.P., 2005.** Responsible use of antibiotics in aquaculture. Food and agriculture organization of the united nations, FAO Fisheries Technical. 469 P.
- Hirahashi, T., Matsumoto, M., Hazeki, K., Saeki, Y., Ui, M. and Seya, T., 2002.** Activation of the



- human innate immune system by *Spirulina*: augmentation of interferon production and NK cytotoxicity by oral administration of hot water extract of *Spirulina plantensis*. *International Immunopharmacology*, 2, 423–434.
- Hironobu, W., Kazuki, O., Asmi, C., Tassakka, T. and Masahiro, S., 2006.** Immunostimulant effects of dietary *Spirulina platensis* on carp, *Cyprinus carpio*. *Aquaculture*, 258, 157 - 163.
- Houston, A.H., 1990.** Components of the haematological response of fishes to environmental temperature change: A review. In "Environmental Physiology of Fishes" (Ali, M. A., ed.) Plenum, New York. pp. 241-298.
- Ibrahim, M.D., Mohamed, F.M. and Marwa, A.I., 2012.** The role of *Spirulina platensis* (*Arthrospira platensis*) in Growth, Immunity of Nile tilapia (*Oreochromis niloticus*) and its resistance to infection. *Journal of Agricultural Science*, 5(6), 5. DOI: 10.5539/jas.v5n6p109.
- James, R., Sampath, K., Thangarathinam, R. and Vasudevan, I., 2006.** Effect of dietary *Spirulina* level on growth, fertility, coloration and leucocyte count in red sword tail, *xiphophorus helleri*. *The Israeli Journal of Aquaculture–Bamidgeh*, 58(2), 97-104.
- Jeney, G. and Anderson, D.P., 1993.** Enhanced immune response and protection in rainbow trout to *Aeromonas salmonicida* bacterin following prior immersion in immunostimulants. *Fish and Shellfish Immunology*, 3(1), 51-58.
- Kalbassi, M., Dorafshan, S., Tavakolian, T., Khazab, M. and Abdolhay, H., 2006.** Karyological analysis of endangered Caspian salmon, *Salmo trutta caspius* (Kessler, 1877). *Aquaculture Research*, 37(13), 1341–1347.
- Khan, Z., Bhadouria, P. and Bisen, P.S., 2005.** Nutritional and therapeutic potential of *Spirulina*. *Current Pharmaceutical Biotechnology*, 6, 373-379.
- Khanzadeh, M., Fereidouni, A.E. and Berenjestanaki, S.S., 2016.** Effects of partial replacement of fish meal with *Spirulina platensis* meal in practical diets on growth, survival, body composition, and reproductive performance of three-spot gourami (*Trichopodus trichopterus*). *Aquaculture International*, 24, 69-84.
- Kiabi, B.H., Abdoli, A. and Naderi, M., 1999.** Status of the fish fauna in the South Caspian basin of Iran. *Zoology in the Middle East*, 18, 57-65.
- Kim, D.D., 2013.** Outdoor mass culture of *Spirulina platensis* in Vietnam. *Journal of Applied Phycology*, 2, 179–181.
- Kop, A. and Durmaz, Y., 2008.** The effect of synthetic and natural pigments on the colour of the cichlids (*Cichlasoma severum* sp., Heckel 1840). *Aquaculture International*, 16, 117-122.
- Kozlenko, R.D.P.M., Ph.D, M.P.H. and Henson, R.H., 1998.** Latest scientific research on spirulina:

- Effects on the AIDS virus, cancer and the immune system. Dana-Farber Cancer Institute and Harvard Medical School, Boston. inspiredliving.com
- Krishnaveni, R., Palanivelu, K. and Velavan, S., 2013.** Effects of probiotics and Spirulina supplementation on haemato-Immunological function of *Catla catla*. *International Journal of Research in Fisheries and Aquaculture*, 3(4), 176-181.
- Kumar, S., Sahu, N.P., Pal, A.K., Dharitri, C., Sona, Y. and Mukherjee., 2005.** Effect of dietary carbohydrate on haematology, respiratory burst activity and histological changes in *L. rohita* juveniles. *Fish and Shellfish Immunology*, 19, 331-344.
- Li, P. and Gatlin, D.M., 2004.** Dietary brewer yeast and prebiotic Grobiotic -A influence growth performance, immune responses and resistances of hybrid striped bass to *Streptococcus iniae* infection. *Aquaculture*, 231, 445-456
- Lu, J. and Takeuchi, T., 2002.** Taste of tilapia, *Oreochromis niloticus*, fed solely on raw spirulina. *Fisheries Science*, 68, 987-988.
- Manoj, G., Venkataraman, L.V. and Srinivas, L., 1992.** Antioxidant properties of Spirulina (*Spirulina platensis*). In: Seshadri C.V., Bai N.J.: Spirulina: National Symposium (India), MCRC, Tharamani, Madras. pp. 48-154.
- Misra, C.K., Das, B.K., Mukherje, S.C. and Pattnaik, P., 2006.** Effect of multiple injections of betaglucon on non-specific immune response and disease resistance in *Labeo rohita* fingerlings. *Fish and Shellfish Immunology*, 20, 305-319.
- Mustafa, M.G., Umino, T. and Nakagawa, H., 1994.** The effect of Spirulina feeding on muscle protein deposition in red sea bream, *Pagrus major*. *Journal of Applied Ichthyology*, 10, 141-145.
- Mustafa, M.G., Takeda, T., Umino, T., Wakamatsu, S. and Nakagawa, H., 1994.** Effects of *Ascomyllum* and *Spirulina* meal as feed additives on growth performance and feed utilization of red sea bream, *Pagrus major*. *J. Fac. Appl. Biol. Sci., Hiroshima Univ*, 33, 125-132.
- Mustafa, M.G. and Nakagawa, H., 1995.** A review: Dietary benefits of algae as an additive in fish feed. *The Israeli Journal of Aquaculture, Bamidgah*, 47, 155-162.
- Nakagawa, H., Sato, M. and Gatlin, D.M., 2007.** Dietary supplements for the health and quality of cultured fish. CABI Publishing, Oxfordshire, UK. 256 P.
- Nandeesh, M.C., Gangadhara, B., Varghese, T.J. and Keshavanath, P., 1998.** Effect of feeding *Spirulina platensis* on the growth, proximate composition and organoleptic quality of common carp, *Cyprinus carpio* L. *Aquaculture Research*, 2, 305-312.
- Nandeesh, M.C., Gangadhara, B., Manissery, J.K. and Venkataraman, L.V., 2001.** Growth performance of two Indian major carps, catla (*Catla catla*) and rohu (*Labeo rohita*) fed diets containing different levels of *Spirulina*

- platensis*. *Bioresource Technology*, 80, 117–120.
- NRC (National Research Council).**, 1993. Nutrient requirements of warmwater fishes and shellfishes, revised edn. National Academy Press, Washington. 225 P.
- Olvera-Novoa, M.A., Dominguez-Cen, L.J., Olivera-Castillo, L. and Martinez-Palacios, C.A., 1998.** Effect of the use of the microalgae *Spirulina maxima* as fish meal replacement in diets for tilapia, *Oreochromis mossambicus* (Peters), fry. *Aquaculture Research*, 29, 709–715.
- Palmegiano, G.B., Gai, F., Dapra, F., Gasco, L., Pazzaglia, M. and Peiretti, P.G., 2008.** Effects of Spirulina and plant oil on the growth and lipid traits of white sturgeon (*Acipenser transmontanus*) fingerlings. *Aquaculture Research*, 39, 587–595.
- Panigrahi, A., Kiron, V., Puangkaew, J., Kobayashi, T., Satoh, S. and Sugita, H., 2005.** The viability of probiotic bacteria as a factor influencing the immune response in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 243, 241–254.
- Pieretti, P.G. and Meineri, G. 2011.** Effects of diets with increasing levels of *Spirulina platensis* on the carcass characteristics, meat quality and fatty acid composition of growing rabbits. *Livest. Sci*, 140:218–224.
- Promya, J. and Chitmanat, C., 2011.** The effects of *Spirulina platensis* and *Cladophora* algae on the growth performance, meat quality and immunity stimulating capacity of the African Sharptooth Catfish (*Clarias gariepinus*). *International Journal of Agriculture and Biology*, 13, 77–82.
- Quillet, E., Faure, A., Chevassus, B., Kreig, F., Harache, Y., Arzel, J., Metailler, R. and Boeuf, G., 1992.** The potential of brown trout (*Salmo trutta* L.) for mariculture in temperate waters. *Icelandic Agricultural Sciences*, 6, 63–76.
- Raa, J., 1996.** The use of immune stimulatory substances in Fish and Shellfish farming. *Reviews in Fisheries Science*, 1(4), 229–288.
- Ragap, H.M., Khalil, R.H. and Mutawie, H.H., 2012.** Immunostimulant effects of dietary *Spirulina platensis* on tilapia *Oreochromis niloticus*. *Journal of Applied Pharmaceutical Science*, 2(2), 26–31.
- Rajabi Islami, H., Arab, N., Assare, R., Rastravan, M.I. and Ebtokari, R., 2016.** Blood parameters of Caspian brown trout (*Salmo trutta caspius*) fingerlings affected by dietary L-ascorbyl-2-polyphosphate. *Iranian Journal of Fisheries Sciences*, 15(3), 1167–1186.
- Regunathan, C. and Wesley, S.G., 2006.** Pigment deficiency correction in shrimp broodstock using Spirulina as a carotenoid source. *Aquaculture Nutrition*, 12, 425–432.
- Ramakrishnan, C.M., Haniffa, M.A., Manohar, M., Dhanaraj, M., Arockiaraj, A.J., Seetharaman, S. and Arunsingh, S.V., 2008.** Effects of probiotics and Spirulina on survival and growth of juvenile common carp *Cyprinus carpio*.

- Israel Journal of Aquacultures Bamidegh*, 60(2), 128–133.
- Rengpipat, S., Rukpratanporn, S., Piyatiratitivorakul, S. and Piamsak, M., 2000.** Immunity enhancement in black tiger shrimp, *Penaeus monodon* by a probiont bacterium. *Aquaculture*, 191, 271–288.
- Ravi, M., Lata, De.S., Azharuddin, S. and Solomon, P., 2010.** The beneficial effects of *Spirulina* focusing on its immunomodulatory and antioxidant properties. *Journal of Dietary Supplements*, 2, 73–83.
- Rook, G.A.W., Steele, J., Umar, S. and Dockrell, H.M., 1985.** A simple method for the solubilisation of reduced NBT, and its use as a calorimetric assay for activation of human macrophages by  $\gamma$ -interferon. *Journal of Immunological Methods*, 82, 161.
- Saurabh, S. and Sahoo, P.K., 2008.** Lysozyme: an important defence molecule of fish innate immune system. *Aquaculture Research*, 39(3), 223–239.
- Sakai, M., 1999.** Current research status of fish immunostimulant. *Aquaculture*, 172, 63–92.
- Salehi-Farsani, A., Soltani, M., Kamali, A. and Shamsaie, M., 2014.** Effect of immune motivator Macrogard and *Spirulina platensis* on some growth, carcass and biochemical indices of stellate sturgeon *Acipenser stellatus*. *Aquaculture, Aquarium, Conservation and Legislation International Journal of the Bioflux Society*, 7, 3.
- Sedgwick, S.D., 1995.** Trout farming handbook, 5th edn. Fishing News Books, Alden Press, Oxford. 169 P.
- Selmi, C., Leung, P.S.C., Fischer, L., German, B., Chen-Yen, Y., Kenny, T.P., Cysewski, G.R. and Gershwin, M.E., 2011.** The effects of *Spirulina* on anemia and immune function in senior citizens. *Cellular and Molecular Immunology*, 8(3), 248–254.
- Sheikhzadeh, N., Heidarieh, M., Karimi Pashaki, A., Nofouzi, K., AhrabFarshbaf, M. and Akbari, M., 2012.** Hilyses® fermented *Saccharomyces cerevisiae*, enhances the growth performance and skin nonspecific immune parameters in rainbow trout (*Oncorhynchus mykiss*). *Fish and Shellfish Immunology*, 32, 1083–1087.
- Shimaa, A.A., 2016.** Effect of *Spirulina platensis* as feed supplement on growth performance, immune response and antioxidant status of mono-sex Nile Tilapia (*Oreochromis niloticus*). *Benha Veterinary Medical Journal*, 30(1), 1–10.
- Siwicki, A. K., Anderson, D. P. 1993 :** Immunostimulation in Fish: Measuring the Effects of Stimulants by Serological and Immunological Methods. U.S. Fish and Wildlife Service, IFI, Poland. 1, 24.
- Schaperclaus, W., Kulow, H. and Schreckenbach, K., 1992.** Fish disease. A.A. Balkema, Rotterdam, the Netherlands. pp. 101–105.
- Shahbazi, S. and Bolhassani, A., 2016.** Immunostimulants: Types and functions. *Journal of Medical*

- Microbiology and Infectious Diseases*, 4(3-4), 45-51.
- Sotoudeh, E., Abedian Kenari, A., Khodabandeh, S. and Khajeh, K., 2015.** Combination effects of dietary EPA and DHA plus alpha-tocopherol: Effects on performance and physiological status of Caspian brown trout (*Salmo trutta caspius*) fry. *Aquaculture Nutrition*, 22(5), 1101-1115.
- Subhadra, B., Lochmann, R., Rawles, S. and Chen, R., 2006.** Effect of dietary lipid source on the growth, tissue composition and hematological parameters of largemouth bass (*Micropterus salmoides*). *Aquaculture*, 255(1), 210-220.
- Tang, H.G., Wu, T.X., Zhao, Z.Y. and Pan, X.D., 2008.** Effects of fish protein hydrolysate on growth performance and humoral immune response in large yellow croaker (*Pseudosciaena crocea* R.). *Journal of Zhejiang University Science*, 9, 684-690.
- Taoka, Y., Hiroto, M., Jae-Yoon, J.O., Min-Jee, J., Sungchul, C.BAI., Won-Jae, L., Kazuya, Y. and Shunsuke, K., 2006.** Growth, stress tolerance and non-specific immune response of Japanese flounder *Paralichthys olivaceus* to probiotics in a closed recirculating system. *Fisheries Science*, 72, 2.
- Teimouri, M., Amirkolaie, A.K. and Yeganeh, S., 2013.** Effect of *Spirulina platensis* meal as a feed supplement on growth performance and pigmentation of Rainbow Trout (*Oncorhynchus mykiss*). *World Journal of Fish and Marine Sciences*, 5(2), 194-202.
- Tietz, N.W., 1986.** Textbook of clinical chemistry, W.B. Saunders, Philadelphia, PA. 1919 P.
- Terry, C.H., Cardinale, J.L. and Smith, S.A., 2000.** Haematology and plasma chemistry reference intervals for cultured tilapia (*Oreochromis hybrid*). *Veterinary Clinical Pathology*, 29, 7-12.
- Tongsiri, S., Mang-Amphan, K. and eerapornpisal, Y., 2010.** Effect of replacing fishmeal with Spirulina on growth, carcass composition and pigment of the mekong giant catfish. *Asian Journal of Science*, 2, 106-110.
- Tort, L., Balasch, J.C. and Mackenzie, S., 2003.** Fish immune system. A crossroads between innate and adaptive response. *Immunologia*, 22(3), 277-286.
- Tukmechi, A., Rahmati Andani, H.R., Manaffar, R. and Sheikhzadeh, N., 2011.** Dietary administration of beta-mercapto-ethanol treated *Saccharomyces cerevisiae* enhanced the growth, innate immune response and disease resistance of the rainbow trout, *Oncorhynchus mykiss*. *Fish Shellfish Immunology*, 30, 923-928.
- Ungsethaphand, T., Peerapornpisal, Y., Whangchai, N. and Sardud, U., 2010.** Effect of feeding *Spirulina platensis* on growth and carcass composition of hybrid red tilapia (*Oreochromis mossambicus* × *O. niloticus*). *Maejo International Journal of Science and Technology*, 4(2), 331-336.

**Vonshak, A., Laorawat, S., Bunnag,**

**B, and Tanticharoen, M., 2014.** The effect of light availability on the photosynthetic activity and productivity of outdoor cultures of *Arthrospira platensis* (Spirulina). *Journal of Applied Phycology*. 26: 1309-1315.

**Watanuki, H., Ota, K., Malina, A.S.,**

**Kato, T. and Sakai, M., 2006.** Immunostimulant effects of dietary *Spirulina platensis* on carp, *Cyprinus carpio*. *Aquaculture*, 258, 157-163.

**Yong-Chin, L., Carina Miranda, T.,**

**Chien-Lun, H., Wen-Ching, T. and Jiann-Chu, C., 2010.** White shrimp *Litopenaeus vannamei* that had received the hot-water extract of *Spirulina platensis* showed earlier recovery in immunity and up-regulation of gene expressions after pH stress. *Fish and Shellfish Immunology*, 29, 1092-1098.

**Zamini, A. and Azimi, A., 2015.** The

effect of *Spirulina platensis* algae in nutrition on blood and immunity parameters of koi (*cyprinus carpio*). *Academie Royale DesScience D Outre-Mer Bulletin DesSeances*, 4(3), 147-151.

**Zeinab, A.K., Aly, M.S., Faiza, A.**

**and Fatma, E.M., 2015.** Effect of *Spirulina platensis* and *Lactobacillus rhamnosus* on growth and biochemical performance of Nile Tilapia (*Oreochromis niloticus*) fingerlings. *International Journal of Current Microbiology and Applied Sciences*, 4(4), 747-763