DOI: 10.22092/ijfs.2019.119703 Hematite-biochemical and immune response of Caspian brown trout (Salmo troutta caspius, Kessler, 1877) juveniles fed different

19(3) 1153-1174

levels of spirulina (Spirulina platensis)

Meshkat Roohani A.¹; Fallahi Kapoorchali M.^{1*}; Abedian Kenari A.²; Sayyad Borani M.¹; Zorriezahra M.J.³

Received: July 2018 Accepted: August 2019

Abstract

The effects of dietary Spirulina platensis on haemato-biochemical and immunity responses of Caspian brown trout (Salmo troutta caspius) juveniles was investigated. For this purpose, diets with five S. platensis inclusion levels (control, 2%FMR (13.2 g kg⁻¹ spirulina in diet), 4%FMR (26.4 g kg⁻¹ spirulina in diet), 6%FMR (39.6 g kg⁻¹ spirulina in diet), and 8%FMR (52.8 g kg⁻¹ spirulina in diet) were prepared. Six hundred juveniles with an average initial weight of 11±1.0 g were assigned to 15 experimental tanks. The experiment lasted for 10 weeks. At the end of the experiment, growth performance, haemato-biochemical parameters including white and red blood cell counts, neutrophils lymphocytes counts, hematocrit, hemoglobin, glucose, albumin, total protein, aspartate amino transferase (AST), alanine amino transferase (ALT), triglyceride, cholesterol, as well as immunity parameters including lysozyme, C3, C4, Immunoglobulin (IgM), ACH50 and respiratory burst activity were assessed. The results indicated that fish fed diets supplemented with 6%FMR and 8%FMR had a significantly higher weight gain (26.13 g and 25.88 g) and specific growth rate (1.74 %bw day⁻¹ and 1.71 %bw day⁻¹) compared with control. Furthermore, 6%FMR and 8%FMR treatments had statistically higher protein efficiency (0.76 and 0.78), lipid efficiency (1.89 and 1.94) and statistically lower feed conversion ratio (2.91 and 2.84) compared to the other treatments respectively (p<0.05). The physiological and immunological factors were improved when fish were fed a high level of S. platensis supplement. S. platensis inclusion also increased activity of Lysozyme C3, C4, IgM and ACH50 and respiratory burst activity and reduced AST and ALT formation. These results indicate that S. platensis supplement is promising for disease prevention in S. trutta caspius juveniles, at an optimum dietary level of 6% in diet.

Keywords: Spirulina, *Salmo trutta caspius*, Growth performances, Haemato-biochemial parameters, Immunity.

¹⁻Inland Waters Aquaculture Research Center, Iranian Fisheries Science Research Institute, Agriculture Research, Education and Extension Organization (AREEO), Bandar Anzali, Gilan, Iran

²⁻Department of Aquaculture, Faculty of Natural Resources and Marine Sciences, Tarbiat Modares University, Noor, Mazandaran, Iran

³⁻Iranian Fisheries Science Research Institute, Agriculture Research, Education and Extension Organization (AREEO). Tehran, Iran

^{*}Corresponding author's Email: m_fallahi2011@yahoo.com

Introduction

performance Growth and disease control are the most important priorities in current aquaculture. Nowadays, use of antibiotics and chemotherapeutics has increased significantly, due to intensification of culture practices and by high prevalence followed infectious diseases (Andrews et al., 2011). Accumulation of antibiotic residues in fish tissues and environment causes human and animal health issues and led to an urgent need for alternative disease preventive substances and use of immune-stimulants in aquaculture (Andrews et al., 2011; Krishnaveni et al., 2013). Several types of stimulants with different mechanisms and functions such as bacterial products, carbohydrates, vaccines. complex immune enhancing drugs, nutritional factors, animal and plant extracts are considered, and their abilities to activate the innate immunity have been studied (Raa, 1996; Sakai, 1999; Shahbazi and Bolhassani, 2016).One supplement source, with ease of production, processing and distribution along with a wide range of macro- and micronutrients of health benefits is spirulina (Ravi et al., 2010). Research has shown that this alga contains a wide variety of compounds such phycobiliproteins, carotenoids, polysaccharides, phycocyanin, unsaturated fatty acids, superoxide dismutase, different vitamins, and other elements which by bring improvement coloration, body growth immunity of fish (Nakagawa et al., 2007). Spirulina (Spirulina platensis), a blue-green filamentous, fresh water and

multi-cell microalgae is gaining more attention from medical scientists as a nutraceutical and source of potential pharmaceuticals. There are several new peer reviewed scientific studies about spirulina's ability to inhibit viral replication, strengthen and enhancing both the cellular and humoral arms of the immune system (Kozlenko et al., 1998; Hirahashi et al., 2002; Andrews et al., 2011). It has been received the most research and public health attention due to its bioactive including compounds vitamins, essential amino acids, minerals, essential fatty acids (gamma linolenic acid), and antioxidant pigments such as phycocyanine (Belay et al., 1996; Regunathan and Wesley, 2006; Ragap et al., 2012); Ceballos et al., 2006). Additionally, spirulina has high protein content (60-70% by dry weight) and is the richest natural source of vitamin B₁₂ (Estrada et al., 2001). This alga contains a whole spectrum of natural mixed carotene and xanthophylls phytopigments (Belay et al., 1996; Estrada et al., 2001; Andrews et al., 2011). It has been confirmed that the addition of small amounts of algae to fish feed can exert pronounced effects on growth, lipid metabolism, body composition and physiological response to stress and disease (Mustafa et al., 1994; Mustafa and Nakagava, 1995; Palmegiano et al., 2008; Abdel-Tawwab and Ahmad, 2009; Teimouri et al., 2013). Spirulina as an immune modulator not only stimulates the immune system, but also strengthens the body's ability to produce new blood cells (Andrews et al., 2011). Important

parts of the immune system, such as bone marrow stem cells, macrophages cells (number of cells phagocytosis), T cells (lymphocyte), NK (non-specific cytotoxic), spleen as well as thymus show tangible activities when treated with spirulina extract (Henrikson, 1998; Watanuki et al., 2006; Tongsiri et al., 2010; Yong-Chin et al., 2010; Shahbazi and Bolhassani, 2016). Recently, spirulina has been speculated to be associated modulation of the host immune system (Hironobu et al., 2006).

Caspian brown trout (Salmo trutta caspius, Kessler 1877), is one of the world's nine subspecies of brown trout (Quillet et al., 1992; Habibi et al., 2013) and attains the greatest size and growth rate of all brown trout (Sedgwick, 1995; Rajabi et al., 2016). This species is a critically endangered anadromous species distributed in southern region of the Caspian Sea. In 1999, this species was declared at risk according to IUCN conditions (Kiabi et al., 1999; Coad, 2000; Kalbassi et al., 2006). Artificial propagation releasing fingerlings to the natural waters is an approach to prevent brown trout's extinction (Rajabi et al., 2016). In addition, in the recent years, this species has attracted interest aquaculture in cage and raceways in intensive culture systems (Kalbassi et al., 2006). S. trutta caspius, like many other species is sensitive to stressors and pathogenic agents. Thus, strong defense mechanisms or immune system needed against pathogens to improve the health of the fish. Despite the importance of Caspian salmon as an endangered species, however, little is known about their nutritional requirements especially in the field of increasing immunity of species and types of immune-stimulants. Therefore, the main objective of this study was to evaluate the effects of S. *platensis* meal as feed additive on immune system, biochemistry and haematology in *S. trutta caspius* as a valuable species.

Materials and methods

Experimental diets

Five artificial diets were formulated using five levels of microalgae S. platensis (Sina microalgae Co., Qeshm, Iran) and in three replicates were examined in a randomized design. The experimental diets were formulated by partially replacing fishmeal in the basal diet (65.89% in control) with spirulina powder at inclusion levels of 0, 2%, 4%, 6% and 8%, respectively. Diet formulations were performed using Lindo software (Lindo copyright, release, 6.11998), and fishmeal and S. platensis powder were used as a protein source. Test diets provided, which were iso-nitrogenous (45-46% crude protein) and iso-caloric (20 MJ kg⁻¹) were analyzed in this regard, only varied in terms of fish meal and S. platensis content. The proximate composition of experimental diets was measured according to the standard methods of Association of Official Analytical Chemists (AOAC, 1995) for moisture, protein, fat, and ash determination. Briefly, Moisture content was estimated by drying the samples to constant weight at 95 °C in a drying oven (GCA, model 18EM. Precision Scientific

Group, Chicago, IL, USA). N content was measured using the Labconco Micro Kjeldahl Apparatus (Labconco Corporation, Kansas, MO, USA) and crude protein was estimated multiplying N content by 6.25.

Lipid content was determined by acetone extraction using the extraction Soxhlet apparatus (Lab-Line Instruments, Inc., Melrose Park, IL, USA) for 16 hours, and ash was determined by combusting dry samples in a muffle furnace (Thermolyne Corporation, Dubuque, IA, USA) at 550 °C for 6 hours. Carbohydrate (i.e., nitrogen-free extract plus fiber) was calculated by the difference:

(100-(protein+fat+ash+moisture)) (Aksnes and Opstvedt, 1998).

Gross energy was calculated using the coefficient of 23.6, 39.5, and 17.2

 g^{-1}) (kJ for protein, fat, and carbohydrates, respectively (NRC. 1993). The experimental formulation and proximate composition are showed in Table1, respectively. Dietary feed ingredients were ground using a laboratory grinder (Philips HR7628, Finland) and then blended into a homogenous doughy matter by adding water and transferred to a meat grinder with a 2 mm-mesh (CGT Company, 2mec, Rome, Italy). After drying, diets were broken by hand to fit fish mouths and placed in a grinding machine for breakdown into small pieces. The resulting diets were then dried in food dryer and stored in plastic bags in a refrigerator at -2 °C until further use.

Table 1: Dietary formulation and proximate composition of experimental diets.

Inquedient (a leaf due weight)		·			
Ingredient (g kg ⁻¹ dry weight)	0%	2%	4%	6%	8%
Fish meal 1	658.9	645.7	632.4	619.4	606.2
Spirulina ²	0	13.2	26.5	39.5	52.7
Wheat flour	115.9	117.8	119.4	120.9	122.4
Fish oil ³	45	45	45	45	45
Soybean oil ⁴	50.3	51.3	52.3	53.3	54.5
Lethitin ⁵	5	5	5	5	5
Filler(sand)	60	57.1	54.6	52.1	49
Mineral mixture ⁶	25	25	25	25	25
Vitamin mixture ⁷	15	15	15	15	15
Anti-oxidant ⁸	0.2	0.2	0.2	0.2	0.2
Choline Chloride ⁹	1	1	1	1	1
Mono calcium phosphate ¹⁰	5	5	5	5	5
Anti fungal ¹¹	2.5	2.5	2.5	2.5	2.5
Lysine ¹²	5	5	5	5	5
Methionine ¹³	5	5	5	5	5
Vitamin C ¹⁴	1	1	1	1	1
Chemical analysis (%)					
protein	45.1	45.2	44.89	45.07	45.1
lipid	17.96	17.75	18.1	18.2	18.1
moisture	7.9	8.02	8.00	7.9	8.1
ash	8.9	8.8	8.7	8.9	9.1
carbohydrate	12.14	11.93	12.21	11.73	11.5
Gross Energy(MJ kg ⁻¹)	19.83	19.73	19.84	19.84	19.8

Herring meal, produced by Mirood, Mazandaran province, Iran.

Produced in Sina micro algae, Qeshm. Hormozgan province, Iran.

Produced by Mirood, Mazandaran province, Iran.

Prepared of Mazandaran aquatic food factory, Sari, Mazandaran province, Iran.

⁶ Mineral premix consisted of (mg kg⁻¹ premix): Man=2600mg, Cu=600mg, Fe=6000mg, Zn=4600mg, Se=100 mg, I=100 mg, Co=50 mg and carrier up to 1 kg.

For Nag. 7 Vitamin premix consisted of (mg kg⁻¹ premix): A=1200000 IU, D3=400000 IU, E= 3000IU, K3=1200mg, C= 5400mg, B1=200mg, H2=200 mg, B2=3360mg, B12=4mg, B9=600mg, B5=9000mg, B7=7200mg

Preparation of the Mazandaran aquatic food factory and its combination including BHT, Etoxyquin , Synergist Propylgallate.

Ocholine Chloride, 60%, Prepared of livestock and poultry pharmaceutical factory Aras Bazar 10.11, 12, 13 Prepared of pharmaceutical factory Aras Taban, Amol, Mazandaran province, Iran.

¹⁴ Prepared of pharmaceutical factory Aras Bazar, Amol, Mazandaran province, Iran. 15 Aqutic vitamin C, production of pharmaceutical factory Aras Bazar, Amol, Mazandaran province, Iran.

Fish rearing

Fish juveniles (n= 600; mean individual initial weight 11±1.0 g) were obtained from Cold Water Breeding Restocking Hatchery, Shahid Bahonar Center in Kelardasht, Mazandaran, Iran, and then were transferred to NIAC multi-purpose co-operative farm fish in Baghbankola village, Amol. Mazandaran, Iran, Fish were allowed to acclimatize for two weeks in indoor fiberglass tanks prior to the experiment and during this period were fed a commercial diet twice daily. After the acclimation period, fish were randomly divided into five equal groups, each comprising three replicates (40 fish per replicate) in 15 separate 400 L fiberglass tanks. The fish were hand-fed three times daily (8:00, 12:00 and 16:00 h) to apparent satiation by visual observation for ten weeks (Sotoudeh et al., 2015) Water temperature, dissolved oxygen and pH were monitored daily (during the experiment, temperature, dissolved oxygen and pH were 15±2 $^{\circ}$ C, 8.3±1 mg L⁻¹ and 7.6 ± 0.3 respectively). Uneaten feed and fecal matter were siphoned off every day. Due to the sensitivity of these juveniles, and to prevent mortality, biometry was performed once at the beginning and once at the end of this period.

Growth parameters

Growth performance was determined and feed utilization was calculated as following (Goytortua-Bores *et al.*, 2006):

Weight gain (WG %)=100×(final body weight-initial body weight)/initial body weight.

Specific growth rate (SGR (%))=100 (ln W2-ln W1)/T; where W1 and W2 are the initial and final weight, respectively, and T is the number of days in the feeding period;

Feed conversion ratio (FCR) = feed consumed (g dry weight)/weight gain (g);

Condition Factor (CF) = Weight/Total length $^{3}\times100$;

Survival rate (%) = (Number of fish at the beginning/Number of fish at the end) $\times 100$.

Sampling

At the end of our experiments, to evaluate haemotological parameters, fish were fasted for 24 hours immediately prior to blood sampling and then six fish from each tank were sampled randomly. To prevent sampling stress. the fish were anesthetized with a stock solution (50 ppm) of clove oil and blood samples were collected quickly (approximately one min/fish) by puncturing the veins in the caudal peduncle with a sterile 5 ML syringe (Esmaeili et al., 2017). The collected blood sample was divided into portions. One portion two was transferred into **Eppendorf** tubes containing heparin anti-coagulant (500 U L⁻¹) for haematological tests and the second portion of blood sample was also transferred into Eppendorf tubes, left to clot at 4 °C and centrifuged at 5000 rpm for 5 min at room for temperature biochemical and Immunological tests.

Haemotology tests

Red blood cells (RBC) and white blood cell (WBC) counts were determined Neubaeur haemocytometer using following the methods of Blaxhall and Daisley (1973). Blood was diluted to 1:200 with Race solution. Five center cells of middle square in Neubauer chamber were used for counting RBC and multiplied by 10,000 (Esmaeili et al., 2017). For counting WBC, four marginal squares in Neubauer chamber were used after blood dilution (1:50) with Race solution and the results were multiplied by 50. Then, blood samples were fixed in methanol and stained using Wright-Giemsa stain for determination of the differential WBC (lymphocyte and neutrophil portions of total WBC) count (Houston, 1990). At least 200 WBCs were counted for differential WBC determinations. Thin slices were used for counting WBC counting was done according to their shape and reported as percentages. To measure the Hemoglobin (Hb) levels obtained were by the Cyanmethemoglibin method (Houston, 1990). In this method, the RBC are haemolized and hemoglobin Cyanmethemoglobin. converted to Twenty microliters of blood samples with 50-ul Drabkin's reagent were mixed and placed in dark place for 10 min to form Cyanemethemoglobin. Then, absorbance was read at 540 nm in a UV with spectrophotometer (Jenway 6800, UK) and finally, hemoglobin value was calculated. Haemotocrit (Ht) was measured using the standard microhaemotocrit method (Subhadra et al., 2006) and reported as percentages.

Microhematocrit capillary tubes were centrifuged at 2500 rpm for 5 min, and then hematocrit value was calculated using graded plate. To calculate others haematological indices, MCV (mean corpuscular volume), MCH (mean corpuscular hemoglobin), **MCHC** (mean corpuscular hemoglobin concentration), the relevant formula (Bain et al., 2011) was used as follows: MCV (fl)=hematocrit/RBC(million per $mm^3)\times10$

MCH (pg)=hemoglobin/RBC(million per mm 3) × 10

MCHC=hematocrit/hemoglobin×100

Blood biochemical parameters

Total protein (modi-biuret method, Tietz, 1986) and Albumin (romocresol green binding method, Doumas and Peters, 1997) assays were performed with Zist Chemistry kits (Zist chemistry Company, Tehran, Iran). Total protein content (at wavelength of 560-520 nm) and Albumin (at wavelength of 630 nm) were determined using an auto analyzer (Thecnicon, RA 1000. system NewYork, NY, USA). Globulin levels of the samples were calculated from the difference in albumin from total protein (Kumar et al., 2005). Total cholesterol, triglyceride and glucose measurements were performed using Pars Azmoon kits commercial (Pars Azmoon Company, Tehran, Iran) according to the company's protocol and using an auto-analyzer (Thecnicon, RA 1000, New York, NY, USA). Briefly, in this method, 10 µl of plasma sample had mixed with 1000 µl of reagent and incubated for 20 min in room temperature. Then, the absorbance of the sample against the blank (10 µl of distilled water mixed with 1000 µl of reagents) had measured at a wavelength of 546 nm. Similar approach had done for other parameters, but with appropriate sample amount and wavelength according each to parameter's protocol (Esmaeili et al., 2017). The Kinetic method was used to measure liver enzymes, aspartate aminotransferase (AST) and alanine aminotransferase (ALT), using Pars Azmoon commercial clinical investigation kits (Parsazmon Company, Tehran, Iran) and were analyzed with an auto analyzer (Technicon RA-1000; Technicon Instruments, New York, NY, USA).

Immunological parameters

The lysozyme activity in serum was determined turbidometrically using the method described by Ellis (1999). In this method, Micrococcus bacteria was used (Sigma, M0508) with phosphate buffer (30 ml, 0.1 M, pH=7) as substrate. For this purpose, activated bacteria were used and cultivated at a concentration in 0.6-0.8 nm adjusted with spectrophotometer (Jenway 6800, UK). Then, 1990 µl of substrate and 10 µl of the sample were added to cuvette. The decrease of absorption at 450 nm each min for a period of 6 min had measured through spectrophotometer (Tukmechi et al., 2011) with chicken egg white (Sigma, A5503) standard. **Turbidometric** method and Pars Azmoon commercial clinical investigation kits (Pars Azmoon Company, Tehran, Iran) were used to measure C3 and C4 serum complement system (Tang et al., 2008). The serum C3 and C4 reacted with the antibodies contained in the kit, and then OD had measured at a wavelength of 340 nm using an auto-analyzer (Thecnicon, RA 1000, New York, NY, USA), compared to the standard in kits and finally, the C3 and C4 amount was calculated based on mg L-1. ACH50 (the activity complement system) was determined according to the indirect ELISA method using a commercial kit (Wielsa, comple 300 total complement functional screen kit, Sweden). The volume of the serum complement producing 50% hemolysis (ACH50) was determined, and the number of ACH50 units in ml was calculated for each experimental fish.

Total immunoglobulin levels were determined according to the method described by Siwicki and Anderson (1993). In this method, analysis of total Ig level in plasma is based on the biuret colorimetric method. First the immunoglobulins were separated from precipitation plasma by with polyethylenic glycol 10,000 (Sigma) and remaining supernatant was read. That number was subtracted from the total protein to give total immunoglobulin. Protein readings from supernatant gave the amount of protein taken out by absorption to polyethylene glycol. To calculate total immunoglobulin, subtracted these readings from total protein on individual samples and compared with standards for calculation of protein. The production of oxygen radicals by

leukocytes was determined by the

reduction of Nitro Blue Tetrazolium

(NBT, Sigma-Aldrich Chemical, St. Louis, MO, USA) according to Rook et al. (1985). Absorbance was converted to NBT units based on a standard curve of NBT diformazan per milliliter of blood.

Statistical analysis

All the data have been analyzed using the SPSS (version 17, SPSS, Richmond, VA, USA) statistical package as described by Dytham (1999). Mean values of all the parameters were subjected to one-way analysis of variance (ANOVA) to study the treatment effect and comparison of any mean values had done by Duncan's Multiple Range Test (DMRT). Comparisons were made at the five percent probability level (p<0.05).

Result

The growth performance of Caspian trout juvenile fed diets containing different levels of spirulina are presented in Table 2. Overall, the optimum growth performance has been

obtained at 6%FMR and 8%FMR treatments, whereas the fish fed control had the lowest growth performances. According to the results, fed supplemented fish diet 6%FMR 8%FMR had and significantly higher weight gain (26.13 g and 25.88 g) and specific growth rate (1.74 %bw day⁻¹ and 1.71 %bw day⁻¹) compared with those fed the control diet (18.18 g and 1.37% bw day⁻¹). Furthermore, 6 %FMR and 8 %FMR treatments had statistically higher protein efficiency (0.76 and 0.78), lipid efficiency (1.89)and 1.94) statistically lower feed conversion ratio (2.91 and 2.84) compared with other treatments (p<0.05). Moreover, there was no significant difference in the condition factor among experimental treatments. In addition, fish fed control, 2 %FMR and 4 %FMR diets exhibited equivalent FCR, protein efficiency and lipid efficiency. Survival rate in all treatments was 100%.

Table 2: Mean growth performance of Salmo trutta caspius fed different levels of Spirulina platensis.

Growth performance	Experimental Diets					
Growth performance	0% control	2%	4%	6%	8%	
$W_1(g)$	11.18±0.23	11.91±1.23	11.16±0.56	10.91±1.61	11.19±0.33	
WG (%)	$62.73\pm12.3^{a*}$	93.93±11.31 ^{ab}	101.41 ± 14.29^{ab}	163.35 ± 67.22^{c}	131.96±35.27 ^{bc}	
SGR (% / day)	0.81 ± 0.13^{a}	1.1 ± 0.1^{ab}	1.16 ± 0.12^{ab}	1.58 ± 0.16^{c}	1.39±0.26bc	
CF	1.33 ± 0.05^{a}	1.28 ± 0.03^{a}	1.38 ± 0.06^{a}	1.27 ± 0.04^{a}	1.31 ± 0.14^{a}	
FCR	3.93 ± 0.74^{b}	3.95 ± 0.49^{b}	2.73 ± 0.06^{a}	2.11 ± 0.49^{a}	2.04 ± 0.56^{a}	
Survival rate (100%)	100	100	100	100	100	

W1: Primary weight WG%= Weight gain

FCR: Feed conservation ratio

SGR: Specific growth ratio

Conditional Factor(CF)

^{*}Values are the least square means±standard errors of the means of triplicate samples. P values determined with oneway ANOVA are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different (p>0.05). Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

The mean haemotology parameters of S. trutta caspius juveniles fed with diets containing different levels platensis are shown in Table 3. Based on obtained results, spirulina had a significant impact on haemotological parameters (RBC, WBC, Hct, Hb, lymphocytes, neutrophils) in S. trutta caspius juveniles. RBC increased with increasing levels of dietary S. platensis to 6%, but in 8%, the rate decreased (p<0.05). WBC, Hct, Hb, differential white blood cell percent, lymphocyte neutrophil increased with and

increasing levels of dietary *S. platensis*. So, the lowest and the highest levels for these parameters (with exception of neutrophils) were observed in the control and 8% treatments, respectively. Other haematological indices such as average volume of red blood cells (MCV), mean hemoglobin in red blood cells (MCH), average concentration of hemoglobin in a red blood cells (MCHC) also have increased with increasing levels of dietary *S. platensis* (p<0.05).

Table 3: Haemotological parameters of Salmo trutta caspius fed with different levels of Spirulina platensis.

Unemetalogical parameters			Experimental Die	ts	
Haemotological parameters	0% control	2%	4%	6%	8%
RBC ($\times 10^6$ n mm ⁻³) WBC (cell ml ⁻¹)	0.77±0.17 ^{a*} 9316±3521.6 ^a	1.22±0.05 ° 10100±1649.24 ^b	1.25±0.16 ^{cd} 11800±2415.7 ^c	1.27±0.09 ^d 13500±2485.4 ^d	1.12±0.15 ^b 15400±2256.5 ^e
Hct (%) Hb (g dl ⁻¹) LYM (% of WBC)	24.66±1.98 ^a 4.32±0.61 ^a 85.5±3.39 ^a	35.83±1.94 b 7.95±0.36 bc 89.5±1.76 b	37.17±1.83 ° 7.59±0.75 ^b 92.66±7.28 °	37.66±0.81 ° 8.16±0.31 ° 98.16±1.16 d	37.83±1.26 ° 8.9±0.51 ^d 98.66±0.81 ^d
Neut (% of WBC) MCV(fl) MCH(pg) MCHC (%)	1.16±0.75 a 280.81±6.96 a 52.37±1.3 a 17.63±1.77 a	1.16±0.75 a 298.76±25.03 ab 56.1±1.72 a 19.98±0.71 b	1.33±0.81 a 300.98±26.45 b 61.23±4.17 a 20.4±1.18 b	1.67±0.81 ° 332.15±17.8 ° 64.57±2.3 ^{ab} 21.68±1.04 ^b	1.5±0.83° 337.21±12.2° 80.13±6.7 b 23.78±2.13°

*Values are least square means \pm standard errors of the means of triplicate samples. p values determined with two-way ANOVA tests are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different (p>0.05). Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

The mean blood biochemical parameters of *S. trutta caspius* juveniles fed with diets containing different levels of *S. platensis* are summarized in Table 4&5. Based on obtained results, with increasing levels of dietary *S. platensis* up to 6%, ALT and AST enzymes decreased (*p*<0.05). Total

protein and globulin decreased and then increased in 8% treatment (p<0.05). Other blood biochemical parameters such as glucose, cholesterol and triglycerides have significantly decreased with increasing levels of dietary S. platensis (p<0.05).

Table 4: Liver enzymes of Salmo trutta caspius fed different levels of Spirulina platensis.

_	Experimental Diets					
Liver enzymes	0% control	2%	4%	6%	8%	
ALT(U L ⁻¹)	323±45.2 ^{d*}	305.4±27.1 ^{bc}	290±20.1 ^b	214.2±28.7 ^a	212.3±16.2 ^a	
AST(U L ⁻¹)	25.4±4.5 ^d	23.7±3.8 ^{cd}	22.9±6.6 ^{bc}	21.3±4.3 ^b	12.2±2.5 ^a	

^{*}Values are least square means \pm standard errors of the means of triplicate samples. p values determined with two-way ANOVA tests are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different (p>0.05). Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

Table 5: Blood biochemical parameters of Salmo trutta caspius fed different levels of Spirulina

	Experimental Diets						
Biochemical factors	0% control	2%	4%	6%	8%		
Total protein (g dl ⁻¹)	2.6±0.4 ^{a*}	2.6±0.4 ^a	2.8±0.7 ^b	3.1±0.3°	4.3±0.5 ^d		
Albumin (g dl ⁻¹)	1.8 ± 0.3^{a}	1.8 ± 0.3^{a}	1.7 ± 0.1^{a}	2 ± 0.3^{a}	2 ± 0.3^{a}		
Globulin (g dl ⁻¹)	0.8 ± 0.1^{a}	0.8 ± 0.1^{a}	1.1 ± 0.2^{b}	1.1 ± 0.3^{b}	2.3 ± 0.3^{c}		
Glucose (mg dl ⁻¹)	172.5 ± 16.3^{e}	176 ± 7.8^{d}	121.5±12.3 ^b	133.1 ± 15.9^{c}	94.5 ± 8.2^{a}		
Cholesterol (mg dl ⁻¹)	438.6 ± 45.1^{d}	412.5±47.3°	401.4 ± 26.2^{b}	346.8 ± 42.2^{a}	394.8 ± 21.6^{b}		
Triglyceride (mg dl ⁻¹)	369.9±5.3 ^e	356.4 ± 40.5^{d}	326.9 ± 45.8^{c}	322.7 ± 58.9^{b}	253.8 ± 61.4^{a}		

^{*}Values are least square means \pm standard errors of the means of triplicate samples. pvalues determined with two-way ANOVA tests are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different (p>0.05). Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

The mean immunity parameters of *S. trutta caspius* juveniles fed with diets containing different levels of *S. platensis* are shown in Table 6 and 7. Based on obtained results, lysozyme enzyme activity increased with increasing dietary *S. platensis* levels up to 6% and then decreased in 8%

Complement C3, C4. treatment. ACH50, free radical oxygen and immunoglobulin were significantly affected by dietary S. platensis levels. The lowest and the highest values for these parameters have been observed in the control and 8% treatments.

Table 6: Lysozyme activity of Salmo trutta caspius fed different levels of Spirulina platensis..

Lysozyme	-	E	Experimental Diet	S	
activity	0% control	2%	4%	6%	8%
After 15 sec (µg ml ⁻¹)	1.44±0.82 ^{a*}	4.3±0.75 ^b	4.95 ± 0.32^{c}	7.33 ± 0.22^{d}	5.87±0.93 ^e
After 180 sec (µg ml ⁻¹)	6.22 ± 0.35^a	6.32 ± 0.38^{a}	6.71 ± 0.28^{b}	8.59±1.06°	7.95 ± 0.44^d

^{*}Values are least square means \pm standard errors of the means of triplicate samples. p values determined with two-way ANOVA tests are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different (p>0.05).Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

Table 7: Immunological parameters of Salmo trutta caspius fed different levels of Spirulina platensis.

Presentation							
Immunological	Experimental Diets						
factor	0% control	6%	8%				
$C_3 \text{ (mg ml}^{-1})$	21.1±3.3 a*	21.1±3.3 ab	23.4±2.5 b	25.7±3.2°	35.7±6.9 ^d		
$C_4 \text{ (mg ml}^{-1})$	13.9±3.6 a	14.7 ± 2.7^{a}	15 ± 2.4^{a}	19.2 ± 1.8^{b}	20.8 ± 1.5^{b}		
ACH50(unit ml ⁻¹)	220.1±3.87 a	225.16 ± 4.9^{b}	228.23±7.1°	234.33 ± 8.73^{d}	247 ± 4.77^{e}		
Igm(mg dl ⁻¹)	183.4 ± 25.4^{a}	183.7 ± 21.7^{a}	194.6±19.2 ^b	321.5 ± 13.6^{c}	334.5 ± 25.8^{d}		
$O^{-}(RLU s^{-1})$	384.96 ± 21.05^{a}	401.5 ± 19.38^{b}	415.51 ± 29.73^{c}	429.28 ± 23.31^{d}	471.03 ± 23.77^{e}		

*Values are least square means \pm standard errors of the means of triplicate samples. p values determined with two-way ANOVA tests are also provided for the main factors and their interactions. Means marked with the same letter are not significantly different (p>0.05).Letters a, b, c, d, and e indicate significant differences in treatments, according to Duncan's test.

Discussion

present results indicated that spirulina significantly improved the growth performance of juvenile Caspian brown trout and the fish meal could be replaced with this microalga until 8% without impairments. During the experimental period, no mortality and disease in fish fed spirulina were observed compared to control. In agreement with our research, many studies have demonstrated a positive impact of spirulina on growth performance (Mustafa et al., 1994; James et al., 2006: Palmegiano et al., 2008; Ramakrishnan et al., 2008; Tongsiri et al., 2010; Ungsethaphand et al., 2010; Promya and Chitmanat, 2011; Teimouri et al., 2013; Khanzadeh et al., 2016; Cao et al., 2018; Gogoi et al., 2018). However, contradictory results have been reported with no beneficial effects of dietary spirulina on growth of common carp Cyprinus carpio (Nandeesha et al., 1998), Catla Catla catla (Nandeesha et al., 2001) and Nile tilapia Oreochromis niloticus (Lu and Takeuchi, 2002). It is likely that for carnivorous fish, this microalga has had better growth performance and better fish muscle quality and our study is in agreement with this hypothesis. Growth improvement to dietary spirulina may be due to the improved feed intake and nutrient digestibility. Some researchers have been reported that spirulina improved intestinal microbial balance, leading to better growth by improving food absorption, digestive enzymes activities and fats transport system (James et al., 2006; Teimouri et al., 2013). In addition, the bio-compounds in spirulina, delayed absorption of dietary nutrients and improved carbohydrate and protein utilization in Moreover, there are several nutrients especially vitamins, minerals, essential amino acids, fatty acids in spirulina, that may be beneficial to fish growth promotion and might activate metabolism and act growth stimulants (Mustafa et al., 1994). Generally, in the current study growth performance in the 6%FMR 8%FMR treatments were higher than the other dietary groups. Differently, similar spirulina content to the 8%FMR (52.8 g kg⁻¹) has been reported to decrease growth in rainbow trout (Oncorhynchus mykiss) (Teimouri et 2013) and Nile tilapia al., (Olvera-Novoa et al., 1998). There are no reports about anti-nutritional factors in spirulina (Vonshak et al., 2014).

However, the slightly lower growth in 8%FMR compared to 6%FMR is due to the lower mineral content such as phosphorous in spirulina, compared to fish meal (Olvera-Novoa *et al.*, 1998). Perhaps the fish meal content in 6%FMR (620 g kg⁻¹ in diet) is optimum for Caspian brown trout growth and higher or lower contents is not enough to reach the highest growth.

There was a significant difference in haemotology parameters among all tested treatments. Our results revealed that S. platensis has positive impact on the health status of S. trutta caspius. Previous studies such as those by Terry et al. (2000) on tilapia, Abdel Tawwab and Ahmad (2009) on tilapia, Yong-Chin et al. (2010) on vannamei shrimp (Litopenaeus vannamei), Andrews et al. (2011) on Nile tilapia, Promya and Chitmanat (2011) on African catfish, Ibrahem et al. (2012) on Nile tilapia, Ragap et al. (2012) on Nile tilapia, Krishnaveni et al. (2013) on catla, Zamini and Azimi (2015) on Carp koi; Salehi Farsani et al. (2014) on stellate sturgeon confirmed that S. platensis has haemotologic positive effects on parameters. In addition, increasing haemotology indices (e.g., RBC, Hb and Hct) of S. trutta caspius with increasing levels of S. platensis can result from a variety of reasons including iron compounds, vitamins also (B12, A and E), and considerable antioxidant capacity contained in S. platensis. This high antioxidant capacity is due to the considerable amount of pigments, particularly Phycocyanobilin (e.g., Allophycocyanin Phycocyanin, and Phycoerythrin). In fact, these pigments are responsible for the removal of peroxide radicals in the body and reduce the rate of haemolysis of RBC by oxidants as well as the ability to significantly increase iron absorption (Kop and Durmaz, 2008; Abdel Tawwab and Ahmad, 2009; Andrews *et al.*, 2011).

Reactions caused by WBC mechanisms of cellular nonspecific or early defense in fish that occur in response to various conditions such as bacterial, viral, fungal, protozoan and parasitic infections, and it is the first indicator to determine the health of every living being (Andrews et al., 2011). Increasing WBC and their differentiate percent with increasing levels of dietary S. platensis is due to stimulating the immune system via bioactive substances contained in the microalgae (Tort, 2003). Furthermore, phytocyanins contained in S. platensis activate types of **WBC** (e.g., macrophages and granulocytes) and are responsible for repairing damaged tissues by pathogens (Selmi et al., 2011). This bioactive substance with induction of cytokine secretion (molecules involved in the immune system by activating macrophages and lymphocytes.) stimulates the production of new WBC marker molecules (Raa, 1996).

Based on these results, levels of liver enzymes (AST and ALT) decreased with increasing levels of *S. platensis*. Based on accepted hypothesis, the increase of these enzymes in animals (including fish), indicate damage to the liver cells (Jeney and Anderson, 1993).

Numerous studies have confirmed the effectiveness of S. platensis to reduce cholesterol and liver protection (Khan et al., 2005; Pieretti and Meineri, 2011; Kim, 2013; Zeinab et al., 2015). One of the most important approaches of antioxidants is liver cell protection. The antioxidant properties of S. platensis have attracted the attention of many researchers. One ofthe latest investigations Manoj et al. (1992) reported that alcoholic extract of S. platensis accelerates oxidation of fat significantly (65%) in comparison to the antioxidant chemicals such (35%),alpha-tocopherol BHA or Butylated Hydroxy Anisole (45%), and beta-carotene (48%).

In this study, serum total protein, albumin and globulin increased with increasing levels of S. platensis. Several studies have been demonstrated the effects of S. platensis on blood biochemical parameters (Schaperclaus et al., 1992; Dunkan and Klesius, 1996; Abdel-Tawwab and Ahmad, Andrews et al., 2011; Hernandez, 2005; Zeinab et al., 2015). Blood serum protein show changes in health conditions affected by internal and factors. Albumin external transporter protein or public transporter of many organic and inorganic ligands, such as thyroxine, bilirubin, penicillin, cortisol, estrogen, free fatty acids, calcium, and magnesium during illness, malnutrition and stress, blood albumin is reduced (Misra et al., 2006; 2011). Glucose, Alexander et al., cholesterol and triglyceride levels fluctuate during protein catabolism and glycogenesis. Serum glucose levels are often referred to as a non-specific marker of stress (Sheikhzadeh *et al.*, 2012). In this study, blood glucose, triglycerides and cholesterol decreased with increasing levels of *S. platensis*. Our results are in agreement with data reported by other researchers (Abdel-Tawwab and Ahmad, 2009; Andrews *et al.*, 2011; Hernandez, 2005; Salehi Farsani *et al.*, 2014).

Long-chain omega-3 fatty acids in the liver by reducing the secretion of lipoproteins (VLDL) can be reduced triacylglycerol levels (Harris et al., 1984). S. platensis is low fat (5% fat) so, 10 gr. of S. platensis has only 36 gr. calorie and no cholesterol. S. platensis particularly in terms of alpha-linolenic acid (36% of the total PUFA) is very rich. The most essential fatty acids in S. platensis that in other food sources are scarce, is gamma linolenic acid. It has been grown under special lighting conditions in the cells, are also able to enrich and enhance it (Khan et al., 2005). Gamma-linolenic acid is a precursor of prostaglandins (PGE) and has special effect on the control of blood cholesterol levels that leads to rapid cell growth. The role of PGE is blood pressure regulation, cholesterol synthesis control and cell survival (Harris et al., 2002).

In this study, we evaluated the immunity status of *S. trutta caspius* fed with different levels of *S. platensis*, using lysozyme, complement C3 and C4, immunoglobulins, oxygen free radical and haemolytic activity of the complement factor. Based on our results, the activity of lysozyme after 15 and 180 seconds with increasing levels

of S. platensis significantly increased (up to 8%). Effect of S. platensis on lysozyme by various researchers have confirmed (Watanuki et al., 2006; Abdel Tawwab and Ahmad, 2009; Andrews et al., 2011; Promya and Chitmanat, 2011; Ragap et al., 2012; Ibrahem et al., 2012; Krishnaveni et al., 2013; Kim, 2013; Shimaa, 2016; Salehi Farsani et al., 2014). Lysozyme by **WBCs** (monocytes, macrophages, neutrophils) and using circulating the blood in different tissues of the body and leads to non-specific released immune organisms (Saurabh and Sahoo, 2008).

The complement system is a part of the immune system that enhances the ability of antibodies and phagocytic cells to clear microbes and damaged cells from an organism, promotes and inflammation, attacks the pathogen's plasma membrane (Claire et al., 2002). Several factors have an impact on immunity, including divalent magnesium ions (especially and calcium), polysaccharides immunogenic, season, sexual maturity, prostaglandins, thromboxane and immunoglobulins (Tort et al., 2003). The activity of the complement system and immunoglobulins is increased using microalgae (Ragap et al., 2012: Krishnaveni et al., 2013; Shimaa, 2016). C3 and C4 high levels are related to the health of the fish. High serum complement activity in fish fed S. platensis could be due to the presence of pigment carotenoids especially beta-carotene. In addition, lysozyme, antibodies and complement factors due adhesion to and colonization with pathogenic microorganisms, prevent infection and pathogenesis. Free radicals of oxygen (O2, H2O and O2-) together with nitric oxides, lysozyme, cytokines and other messenger molecules, (prostaglandins, leukotrienes and thromboxanes) increased phagocytic and Pynocytic capacity of macrophage cells (Taoka et Measuring any of these al., 2006). factors can be effective in determining the activation of macrophages. Several studies have demonstrated a positive impact of immune-stimulants such as probiotics and micro-algae on amount of oxygen free radicals and phagocytic activity (Dunkan and Klesius, 1996; Rengpipat et al., 2000; Li and Gatlin, 2004; Panigrahi et al., 2005; Taoka et al., 2006; Watanuki et al., 2006; Abdel-Tawwab and Ahmad, 2009; Andrews et al., 2011; Ragap et al., 2012; Kim, 2013; Ibrahem et al., 2012). Oxidative stress is the accumulation of ROS (Reactive Oxygen Species, ROS). It may be an important factor in the process of developing a variety of The immune system is diseases. vulnerable to oxidative damage. S. platensis also contains other pigments Phycocyanobilin including Phycocyanin, Allophycocyanin Phycoerythrin mainly that are responsible for antioxidant activity and the removal of peroxide radicals in the body (Ragap et al., 2012)

The results of different impacts levels of *S.platensis* on immune and haemotology parameters in *S.trutta caspius* show that *S.platensis* may be beneficial in improve growth, immunity levels in Caspian Sea salmon and create

resistance to pathogens. These results indicate that *S. platensis* supplementation is promising for disease prevention in *S. trutta caspius* juveniles and can substitute for up to 8% of the fish meal in the diet, although the optimum performance is obtained with 6% substitution

References

- Abdel-Tawwab, Μ. and Ahmad, 2009. Live M.H., Spirulina (Arthrospira platensis) as a growth and immunity promoter for Nile tilapia, Oreochromis niloticus (L.), challenged pathogenic with Aeromonas hydrophila. Aquaculture Research, 40, 1037-1046.
- Aksnes, A. and Opstvedt, J., 1998. Content of digestible energy in fish feed ingredients determined by the ingredient-substitution method. *Aquaculture*, 161, 45–53.
- Alexander, C., Sahu, N.P., Pal, A.K. and Akhtar, M.S., 2011.

 Haematoimmunological and stress responses of Labeo rohita (Hamilton) fingerlings: Effect of rearing temperature and dietary gelatinized carbohydrate. Journal of Animal physiology and Animal Nutrition, 95, 653-663.
- Andrews, S.R., Sabu, N.P., Pal, A.K., Mukherjee, S.C. and Kumar, S., 2011. Yeast extract, brewer's yeast and Spirulina in diets for *Labeo rohita* fingerlings affect haematoimmunological responses and survival following *Aeromonas hydrophila* challenge. *Research in Veterinary Science*, 91, 103–109.

- AOAC, 1995. Official methods of analysis. 16th ed. Association of Official Analytical Chemists. Gaithersburg, MD, USA. 2200 P.
- Bain, B., Bates, I. and Laffan, M.,2011. Dacie and Lewis PracticalHaematology. 12th Edition,Published Edinburgh ChurchillLivingstone.
- Belay, A., Kato, T. and Ota, Y., 1996. Spirulina (*Arthrospira*): Potential application as an animal feed supplement. *Journal of Applied Phycology*, 8, 303-311.
- **Blaxhall, P.C. and Daisley, K.W.,** 1973. Routine haematological methods for use with fish blood. *Journal of Fish Biology*, 5(6), 771–781.
- Cao, S.P., Zou, T., Zhang, P.Y., Han, D., Jin, J.Y., Liu, H.K., Yang, Y.X., Zhu, X.M. and Xie, S.Q., 2018. Effects of dietary fishmeal replacement with *Spirulina platensis* on the growth, feed utilization, digestion and physiological parameters in juvenile gibel carp (*Carassis auratus gibelio* var. CAS III). *Aquaculture Research*, 49, 1320-1328.
- Ceballos, J.B., Villareal, H., Garcia, T., Perez-Jar, L. and Alfonso, E., 2006. Effect of Spirulina platensis meal as feed additive on growth, survival and development in Litopenaeus schmitti shrimp larvae. Revista de Investigaciones Marinas, 26(3), 235-241.
- Claire, H., Holland, M. and Lambris, J.D., 2002. The complement system in teleosts. *Fish and Shellfish Immunology*, 12, 399–420.

- **Coad, B.W., 2000**. Criteria for assessing the conservation status of taxa (as applied to Iranian freshwater fishes). *Biologia*, 55,539-557.
- **Doumas, B.T. and Peters, J.r.T.,** 1997. Serum and urine albumin: a progress report on their measurement and clinical significance. *Clinica Chimica Acta*, 258(1), 3-20.
- Dunkan, P.L. and Klesius, P.H., 1996. Effects of feeding spirulina on specific and non-specific immune responses of channel catfish. *Journal of Aquaculture Animal Health*, 8, 308-313.
- **Dytham, C., 1999**. Choosing and using statistics: A biologist's guide. *Journal of Ecology*, 87(4), 734–735.
- **Ellis, A.E., 1999.** Lysozyme assays. Techniques in Fish Immunology. pp. 101-103.
- Esmaeili, M., Abedian Kenari, A. and Rombenso, A.N., Immunohematological status under acute ammonia stress of juvenile rainbow trout (Oncorhynchus mykiss Walbaum, 1792) fed garlic (Allium sativum) powder-supplemented meat meal-based and bone feeds. Clinical Comparative and Pathology, DOI: 10.1007/s00580-017-2457-8.
- **Estrada, J.P., Bescós, P.B. and Del Fresno, A.V.** 2001. Antioxidant activity of different fractions of *Spirulina platensis* protean extract. *Il farmaco*, 56, 497-500.
- Gogoi, S., Mandal, S. and Patel, A., 2018. Effect of dietary Wolffia arrhiza and Spirulina platensis on growth performance and pigmentation of Queen loach Botia

- dario (Hamilton, 1822). Aquaculture Nutrition, 24, 285-291.
- Goytortua-Bores, E., Civera-Cerecedo, R., Rocha-Meza, S. and A., Green-Yee, 2006. **Partial** of replacement red crab (Pleuroncodes planipes) meal for fish meal in practical diets for the white shrimp Litopenaeus vannamei, Effects on growth and in vivo digestibility. Aquaculture, 256, 414-422.
- Habibi, E., Kalbassi, M., Hosseini, S. and Qasemi, S., 2013. Feasibility of Identification of Fall and Spring Migrating Caspian trout (Salmo trutta caspius) by Using AFLP Molecular Marker. Turkish Journal of Fisheries and Aquatic Sciences, 13, 241-248
- Harris, W.S., Connor, W.E., Inkeles, S.B. and Illinworth, D.R., 1984. Dietary omega-3 fatty acids prevent carbohydrate-induced hypertriglyceridemia. Metabolism 33(11), 1016-1019.
- Harris, S.G., Josue, P., Koumas, L., Denise, R. and Phipps, R.P., 2002. Prostaglandins as modulators of immunity. *TRENDS in Immunology*, 23, 3.
- Henrikson, R., 1998. Earth food Spirulina. California/USA. Ronore Enterprises. 180 P.
- Hernandez, S.P., 2005. Responsible use of antibiotics in aquaculture. Food and agriculture organization of the united nations, FAO Fisheries Technical. 469 P.
- Hirahashi, T., Matsumoto, M., Hazeki, K., Saeki, Y., Ui, M. and Seya, T., 2002. Activation of the

- human innate immune system by Spirulina: augmentation of interferon production and NK cytotoxicity by oral administration of hot water extract of *Spirulina plantesis*. *International Immunopharmacology*, 2, 423–434.
- Hironobu, W., Kazuki, O., Asmi, C., Tassakka, T. and Masahiro, S., 2006. Immunostimulant effects of dietary *Spirulina platensis* on carp, *Cyprinus carpio*. *Aquaculture*, 258, 157 163.
- Houston, A.H., 1990. Components of the haemotological response of fishes to environmental temperature change: A review. In "Environmental Physiology of Fishes" (Ali, M. A., ed.) Plenum, New York. pp. 241-298.
- Ibrahem, M.D., Mohamed, F.M. and Marwa, A.I., 2012. The role of Spirulina platensis (Arthrospira platensis) in Growth, Immunity of Nile tilapia (Oreochromis niloticus) and its resistance to infection. Journal of Agricultural Science, 5(6), 5. DOI: 10.5539/jas.v5n6p109.
- James, R., Sampath, K., Thangarathinam, R. and Vasudevan, I., 2006. Effect of dietary Spirulina level on growth, fertility, coloration and leucocyte count in red sword tail, xiphophorus helleri. The Israeli Journal of Aquaculture–Bamidgeh, 58(2), 97-104.
- Jeney, G. and Anderson, D.P., 1993.

 Enhanced immune response and protection in rainbow trout to *Aeromonas salmonicida* bacterin following prior immersion in

- immunostimulants. *Fish* and *Shellfish Immunology*, 3(1), 51-58.
- Kalbassi, M., Dorafshan, Tavakolian, T., Khazab, M. and Abdolhay, H., 2006. Karyological analysis of endangered Caspian salmon, Salmo trutta caspius (Kessler. 1877). Aquaculture Research, 37(13), 1341–1347.
- Khan, Z., Bhadouria, P. and Bisen, P.S., 2005. Nutritional and therapeutic potential of Spirulina. *Current Pharmaceutical Biotechnology*, 6, 373-379.
- Khanzadeh, M., Fereidouni, A.E. and Berenjestanaki, S.S., 2016. Effects of partial replacement of fish meal with *Spirulina platensis* meal in practical diets on growth, survival, body composition, and reproductive performance of three-spot gourami (*Trichopodus trichopterus*). *Aquaculture International*, 24, 69-84.
- Kiabi, B.H., Abdoli, A. and Naderi, M., 1999. Status of the fish fauna in the South Caspian basin of Iran. Zoology in the Middle East, 18, 57-65.
- **Kim, D.D., 2013.** Outdoor mass culture of *Spirulina platensis* in Vietnam. *Journal of Applied Phycology*, 2, 179–181.
- Kop, A. and Durmaz, Y., 2008. The effect of synthetic and natural pigments on the colour of the cichlids (*Cichlasom aseverum* sp., Heckel 1840). *Aquaculture International*, 16, 117-122.
- Kozlenko, R.D.P.M., Ph.D, M.P.H. and Henson, R.H., 1998. Latest scientific research on spirulina:

- Effects on the AIDS virus, cancer and the immune system. Dana-Farber Cancer Institute and Harvard Medical School, Boston. inspiredliving.com
- Krishnaveni, R., Palanivelu, K. and Effects of S., 2013. Velavan, probiotics and **Spirulina** supplementation on haemato-Immunological function of Catla International Journal of catla. Research in **Fisheries** and Aquaculture, 3(4), 176-181.
- Kumar, S., Sahu, N.P., Pal, A.K., Dharitri, C., Sona, Y. Mukherjee., 2005. Effect of dietary carbohydrate haematology, on burst activity respiratory and histological changes in L. rohita juveniles. Fish and Shellfish Immunology, 19, 331-344.
- Li, P. and Gatlin, D.M., 2004. Dietary brewer yeast and prebiotic Grobiotic -A influence growth performance, immune responces and resistances of hybrid striped bass to *Streotococcus iniae* infection. *Aquaculture*, 231, 445-456
- Lu, J. and Takeuchi, T., 2002. Taste of tilapia, *Oreochromis niloticus*, fed solely on raw spirulina. *Fisheries Science*, 68, 987–988.
- Manoj, G., Venkataraman, L.V. and Srinivas, L., 1992. Antioxidant properties of Spirulina (*Spirulina platensis*). In: Seshadri C.V., Bai N.J.: Spirulina: National Symposium (India), MCRC, Tharamani, Madras. pp. 48–154.
- Misra, C.K., Das, B.K., Mukherje, S.C. and Pattnaik, P., 2006. Effect of multiple injections of betaglucan

- on non-specific immune response and disease resistance in *Labeo* rohita fingerlings. Fish and Shellfish Immunology, 20, 305-319.
- Mustafa, M.G., Umino, T. and Nakagawa, H., 1994. The effect of Spirulina feeding on muscle protein deposition in red sea bream, *Pagrus major*. *Journal of Applied Ichthyology*, 10,141–145.
- Mustafa, M.G., Takeda, T., Umino, T., Wakamatsu, S. and Nakagawa, H., 1994. Effects of *Ascophyllum* and *Spirulina* meal as feed additives on growth performance and feed utilization of red sea bream, Pagrus major. J. Fac. Appl. Biol. Sci., Hiroshima Univ, 33, 125-132.
- Mustafa, M.G. and Nakagawa, H., 1995. A review: Dietary benefits of algae as an additive in fish feed. *The Israeli Journal of Aquaculture, Bamidgeh*, 47,155-162.
- Nakagawa, H., Sato, M. and Gatlin, D.M., 2007. Dietary supplements for the health and quality of cultured fish. CABI Publishing, Oxfordshire, UK. 256 P.
- Nandeesha, M.C., Gangadhara, B., Varghese, T.J. and Keshavanath, P., 1998. Effect of feeding *Spirulina platensis* on the growth, proximate composition and organoleptic quality of common carp, *Cyprinus carpio* L. *Aquaculture Research*, 2, 305–312.
- Nandeesha, M.C., Gangadhara, B., Manissery, J.K. and Venkataraman, L.V., 2001. Growth performance of two Indian major carps, catla (*Catla catla*) and rohu (*Labeo rohita*) fed diets containing different levels of *Spirulina*

- platensis. Bioresource Technology, 80,117–120.
- NRC (National Research Council)., 1993. Nutrient requirements of warmwater fishes and shellfishes, revised edn. National Academy Press, Washington. 225 P.
- Olvera-Novoa, M.A., Dominguez-Cen, L.J., Olivera-Castillo, L. and Martinez-Palacios, C.A., 1998. Effect of the use of the microalgae Spirulina maxima as fish meal replacement in diets for tilapia, Oreochromis mossambicus (Peters), fry. Aquaculture Research, 29, 709-715.
- Palmegiano, G.B., Gai, F., Dapra, F., Gasco, L., Pazzaglia, M. and Peiretti, P.G., 2008. Effects of Spirulina and plant oil on the growth and lipid traits of white sturgeon (Acipenser transmontanus) fingerlings. Aquaculture Research, 39, 587–595.
- Panigrahi, A., Kiron, V., Puangkaew, J., Kobayashi, T., Satoh, S. and Sugita, H., 2005. The viability of probiotic bacteria as a factor influencing the immune response in rainbow trout (*Oncorhynchus mykiss*). Aquaculture, 243, 241-254.
- Pieretti, P.G. and Meineri, G. 2011. Effects of diets with increasing levels of Spirulina platensis on the carcass characteristics, meat quality and fatty acid composition of growing rabbits. Livest. Sci, 140:218-224.
- Promya, J. and Chitmanat, C., 2011.

 The effects of *Spirulina platensis* and *Cladophora* algae on the growth performance, meat quality and

- immunity stimulating capacity of the African Sharptooth Catfish (*Clarias gariepinus*). *International Journal of Agriculture and Biology*, 13, 77-82.
- Quillet, E., Faure, A., Chevassus, B., Kreig, F., Harache, Y., Arzel, J., Metailler, R. and Boeuf, G., 1992. The potential of brown trout (Salmo trutta L.) for mariculture in temperate waters. *Icelandic Agricultural Sciences*, 6, 63–76.
- Raa, J., 1996. The use of immune stimulatory substances in Fish and Shellfish farming. *Reviews in Fisheries Science*, 1(4), 229–288.
- Ragap, H.M., Khalil, R.H. and Mutawie, H.H., 2012. Immunostimulant effects of dietary Spirulina platensis on tilapia Oreochromis niloticus. Journal of Applied Pharmaceutical Science, 2(2), 26-31.
- Rajabi Islami, H., Arab, N., Assare, R., Rastravan, M.I. and Ebtekari, R., 2016. Blood parameters of Caspian brown trout (*Salmo trutta caspius*) fingerlings affected by dietary L-ascorbyl-2-polyphosphate. *Iranian Journal of Fisheries Sciences*, 15(3), 1167-1186.
- **Regunathan, C. and Wesley, S.G., 2006.** Pigment deficiency correction in shrimp broodstock using Spirulina as a carotenoid source. *Aquaculture Nutrition*, 12, 425–432.
- Ramakrishnan, C.M., Haniffa, M.A., Manohar, M., Dhanaraj, M., Arockiaraj, A.J., Seetharaman, S. and Arunsingh, S.V., 2008. Effects of probiotics and Spirulina on survival and growth of juvenile common carp *Cyprinus carpio*.

- *Israel Journal of Aquacultures Bamidgeh*, 60(2), 128–133.
- Rengpipat, S., Rukpratanporn, S., Piyatiratitivorakul, S. and Piamsak, M., 2000. Immunity enhancement in black tiger shrimp, *Penaeus monodon* by a probiont bacterium. *Aquaculture*, 191, 271–288.
- Ravi, M., Lata, De.S., Azharuddin, S. and Solomon, P., 2010. The beneficial effects of Spirulina focusing on its immunomodulatory and antioxidant properties. *Journal of Dietary Supplements*, 2, 73–83.
- Rook, G.A.W., Steele, J., Umar, S. and Dockrell, H.M., 1985. A simple method for the solubilisation of reduced NBT, and its use as a calorimetric assay for activation of human macrophages by y-interferon. *Journal of Immunological Methods*, 82, 161.
- Saurabh, S. and Sahoo, P.K., 2008.

 Lysozyme: an important defence molecule of fish innate immune system. *Aquaculture Research*, 39(3), 223–239
- **Sakai, M., 1999.** Current research status of fish immunostimulant. *Aquaculture*, 172, 63-92.
- Salehi-Farsani, A., Soltani, M., Kamali, A. and Shamsaie, M., **2014.** Effect of immune motivator Macrogard and Spirulina platensis carcass on some growth, biochemical indices of stellate stellatus. sturgeon Acipenser Aquaculture, Aquarium, Legislation Conservation and International Journal of the Bioflux Society, 7, 3.

- **Sedgwick, S.D., 1995.** Trout farming handbook, 5th edn. Fishing News Books, Alden Press, Oxford. 169 P.
- Selmi, C., Leung, P.S.C., Fischer, L., German, B., Chen-Yen, Y., Kenny, T.P., Cysewski, G.R. and Gershwin, M.E., 2011. The effects of Spirulina on anemia and immune function in senior citizens. *Cellular and Molecular Immunology*, 8(3), 248–254.
- Sheikhzadeh, N., Heidarieh, Karimi Pashaki, A., Nofouzi, K., AhrabFarshbaf, M. and Akbari, M., 2012. Hilyses® fermented Saccharomyces cerevisiae, enhances the growth performance and skin nonspecific immune parameters in rainbow trout (Oncorhynchus mykiss). Fish and Shellfish Immunology, 32, 1083-1087.
- Shimaa, A.A., 2016. Effect of platensis Spirulina feed supplement on growth performance, immune response and antioxidant status of mono-sex Nile Tilapia (Oreochromis niloticus). Benha *Veterinary Medical Journal*, 30(1), 1-10.
- Siwicki, A. K., Anderson, D. P. 1993:
 Immunostimulation in Fish:
 Measuring the Effects of Stimulants
 by Serological and Immunological
 Methods. U.S. Fish and Wildlife
 Service, IFI, Poland. 1, 24.
- Schaperclaus, W., Kulow, H. and Schreckenbach, K., 1992. Fish disease. A.A. Balkema, Rotterdam, the Netherlands. pp. 101-105.
- **Shahbazi, S. and Bolhassani, A., 2016.** Immunostimulants: Types and functions. *Journal of Medical*

- *Microbiology and Infectious Diseases*, 4(**3-4**), 45-51.
- Sotoudeh, E., Abedian Kenari, A., Khodabandeh, S. and Khajeh, K., 2015. Combination effects of dietary EPA and DHA plus alpha-tocopherol: Effects on performance and physiological status of Caspian brown trout (*Salmo trutta caspius*) fry. *Aquaculture Nutrition*, 22(5), 1101-1115.
- Subhadra, B., Lochmann, R., Rawles, S. and Chen, R., 2006. Effect of dietary lipid source on the growth, tissue composition and hematological parameters of largemouth bass (*Micropterus salmoides*). Aquaculture, 255(1), 210-220.
- Tang, H.G., Wu, T.X., Zhao, Z.Y. and Pan, X.D., 2008. Effects of fish protein hydrolysate on growth performance and humoral immune response in large yellow croaker (Pseudosciaena crocea R.). Journal of Zhejiang University Science, 9, 684-690
- Taoka, Y., Hiroto, M., Jae-Yoon, J.O., Min-Jee, J., Sungchul, C.BAI., Won-Jae, L., Kazuya, Y. and Shunsuke, K., 2006. Growth, stress tolerance and non-specific immune response of Japanese flounder *Paralichthys olivaceus* to probiotics in a closed recirculating system. *Fisheries Science*, 72, 2.
- **Teimouri, M., Amirkolaie, A.K. and Yeganeh, S., 2013.** Effect of Spirulina platensis meal as a feed supplement on growth performance and pigmentation of Rainbow Trout (Oncorhynchus mykiss). World

- Journal of Fish and Marine Sciences, 5(2), 194–202.
- **Tietz, N.W., 1986**. Textbook of clinical chemistry, W.B. Saunders, Philadelphia, PA. 1919 P.
- Terry, C.H., Cardinale, J.L. and Smith, S.A., 2000. Haemotology and plasma chemistry reference intervals for cultured tilapia (*Orechromis hybrid*). *Veterinary Clinical Pathology*, 29, 7–12.
- Tongsiri, S., Mang-Amphan, K. and eerapornpisal, Y., 2010. Effect of replacing fishmeal with Spirulina on growth, carcass composition and pigment of the mekong giant catfish. *Asian Journal of Science*, 2, 106–110.
- **Tort, L., Balasch, J.C. and Mackenzie, S., 2003.** Fish immune system. A crossroads between innate and adaptive response. *Inmunologia*, 22(3), 277-286.
- Tukmechi, A., Rahmati Andani, H.R.. Manaffar. R. and Sheikhzadeh, N., **2011.** Dietary administration of beta-mercapto-Saccharomyces ethanol treated cerevisiae enhanced the growth, innate immune response and disease resistance of the rainbow trout, Oncorhynchus mykiss. Fish Shellfish Immunology, 30, 923–928.
- Ungsethaphand, T., Peerapornpisal, Y., Whangchai, N. and Sardsud, U., 2010. Effect of feeding *Spirulina platensis* on growth and carcass composition of hybrid red tilapia (*Oreochromis mossambicus* × O. niloticus). Maejo International Journal of Science and Technology, 4(2), 331–336.

- Vonshak, A, Laorawat, S., Bunnag, B, and Tanticharoen, M. 2014. The effect of light availability on the photosynthetic activity and productivity of outdoor cultures of *Arthrospira platensis* (Spirulina). Journal of Applied Phycology. 26: 1309-1315.
- Watanuki, H., Ota, K., Malina, A.S., Kato, T. and Sakai, M., 2006. Immunostimulant effects of dietary *Spirulina platensis* on carp, *Cyprinus carpio*. *Aquaculture*, 258, 157-163.
- Yong-Chin, L., Carina Miranda, T., Chien-Lun, H., Wen-Ching, T. and Jiann-Chu, C., 2010. White shrimp Litopenaeus vannamei that had received the hot-water extract of Spirulina platensis showed earlier recovery in immunity and upregulation of gene expressions after pH stress. Fish and Shellfish Immunology, 29, 1092-1098.
- Zamini, A. and Azimi, A., 2015. The effect of *Spirulina platesis* algae in nutrition on blood and immunity parameters of koi (cyprinus carpio). Academie Royale DesScience D Outre-Mer Bulletin DesSeances, 4(3), 147-151.
- Zeinab, A.K., Aly, M.S., Faiza, A. and Fatma, E.M., 2015. Effect of Spirulina platensis and Lactobacillus rhamnosus on growth and biochemical performance of Nile Tilapia (Oreochromis niloticus) fingerlings. International Journal of Current Microbiology and Applied Sciences, 4(4), 747-763