[DOR: 20.1001.1.15622916.2020.19.3.31.7]

Morphometric and molecular characterization of Dactylogyrus lamellatus isolated from farmed Grass carp, Ctenopharyngodon idella (Valenciennes, 1844), in Guilan Province, Iran

Daghigh Roohi J.¹; Dalimi A.^{2*}; Pourkazemi M.³; Ghasemi M.¹; Shamsi S.⁴

Received: July 2018 Accepted: April 2019

Abstract

Grass carp is an herbivorous species, which is actually popular in Iran. Monogenean parasites, particularly members of the family Dactylogyridae, have been one of the main causes of mortalities in Iranian fish farms and therefore they have been subject of many studies in Iran. The main aim of the present study was to describe morphological and molecular characteristics of monogenean parasites recovered from the farmed grass carp (*Ctenopharyngodon idella*) in the Guilan Province during 2015 to 2016. A total of 80 grass carps have been examined for infestation with monogenean parasites. 10923 *Dactylogyrus* species were recovered from 95% of the farmed grass carp. The parasite was first classified based on their morphological characteristics and identified as *Dactylogyrus lamellatus*. The parasite was then subjected to PCR and sequencing of the 28S rDNA gene. The phylogenetic tree showed that *D. lamellatus* genetically was similar to those previously reported from other countries and Mashhad farms in Iran.

Keywords: Grass carp, *Dactylogyrus lamellatus*, 28S rDNA, Guilan, Iran.

¹⁻Inland water Aquaculture Research Center, Iranian Fisheries Sciences Research Institute, Agricultural Research Education and Extension Organization (AREEO), Bandar Anzali, Iran

²⁻Department of Parasitology, Faculty of Medical Sciences, Tarbiat Modares University, P.O.Box: 14115-331, Tehran, Iran

³⁻Iranian fisheries Science Research Institute, Agricultural Research Education and Extension Organization (AREEO), Tehran, Iran

⁴⁻School of Animal and Veterinary Sciences; Graham Centre for Agricultural Innovations, Charles Sturt University, NSW 2650, Australia

^{*}Corresponding author's Email: dalimi_a@modares.ac.ir

Introduction

Grass carp is a large member of the carp and minnow family (Cyprinidae) and is native to south eastern Russia and north western China. This herbivorous been species has deliberately introduced into many countries for vegetation control (Cudmore and purposes Mandrak. 2004) and also for aquaculture due to rapid growth rate (Zolfinejad et al., 2017). In Iran, this species was intentionally released for vegetation control in Anzali lagoon and introduced to the Guilan province for polyculture fish farming about 40 years ago. Today, this fish is artificially propagated in fish hatcheries of Guilan province and was sent to the whole of the country for cultivation.

Class Monogenea is a group of parasitic worms commonly found in fishes. These parasites feed of mucus and epithelial cells of the skin and gills and sometimes of the blood (Chaudhary al., 2013). Infestation with etmonogenean parasites may lead to serious hyperplasia of the gill filaments epithelium. impairs respiratory function, negatively affects growth, and even associates with high mortalities, especially in small carps (Lu et al., 2012; Jiang et al., 2013; Tu et al., 2015). Dactylogyrus represents most dominant genus among the Monogenea. These are common ectoparasites usually attached to the gills of freshwater fish of the family Cyprinidae (Gibson et al., 1996; Woo 2006). The fish heavily infected with Dactylogyrus spp. is susceptible to secondary infestations that can play a significant role in the pathogenicity of these parasites, which result in considerable economic losses (Dove and Ernst, 1998; Woo *et al.*, 2002; Reed *et al.*, 2009; Tu *et al.*, 2015).

Thus far, grass carp is known to be parasitized by at least two species of Dactylogyrus (Shamsi et al., 2009; Daghigh Roohi et al., 2016) in both pond and natural lake habitats in Iran. Identification of the Monogenea in Iran often been based morphological basis. Monogenean identification solely based on morphological features can be problematic and ambiguous for closely related species. For effective diagnosis, treatment and control of parasitic diseases, the accurate identification of parasites is necessary (Mcmanus and Bowles, 1996). Recently, molecular provided markers have useful alternative to the identification and validation of monogenean (Cunningham et al., 1995; Tu et al., 2015). Our knowledge on genetic character of Iranian monogenea is poor and little is known about molecular of aspects these parasites (Mozhdeganlou et al., 2011; Omidzahir et al., 2012; Mousavi et al., 2013; Omidzahir et al., 2015). In the present study, Dactylogyrids that recovered from cultured grass carp in the Caspian described region are morphologically, then their genetic characterization was done based on sequences of 28SrDNA gene.

Materials and methods

Fish collection

This work was carried out from April 2015 to November 2016 in Guilan Province, North of Iran (Fig. 1). A total of 80 grass carp (Ctenopharyngodon idella) were collected from 10 fish farms. Fish were one summer old with the average total length of 11.38±2.84 an average weight cm and 15.89±12.69 g. The fish were placed in plastic tanks with water obtained from the collection sites and transferred to the Parasitology Laboratory of National Inland Water Aquaculture Research Center in Anzali where they were randomly distributed into several 100-L aquariums containing water from the fish farm and investigated within 72 hours. Fish were examined for the presence of Dactylogyrids parasites under a dissection microscope (4 to 40 magnifications). Live parasites were picked up from the gills scraping by using small needles. Then mounted in glycerine-jelly for morphological investigations (Gussev, 1983).



Figure 1: Distribution of studied ponds in Guilan Province, Iran.

Parasites collection and morphological identification

Monogenean parasites were recovered from gill filaments of Ctenopharyngodon idella. **Parasite** identification was performed based on the character of copulatory organ, the haptoral sclerites like anchors, bars and hooks. Photographs of mounted captured parasites were at magnifications of 20X and 40X using a Nikon **Eclipse** 50i compound microscope with Nikon Digital Sight DS-L1 image analysis software and Nikon Digital Sight DS-SM camera. For morphometric analysis, a total of 7 point to point measurements were made on Dactylogyrus specimen photographs, using Image J software. Drawings were made with the aid of a drawing tube Nikon Y-IDT Japan. Finally, drawing of specimens and their measurement data were compared by parasites identification (Bykhovskayakeys Pavlovskaya et al., 1962; Gussev, 1985). After species identification, the same parasite specimen was stored in %75 ethanol tube for molecular analysis.

Molecular Studies

DNA extraction and PCR

The genomic DNA was extracted from one specimen using the YTA Genomic DNA Extraction Mini Kit (Yekta Tajhiz according Azma, Iran) to the manufacturer's manual. The 28S rDNA region of specimens was amplified using primers (forward. 5'TCTAGTAACGGCGAGTGAACG-3') (Chiary et al., 2014) and the modified (5'reverse primer

GTGGGAAGGTCTACCTCAGC-3').

Each amplification reaction performed in a final volume of 25µl containing 12.5 µl 2× Taq Mastermix, 1 ul of each primer (Macrogen, South Korea), and 2 µl of genomic template DNA. The amplification was carried out in a thermocycler (BIO-RAD, USA) under the following conditions: after an initial denaturation at 94 °C for 3 min, 35 cycles at 94 °C for 30 (denaturation), 59 °C for 30 sec (annealing), 72 °C for 1 min (extension) with a final extention at 72 °C for 10 min. The PCR products were analysed on 1.5% agarose gel and visualized under a UV illuminator, and the results were recorded.

DNA Sequencing

Purified fragments of PCR products were sequenced from both forward and reverse sites of each PCR product. Sequencing was carried out using the same primers as used for PCR amplification (Macrogen, South Korea).

Phylogenetic analysis

28SrDNA sequences of the *Dactylogyrus lamellatus* specimens in the present study were aligned separately using the Clustal W software and then manually adjusted to perform the phylogenetic analysis. Gaps and ambiguously aligned regions were removed. The phylogenetic tree was built using Mega 6.0 by the UPGMA method.

Statistical calculation and analysis

In this study, all calculations were performed using the following formulas (Bush *et al.*, 1997):

Prevalence=Number of infected hosts /Total of hosts×100

Mean intensity=Total number of parasites / Number of infected hosts

Mean abundance=Total number of a particular parasite/Total number of infected and uninfected hosts

Dominance=Total number of a particular parasite/Total number of parasites × 100

The mean intensity of infestation and abundances of Monogenean among different seasons was tested by the Kruskal-Wallis test (KW, multiple comparisons) and Mann-Whitney test. The results were considered significant at the 95% level (p<0.05). Computations were performed using the SPSS.15 programme.

Results

Totally, 10923 *Dactylogyrus* parasites were collected from 80 pieces of grass carp. Morphological examination of taxonomically important features such as the shape of the anchors, connective bar and the shape of a copulatory organ (Fig. 2) as well as detailed morphometric (Table 1) revealed that the only one species is identified as *D. lamellatus* in the present study.

The selected rDNA region was successfully amplified and sequenced. The sequence was deposited in GenBank with accession number MG657261. The 28SrDNA sequence size for *D. lamellatus* was 624 base pairs (bp) (Fig. 3). To build a

phylogenetic tree we used of different 28SrDNA sequences, which are brought in Table 2.

Table 1: Morphometrics (µm) of Dactylogyrus lamellatus recovered from Ctenopharyngodon idella.

Species	Dactylogyrus lamellatus		
Characters	(n= Present Study	10) Bykhovskaya Pavlovskaya	
		(1962)	
Body length	320.55 ± 80.92	Up to 480	
	(221.48-444.63)		
Body width	71.57 ± 21.23	Up to 110	
	(44.58-94.76)		
Length of marginal hooks	27.12±3.16	21-31	
	(20.62-32.47)		
Length of median hooks	35.59 ± 2.02	35-41	
	(32.08-37.64)		
Connecting bar	26.95±1.91	28-30	
Length	(24.04-29.11)		
Connecting bar	3.07 ± 0.42	Up to 4	
Width	(2.57-3.82)	_	
The total length of the	42.68±3.77	About 50	
copulatory organ	(35.78-46.86)		

n = number of specimens measured and range in size given in brackets.




Figure 2: Drawing of *Dactylogyrus lamellatus*: the opisthohapthoral central hook complex (A); copulatory organ (B).




Figure 3: Agarose gel electrophoresis of PCR amplification products from Dactylogyrus lamellatus in the present study following exposure to UV light.

Table 2: List of *Dactylogyrus* species used in this study with their host species, Genbank accession numbers and locality to build a phylogenetic tree.

Taxa	Host name	GenBank Accession number	Locality of collection
Dactylogyrus extensus	Cyprinus carpio	AJ969944	Zcech Republic
Dactylogyrus nminutus	Cyprinus carpio	MF926269	Guilan, Iran
Dactylogyruns nachmerowi	Cyprinus carpio	MF979966	Guilan, Iran
Dactylogyrus vastator	Cyprinus carpio	MF928712	Guilan, Iran
Dactylogyrus aristichthys	Aristichthys nobilis	MH023397	Guilan, Iran
Dactylogyrus nobilis	Aristichthys nobilis	MH023399	Guilan, Iran
Dactylogyrus hypophthalmichthys	Aristichthys molitrix	EF100532	China
Dactylogyrus suchengtaii	Aristichthys molitrix	MG825765	Guilan, Iran

Table 2 continued:			
Dactylogyrus anchoratus	Cyprinus carpio	JX524546	Mashhad, Iran
Dactylogyrus inexpectatus	Carassius auratus	AJ969945	Zcech Republic
Dactylogyrus quanfami	Cirrhinus moliorella	EF100536	China
Dactylogyrus labei	Catla catla	JX566720	India
Dactylogyrus sphyrna	Rutilus rutilus	AJ969943	Zcech Republic
Dactylogyrus cryptomeres	Gobio gobio	AJ969947	Zcech Republic
Dactylogyrus lamellatus	Ctenopharyngodon idella	AY307019	China
Dactylogyrus lamellatus	Ctenopharyngodon idella	MG657261	Guilan, Iran •
Dactylogyrus lamellatus	Ctenopharyngodon idella	KY039154	Mashhad, Iran
Dactylogyrus lamellatus	Ctenopharyngodon idella	EF100533	China
Dactylogyrus lamellatus	Ctenopharyngodon idella	AG969948	Zcech Republic
Dactylogyrus pekinensis	Megalobrama amblycephala	EF100535	China
Dactylogyrus petruschewskyi	Megalobrama amblycephala	AY548927	China
Tetraonchus monenteron	Esox lucius	AJ969953	Zcech Republic

(• This study)

In the present study, 95% of the fish were found to be infected with D. lamellatus. The highest rate of infestation has been observed with an average of 505 parasites per fish in summer. In Table 3 the prevalence, mean intensity, range and total of D. lamellatus in grass carp in different seasons is compared (Mann-Whitney test, p<0.05). As shown in Table 3 this parasite was present during all seasons

and cultivation period, however, the mean intensity was significantly higher in summer than other seasons (p<0.05). The low intensity of this parasite was observed in autumn.

Phylogenetic trees showed 100% similarity between our isolates and other *Dactylogyrus lamellatus* specimens registered in GenBank (Fig. 4).

Table 3: The prevalence, mean intensity, range and total number of *Dactylogyrus lamellatus* in *Ctenopharingodon idella*, in the present study and different seasons.

Parameters Season	Prevalence (%)	Mean±SE	Range	Total
Spring N=47	100	98.17±66.04°	3-836	4614
Summer N=12	100	505±146.51°	61-1527	6060
Autumn N=21	80.95	14.64±5.92 ^b	1-101	249

N: number of examined fish, SE: Standard Error, * Data with different letters in a column are statistically significant at the p<0.05 level




Figure 4: Phylogenetic tree constructed by UPGMA analysis based on 28S rDNA for selected species of *Dactylogyrus*, with *Tetraonchus monenteron* used as the outgroup.

Discussion

Result indicated that only one species of monogenean, *D. lamellatus*, was recovered from *Ctenopharyngodon idella* in the Guilan province fish farms. The parasite identification was performed by morphometric and molecular analysis.

Grass carp is readily infested by several monogeneans, among them D. lamellatus is the most harmful parasite, which is spread throughout the country and causes high mortality in fingerlings (Jalali and Molnar, 1990). This survey demonstrated, D. ctenopharyngodonis as another *Dactylogyrus* on grass carp which introduced from China in 1991 and has been reported from some of the water resources in the country (Shamsi et al., 2009; Daghigh Roohi et al., 2016) has not been spread out in fish ponds yet. The study of grass carp parasites in Zarrineh rud River showed that 11.87% of the fish were infested with D. lamellatus, whereas in this study, the rate of fish infestation was determined about 95% due to the high density of host fish in these ponds. As mentioned in table 3 the mean intensity of this parasite in Guilan fish ponds was also higher than natural waters like Anzali wetland in Guilan province or Khandaghloo Dam Lake in Zanjan province (Pazooki et al., 2006; Dghigh Roohi et al., 2016). Phylogenetic tree showed that D. lamellatus is genetically identical to those previously reported from other countries and also other parts of the country like Mashhad region which reported by Ahmadi et al. (2017). On the other hand, phylogenetic tree shows that lamellatus is closely related to parasites of other Chinese carp like big head and silver carp. Because of grass carp is an alien fish, which introduced to Iran for cultivation so, we can conclude its private parasite, D. lamellatus probably has introduced to Iran due to lack of proper quarantine for their host. as suggested by authors a widely distributed species has

the ability to adapt itself to a broad range of climates.

As shown in Table 3, this parasite infestation was significantly higher in summer than other seasons. It seems that the parasite intensity significantly depends on the water temperature. According to the previous studies, the time of D. lamellatus sexual maturation depends on the temperature of water. It is indicated that the egg production in D. lamellatus began 8, 6 and 4 days after infection at temperatures of 17-19 $^{\circ}\mathrm{C}$ and 22-26 °C , 20-24 respectively (Molnar, 1971).

Therefore, due to the temperature conditions in Guilan province, the different seasonal infection intensity, which obtained in our study, is justifiable.

This study provides further insights into the biodiversity and taxonomic status of monogenean parasites in Iran. Iran is a country with various zoogeographical regions, resulting in a high diversity of monogenean parasites reported from the country. Therefore, there is a huge potential for further work on the genetic character of monogenean fauna in the region and to understand the taxonomic relationship between Iranian monogenea with those from the rest of the world.

Acknowledgements

The present work is part of PhD. thesis, supported financially by Iranian fisheries Science Research Institute, Agricultural Research Education and Extension Organization (AREEO), Tehran, Iran. The author wishes to thank all the personnel of the

Parasitology Department of Medical Sciences Faculty of Tarbiat Modares University for their kind assistance.

Conflict of interest statement

We declare that we have no conflict of interest.

Reference

Ahmadi, A., Borji, H., Naghibi, A., Nasiri, M.R. and Sharifiyazdi, H., 2017. Morphologic and molecular (28SrDNA) characterization of **Dactylogyrus** spp. in Cyprinus carpio and Ctenopharyngodon idella in Mashhad, Iran. Canadian Journal of Comparative *Medicine*, 81(4), 280-284.

Bush, A.O., Lafferty, K.D., Lotz, J.M. and Shostak, A.W., 1997.

Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *Journal of Parasitology*, 83(4), 575-583.

Bykhovskaya-Pavlovskaya, I.E., Gussev, A.V., Dubinina, M.N., Izyumova, N.A., Smirnova, T.S., Sokolovskaya, A.L., Schtein, G.A., Shulman, S.S. and Epshtein, V.M., 1962. Key to parasites of freshwater fishes of the USSR. Academy of Science of the USSR, Zoological Institute. 919 P.

Chaudhary, A., Verma, C., Shobhna, M.V. and Singh, H.S., 2013. A review of Monogenean diversity in India: Pathogens of fish diseases. *Journal of Coastal Life Medicine*, 1, 151-168.

DOI:10.12980/JCLM.1.20133D130

- Chiary, H.R., Chaudhary, A., Singh, H.S. and Goswami, U.C., 2014.

 Molecular characterization of two non-native species of *Dactylogyrus* (Monogenea: Dactylogyridae) recovered from introduced hosts in India. *BioInvasions Record*, 3, 297-300. DOI:10.3391/bir.2014.3.4.12
- Cudmore, B. and Mandrak, N.E., 2004. Biological Synopsis of Grass Carp (*Ctenopharyngodon idella*). Canadian Manuscript Report of Fisheries and Aquatic Sciences 2705. 44 P.
- McGillivray, Cunningham, C.O., D.M. and MacKenzie, K., 1995. Phylogenetic analysis of salaris *Gyrodactylus* Malmberg, 1957 based on the small subunit (18S)ribosomal **RNA** gene. Molecular and Biochemical Parasitology, 71, 139-142
- Daghigh Roohi, J., Sattari, M., Masoumian, M., Nezamabadi, H., Mirhasheminasab. S. F.. Asgharnia, M., Ghorbanpour, N., M., Nahrvar. Rufchaei. Ramezani, B., Musavi, A., Abbasi, K., Rastin, R., Mahisefat, F. and Syaddokht, J., 2016. Study of parasites occurence and intensity in fishes of Anzali lagoon. Iranian Fisheries Science Research Institute. Final report of the project No. 48174. 36 P (In Persian).
- Dove, A.D. and Ernst, I., 1998.

 Concurrent invaders—four exotic species of Monogenea now established on exotic freshwater fishes in Australia. *International Journal for Parasitology*, 28, 1755-1764.

- Gibson, D.I., Timofeeva, T.A. and Gerasev, P.I., 1996. A catalogue of the nominal species of the monogenean genus *Dactylogyrus* Diesing, 1850 and their host genera. *Systematic Parasitology*, 35, 3-48.
- Gussev, A.V., 1983. Methods for the collection and processing fish parasites monogenean material Nauka, Leningrad, USSR, 48 P. (In Russian).
- Gussev, A.V., 1985. Identification key to parasites of freshwater fish of USSR vol. II. Nauka, Leningrad, (In Russian). pp.10-424.
- Jalali, B. and Molnar, K., 1990.

 Occurrence of monogeneans on freshwater fishes in Iran.

 Dactylogyrus spp. on cultured Iranian fishes. Acta Veterinaria Hungarica, 38, 239-242.
- Jiang, B., Chi, C., Fu, Y.W., Zhang, Q.Z. and Wang, G.X., 2013. In vivo anthelmintic effect of flavonol rhamnosides from *Dryopteris crassirhizoma* against *Dactylogyrus intermedius* in goldfish (*Carassius auratus*). *Parasitology Research*, 112, 4097-4104. DOI:10.1007/s00436-013-3600-3
- Lu, C., Zhang, H.Y., Ji, J. and Wang, G.X., 2012. In vivo anthelmintic activity of *Dryopteris crassirhizoma*, *Kochia scoparia*, and *Polygala tenuifolia* against *Dactylogyrus intermedius* (Monogenea) in goldfish (*Carassius auratus*). *Parasitolology Research*, 110,1085-1090. DOI:10.1007/s00436-011-2592-0
- Mcmanus, D.P. and Bowles, J., 1996.

 Molecular genetic approaches to parasite identification: their value in

- diagnostic parasitology and systematic. *International Journal for Parasitology*, 96(7), 687-704.
- Molnar, K., 1971. Studies on gill of the parasitosis grass (Ctenopharyngodon idella) caused **Dactylogyrus** by lamellatus Achmerow, 1952. 1. Morphology and Biology of Dactylogyrus lamellatus. Acta Veterinaria Academiae Scientiarum Hungricae, *Tomus*, 21(**2-3**), 267-289.
- Mousavi, H.E., Omidzahir, S., Shayan, Soltani. M., P., Ebrahimzadeh, E., Mousavi, S. Hoseini. 2013. and M., Morphometrical and molecular characterization of Gyrodactylus cichlidarum (Gyrodactylidae) from Astronotus ocellatus (Cichlidae) in Comparative Iran. Clinical 1093-1097. Pathology, 22, DOI:10.1007/s00580-012-1534-2
- Mozhdeganlou, Z., Ebrahimzadeh, M., Shayan, P., Soltani, M., Ebrahimzadeh, E. and Rostami, M., 2011. Detection of single Dactylogyrus spp. in DNA extracted from infected gill tissue of fishes using polymerase chain reaction. Iranian Journal of Veterinary Research, 5, 77-144.
- Omidzahir, S., Ebrahimzadeh Mousavi, H.A., Soltani, M., Shayan, P., Ebrahimzadeh, E. and Hoseini, M., 2012. Identification of Gyrodactylus gurleyi in Carassius auratus using morphometric and molecular characterization. Iranian Journal of Veterinary Medicine, 6, 41-46.

- Omidzahir, S., Mousavi, H., Shayan, P., Abkooh, E. and Mahmoodzadeh, H., 2015. Morphometric and molecular analysis of *Gyrodactlus kobayashi* in *Carassius auratus* (Linnaeus, 1758). *Journal of Veterinary Research*, 70, 425-432.
- Pazooki, J., Jalali Jafari, B. and Ghobadian, M., 2006. Monogenean species from freshwater fishes of Zanjan province, Iran. *Iranian Journal of Fisheries Sciences*, 6 (1), 103-112.
- Reed, P., Francis-Floyd, R., Klinger, R. and Petty, D., 2009.

 Monogenean parasites of fish. Fisheries and Aquatic Sciences University of Florida UF, FA28, USA. pp. 1-4.
- Shamsi, S., Jalali, B. and Aghazadeh Meshgi, M., 2009. Infection with *Dactylogyrus* spp. among introduced cyprinid fishes and their geographical distribution in Iran. *Iranian Journal of Veterinary Research*, 10, 70-74.
- Tu, X., Ling, F., Huang, A. and Wang, G., 2015. The first report of Dactylogyrus formosus Kulwiec, 1927 (Monogenea: Dactylogyridae) from goldfish (Carassius auratus) in central China. Parasitology Research, DOI: 10.1007/s00436-015-4474-3
- Woo, P., Bruno, D. and Lim, L., 2002. Diseases and disorders of finfish in cage culture. 365 hal. CABI, Malaysia. 353 P.
- Woo, P., 2006. Fish diseases and disorders: protozoan and metazoan

infections. vol 1. 2th edition. CABI, UK. 791 P.

Zolfinejad, K., Khara, H. and Filizadeh, Y., 2017. Food preference and growth of grass carp, *Ctenopharyngodon idella* (Cuvier and Valenciennes, 1844) fed some aquatic and terrestrial plants. *Iranian Journal of Fisheries Sciences*, 16(4), 1278-1286.