Downloaded from jifro.ir on 2025-11-18

# **Research Article**

# Diatom community structure along physico-chemical gradients in Southern Caspian Sea (Noor shore)

Zarei Darki B.1\*; Bigham S.2; Patimar R.3

Received: December 2019 Accepted: March 2020

#### Abstract

Diatom species diversity and cell abundance are influenced by environmental factors. The effect of physico-chemical factors on interseasonal dynamics of planktonic diatom community structure was examined along Noor shore of Caspian Sea to determine which of the variables impact more on the diatom diversity, distribution and cell number. Overall, 144 samples were collected during four sampling campaigns along four transects of 2 km long each with three sampling sites in 2014 (February, May, July and November). In total, 62 diatom species were identified, of which 27 species were typical marine or brackish-water and the rest were marine-freshwater or freshwater. In the present study, diatom species diversity was highest in winter. As a result, Shannon-Wiener Index was 3.5 in winter, while in spring it was reduced to 0.8. Among the revealed species, the centric Thalassiosira caspica was the most abundant (up to 2.75×10<sup>6</sup> cells L<sup>-1</sup>). Comparison of species and their abundances between coastal, middle and offshore stations, as well as all stations as a whole, showed that they differed (Test  $\chi$  2, p<0.05) except for *Navicula* sp. (in summer) and *Ulnaria ulna* (in autumn and winter). Canonical correspondence analysis results showed that in particular temperature and salinity, in the second place transparency, phosphates and to a lesser extent dissolved oxygen and pH can affect diatom species diversity and abundance. Diatoms Thalassiosira fasciculata, T. caspica and Melosira caspica may be recommended as biological indicators along the shore of Caspian Sea.

**Keywords:** Abundance, Bacillariophyceae, Caspian Sea, CCA analysis, Diatom diversity, Phytoplankton

<sup>1-</sup>Department of Marine Biology, Faculty of Marine Sciences, Tarbiat Modares University, Noor, Mazandaran Province, Iran

<sup>2-</sup>Department of Marine Science and Technology, University of Hormozgan, Bandar Abas, Iran

<sup>3-</sup>Department of Fisheries, Faculty of Natural Resources, Gonbad Kavous University, Gonbad Kavous, Iran

<sup>\*</sup>Corresponding author's Email: zareidarki@modares.ac.ir

## Introduction

The Caspian Sea has particular natural conditions, rich natural resources and an important geopolitical role in the surrounding regions (Kostianoy and Kosarev, 2005). More than 700 species and infraspecific taxa of algae have been recorded in the Caspian Sea (Guseynov et al., 2015). Among them diatoms the most diverse are (Yablonskaya, 2007). They play a significant role world-wide, responsible for primary production in the pelagic zone (Charles, 1985). Furthermore, diatoms are siliceous algae that are sensitive to chemical conditions (Mann, 1999); therefore they frequently used as biological indicators to evaluate the water quality (Hall and Smol, 1992; De la Rey et al., 2004; Ponmanickam et al., 2007).

Same as other algae, diatom species diversity, cell abundance and biomass are influenced by environmental factors (Habib et al., 1997; Naz and Türkmen, 2005: da Silva et al.. 2005). Temperature, light intensity, of nutrients, concentration phytoplankton density, population antagonistic relations aquatic hydrological climate and plants. changes and site depth are variables that can affect diversity and distribution of phytoplankton communities, especially diatoms (Onyema, 2008).

Presently salt composition of Caspian Sea is a result of changes in the originally oceanic water influenced by continental flows that occurred up after separating the Caspian Sea from the World Ocean (Kostianoy and Kosarev, 2005). The effect of seasonal changes

and small rivers that enter it are considered the main factors in determining pH in Southern Caspian Sea (Tuzhilkin and Kosarev, 2005). Furthermore, the river flow results in increasing phytoplankton diversity and decreasing water transparency (Bagheri al.. 2012). Phytoplankton et distribution, especially diatoms, is not uniform, primarily due to the water temperature. For example, in northern Caspian Sea, in winter when it is ice covered, the growth of most algae is halted. However, in middle Southern parts of the sea, algal growth continues at different growth rates (Tuzhilkin and Kosarev, 2005). During winter. inorganic phosphorus accumulated mainly within surface of the Caspian Sea due to increasing freshwater inflow and reduction in consumption. The temperature and photosynthesis increase, especially in summer, resulting in consumption of inorganic phosphorus (Nausch et al., 2008).

Taking into account all these conditions presently occurring in the Caspian Sea, it is important to determine variables that cause diatom distribution. The aim of this study was to follow the annual dynamics of the planktonic diatom structure under the influence of physico-chemical factors on the Noor coast in Southern Caspian Sea.

# Materials and methods

Four transects and 12 sampling sites were selected along Noor shore in Southern Caspian Sea (Fig. 1, Table 1).




Figure 1: Phytoplankton sampling transects (A, B, C, D) each bearing three sites (1, 2, 3) at Noor, Southern Caspian Sea.

Table 1: Geographic coordinates and depths at the investigated sites in February, May, July and November 2014.

Transect		Site	Depth	Northern latitude	Eastern longitude		
1	Sabze-rud River		A1	0.5	36°34'55	52°59'24	
			A2	5.0	36°34'40	52°59'82	
			A3	12.0	36°35'99	52°59'38	
	Noor River		B1	0.5	36°35'18	52°01'07	
2			B2	5.0	36°35'82	52°01'98	
			B3	12.0	36°35'37	52°01'02	
	Tarbiat University	M. 1	C1	0.5	36°35'22	52°02'05	
3		Modares	C2	5.0	36°35'62	52°02'11	
			C3	12.0	36°35'90	52°02'12	
			D1	0.5	36°35'74	52°05'89	
4	Lavij River		D2	5.0	36°35'95	52°05'72	
			D3	12.0	36°36'21	52°05'08	

The transects, which were nearly two kilometer in length, were placed at right angles to the coast and parallel to the Sabzehrood River (A), Noor River (B), Lavidjrood River (D), and Tarbiat Modares University (C). Sampling was carried out by a boat, seasonally from 2014 (May, July and November) to

February 2015. Both qualitative and quantitative phytoplankton samples were taken with a plankton net (55  $\mu$ m mesh size) and a Ruttner's bathometer (2 liter in volume), respectively. Overall, 144 algal samples were collected. In order to study the affecting factors on diatom dynamics, some

physico-chemical factors such as pH, temperature, transparency, oxygen saturation (O %), dissolved oxygen (DO), salinity, electrical conductivity (EC), phosphates were measured. Water transparency was estimated by a Secchi disk.

Phytoplankton samples were transported to the laboratory in dark cold containers as quickly as possible (Fathi and Flower, 2005). They were by a settling condensed method (Wasser et al., 1989) and quantitatively analyzed with a Nikon-Eclipse E200 microscope in laboratory of the Marine Biology Department, Tarbiat Modares University, Iran. To identify diatom species, before microscopic analysis, the diatoms were processed for the purpose of deleting organic contents of the cell by cold methods using 30% sulfuric acid with potassium permanganate (Proshkina-Lavrenko Lange-Bertalot 1974, 2001). Furthermore. specialized taxonomic references were used (Proshkina-Lavrenko Makarova 1968: and Proshkina-Lavrenko, 1974; Krammer and Lange-Bertalot, 1991; Lange-Bertalot, 2001).

After sampling, chlorophyll-a was immediately measured. For extraction, acetone and citric acid were used and results were interpreted using a BioTek ELISA microplate reader (model 264852) at wavelengths of 630, 647, 664 nm (Lorenzen, 1967).

To analyze the diatom species diversity, Shannon-Wiener Index (SWI) was estimated using PRIMER 5.0

program. To determine differences among seasons and stations, SWI was applied with SPSS 20 program. Chisquare test  $(\chi 2)$  was used to compare species and cell abundances among stations and seasons. To determine the between diatoms relationship variables, environmental canonical correspondence analysis (CCA) was applied using CANOCO 4.5 software (including ANOVA).

### **Results**

Average values of physico-chemical factors during the study period are presented in Table 2. Moreover, chlorophyll-a concentrations were measured along transects during the investigated time (Fig. 2).

In total 62 species of diatoms belonging to 30 genera were recorded (Table 3). Among them, 27 species were typically marine or brackish-water and the rest (>50%) were brought by river runoff. Diatom species diversity varied with the seasons and sampling sites fluctuations. The maximum species diversity (41 species) was observed in winter, and the pennate Nitzschia sp. had the highest abundance  $(2.25\times10^6 \text{ cells L}^{-1})$ . The lowest diatom species diversity was recorded in spring (14 species); the centric Thalassiosira caspica was the most abundant at transect A  $(2.75\times10^6 \text{ cells } \text{L}^{-1})$ . In summer, among 20 recorded species, centric the Chaetoceros muelleri the maximum abundance reached  $(2.00\times10^6 \text{ cells } \text{L}^{-1})$  at station A2. In autumn, 34 diatom species were found in all stations. *Thalassionema* nitzschioides  $(11\times10^6 \text{ cells } \text{ L}^{-1})$ , Nitzschia sp.  $(5.25\times10^6 \text{ cells } \text{ L}^{-1})$ ,

Pseudosolenia calcar-avis  $(6.75 \times 10^6)$  cells L<sup>-1</sup>) were the most numerous, especially at stations B1, C1 and D1.

Table 2: Mean values and ranges of physico-chemical factors in coastal water along Noor shore in Southern Caspian Sea over the studied period.

		Physical and chemical factor, Mean ± SD							
Season	S	pН	DO, mg L <sup>-1</sup>	Ο%	Salinity, ‰	T, °C	EC	Ph	
	A	8.16±0.02	11.76±0.16	106.76 ±0.11	10.96±0.63	10.40±0.41	18.51±0.67	0.10±0.09	
Winter	В	8.19±0.03	10.56±0.78	97.66±4.87	10.97±0.99	10.90±0.17	$14.43\pm1.84$	$0.12\pm0.01$	
	C	$8.22 \pm 0.03$	9.73±0.47	91.53±7.72	11.60±0.12	10.96±0.11	$18.81 \pm 0.24$	$0.09\pm0.06$	
	D	$8.30 \pm 0.02$	9.15±0.22	83.93±1.40	$11.00 \pm 1.07$	11.10±0.49	$17.46 \pm 1.50$	$0.10\pm0.02$	
	A	$8.46 \pm 0.22$	$7.50\pm0.26$	91.90±4.06	11.23±0.80	$24.60\pm0.25$	18.37±0.2.	$0.11 \pm 0.04$	
Carina	В	8.65±0.04	7.53±0.20	91.06±1.92	11.10±0.18	25.40±0.10	18.51±0.51	$0.10\pm0.01$	
Spring	C	$8.68 \pm 0.05$	$7.50\pm0.17$	94.03±1.92	11.43±0.09	26.16±0.58	18.64±0.17	$0.09\pm0.03$	
	D	$8.66 \pm 0.03$	$7.56\pm0.15$	92.60±1.35	11.20±0.09	$25.40\pm0.25$	$18.35 \pm 0.25$	$0.09\pm0.02$	
	A	$8.38 \pm 0.09$	4.56±0.87	57.50±11.4	11.60±0.01	27.90±0.63	18.94±0.03	$0.46\pm0.24$	
Summer	В	8.57±0.01	6.60±2.02	75.50±1.67	11.60±0.01	27.60±0.60	19.34±0.49	$0.20\pm0.03$	
Summer	C	8.67±0.03	5.50±0.30	71.90±3.58	11.84±0.33	$28.30\pm0.17$	$2.64\pm1.48$	$0.19\pm0.12$	
	D	$8.68 \pm 0.03$	$6.03\pm0.05$	77.26±0.40	$11.70\pm0.03$	$28.00 \pm 0.37$	$19.09\pm0.04$	$0.24 \pm 0.14$	
	A	$8.60\pm0.24$	13.04±0.80	107.93±1.18	11.27±0.05	14.70±0.55	18.36±0.13	$0.20\pm0.14$	
Autum	В	8.48±0.16	7.96±0.58	75.50±1.67	11.10±0.16	11.70±0.41	18.19±0.32	$0.08\pm0.06$	
Autumn	C	8.66±0.06	12.44±2.14	$107.16 \pm 2.01$	11.14±0.16	$14.70\pm0.15$	18.05±0.37	$0.09\pm0.03$	
	D	$8.58 \pm 0.13$	10.25±0.32	$101.53 \pm 0.40$	10.67±0.67	$15.00 \pm 0.15$	17.42±1.51	$0.09\pm0.02$	

Notes: S, site; SD, standard deviation; DO, dissolved oxygen; O%, oxygen saturation; T, temperature; EC, electrical conductivity; Ph, phosphates.




Figure 3: Variation of Shannon-Wiener Index applied to planktonic diatomic species diversity at different sampling stations (1 to 3) along transects (A to D) in different seasons.

Table 3: Planktonic diatom species composition in coastal waters along Noor shore in Southern Caspian over the studied period.

N	Taxa		Season			
			S	A	W	
1	Amphora ovalis (Kützing) Kützing 1844	-	-	+	+	
2	Amphora sp.	+	+	-	+	
3	Aulacoseira granulate (Ehrenberg) Simonsen 1979	-	-	+	+	
4	Chaetoceros muelleri Lemmermann 1898*	-	+	+	+	
5	Chaetoceros peruvianus Brightwell 1856*	-	-	+	-	
6	Chaetoceros subtilis Cleve 1896*	-	+	-	-	
7	Chaetoceros throndsenii Marino, Montresor & Zingone 1991*	-	-	+	+	
8	Chaetoceros sp	-	-	+	-	
9	Cocconeis placentula Ehrenberg 1838	-	-	+	+	
10	Coscinodiscopsis jonesiana (Greville) E. A. Sar & I. Sunesen in Sar,	+	_	_	_	
	Sunesen & Hinz 2008 *					
11	Coscinodiscus gigas Ehrenberg 1843*	-	-	-	+	
12	Coscinodiscus sp. Gran & Angst 1931	_	-	+	_	
13	Cyclotella meneghiniana Kützing 1844	_	+	+	+	
14	Cyclotella sp.	-	-	+	-	
15	Cymbella sp.	+	-	_	+	
16	Diatoma vulgaris Bory de Saint-Vincent 1824	_	-	+	+	
17	Diploneis interrupta (Kützing) Cleve 1894*	+	-	_	-	
18	Diploneis papula (A. W. F. Schmidt) Cleve 1894*	_	+	_	-	
19	Diploneis smithii (Brébisson) Cleve 1894*	_	-	+	+	
20	Epithemia turgida (Ehrenberg) Kützing 1844	_	+	_	_	
21	Fragilaria capucina Desmazières 1830	+	+	+	+	
22	Gyrosigma attenuatum (Kützing) Rabenhorst 1853	_	+	+	+	
23	Gyrosigma sp.	_	-	-	_	
24	Halamphora coffaeiformis (C. Agardh) Mereschkowsky 1903	_	-	+	_	
25	Halamphora veneta (Kützing) Levkov 2009	_	-	+	-	
26	Hantzschia sp.	_	-	_	+	
27	Melosira juergensii var. caspica A. Henckel*	-	-	-	+	
28	Melosira lineata (Dillwyn) C. Agardh 1824*	-	+	-	-	
29	Melosira moniliformis C. Agardh 1824*	_	-	+	-	
30	Melosira sp.	_	-	_	+	
31	Melosira varians C. Agardh 1827	-	-	+	+	
32	Navicula cincta (Ehrenberg) Ralfs 1861*	_	-	+	+	
33	Navicula cryptocephala Kützing 1844*	+	+	+	-	
34						
	Navicula erifuga Lange-Bertalot 1985*	-	-	-	+	
35	Navicula radiosa Kützing 1844	_	_	_	+	
36	Navicula sp.	+	_	+	_	
37	Nitzschia acicularis (Kützing) W. Smith 1853*	_	_	_	+	
38	Nitzschia frustulum (Kützing) Grunow 1880*	_	_	_	+	
39	Nitzschia longissima (Brébisson) Ralfs 1861*	_	_	_	+	
40	Nitzschia palea (Kützing) W. Smith 1856	_	_	_	+	
41	Nitzschia reversa W. Smith 1853	_	_	+	+	
42	Pinnularia sp.	_	_	+	_	
43	Planothidium vanheurckii (Grunow in Van Heurck) E. W. Thomas, Van					
-	de Viiver & Kociolek 2015	-	-	-	+	
44	Pseudosolenia calcar-avis (Schultze) B. G. Sundström 1986*	-	+	+	_	
45	Pseudo-nitzschia seriata (Cleve) H. Peragallo 1899*	+	+	+	+	
46	Pseudo-nitzschia sp.	+	+	+	+	
47	Rhizosolenia sp.	-	+	+	+	

**Table 3 (Continued):** 

N	Taxa	Season			
		Sp	S	A	W
48	Rhoicosphenia abbreviata (C. Agardh) Lange-Bertalot 1980	-	-	+	+
49	Rhoicosphenia sp.	-	-	+	+
50	Skeletonema costatum (Greville) Cleve 1873*	-	+	-	-
51	Skeletonema subsalsum (A. Cleve) Bethge 1928*	-	-	+	+
52	Skeletonema sp.	-	-	+	-
53	Surirella ovalis Brébisson 1838	-	-	-	+
54	Synedra sp.	+	+	-	+
55	Tabularia fasciculata (C. Agardh) D. M. Williams & Round 1985*	+	-	-	-
56	Tabularia sp.	-	+	-	+
57	Thalassionema nitzschioides (Grunow) Mereschkowsky 1902*	+	+	+	+
58	Thalassionema sp.	-	-	-	+
59	Thalassiosira caspica I. V. Makarova 1957*	+	+	+	+
60	Thalassiosira coronifera Proshkina-Lavrenko*	-	-	-	+
61	Thalassiosira sp. Cleve P. T. 1873	-	-	-	+
62	Ulnaria ulna (Nitzsch) Compère 2001*	+	+	+	+

<sup>\*</sup> Marine species including not only typical marine species, but also species of brackish water ecotypes Notes: Sp, spring; S, summer; A, autumn; W, winter

Shannon-Wiener Index (SWI) different seasons reached a maximum value of 3.5 at station A1 in winter, whereas the minimum value was 0.8 at station D3 in spring (Fig. 3). Analysis of variance showed significant differences in SWI among seasons \*\*\*, ANOVA, F=26.805). (p < 0.05)Mean SWI values for transects ranged between 1.2 and 3.2.

A comparison of species and their abundances among coastal, middle and offshore stations, as well as all stations as a whole, showed that they significantly differed (Test  $\chi$  2, p<0.05) except for *Navicula* sp. (in summer) and *Ulnaria ulna* (in autumn and winter).

Based on CCA results, the eigenvalue of axis 1 (0.23-0.44) explained from 21.6 to 30.3% and the eigenvalue of axis 2 (0.23-0.32) interpreted 41.4-51.4%, indicating a relationship between diatom abundance and water temperature, transparency,

salinity, EC, oxygen saturation, DO, pH as well as phosphates in different seasons (Fig. 4). Interestingly, the indices species-environment correlations of both axes were 100%. In spring Tabularia fasciculata, Amphora sp, and Th. caspica were positively at correlation with DO offshore stations. The abundance of Pseudonitzschia seriata, Ulnaria ulna, Synedra **Diploneis** interrupta, and sp, Coscinodiscopsis ionesiana were correlated with the water temperatures, which had a significant impact on their presence in the water column. The pH did not have a strong impact on the species abundance. Chlorophyll-a was correlated with *Navicula* sp., Thalassionema cryptocephala and nitzschioides.

In summer, *Rhizosolenia* sp. and *Tabularia* sp. were associated with salinity at middle stations. *Pseudonitzschia seriata* and *Melosira caspica* were correlated positively with oxygen

saturation and DO. Flagilaria capucina, Melosira lineata and Epithemia turgida were principally responsible for chlorophyll-a at stations nearest to the coast. Diploneis papula, Cyclotella meneghiniana, Skeletonema costatum, Thalassionema nitzschioides, Synedra

sp. and *Amphora* sp. were associated with phosphates at these coastal stations. *Gyrosigma attenuatum* was positively correlated with transparency. Temperature directly influenced *Chaetoceros muelleri* and *C. subtilis* at offshore stations.




Figure 3: Variation of Shannon-Wiener Index applied to planktonic diatomic species diversity at different sampling stations (1 to 3) along transects (A to D) in different seasons.

In autumn, the correlation between diatom abundance and physicochemical factors were not ascertained for many species. However, Diatoma vulgaris was positively related to phosphates; Nitzschia reversa was positively correlated with temperature; Chaetoceros muelleri was moderately associated with pH; and Th. caspica and Gyrosigma attenuatum were correlated with transparency. Thalassionema nitzschioides, Pseudonitzschia sp, Coscinodiscus sp., moniliformis, Melosira Chaetoceros throndsenii, Chaetoceros sp. were positively correlated with salinity. Cyclotella sp. and Ulnaria ulna were positively correlated with chlorophyll-a at coastal stations.

winter, Diatoma vulgaris was positively correlated with oxygen. When temperature increased, abundance of some species. for instance. Aulacoseira granulata, Melosira caspica, M. varians. throndsenii, Pseudo-Chaetoceros nitzschia sp., increased. Moreover, the abundance of Coscinodiscus gigas, meneghiniana, Cyclotella Nitzschia reversa, Rhizosolenia sp., Rhoicosphenia abbreviata, Thalassionema sp. and Thalassiosira decreased when the salinity decreased. Thalassiosira coronifera Th. caspica were positively correlated with transparency.




Figure 4: CCA biplot between planktonic diatom abundance and physico-chemical variables in four seasons at Noor in Southern Caspian Sea.

Notes: CS, coastal station; MS, middle station; OS, off-shore station; DO, dissolved oxygen; EC, electrical conductivity; O, oxygen saturation; Phosph, phosphates; Sal, salinity; Trans, transparency; Temp, temperature; AO, Amphora ovalis; ASp, Amphora sp.; MG, Aulacoseira granulate; ChM, Chaetoceros muelleri; ChP, Ch. peruvianus; ChS, Ch. subtilis; ChT, Ch. throndsenii; ChSp, Chaetoceros sp.; CP, Cocconeis placentula; CJ, Coscinodiscus jonesiana; CG, C. gigas; CSP, Coscinodiscus sp.; CM, Cyclotella meneghiniana; CySP, Cyclotella sp.; CySP1, Cymbella sp.; DV, Diatoma vulgaris; DI, Diploneis interrupta; DP, D. papula; DS, D. smithii; ET, Epithemia turgida; FC, Fragilaria capucina; GA, G. attenuatum; GSp, Gyrosigma sp.; AC, Halamphora coffeiformis; AV, H. veneta; HSp, Hantzschia sp.; MC, Melorira caspica; ML, M. lineata; MM, M. moniliformis; MSp, Melosira sp.; MV, M. varians; NC, Navicula cincta; NCr, N. cryptocephala; NL, N. erifuga; NR, N. radiosa; NSp, Navicula sp.; NA, Nitzschia acicularis; NF, N. frustulum; NL, N. longissima; NP, N. palea; NR, N. reversa; PSp, Pinnularia sp.; PV, P. vanheurckii; PS, Pseudo-nitzschia seriata; PsSp, Pseudo-nitzschia sp.; RC, Rhizosolenia calcar-avis; RSp, Rhizosolenia sp.; RSp, Rhoicosphenia sp.; RA, R. abbreviata; SC, Skeletonema costatum; SS, S. subsalsum; SSp, Skeletonema sp.; SO, Surirella ovalis; SySp, Synedra sp.; TF, T. fasciculata; TSp, Tabularia sp.; TN, Th. nitzschioides; TSp, Thalassionema sp. TC, Th. caspica; TCo, Th. coronifera; TSp, Thalassiosira sp.; SU, Ulnaria ulna.

### **Discussion**

Study of the structure and function of phytoplankton communities in a marine ecosystem is important for understanding biogeochemical and ecological processes and for identifying common distributional patterns that might suggest underlying mechanisms controlling them (Goericke, McFarland. 2014). In total 306 planktonic diatom species belonging to 78 genera, 41 families and 22 orders have been recorded in the Caspian Sea Jafarova. (Karayeva and 2007: Karayeva and Bukhliyarova, 2010). During the present study, 62 diatom species were identified. Among them, Th. caspica had the highest abundance  $(2.75\times10^6 \text{ cell } \text{L}^{-1})$  in spring, and the lowest diatom species diversity was observed at this time of the year. The decrease in diatom diversity and mass proliferation of a single species in the study area in spring may be related to river discharge, including domestic, industrial and agricultural wastewater (Jamshidi and Bin Abu Bakar, 2011). In winter, diatom species diversity was highest. SWI varied between 0.8 and 3.5, suggesting an unstable diatom community structure during the year (Tahami et al., 2012). Interseasonal changes in water physicochemical variables may affect the structure of phytoplankton communities (Liu et al., 2010). The most important factor among them is temperature (Kideys et al., 2005; Abrantes et al., 2006). Salmanov (1987) and Resende et (2007) reported that seasonal changes in temperature are considered

main factor controlling the phytoplankton species composition and structure. Algal distribution in the Caspian Sea is also dependent on temperature. Eurythermal algae such as Fragilaria capucina, Pseudo-nitzschia seriata, Thalassionema nitzschioides, Thalassiosira caspica and Ulnaria ulna live in Southern Caspian, and most diatoms living in Southern part of the sea are often found in the Middle Caspian (Karayeva and Makarova, 1973). When water temperature decreases to 8°C throughout Southern Caspian Sea, diatom diversity and cell abundance increase and vice versa (Salmanov, 1987). Their species diversity decreases when water temperature increases to 30°C (Bagheri et al., 2012). The present results also showed that changes in temperature affected the species appearance and proliferation, especially those of Chaetoceros muelleri, C. subtilis, C. throndsenii. Coscinodiscopsis jonesiana, Diatoma vulgaris, Diploneis interrupta, Gyrosigma attenuatum, Aulacoseira granulata, Melosira caspica, M. varians, Nitzschia cincta, N. acicularis, N. frustulum, N. palea, N. Pseudo-nitzschia reversa, seriata, Ulnaria ulna, Th. caspica, and T. coronifera.

Salinity is also important. During the study period it ranged from 9.8 to 19.1. Salinity varied greatly between winter and summer (Khosropanah *et al.*, 2011) and with distance from the coast and also with depth (Guliyev *et al.*, 2014). In general aquatic organisms inhabiting Caspian Sea grow within four basic and

intermediate three salinity zones. having specific salt composition at all barrier salinities (Plotnikov and Aladin, 2011). In the study zone, freshwater occupied areas were around Sabzehrood River, Noor River and Lavidirood River estuaries. Freshwater and brackish-water diatom species contributed more than half the total diatom species richness. being especially numerous in spring and summer. The number of brackish and marine species increased with distance from the shore and with increasing EC, and the freshwater ones, in contrast decreased. Overall, based on CCA species marine (Thalassionema nitzschioides. Melosira moniliformis. Chaetoceros throndsenii Coscinodiscus gigas) were positively correlated with salinity and EC, while freshwater species (Cyclotella meneghiniana, Nitzschia reversa, Rhoicosphenia abbreviata) were negatively correlated with these variables. In particular, the salinity influence was found to be higher at middle stations of each transect where the river algae invaded brackish-marine waters and a significant decrease in phytoplankton growth can be observed (Lionard et al., 2005).

Presence of oxygen in water allows aquatic organisms to live. Measurements of DO helps to estimate the health of water ecosystems and the amount of organic pollutants in water bodies (Xie *et al.*, 2003). Anthropogenic stress in Caspian Sea has resulted in decrease of Dissolved

Oxygen (Kostianov and Kosarev, 2005; Hadjizadeh Zaker, 2007). Furthermore, the lagoons and coastal areas in Southern Caspian Sea are constantly polluted by human activities (Kideys 2003; and Moghim, CEP. 2006: Kopelevich et al., 2008). Therefore, reliable indicators are needed for rapid determination of water quality. The diatoms Thalassiosira fasciculata, Th. caspica and Melosira caspica may be recommended as biological indicators because they were positively correlated with DO in our study.

Among other species, it is notable that G. attenuatum, Th. caspica, and Th. coronifera are sensitive to transparency. In the Caspian Sea, transparency can decrease because of fluctuations in the water level and as a result destruction of coastal soil, the inflow river drifts as well as significant growth of hydrobionts (Nasrollahzadeh Saravi and Hosseini, 2004; Plotnikov and Aladin, 2011; Salmanov et al., 2015). Therefore, chlorophyll-a represents an unmeasured potential for influencing DO and also allows indirect assessment of ecological conditions in a water body (Miltner, 2010; Jamshidi and Bin Abu Bakar, 2011). In Caspian Sea mean chlorophyll-a content in the surface layer increased since 2003 (Jamalomidi, 2013). High chlorophyll-a values were observed in our study, especially at the coastal stations in autumn. In spring, chlorophyll-a was negatively correlated with water transparency. Study of chlorophyll-a distribution in some lakes showed significant hyperbolic

relationship between transparency and algal biomass (Canfield and Hodgson, 1983; Hosseini and Ordog, 1995).

To a greater extent this is due to increasing concentrations of nitrates and phosphates caused by discharge of untreated domestic sewage and pesticides from agricultural activities into rivers (Khosropanah et al., 2011). It is known that some diatoms are not able to dominate when phosphorus is deficient, even if silicate and nitrate are in excess (Katiyar et al., However, some of them can apply physiological and molecular strategies such as phosphorus scavenging or recycling as well as adjusting cell growth to adapt to limiting phosphorus concentrations (Brembu et al., 2017). According to the conducted research, phosphates were responsible for growth of Diatoma vulgaris, Diploneis papula, Cyclotella meneghiniana, Skeletonema costatum, Thalassionema nitzschioides, Synedra sp. and Amphora sp. at coastal stations.

Thus the present study showed that temperature, salinity, transparency, phosphates, and to a lesser extent DO and pH, can affect planktonic diatom species diversity in the Southern Caspian. In some cases it was not possible to determine the relationships between species and physico-chemical factors. However, one thing absolutely certain: ecological conditions of Caspian Sea along the Noor shore are unfavorable because of anthropogenic activities and need to be improved to preserve its unique aquatic species diversity.

# Acknowledgments

The authors express their gratitude and appreciation to Tarbiat Modares University for providing funding and research facilities. We also thank laboratory assistant of the Faculty of Marine Sciences for help.

#### References

Abrantes, N., Antunes, S.C., Pereira, M.J. and Gonçalves, F., 2006. Seasonal succession of cladocerans and phytoplankton and their interactions in a shallow eutrophic lake (Lake Vela, Portugal). *Acta Oecologica*, 29(1), 54-64. DOI: 10.1016/j.actao.2005.07.006.

Bagheri, S., Mansor, M., Türkoğlu, M., Makaremi, M., Wan Omar, W.M. and Negarestan, H., 2012. Phytoplankton species composition and abundance in the southwestern Caspian Sea. *Ekoloji*, 21(83), 32-43. DOI: 10.5053/ekoloji.2012.834.

Brembu, T., Mühlroth, A., Alipanah, L. and Bones, A.M., 2017. The effects of phosphorus limitation on metabolism carbon in diatoms. Philosophical Transactions of the Royal Society *B*: **Biological** 372(1728), 20160406. Sciences, Published online, DOI: 10.1098/ rstb.2016.0406.

Canfield Jr., D.E. and Hodgson, L.M., 1983. Prediction of Secchi disc depths in Florida lakes: Impact of algal biomass and organic color.

- *Hydrobiologia*, 99(**1**), 51-60. DOI: 10.1007/BF00013717.
- CEP, 2006. Identification of the extent and causes of cyanobacterial bloom in September-October 2005 and development of the capacity for observation and prediction of HAB in the Southern Caspian Sea using Remote Sensing Technique. Caspian Environment Programme 2006.
- Charles, D.F., 1985. Relationships between surface sediment diatom assemblages and lakewater characteristics in Adirondack lakes. *Ecology*, 66(3), 994-1011. DOI: 10.2307/1940561.
- da Silva. **C.A..** Train. S. and L.C.. Rodrigues. 2005. Phytoplankton assemblages in a **Brazilian** subtropical cascading reservoir system. Hydrobiologia, 537(1), 99-109. DOI: 10.1007/ s10750-004-2552-0.
- **De la Rey, P.A., Taylor, J.C., Laas, A., Van Rensburg, L. and Vosloo, A., 2004.** Determining the possible application value of diatoms as indicators of general water quality: A comparison with SASS 5. *Water SA*, 30(3), 325-332. DOI: 10.4314/wsa.v3013.4080.
- Fathi, A.A. and Flower, R.J., 2005. Water quality and phytoplankton communities in Lake Qarun (Egypt). *Aquatic Science*, 67(3), 350-362. DOI: 10.1007/s00027-005-0777-2.
- Goericke, R., 2011. The structure of marine phytoplankton communities patterns, rules and mechanisms. *CalCOFI Reports*, 52, 182-197.

- J.A., S.R., Gulivev. Hadjiveva, Ghorbanpour. Sh. and Aminbekov, A., 2014. Study of vertical distribution of heavy metals in the border-bottom water in the area of the Absheron Peninsula. International **Journal** of Basic Sciences and Applied Research, 3(9), 606-610.
- Guseynov, M.K., Guseinov, K.M. and Gasanova, A.SH., 2015. Caspian biological resources. *South of Russia: ecology, development*, 10(2), 38-53. DOI: 10.18470/1992-1098-2-38-53.
- Habib. O.A., Tippett, R. and 1997. Seasonal Murphy, K.J., changes in phytoplankton community structure in relation to physico-chemical factors in Loch Lomond, Scotland. Hydrobiologia, 350, 63-79. DOI: 10.1023/A:1003037012226.
- Hadjizadeh Zaker. N., 2007. Characteristics and seasonal variations of dissolved oxygen. **International Journal** of Environmental Research, 1(4), 296-301.
- Hall, R.I. and Smol, J.P., 1992. A weighted-averaging regression and calibration model for inferring total phosphorus concentration from diatoms **British** Columbia Lakes. Freshwater (Canada) Biology, 27(3), 417-434. DOI: 10.1111/j.1365-2427.1992.tb00551.x.
- Hosseini, S.A. and Ordog, V., 1995. Relationships of chlorophyll a with

- some physical and chemical parameters in fish ponds. *Aquaculture Hungarica*, 8, 64-66.
- Jamalomidi, M., 2013. Temporal changes of surface chlorophyll in south of Caspian Sea based on data gained by MODIS of Aqua satellite. *International Journal of Agriculture and Crop Sciences*, 5(12), 1269-1275.
- Jamshidi, S. and Bin Abu Bakar, N., 2011. A study on distribution of chlorophyll-a in the coastal waters of Anzali Port, South Caspian Sea. *Ocean Science Discussions*, 8(1), 435-451. DOI: 10.5194/ods-8-435-2011.
- **Karayeva, N.I. and Bukhliyarova, L.N., 2010.** An addition to the Bacillariophyta flora of the Caspian Sea. *Algologia*, 20(4), 471-481, in Russian.
- Karayeva, N.I. and Jafarova, S.K., 2007. The main directions of algological researches at the Institute of Botany, National Academy of Sciences of Azerbaijan, Izvestie NANA. Seria Biologicheskie Nauki, 5(6), 11-20.
- **Karayeva, N.I. and Makarova, I.V., 1973.** Specific features and origin of the Caspian Sea diatom flora. *Marine Biology*, 21, 269-275. DOI: 10.1007/BF00381083.
- Katiyar, D., Lall, A.M. and Singh, B., 2010. Effect of phosphate on growth of diatoms. *Indian Journal of Scientific Research*, 1(2), 103-106.
- Khosropanah, N., Nejatkhah Manavi, P., Koohilay, S. and Naseri, M.T.,

- **2011.** Variations in nitrate and phosphate contents of waters in the southwest Caspian Sea. *Journal of the Persian Gulf*, 2(5), 27-33.
- **Kideys, A.E. and Moghim, M., 2003.**Distribution of the alien ctenophore *Mnemiopsis leidyi* in the Caspian Sea in August 2001. *Marine Biology*, 142(1), 163-171. DOI: 10.1007/s00227-002-0932-4.
- **Kideys, A.E., Soydemir, N., Eker, E., Vladymyrov, V., Soloviev, D. and Melin, F., 2005.** Phytoplankton distribution in the Caspian Sea during March 2001. *Hydrobiologia*, 543(1), 159-168. DOI: 10.1007/s10750-004-6953-x.
- Kopelevich, O.V., Burenkov, V.I. and Sheberstov, S.V., 2008. Biodiversity. Case studies of optical remote sensing in the Barents Sea, Black Sea and Caspian Sea. In: *Remote Sensing of the European Seas.* Barale V. and Gade M. editors. Springer, Heidelberg, Germany: 53-66.
- Kostianoy, A. and Kosarev, A.N., 2005. The Caspian Sea Environment. Springer Science and Business Media, Berlin, Germany, 271P.
- Krammer, K. and Lange-Bertalot, H., 1991. Süβwasserflora von Mitteleuropa, Bd 2/3: Bacillariophyceae 3. Teil Centrales, Fragilariaceae, Eunotiaceae. Spektrum Akademischer Verlag, Berlin/Heidelberg, Germany, 598P.
- Lange-Bertalot, H., 2001. Diatoms of Europe, Volume 2: Navicula sensu

- strict; 10 genera separated from Navicula sensu lato; Frustulia. A.R. Gantner Verlag K.G., Koenigstein, Germany, 526P.
- Lionard, M., Muylaert, K., Van Gansbeke, D. and Vyverman, W., 2005. Influence of changes in salinity and light intensity on growth of phytoplankton communities from the Schelde River and estuary (Belgium/The Netherlands). *Hydrobiologia*, 540(1), 105-115. DOI: 10.1007/s10750-004-7123-x.
- Liu, C., Liu, L. and Shen, H., 2010.
  Seasonal variations of phytoplankton community structure in relation to physico-chemical factors in Lake Baiyangdian, China. *Procedia Environmental Sciences*, 2, 1622-1631.

  DOI: 10.1016/j.proenv.2010.10.173.
- **Lorenzen, C.J., 1967.** Determination of chlorophyll and phaeo-pigments: spectrophotometric equations. *Limnology and Oceanography*, 12, 343-346. DOI: 10.4319/lo.1967.12.2.0343.
- **Mann, D.G., 1999.** The species concept in diatoms. *Phycologia*, 38(**6**), 437-495. DOI: 10.2216/ 10031-38-6-437.1.
- McFarland, M.N., 2014. Impact of phytoplankton community structure and function on marine particulate optical properties. Ph.D Theses, University of Rhode Island, Kingston, USA. DOI: 10.23860/diss-mcfarland-malcom-2014.
- Miltner, R.J., 2010. A method and rationale for deriving nutrient criteria

- for small rivers and streams in Ohio. *Environmental Management*, 45(4), 842-855. DOI: 10.1007/s00267-010-9439-9.
- Nasrollahzadeh Saravi, H. and Hosseini, S.A., 2004. Correlation between changes of chlorophyll-a and transparency in the Southern Caspian Sea. *Iranian Scientific Fisheries Journal*, 13(1), 191-200, in Persian. DOI: 10.22092/ISFJ.2004.113727.
- Nausch, M., Nausch, G., Wasmund, N. and Nagel, K., 2008. Phosphorus pool variations and their relation to Cyanobacteria development in the Baltic Sea: A three-year study. *Journal of Marine Systems*, 71(1-2), 99-111. DOI: 10.1016/j.jmarsys.2007.06.004.
- Naz, M. and Türkmen, M., 2005. Phytoplankton biomass and species composition of Lake Gölbaşi (Hatay-Turkey). *Turkish Journal of Biology*, 29(1), 49-56. DOI: 10.12714/egejfas.2006.23.1.5000156798.
- Onyema, I.C., 2008. A checklist of phytoplankton species of the Iyagbe Lagoon, Lagos. *Journal of Fisheries and Aquatic Science*, 3(3), 167-175. DOI: 10.3923/ifas.2008.167.175.
- Plotnikov, I.S. and Aladin, N.V., 2011. An overview of hybrid marine and lacustrine seas and saline lakes of the world. *Lakes and Reservoirs:* Research and Management, 16(2), 97-108. DOI: 10.1111/j.1440-1770.2011.00452.x.
- Ponmanickam, P., Rajagopal, T., Rajan, M.K., Achiraman, S. and

- Palanivelu, K., 2007. Assessment of drinking water quality of Vembakottai Reservoir, Virudhunagar district, Tamil Nadu. *Journal of Experimental Zoology*, 10(2), 485-488.
- Proshkina-Lavrenko, A.I. and Makarova, I.V., 1968. Plankton algae of the Caspian Sea, Nauka Press, Leningrad, Russia, in Russian, 295P.
- Proshkina-Lavrenko, A.I., 1974.

  Diatoms of the USSR: Fossil and Recent. Nauka Press 1(1),
  Leningrad, Russia, in Russian, 403P.
- Resende, P., Azeiteiro, U.M., Gonçalves, F. and Pereira, M.J., 2007. Distribution and ecological preferences of diatoms and dinoflagellates in the west Iberian coastal zone (North Portugal). *Acta Oecologica*, 32(2), 224-235. DOI: 10.1016/j.actao.2007.05.004.
- **Salmanov, M.A., 1987.** The role of microflora and phytoplankton in the production processes of the Caspian Sea. Nauka Press, Moscow, Russia, 214P.
- Salmanov, M.A., Guseynov, A.T. and Gasanov, G.F., 2015. Changes in transparency of water and reduction of biomass of benthic communities due to the damage to the banks and the overall pollution in western part of the South Caspian Sea. *South of Russia: Ecology, Development*, 10(2), 32-37. DOI: 10.18470/1992-1098-2015-2-32-37.
- Tahami, F.S., Mazlan, A.G., Negarestan, H., Najafpour, S.,

- Wan Muda, W.M.L. and Najafpour, G.D., 2012. Phytoplankton combination in the southern part of Caspian Sea. *World Applied Sciences Journal*, 16(1), 99-105.
- **Tuzhilkin, V.S. and Kosarev, A.N., 2005.** Thermohaline structure and general circulation of the Caspian Sea waters. In: *The Caspian Sea Environment*. Kostianoy, A. and Kosarev, A. editors. The handbook of environmental chemistry, vol 5p. Springer, Berlin, Germany: 33-57. DOI: 10.1007/698-5-003.
- Wasser, S.P., Kondratieva, N.V., N.P., Masyuk, Palamar'-Mordvintseva, G.M., Vetrova, Z.I., Kordyum, E., Moshkova, N.A., Prikhotkova, L.P., Stupina, V.V., Tsarenko, P., Unger, V.P.. Radchenko, M.I., Vinogradova, O., Bukhtiyarova, L. 1989. Razumna, L.F., Algae Reference Book. Naukova Dumka Publisher, Kiev, Ukraine, in Russian, 608P.
- Xie, L.Q., Xie, P. and Tang, H.J., 2003. Environment of dissolved phosphate release from sediment to lake water by Microcystis biomass An enclosure experiment in a hyper eutrophic, subtropical Chinese lake. *Environmental Pollution*, 122(3), 391-399. DOI: 10.1016/s0269-7491(02)00305-6.
- Yablonskaya, E.A., 2007. Biology of the Caspian Sea. VNIRO, Moscow, Russia, in Russian, 142P.