Downloaded from jifro.ir on 2025-11-18

Research Article

Effects of using enriched copepod with microalgae on growth, survival, and proximate composition of giant freshwater prawn (*Macrobrachium rosenbergii*)

Rasdi N.W.^{1,2}*; Ikhwannuddin M.³; Syafika C.A.¹; Azani N.^{1,3}; Ramli A.¹

Received: March 2020 Accepted: May 2020

Abstract

Copepods are suitable to be used as a starter feed for prawn or fish larvae in hatchery. Nowadays, farmers are still having problems sustaining growth of Macrobrachium rosenbergii production since the use of commercial diets such as micro-pallet. This study was carried out for 30 days to determine the best enrichment formula for copepods. Five dietary treatments with triplicates each were used in this study to test efficacy of enriched copepod (Oithona sp.) to M. rosenbergii larvae. The enrichments used were Tetraselmis sp., Nannochloropsis sp., mixed algae (Tetraselmis sp. and Nannochloropsis sp.), yeast (Saccharomyces cerevisiae) and micro-pallet (Dindings Shrimp Feed 2203) diet that acts as the control. Copepods were enriched twice per day (10:00 a.m. and 4:00 p.m.) at 500 mg L⁻¹ ¹ concentration for each diet. Postlarvae were fed at 6-7% of body mass per day. The growth, survival and proximate analysis of M. rosenbergii larvae were conducted to compare effectiveness of each consecutive diet. Higher specific growth rate result of M. rosenbergii was observed when fed with copepod enriched with mixed algae 4.96±0.02% and the lowest was when fed with copepod enriched with Nannochloropsis sp. 4.58±0.03%. The greatest survival rate for M. rosenbergii was seen when fed with copepod enriched with mixed algae 95.16±1.04% compared to that fed with copepod enriched with micropallet 89.16±1.89%. Among used enrichments, mixed algae produced the best result compared with other enrichment, especially in the proximate analysis of the M. rosenbergii larvae. Mean value of protein, lipid, moisture and ash of M. rosenbergii were found to be $70.45\pm1.22\%$, $8.55\pm1.11\%$, $15.30\pm0.16\%$, and $9.05\pm0.59\%$, respectively. The results indicated that it is highly recommended to use copepod-enriched with mixed algae to feed giant freshwater prawn larvae (PL5 and above).

Keywords: Copepod, Giant freshwater prawn, Enrichment, Proximate analysis, Microalgae

¹ Faculty of Fisheries and Food Sciences, University Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia.

²⁻Institute of Tropical Biodiversity and Sustainable Development, University Malaysia Terengganu, 21300 Kuala Nerus, Terengganu, Malaysia.

³⁻Institute of Tropical Aquaculture and Fisheries, University Malaysia Terengganu, 21300 Kuala Nerus, Terengganu, Malaysia.

^{*} Corresponding author's Email: nadiah.rasdi@umt.edu.my

Introduction

About 80% of zooplankton represents copepods and it is food source for many marine fish larvae in the ocean. Due to several factors that are important for and reproduction copepod growth related food quality and environmental conditions, copepods are difficult to be mass cultured. Food quality is the main factor regulating copepod nutrition and production (Drillet et al., 2011). Copepods are zooplanktons that can be enriched by providing better nutrition and are very suitable to use as starter feed to marine or freshwater fish and shellfish in hatchery (Rasdi and Oin, 2018a: Rahmati et al., 2020). This makes copepods a suitable group as live feed in aquaculture.

Copepods are live foods containing high levels of fatty acids needed by many larvae of marine fish species since it is the best primarily consumed food by small larvae of aquatic organisms in the ocean. Copepod is superior in nutritional terms compared with other live foods species (Ajiboye et al., 2011). Nutritional content of copepod largely depends on its culture medium, especially on the quality of feed provided to them (Agadjihouede et 2014). The dry-weight al.. concentration of protein is much higher in copepods than in other zooplankton species which is very important for growth and survival of larvae. Other live foods even after enrichment are still incomparable to copepods in terms of nutritional quality (Hamre et al., 2013).

In general, Artemia gives low growth and survival rate compared to marine copepod that boost growth and survival of tiger shrimp Penaeus monodon (Ananthi et al., 2011). Marine copepods are nutritionally superior for marine fish and crustacean larvae, and are valuable source of proteins, lipids, carbohydrates, amino acids, fatty acids and enzymes (amylase, protease, exonuclease and esterase), which are essential for larval digestion (Kumar et al., 2017). Artemia nauplii have a low nutritional content compare to copepod even though Artemia is easy to produce in large numbers. Previous studies have compared the growth and survival of seahorse fed with enriched-copepod and enriched-Artemia with Super Selco. Growth and survival of seahorses were significantly higher when fed copepod nauplii (Zhang et al., 2015). Based on Payne and Rippingale (2000), Artemia nauplii showed a negative result and copepod nauplii were well digested by juvenile seahorses compared to Artemia nauplii. Enriching copepods with high HUFAs make them more suitable for feeding young fish or prawn larvae (Sargent et al., 1997; Naman et al., survival М. 2021). Hence, rosenbergii may improve by feeding them enriched-copepod.

M. rosenbergii post-larvae culture is still an important and recent activity in many countries, where there is little research on giant freshwater prawn nutrition and methods to improve the growth and survival rates through enrichment and production of live food. The latest study has shown that highly

unsaturated fatty acids (HUFA) are very important for many fish species (Chang Southgate, 2001). Nutritional and composition and environmental conditions are factors that affect growth and survival rate of fish larvae in many cases. Production of digestive enzymes develops an efficient digestive capacity in the fish body (Ikpegbu et al., 2011). Availability of high nutrition, suitable consumed diets and efficiently digested diet would promote the success of larval rearing in a particular fish species (Giri et al., 2002). In general, higher proteins, lipids and highly unsaturated fatty acids are required for early stages of T. tambroides for growth, survival and neural development. The study on the feeding of freshwater larvae using Artemia, Moina and Daphnia have been done due to their height in protein sources for larval development, and their broad spectrum of digestive enzymes and fatty acids (Macedo and Pinto-Coelho, 2001; Yuslan et al., 2021). Independently of their high nutritional value, other than easy to digest, live feed is the best starter feed as it is easy to detect and capture by fish or prawn larvae due to its swimming movements in the water column (Conceição et al., 2010).

It is important to use live food to improve the success of larval fish rearing. In most cases, live food such as copepod, *Artemia*, and rotifers provide adequate nutrients for finfish and crustacean larvae (Drillet *et al.*, 2008). However, for the good improvement of growth and survival of larvae reared in

the hatchery, commercially live used food nutrition such as rotifer and *Artemia* are still not enough for the size and nutrition required by certain species of fish and prawn larvae to enhance their growth performance (Solgaard *et al.*, 2007). More specifically, low survival and high deformity will occur with inadequate nutritional supply provided during these early stages of development (Rajkumar and Kumaraguru Vasagam, 2006).

At present, live feeds such as Artemia and rotifers have been used widely in the culture of fish larvae where enrichment of Artemia and rotifers is commonly done using the oil emulsion method containing various amounts of DHA ethyl ester and EPA ethyl ester (Samat et al., 2020). However, some fish larvae require specific nutrients such as a higher content of PUFA. Even after artificial enrichment using emulsified oils and other common enrichment products. rotifers and Artemia are nutritionally inadequate and cannot provide adequate nutrition for some fish larvae compared to enriched copepod (Rasdi et al., 2016).

The consumption of copepods can improve the survival of prawn larvae since copepod is better regarding its appropriate size for ease of consumption by prawn larvae (Drillet *et al.*, 2006). The nutritional content, digestibility, and size of the offered food are the factors for successful rearing of early development stages of fish or prawn larvae because proper

nutrition is considered as the prime factor in providing better digestion, growth and survival of fish in the ocean (Lavens and Sorgeloos, 1996). The aim of this study was to determine the best enrichment formula for cyclopoid copepod (*Oithona* sp.) that could stimulate better growth and survival to *M. rosenbergii* post-larvae. Besides, this study was designed to evaluate the proximate composition of copepod and *M. rosenbergii* post-larvae after feeding with enriched-copepod.

Materials and methods

Sampling and copepod stock culture Oithona sp. was sampled from the lagoon of Setiu Wetland, located in the state of Terengganu, Malaysia (5'37''30.64N, 102'47"15.08E). Habitat condition factors, such as temperature and salinity of sampling area were recorded. Normal habitat condition for optimal temperature being at 26-27°C and the optimal salinity at 24-26 ppt (Kassim and Busra, 2012) were maintained until the end of the experiment. The copepod was cultured and sustained at hatchery of Universiti Malaysia Terengganu (UMT). As a stock culture, the copepod was first cultured in a 1000L tank. Daily observation and feeding were provided. Apart from that, once in two days, 30% of the water was changed in order to avoid any contamination and toxicities from food given.

Experimental design and diets preparation

The diets for copepod were prepared. These adult copepods were isolated from stock and transported to other culture tanks (500 L) and fed for one generation with the specified food before data was collected to remove any previous consequences of regiments on copepods (Rasdi and Oin, 2018b). The copepod culture was closely monitored for egg reproduction and hatching of new nauplii. Next, the copepod was harvested and placed into four enrichment tanks for 6 hours before fed to the prawn larvae (Esmaeili Fereidouni et al., 2013). Copepodites of the copepod were enriched using four different treatments. Four treatments consisted of **Tetraselmis** sp., Nannochloropsis sp., mixed algae (Tetraselmis sp., and Nannochloropsis sp.), yeast (Saccharomyces cerevisiae) and micro-pallet (Dindings Shrimp Feed, 2203) diet that acted as the control (Lananan et al., 2013; Kasan et al., 2020). Microalgae were cultured at plankton lab of UMT and were upscaled and maintained. The quality of the medium (Conway medium) used for cultivation was to determine the growth performance of microalgae (Lam and Lee, 2012). The concentration of the used algal diets was 500 mg L⁻¹ which is equivalent to 2×10^7 algal cells mL⁻¹ (Kassim and Busra, 2012). For mixed algae, which are Tetraselmis sp. + Nannochloropsis sp., 1:1 ratio was used. Copepods were fed yeast at the concentration of 500 mg L⁻¹ (Paray and Al-Sadoon, 2016).

Culture of M. rosenbergii post-larvae Post-larvae of M. rosenbergii (PL5) were purchased from a hatchery in Aquatrope UMT, located in Kuala Terengganu, Malaysia. The post-larvae were transported to the hatchery for test feeding. For feeding experiments, 200 M. rosenbergii post-larvae (PL5) were cultured in each 25L aquarium with 20L of water. Gentle aeration was provided for each tank using the aeration system that was prepared in the hatchery. Five dietary treatments with three replicates each were used in this study to test efficacy of the enrichedcopepods to the prawn larvae. 50 adult copepods (individual(s) /larvae) were fed twice per day (10:00 a.m. and 4:00 p.m.) with different diets for 30 days (Chakraborty et al., 2001; Redzuari et al., 2012). The water exchange was done for 50% every week and the water quality were maintained at the temperature of 26 -27°C, salinity of 0-6 ppt and pH of 3-5. The post-larvae were fed at 6-7% of body mass per day (Balaji *et al.*, 2002).

Growth and survival of M. rosenbergii post-larvae analysis

The body weight of post-larvae was measured every six days. Ten post-larvae were sampled randomly for every treatment and replicate. Samples were weighed to calculate the growth rate of post-larvae. Specific Growth Rate (SGR) was calculated using the formula: (ln Wt – ln Wi x 100) / t. Where, Wt=final weight (g), Wi=initial weight (g), t=time (days).

For the survival rate, mortality of postlarvae was recorded every day. Survival rate was calculated using the formula:

Survival rate = [(Total number of surviving fish) / Total number of initial prawn larvae density in the tank] x 100% (Mandal, 2010).

Proximate analysis of copepod and M. rosenbergii post-larvae

Weight of the sample was measured. Samples were packed in separate polyethene bags, labelled and stored in freezer at -20°C for further laboratory analyses. Protein, lipid, moisture and ash were analysed at Department of Fisheries, laboratory of Universiti Malaysia Terengganu. Methods used are as follows:

Estimation of Protein:

Total protein content of fish was identified by Micro-Kjeldhal method (Kirk *et al.*, 1991).

Percentage of protein was calculated by multiplying percentage of N2 (%) with an empirical factor of 6.25. Protein (%) = % of total amount of N2×6.25.

Estimation of Lipid:

Total lipid content of prawn was estimated by method of Bligh and Dyer (1959).

Percentage of lipid calculated by: Percentage of fat (%)=(weight of the extract/weight of the sample) \times 100.

Estimation of moisture:

Moisture content of the prawn, *Macrobrachium rosenbergii* was estimated using hot air oven method (Jain and Singh, 2000).

Percentage calculated by: Percentage (%) of moisture=(weight loss/actual weight of the sample)×100.

Estimation of Ash:

Ash content of the prawn was evaluated by the method of Association of Official Analytical Chemists (AOAC, 1991).

Percentage of ash content was estimated by: Ash (%)=weight of ash/weigh of sample × 100. *Data analysis*

Data were presented as mean±standard deviation (SD). All the data were collected throughout the experiment and analysed using a one-way analysis of variance (ANOVA). This was in order to compare growth, survival and composition proximate of M. rosenbergii post-larvae and compare the effectiveness of each consecutive Differences were diet. considered significant at p<0.05 level. When the main treatment effect was significant,

post-hoc comparisons were made using Tukey's test. All data were tested for normality, homogeneity and independence to satisfy assumptions for ANOVA.

Results

Specific Growth Rate

Figure 1 shows the overall specific growth rate of M. rosenbergii postlarvae in each treatment after being cultured for 30days. The highest growth rate of *M. rosenbergii* post-larvae occurred when larvae fed with copepod enriched with mix algae (4.96±0.02%), compared with other enrichments such as Tetraselmis sp. (4.75±0.16%), yeast $(4.79\pm0.05\%)$ micro-pallet and $(4.66\pm0.16\%)$. Nannochloropsis (4.58±0.03%) showed the lowest larvae growth. However, all treatments were significantly different among each other (p<0.05).

Survival Rate

There was significant difference among the dietary treatments (p<0.05). M. rosenbergii post-larvae fed with copepod enriched mix algae showed the highest survival rate compared to other diets of *Tetraselmis* sp. (93.33 \pm 2.51%), Nannochloropsis sp. (93.00 \pm 1.03%) and yeast (90.66 \pm 1.75%, Fig. 2).




Figure 1: Specific Growth Rate (SGR) of *Macrobrachium rosenbergii* post-larvae fed with different enriched copepods and diets. Different small letters indicate significant difference among treatment, error bars show standard deviation.




Figure 2: Survival rate of *Macrobrachium rosenbergii* post-larvae fed with different enriched copepods and diets. Different small letters indicate significant difference among treatments, error bars show standard deviation.

However, the lowest survival rate was recorded when M. rosenbergii being fed with copepod enriched micro-pallet (89.16±1.89%). Therefore, all treatments were significantly different among each other (p<0.05).

Proximate Analysis of M. rosenbergii Post-larvae Table 1 shows the results for proximate analysis of M. rosenbergii post larvae. Post-larvae fed with copepod enriched with mix algae showed the highest content of protein, lipid, moisture and ash $(70.45\pm1.22\%, 8.55\pm1.11\%, 15.30\pm0.16\%$ and $9.05\pm0.59\%)$, respectively.

Table 1: Body composition of *M. rosenbergii* fed with different enrichment. Different small letters indicate significant difference among treatments.

Body Composition	Enrichment	Mean±SD (%)
Protein	Tetraselmis sp.	65.66 ± 1.09 ^a
	Nannochloropsis sp.	$67.62 \pm 0.78^{a,b}$
	Mix algae (<i>Tetraselmis</i> sp. + <i>Nannochloropsis</i> sp.)	70.45 ± 1.22 °
	Yeast	66.28 ± 0.92 a
	Micro pellet	69.97 ± 2.18 °
Lipid	Tetraselmis sp.	$7.82 \pm 1.00^{a, b}$
	Nannochloropsis sp.	6.21 ± 0.63^{a}
	Mix algae (<i>Tetraselmis</i> sp. + <i>Nannochloropsis</i> sp.)	8.55 ± 1.11^{b}
	Yeast	6.53 ± 0.45 a,b
	Micro pellet	$8.43 \pm 0.52^{\ b}$
Moisture	Tetraselmis sp.	10.54 ± 1.08 a
	Nannochloropsis sp.	13.85 ± 6.48 a
	Mix algae (<i>Tetraselmis</i> sp. + <i>Nannochloropsis</i> sp.)	15.30 ± 0.16 a
	Yeast	10.53 ± 0.92 a
	Micro pellet	14.42 ± 0.97 a
Ash	Tetraselmis sp.	8.60 ± 0.32^{a}
	Nannochloropsis sp.	8.15 ± 0.64^{a}
	Mix algae (<i>Tetraselmis</i> sp. + <i>Nannochloropsis</i> sp.)	9.05 ± 0.59^{a}
	Yeast	8.44 ± 0.24^{a}
	Micro pellet	8.86 ± 1.44^{a}

Proximate Analysis of Diets, Unenriched Copepod, and Enriched Copepod.

Based on previous studies, the proximate of different used feeding were tabulated in order to compare with the results of the present study. According to Table 2, micro pellet (53.8%) contained the highest protein content compared with other diets, followed bv yeast (46.11%). Nannochloropsis sp. (35.8%), mix diet of microalgae (33.7%), and Tetraselmis sp. (25.7) respectively. Previous results on the proximate analysis of Oithona sp. (Table 3), showed greater variations in protein content with 69.24%. The lipid content was 15.36% and moisture and ash contents were 83.07% and 3.96% respectively (Santhanam and Perumal, 2012). However, in observation (Table 4), protein and lipid contents of enriched copepod were comparatively higher than unenriched copepod. The proximate analysis of copepod observed that copepod enriched with mix algae contained the highest content of protein and lipid (71.54±1.03% and $10.48\pm2.22\%$) respectively. Compared to other diets, micro pellet showed the lowest content of protein, 63.15±0.52%, and lipid, Therefore, $6.05\pm4.55\%$. the body composition of copepod depends on the type of provided enrichment (p>0.05, Table 4).

Table 2: Proximate analysis of different enrichments used to enrich copepod, Oithona sp.

Body composition	Protein	Lipid	Moisture	Ash	Reference
Tetraselmis sp.	25.7	9.4	nr	nr	
Nannochloropsis sp.	35.8	11.7	nr	nr	Arkronrat <i>et al.</i> , 2016
Mix diet (<i>Tetraselmis</i> sp. + <i>Nannochloropsis</i> sp.)	33.7	10.8	nr	nr	
Yeast	46.11	1.51	4.37	10.20	Olvera-Novoa <i>et</i> al., 2002
Micro pellet	53.8	30.5	nr	8.4	Rust et al., 2015

^{*}nr: data was not recorded

Table 3: Body composition of unenriched cyclopoid copepod, Oithona sp.

Body Composition	Mean ± SD (%)	Reference	
Protein	69.24 ± 0.070		
Lipid	5.36 ± 0.103	Conthonorm and Dominal 2012	
Moisture	83.07 ± 0.075	Santhanam and Perumal, 2012	
Ash	3.962 ± 0.081		

Table 4: Body composition of copepod-enriched with different enrichments. Different small letters indicate significant difference among treatment (p<0.05).

Body Composition	Enrichment	Mean±SD (%)
Protein	Tetraselmis sp.	66.36 ± 2.47 a,b
	Nannochloropsis sp.	$67.89 \pm 1.20^{\text{ b,c}}$
	Mix algae (<i>Tetraselmis</i> sp. + <i>Nannochloropsis</i> sp.)	$71.54 \pm 1.03^{\text{ b,c}}$
	Yeast	68.95 ± 0.52 °
	Micro pellet	63.15 ± 0.31^{a}
Lipid	Tetraselmis sp.	8.55 ± 1.11^{a}
	Nannochloropsis sp.	$7.06 \pm 6.90^{\text{ a}}$
	Mix algae (<i>Tetraselmis</i> sp. + <i>Nannochloropsis</i> sp.)	10.48 ± 2.22^{a}
	Yeast	6.52 ± 0.45^{a}
	Micro pellet	6.05 ± 4.55 a
Moisture	Tetraselmis sp.	10.54 ± 1.08 a,b
	Nannochloropsis sp.	13.85 ± 6.48 a,b
	Mix algae (<i>Tetraselmis</i> sp. + <i>Nannochloropsis</i> sp.)	$8.99 \pm 3.58^{\text{ a}}$
	Yeast	10.53 ± 0.92 a,b
	Micro pellet	$14.42 \pm 0.97^{\ b}$
Ash	Tetraselmis sp.	11.90 ± 5.59^{a}
	Nannochloropsis sp.	4.08 ± 0.31^{a}
	Mix algae (<i>Tetraselmis</i> sp. + <i>Nannochloropsis</i> sp.)	9.13 ± 0.93^{a}
	Yeast	7.25 ± 3.08^{a}
	Micro pellet	9.08 ± 1.30^{a}

Discussion

Post-larval stages of many shrimp require feed with high species nutritional content for development and continuous rearing in a hatchery and it is important to produce live food with high nutritional quality for feeding. The nutritional requirements of the copepods as a starter diet (Singh et al., 2019) for Macrobrachium rosenbergii have to be fully understood in order to sustain growth and ability to be produced on massive scale. The results showed that microalgae were the best enrichment for copepod to improve the growth and survival rate of *Macrobrachium rosenbergii* postlarvae.

From the results (Fig. 1), it is observed that the Specific Growth Rate (SGR)

and survival rate of post-larvae of M. rosenbergii were better fed with copepod-enriched with mix algae compared with other enrichments and such **Tetraselmis** as sp., Nannochloropsis sp., and yeast. The provided diets affected Specific Growth Rate (SGR) of M. rosenbergii postlarvae (P=0.03, Fig. 1). Apart from all treatments, micro pellet showed the lowest growth of post-larvae. To improve the nutritional content of copepod, algae are primary food for the advancement of copepod (Rasdi et al., 2018). Several reports have shown P. monodon larvae fed with a mixed diet ofmuelleri Chaetoceros and Thalassiosira suecica performed well (D'Souza and Blackburn, 2013). Consistent with the literature, this research found that the survival and development of M. rosenbergii postlarvae were better on copepod-enriched mixed algae as a diet.

From the results (Fig. 2) it is clear that post-larvae of M. rosenbergii had high survival rate when fed with copepod-enriched mix algae. marine algae Tetraselmis sp. and Nannochloropsis sp. are generally high in EPA that can enhance growth and survival rate of many species (Rasdi and Qin, 2015). This study supports evidence from previous observations by De Araujo and Valenti (2007) that suggest that use of Artemia-enriched with microalgae during larval development of M. americanum can enhance growth and improves survival due to better nutrition after enrichment. As stated by Rasdi and Qin (2018a),

improvement of the nutritional content of copepod enriched with microalgae can be used to increase the survival rate of post-larvae in aquaculture. Besides, the lowest result for survival rate was post-larvae feed with micro-pallet. The digestibility and nutritional quality of formulated diets were not as efficient as a starter food for fish larvae (Carneiro et al 2003). The nutritional composition in pellet was not enough to cover the required levels by larvae for growth. The protein composition in a micro-pallet was lower than the content of protein in enriched-copepods such as Tetraselmis sp. and Nannochloropsis sp. Therefore, use of pallet as starter feeding to prawn larvae will also indirectly increase ammonia level in the water (Coutteau and Mourente, 1997). High ammonium content in the water can be toxic for aquatic animals. It is difficult for aquatic organisms to sufficiently excrete the toxicant, leading to toxic build up in internal tissues and blood that can cause mortality.

Proximate analysis was done to compare the body composition of *M. rosenbergii* post-larvae after fed with differently enriched copepod. Post-larvae fed with copepod that was enriched with mix algae showed highest percentage of protein compared to other enrichments. Then, the post-larvae fed with copepod-enriched *Tetraselmis* sp. showed the lowest percentage of protein. Previous early studies on the rearing of *M. rosenbergii* showed an optimum dietary protein level ranged from 15 to 80% (Mai *et al.*, 2009). It is supported by Ben-Amotz *et al.*, (1987)

that studied on Penaeus japonicus. They concluded that the diet of less than 10% protein was low in HUFA content that supported growth and survival of Penaeus japonicus larvae. Protein content below 15% can cause mortality during rearing of post-larvae as protein is the first and foremost factor that has to be considered for formulating a fish feed. However, excess protein in the diet may also inhibit growth (Lim et al., 2003). The excess protein would be metabolized by the shrimp as a source of energy, and nitrogen will be excreted as ammonia. These studies are in agreement with those obtained by the previous, 70% content of protein in body composition of M. rosenbergii is regarded to be the best result when post-larvae fed with copepod-enriched mix algae.

Based on a previous study of Hilton et al. (1984), the freshwater prawn requires 6% to 10% lipid. Furthermore, based on that study, they also observed that lower lipid content (>10%) can causes reduction in growth for postlarvae of M. rosenbergii. Therefore, based on proximate analysis that is done, the shrimp that was fed with high lipid content food sources had significantly more lipid and carbohydrate than others. In addition, this is also supported by findings, which report that proximate analysis of P. monodon larvae fed with mix algae that contain high lipid content produce highest lipid and carbohydrate content (D'Souza and Loneragan, 1999). Protein and lipid were found to be the major biochemical component in the cultured copepod as a feeding for larvae and crustaceans (Santhanam et al., 2004). Copepods fed with manipulated diets were found to contain remarkable biochemical characteristics (Eviemo et and were capable of 2003) stimulating appetites and feeding in larvae (Olivotto et al., 2011). Based on the result (Table 4), copepod fed with single diet of Nannochloropsis sp. showed the lowest percentage of lipid relatively produce content, population growth and reproductive performance, which was mainly due to derived low DHA and other HUFAs content of copepods (Rasdi and Qin, 2015). Therefore, the best result for lipid is prawn fed with copepodenriched mix algae (Tetraselmis sp. + Nannochloropsis sp.). Copepodenriched mix algae contain high in lipid. Thus, it makes prawn post-larvae absorb enough lipids for their energy and growth. Roughly the enrichment of copepod using mix algae suitable to use in aquaculture, increased the lipid content for improvement of growth and survival.

In the study, the moisture and ash content were maximum in body composition of post-larvae fed with copepod-enriched algae. mix Meanwhile, the minimum percentage of moisture was obtained in post-larvae copepod-enriched fed with yeast. Therefore. for metabolic process, minerals such as, calcium, phosphorus, magnesium, sodium, potassium, chloride, and sulfur are inorganic elements required in the body. Calcium is required for exoskeleton formation, muscle contraction, and osmoregulation (Moss *et al.*, 2019). Shrimp are able to absorb calcium directly from the water, and shrimps living in seawater do not need calcium supplements in the diet (Davis and Gatlin, 1996).

In conclusion, growth, survival and proximate composition of M. rosenbergii post-larvae were affected by enrichment of copepod and provided mix diets. From the result, mix algae (Tetraselmis sp. + Nannochloropsis sp.) were the best enrichment for copepod to increase the growth and survival rate of M. rosenbergii post-larvae, and for obtaining a better feed conversion ratio for larvae rearing. It is followed by Tetraselmis sp., Nannochloropsis sp., micro-pallet and yeast which showed the slowest growth and survival for M. rosenbergii post-larvae. The micropallet showed the lowest survival rate on M. rosenbergii post-larvae among the dietary treatments. In general, feeding on live microalgae and yeast can improve the mass culture of However, selection copepods. suitable microalgae species that are high in nutrients is important, as the nutritional profile differs among algal species. Many types of algae have been offered to copepods as food, but there is a need to identify algae with high content of essential nutrients, such as fatty acids (Rasdi and Qin, 2018c). Copepods fed microalgae have shown positive results on growth, reproduction biochemical and composition of copepods depending on the algal

nutritional content fed to copepods. The mix algae used as enrichment to copepod can reduce the cost in aquaculture since it can increase the growth and survival of larvae rearing. However, other enrichment and diet still can be used in larvae rearing culture.

Acknowledgment

Writing of this article was supported by Fundamental Research Grant Scheme (FRGS) provided by Ministry of Higher Education of Malaysia under grant Vote No. 59530, in order to generate new ideas and methodology for development of live food nutrition in Malaysian aquaculture industry.

References

Agadjihouede, H., Montchowui, E., Montcho, S.A., Bonou, C.A. and Laleye, P.A., 2014. Growth and development of three species of the zooplankton (Brachionus calyciflorus, Moina micrura and Thermocyclops sp.) breeding on poultry dropping in mixed condition in tanks. International Journal of Fisheries and Aquatic Studies, 2(1D), 189-196.

Ajiboye, O.O., Yakubu, A.F., Adams, T.E., Olaji, E.D. and Nwogu, N.A., 2011. A review of the use of copepods in marine fish larviculture. *Reviews in Fish Biology and Fisheries*, 21, 225-246. DOI: 10.1007/s11160-010-9169-3.

Ananthi, P., Santhanam, P., Nandakumar, R., Ananth, S., Jothiraj, K., Dinesh Kumar, S.,

- Balaji Prasath. В. and Jayalakshmi, T., 2011. Production and utilization of marine copepods as live feed for larval rearing of tiger shrimp Penaeus monodon special emphasis on astaxanthin enhancement. Indian Journal of *Natural Sciences*, 2(8), 494-503. DOI:10.1.1.734.9130
- AOAC, 1991. Official methods of analysis of the association of official analytical chemists Volume 2, 15th edition. Helrich K. editor. Association of Official Analytical Chemists, Inc. Arlington, Virginia, USA. pp. 685-1298.
- Arkronrat, W., Deemark, P. and Oniam. V.. 2016. Growth performance and proximate composition of mixed cultures of marine microalgae (Nannochloropsis Tetraselmis sp.) with and monocultures. Songklanakarin Journal of Science and Technology, 38(1), 1-5.
- Balaji, K., Sahu, N.P. and Tripathi, S.D., 2002. Effect of different protein energy ratio in isocaloric diet on growth and survival of post larvae of *Macrobrachium rosenbergii* (De Man). Aquaculture Nutrition Workshop Proceeding, 7-10.
- Ben-Amotz, A., Fishler, R. and Schneller, A., 1987. Chemical composition of dietary species of marine unicellular algae and rotifers with emphasis on fatty acids. *Marine Biology*, 95, 31-36. DOI: 10.1007/BF00447482.

- Bligh, E.G. and Dyer, W.J., 1959. A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology*, 37(8), 911-917. DOI: 10.1139/o59-099.
- Carneiro, P.C.F., Mikos, J.D., Schorer, M. Campagnoli, O.F.P.R. and Bendhack, F., 2003. Live and formulated diet evaluation through initial growth and survival of jundiá larvae, *Rhamdia quelen. Scientia Agricola*, 60(4), 615-619. DOI: 10.1590/S0103-90162003000400002.
- Chakraborty, D., Gopikrishna, M. and Altaff, K., 2001. Study on the growth in postlarvae of *Macrobrachium rosenbergii* (de Man) on a copepod and a cladocerans live food. *Journal of Aquatic Biology*, 16(1-2), 81-85.
- Chang, M. and Southgate, P.C., 2001. Effects of varying dietary fatty acid composition on growth and survival of seahorse, *Hippocampus* sp., juveniles. *Aquarium Science and Conservation*, 3, 205-214. DOI: 10.1023/A:1011363807074.
- Conceição, L.E.C., Yúfera, M., Makridis, P., Morais, S. and Dinis, M.T., 2010. Live feeds for early stages of fish rearing. *Aquaculture Research*, 41(5), 613-640. DOI: 10.1111/j.1365-2109.2009.02242. x.
- Coutteau, P. and Mourente, G., 1997. Lipid classes and their content of n-3 highly unsaturated fatty acids (HUFA) in *Artemia franciscana* after hatching, HUFA-enrichment and

- subsequent starvation. *Marine Biology*, 130(1), 81-91. DOI: 10.1007/s002270050227.
- D'souza, F.M.L. and Loneragan, N.R., 1999. Effects of monospecific and mixed-algae diets on survival, development and fatty acid composition of Penaeid prawn (Penaeus spp.) larvae. Marine Biology, 133(4), 621-633. DOI: 10.1007s002270050502.
- D'souza, M.R. and Blackburn, S.I.,
 2013. Live microalgae as feeds in aquaculture hatcheries. In: Advances in aquaculture hatchery technology.
 Brown M.R. and Blackburn S.I. editors, Woodhead Publishing,
 Cambridge, UK. 117-156, 157e-158e. DOI: 10.1533/9780857097460.1.117.
- **Davis, D.A. and Gatlin, D.M.III., 1996.** Dietary mineral requirements of fish and marine crustaceans. *Reviews in Fisheries Sciences*, 4(1), 75-99. DOI: 10.1080/10641269609388579.
- **De Araujo, M.C. and Valenti, W.C., 2007.** Feeding habit of the Amazon River prawn *Macrobrachium amazonicum* larvae. *Aquaculture*, 265(**1-4**), 187-193. DOI: 10.1016/j.aquaculture.2007.01.016.
- Drillet, G., Jørgensen, N.O.G., Sørensen, T.F., Ramløv, H. and Hansen, B.W., 2006. Biochemical technical observations and supporting the use of copepods as live feed organisms in marine larviculture. Aquaculture Research, DOI: 37(8), 756-772. 10.1111/j.1365-2109. 2006.01489.x.

- Drillet, G., Jepsen, P.M., Højgaard, J.K., Jørgensen, N.O.G. and Hansen, B.W., 2008. Strain-specific vital rates in four *Acartia tonsa* cultures II: life history traits and biochemical contents of eggs and adults. *Aquaculture*, 279(1-4), 47-54. DOI: 10.1016/j.aquaculture. 2008.04.010.
- Drillet, G., Frouël, S., Sichlau, M.H., Jepsen, P.M., Højgaard, J.K., Joarder, A.K. and Hansen, B.W., 2011. Status and recommendations on marine copepod cultivation for use as live feed. *Aquaculture*, 315(3-4), 155-166. DOI: 10.1016/j.aquaculture.2011.02.027.
- Esmaeili Fereidouni, A., Fathi, N. and Khalesi, M.K., 2013. Enrichment of *Daphnia magna* with canola oil and its effects on the growth, survival and stress resistance of the *Caspian kutum (Rutilus frisii kutum)* larvae. *Turkish Journal of Fisheries and Aquatic Sciences*, 13(1), 119-126. DOI: 10.4194/1303-2712-v13-1-15.
- **Evjemo, J.O., Reitan, K.I. and Olsen, Y., 2003.** Copepods as live food organisms in the larval rearing of halibut larvae (*Hippoglossus hippoglossus* L.) with special emphasis on the nutritional value. *Aquaculture*, 227(1), 191-210. DOI: 10.1016/S0044-8486(03)00503-9.
- Giri, S.S., Sahoo, S.K., Sahu, B.B., Sahu, A.K., Mohanty, S.N., Mukhopadhyay, P.K. and Ayyappan, S., 2002. Larval survival and growth in *Wallago attu* (Bloch and Schneider), Effects of light,

- photoperiod and feeding regimes. *Aquaculture*, 213(**1-4**), 151-161. DOI: 10.1016/S0044-8486(02)00012-1.
- Hamre, K., Yúfera, M., Rønnestad, I., Boglione, C., Conceição, L.E.C. and Izquierdo, M., 2013. Fish larval nutrition and feed formulation: knowledge gaps and bottlenecks for advances in larval rearing. *Reviews in Aquaculture*, 5(s1), S26-S58. DOI: 10.1111/j.1753-5131. 2012.01086.x.
- Hilton, J.W., Harisson, K.F. and Silnger, S.J., 1984. A semi-purified test diet for *Macrobrachium rosenbergii* and the lack of need for supplemental lecithin. *Aquaculture*, 37(3), 209-215. DOI: 10.1016/0044-8486(84)90153-4.
- **Ikpegbu, E., Nlebedum, U., Nnadozie, O. and Agbakwuru, I., 2011.** Fast green FCF or Ehrlich's hematoxylin as counterstain to periodic acid-Schiff reaction: A comparative study. *Histologic*, 54(2), 29-30.
- Jain, P.C. and Singh, P., 2000. Moisture determination of jaggery in microwave oven. *Sugar Tech*, 2, 51-52. DOI: 10.1007/BF02945760.
- Kasan, N.A., Hashim, F.S., Haris, N., Zakaria, M.F., Mohamed, N.N., Abd Wahid, M.E., Katayama, T., Takahashi, K. and Jusoh, M., 2020. Isolation of freshwater and marine indigenous microalgae species from Terengganu water bodies for potential uses as live feeds aquaculture in industry. International Aquatic Research,

- 12(**1**), 74-83. DOI: 10.22034/IAR(20).2020.671730.
- Kassim, Z. and Busra, I., 2012.
 Culture of harpacticoid copepods: understanding the reproduction and effect of environmental factors. In: *Aquaculture*, Muchlisin Z. editor. InTech Open Access Book. pp. 343-357. DOI: 10.5772/28373. Available from: https://www.intechopen.com/books/aquaculture/c opepods-in-aquaculture.
- **Kirk, R.S., Sawyer, R. and Egan, H., 1991.** Pearson's composition and analysis of foods, 9th edition. Addison-Wesley Longman Ltd, Boston, USA. 708 P.
- Kumar, S.D., Santhanam, P., Ananth, S., Kavivarasan, M., Nithya, P., Dhanalakshmi, B., Park, M.S. and Kim, M.K., 2017. Evaluation of suitability of wastewater-grown microalgae (Picochlorum maculatum) and copepod (Oithona rigida) as live feed for white leg shrimp Litopenaeus vannamei postlarvae. Aquaculture International, 393-411. DOI: 25(1), 10. 1007/s10499-016-0037-6.
- Lam, M.K. and Lee, K.T., 2012.

 Potential of using an organic fertilizer to cultivate *Chlorella vulgaris* for biodiesel production. *Applied Energy*, 94, 303-308. DOI: 10.1016/j.apenergy. 2012.01.075.
- Lananan, F., Jusoh, A., Ali, N., Lam, S.S. and Endut, A., 2013. Effect of Conway medium and f/2 medium on the growth of six genera of South China Sea marine microalgae.

- *Bioresource Technology*, 141, 75-82. DOI:
- 10.1016/j.biortech.2013.03.006.
- Lavens, P. and Sorgeloos, P., 1996.

 Manual on the production and use of live food for aquaculture. FAO Fisheries Technical Paper No. 361.

 Food and Agriculture Organization of the United Nations, Rome, Italy.
- **Lim, L.C., Dhert, P. and Sorgeloos, P., 2003.** Recent developments in the application of live feeds in the freshwater ornamental fish culture. *Aquaculture*, 227(1-4), 319-331. DOI: 10.1016/S0044-8486(03)00512-x.
- C.F. and Pinto-Coelho, Macedo, **R.M.**, 2001. Nutritional status response of Daphnia laevis and Moina micrura from a tropical reservoir to different algal diets: Scenedesmus quadricauda and Ankistrodesmus gracilis. Brazilian Journal of Biology, 61(4), 555-562. DOI: 10.1590/ s1519-69842001000400005.
- Mai, M.G., Engrola, S., Morais, S., Portella, M.C., Verani, J.R., Dinis, M.T. and Conceição, L.E.C., 2009. Co-feeding of live feed and inert diet from first-feeding affects *Artemia* lipid digestibility and retention in Senegalese sole (*Solea senegalensis*) larvae. *Aquaculture*, 296(3-4), 284-291. DOI: 10.1016/j.aquaculure.2009.08.024.
- Mandal, S.C., Sahu, N.P., Kohli, M.P.S., Das, P., Gupta, S.K. and Munilkumar, S., 2010.

 Replacement of live feed by formulated feed: effect on the growth

- and spawning performance of Siamese fighting fish (*Betta splendens*, Regan, 1910). *Aquaculture Research*, 41(**11**), 1707-1716. DOI: 10.1111/j.1365-2109.2010.02564.x.
- Moss, A.S., Ishikawa, M., Koshio, S., Yokoyama, S. and Dawood, M.A.O., 2019. Effects of different levels of marine snail shells in the diets of juvenile kuruma shrimps *Marsupenaeus japonicus* as a source of calcium. *North American Journal of Aquaculture*, 81(1), 55-66. DOI: 10.1002/naaq.10066.
- Naman, N., Kassim, Z. and Rasdi, N.W., 2021. The effect of copepod enriched-vegetable based diet on Giant **Tiger** Prawn (Penaeus monodon) post-larvae. In IOPConference Series: Earth and Environmental Science. 674(1). 012081. **IOP** Publishing. DOI: 10.1088/1755-1315/674/1/012081.
- Olivotto, I., Planas, M., Sinoes, N., Holt, G.J., Avella, M.A. and Calado, R., 2011. Advances in breeding and rearing marine ornamentals. *Journal of the World Aquaculture Society*, 42(2), 135-166. DOI: 10.1111/j.1749-7345.2011.00453.x.
- Olvera-Novoa, M.A., Martínez-Palacios, C.A. and Olivera-Castillo, L., 2002. Utilization of torula yeast (*Candida utilis*) as a protein source in diets for tilapia (*Oreochromis mossambicus* Peters) fry. *Aquaculture Nutrition*, 8(4), 257-264. DOI: 10.1046/j.1365-2095. 2002.00215.x.

- Paray, B.A. and Al-Sadoon, M.K., 2016. Utilization of organic manure for culture of cladocerans, *Daphnia carinata*, *Ceriodaphnia cornuta* and copepod, *Thermocyclops decipiens* under laboratory conditions. *Indian Journal of Geo-Marine Sciences*, 45(3), 399-404.
- Payne, M.F. and Rippingale, R.J., 2000. Rearing West Australian seahorse. Hippocampus subelongatus, juveniles on copepod enriched nauplii and Artemia. Aquaculture, 188(3-4), 353-361. DOI: 10.1016/s0044-8486(00)00349-5.
- Rahmati. R., Esmaeili Fereidouni, A., Rouhi, A. and Agh, N., 2020. Effects of different diets population growth and fatty acids composition in cyclopoid copepod, Acanthocyclops (Mirabdullayev and Defaye, 2002): A potential supplementary live food for freshwater fish larvae. Iranian Journal of Fisheries Sciences, 19(3), 1447-1462. DOI: 10.22092/ijfs.2019.120729.
- **Rajkumar, M. and Kumaraguru Vasagam, K.P., 2006.** Suitability of the copepod, *Acartia clausi* as a live feed for Seabass larvae (*Lates calcarifer* Bloch), Compared to traditional live-food organisms with special emphasis on the nutritional value. *Aquaculture*, 261(2), 649-658. DOI:
- Rasdi, N.W. and Qin, J.G., 2015. Effect of N: P ratio on growth and

10.1016/j.aquaculture.2006.08.043.

- chemical composition of *Nannochloropsis oculata* and *Tisochrysis lutea. Journal of Applied Phycology*, 27, 2221-2230. DOI: 10.1007/s10811-014-0495-z.
- **Rasdi, N.W., Qin, J.G. and Li, Y., 2016.** Effects of dietary microalgae on fatty acids and digestive enzymes in copepod *Cyclopina kasignete*, a potential live food for fish larvae. *Aquaculture Research*, 47(**10**), 3254-3264. DOI: 10.1111/are.12778.
- Rasdi, N.W., Suhaimi, H., Yuslan, A., Sung, Y.Y., Ikhwanuddin, M., Omar, S.S., Qin, J.G., Kassim, Z. and Yusoff, F.M., 2018. Effect of mono and binary diets on growth and reproduction of cyclopoid copepod. Aquaculture, Aquarium, Conservation and Legislation Bioflux, 11(5), 1658-1671.
- Rasdi, N.W. and Qin, J.G., 2018a. Improvement of copepod nutritional quality as live food for aquaculture: A review. *Aquaculture Research*, 47(1), 1-20. DOI: 10.1111/are.12471.
- Rasdi, N.W. and Qin, J.G., 2018b.
 Copepod supplementation as a live food improved growth and survival of Asian seabass *Lates calcarifer* larvae. *Aquaculture Research*, 49(11), 3606-3613. DOI: 10.1111/are.13828.
- Rasdi, N.W. and Qin, J.G., 2018c.
 Impact of food type on growth, survival and reproduction of the cyclopoid copepod *Cyclopina kasignete* as a potential live food in aquaculture.

 Aquaculture

- *International*, 26(**5**), 1281-1295. DOI: 10.1007/s10499-018-0283.x.
- Redzuari, A., Azra, M.N., Abol-Munafi, A.B., Aizam, Z.A., Hii, Y.S. and Ikhwanuddin, M., 2012. Effects of feeding regimes on survival, development and growth of blue swimming crab, *Portunus pelagicus* (Linnaeus, 1758) larvae. *World Applied Sciences Journal*, 18(4), 472-478. DOI: 10.5829/idosi.wasj.2012.18.04.313.
- **M.B.**, Barrows, Rust, F.T., Drawbridge, M., Hart, **E.R.**, Stuart, K., Webb, K., Barnett, Nicklason, P.M. H.J., and Johnson, R.B., 2015. Development and characterization of several open formula reference diets for marine fish larvae. Bulletin of Fisheries Research Agency, 40, 75-77.
- Samat, N.A., Yusoff, F.M., Rasdi, N.W. and Karim, M., 2020. Enhancement of live food nutritional status with essential nutrients for improving aquatic animal health: A review. *Animals*, 10(12), 2457. Published online, DOI: 10.3390/ani10122457.
- Santhanam, P., Perumal, P. and Rajkumar, M., 2004. Effects of feeding Artemia on growth and survival of P. monodon larvae.

 Journal of Applied Fisheries and Aquaculture, 4(2), 42-46.
- Santhanam, P. and Perumal, P., 2012. Evaluation of the marine copepod *Oithona rigida* Giesbrecht as live feed for larviculture of Asian seabass *Lates calcarifer* Bloch with special

- reference to nutritional value. *Indian Journal of Fisheries*, 59(2), 127-134.
- Sargent, J.R., McEvoy, L.A. and Bell, 1997. J.G., Requirements, presentation and sources polyunsaturated fatty acids in marine fish larval feeds. Aquaculture, 155(1-4), 117-127. DOI: 10.1016/S0044-8486(97)00122-1.
- Singh, K., Munilkumar, S., Sahu, N.P., Das, A. and Devi, G.A., 2019. Feeding HUFA and vitamin C-enriched *Moina micrura* enhances growth and survival of *Anabas testudineus* (Bloch, 1792) larvae. *Aquaculture*, 500, 378-384. DOI: 10.1016/j.aquaculture.2018.09.049.
- Solgaard, G., Standal, I.B. and Draget, K.I., 2007. Proteolytic activity and protease classes in the zooplankton species Calanus finmarchicus. Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology, 147(3), 475-481. DOI: 10.1016/j.cbpb.2007. 02.014.
- Yuslan, A., Najuwa, S., Hagiwara, A., Ghaffar, M.A., Suhaimi, H. and Rasdi, N.W., 2021. Production performance of *Moina macrocopa* (Straus 1820) (crustacea, cladocera) cultured in different salinities: the effect on growth, survival, reproduction, and fatty acid composition of the neonates. *Diversity*, 13(3), 105. Published online, DOI:10.3390. d13030105.
- Zhang, D., Lin, T. and Liu, X., 2015. A comparison of growth, survival, and fatty acid composition of the lined seahorse, *Hippocampus erectus*, juveniles fed enriched *Artemia* and a calanoid copepod, *Schmackeria dubia*. *Journal of the World Aquaculture Society*, 46(6), 608-616. DOI: 10.1111/jwas.12233.