Downloaded from jifro.ir on 2025-11-02

# **Research Article**

# Optimum dietary protein requirement for maximum growth performance of Caspian kutum (*Rutilus frisii*) juveniles

# Zahmatkesh A.1\*; Karimzadeh K.2

Received: March 2021 Accepted: November 2021

#### **Abstract**

A 12-weeks feeding trial was conducted to estimate the optimum dietary protein level of Caspian kutum white fish as one of the highly important commercial species in the south of the Caspian Sea. The fish juveniles (0.59±0.05g body weight) were randomly distributed into 21 aquariums (30 fish per aquarium). The fish were fed with seven experimental diets containing 28, 32, 36, 40, 44, 48, and 52 % crude protein (CP).

Results showed that dietary protein levels significantly (p<0.05) affected growth performance and feed utilization of Caspian kutum juveniles. The fish fed a diet with 40% CP showed higher survival rate (p<0.05) than other groups. Feeding the fish with diets at 40-44% CP improved the weight gain (WG) and specific growth rate compared to 28%, 32%, and 52% CP diets. The feed conversion ratio and protein productive value were promoted by increasing the dietary CP level, while the protein efficiency ratio was decreased in diets containing high CP levels (44-52%). The fish receiving the protein supplemented diets showed higher protein and lipid content than the initial fish population. The results indicated that dietary CP level only affected the body protein content of the Caspian kutum juveniles (p<0.05). A broken-line regression calculated the protein requirement at 42.85-43.23 % based on the WG and PPV (protein productive value).

**Keywords:** Body composition, Caspian kutum, Dietary crude protein, Growth, Feed efficiency

<sup>1-</sup>Aquaculture Department, Guilan Agricultural and Natural Resources Research and Education Center, Agricultural Research, Education and Extension Organization (AREEO), Rasht, Iran.

<sup>2-</sup>Department of Marine Biology, Lahijan Branch, Islamic Azad University, Lahijan, Iran.

<sup>\*</sup>Corresponding author's Email: aszahmat@yahoo.co.uk

# Introduciton

aquaculture industry is producing more than 114 million tons of various fisheries products (FAO, 2020) playing a considerable role in the supply of the required human food as well as contributing to food security in the world. Introducing new fish species to the fish farm can improve the supply, diversify aquaculture products and help promote the industry's efficacy in maintaining food security. The Caspian white fish or kutum (Rutilus frisii Kutum Kaminski, 1901) is a commercial fish of the Caspian Sea (Samavat et al., 2019) which is spread over a wide area ranging from the Kura river in Azerbaijan up to the Atrak river in Turkmenistan as well as in the Iranian side of the Caspian Sea (Mirzajani et al., 2016).

Like other fish under cultural conditions, the Caspian kutum need a balanced diet consisting of three basic nutritional elements such as proteins, lipids and carbohydrates (Craig and Helfrich, 2009; Hosseini Shekarabi et al., 2021a). Thus, the use of a diet comprising optimum protein level constitutes a major portion of this species' successful rearing and production plan. Protein is an important element that triggers the formation of different organs and tissues (Kaushik et al., 1995; NRC. 2011; Hosseini Shekarabi et al., 2021 a, b). Also it can help improve growth rate particularly among carnivorous fish (McGoogan and Gatlin, 1999). Therefore, it is imperative to secure adequate crude protein (CP) content within the fish diet (Siddiqui and Khan, 2009; Akpinar et al., 2012). Since

the protein requirements of different fishes might vary according to fish age and species (Bowen, 1987; NRC, 2011), the optimum protein contents in the diet of various Cyprinid species have already been specified (Chang-an *et al.*, 2014; Jin *et al.*, 2015; Yun *et al.*, 2015).

A limited number of studies on the effect of macronutrients, especially dietary CP levels, on the growth and body composition of the Caspian kutum fish juveniles, reported that the proper dietary protein is 35% for this species (Neverian et al., 2005; Mahmoodi et al., 2013). Indeed, an inadequate level of CP in fish practical diets can cause low fish farming performance, increased production cost and decline in water al..(Neverian quality et2008a: Falahatkar et al., 2012). Therefore, the present study has been undertaken to assess the effects of dietary protein levels on the growth, feed utilization and body composition of kutum juveniles, together with an estimation of the required protein level in diets using the broken line regression method.

## Material and methods

Diets preparation

Seven semi-purified diets containing graded levels of 28, 32, 36, 40, 44, 48, and 52% crude protein were formulated by Lindo Ver.6.1 (Neverian *et al.*, 2005; Zahmatkesh and Karimzadeh, 2018) (Table 1). Gelatin, egg white and fish meal were used as protein sources in the diets. Upon weighing the ingredients, they were thoroughly mixed, added with distilled water to produce a paste form. The paste material was pelletized

through a meat chopper machine with apertures 2mm in diameter. The dough obtained was dried in the oven at 60°C over-night. The dough was first formed into small-sized pellets and passed

through sieves with mesh sizes ranging from 1 to 1.5 mm. The pellets were kept at -20°C until used.

Table 1: Formulation and proximate composition (% dry weight) (mean±SD) of experimental diets.

Ingredients	Dietary protein level (%)							
	28	32	36	40	44	48	52	
Gelatin <sup>a</sup>	7	7	7	7	7	7	7	
Egg albumin b	0.56	5.98	11.42	16.86	22.28	27.78	33.14	
Herring fish	20	20	20	20	20	20	20	
meal <sup>c</sup>								
Baker yeast d	10	10	10	10	10	10	10	
Wheat flour e	36.42	30.96	25.48	20	14.54	9	3.58	
Potato starch f	5	5	5	5	5	5	5	
Dry milk <sup>g</sup>	4	4	4	4	4	4	4	
Fish oil h	4	4	4	4	4	4	4	
Sunflower oil i	7.42	7.46	7.5	7.54	7.58	7.62	7.68	
Mineral premix k	2.5	2.5	2.5	2.5	2.5	2.5	2.5	
Vitamin premix <sup>1</sup>	3	3	3	3	3	3	3	
Antioxidant <sup>m</sup>	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
Proximate composition								
Dry matter	92.3±1.0	92.4±1.1	92.2±0.9	$92.6 \pm 0.8$	92.4±1.1	92.5±0.8	92.4±1.1	
Crude protein	$28.4 \pm 0.7$	$31.8\pm0.9$	$35.7 \pm 0.2$	$40.5\pm0.9$	44.6±0.6	$48.0\pm1.1$	$51.8\pm0.4$	
Crude fat	16.8±0.9	$16.9 \pm 0.5$	$16.8 \pm 0.4$	$16.9\pm0.2$	$17.0\pm0.3$	$16.9\pm0.4$	$16.9 \pm 0.2$	
ash	$7.7 \pm 0.4$	$7.9 \pm 0.5$	$8.6 \pm 0.2$	$8.6 \pm 0.2$	$8.7 \pm 0.4$	$9.1\pm0.3$	$9.4\pm0.2$	
Crude fiber	$5.1\pm0.2$	$4.2\pm0.2$	$2.3\pm0.2$	$2.3\pm0.1$	$1.8\pm0.3$	$3.9\pm0.4$	$1.3\pm0.2$	
NFE n	$34.3 \pm 0.5$	31.6±1.3	$29.1\pm0.9$	$24.3 \pm 2.1$	$20.4\pm1.0$	$14.7 \pm 2.4$	$13.2 \pm 1.1$	
Energy (kcal/g)	$3.37 \pm .09$	$3.43 \pm 0.0$	$3.47 \pm 0.0$	$3.52\pm0.1$	$3.59\pm0.0$	$3.58\pm0.1$	$3.64\pm0.1$	
		3	6	0	9	0	8	

<sup>&</sup>lt;sup>a</sup> Supplied by JAHANSHIMI Co., Ltd. (Tehran, Iran), 950 g/kg crude protein.

<sup>&</sup>lt;sup>b</sup> Supplied by GOLPOODR Co., Ltd. (Golestan, Iran), 800 g/kg crude protein.

<sup>&</sup>lt;sup>c</sup> Supplied by PARSKILKA Co., Ltd. (Mazandran, Iran), 600 g/kg crude protein, 110 g/kg crude lipid, 105 g/kg ash, 101 g/kg moisture.

<sup>&</sup>lt;sup>d</sup> Supplied by PURETOP Co., Ltd. (Khozestan, Iran), 460 g/kg crude protein.

<sup>&</sup>lt;sup>e</sup> Supplied by AZARGANDOM Co., Ltd. (Guilan, Iran).

f Supplied by TAKALVAND Co., Ltd. (Hamadan, Iran).

g Supplied by NUTRICIA Co., Ltd. (Netherlands), 194 g/kg crude protein, 247 g/kg crude lipid.

h Supplied by PARSKILKA Co., Ltd. (Mazandran, Iran).

<sup>&</sup>lt;sup>i</sup> Supplied by BAHAROIL Co., Ltd. (Tehran, Iran).

k Mineral premix (mg/kg of diet): MgSO4·7H2O, 3600; KI, 20; FeSO4·H2O, 5200; ZnSO4·H2O, 3600; CuSO4·5H2O, 500; Na2Se2O3, 0.2; MnSO4·H2O, 3600; CoCl2·6H2O, 15. Supplied by ISFAHAN MOKAMMEL Co., Ltd. (Isfahan, Iran).

<sup>&</sup>lt;sup>1</sup> Vitamin premix (mg/kg of diet): retinyl acetate (2800 IU/mg), 40; cholecalciferol, 0.6; dl-\_-tocopheryl acetate, 600; menadione, 60; thiamine hydrochloride,160; riboflavin, 220; pyridoxine hydrochloride, 160; vitamin B12, 0.4; ascorbic acid, 1000; folic acid, 20; biotin, 2; niacin, 600; calcium d-pantothenate, 640; inositol, 500. Supplied by ISFAHAN MOKAMMEL Co., Ltd. (Isfahan, Iran).

<sup>&</sup>lt;sup>m</sup> BHT. Supplied by JAHANSHIMI Co., Ltd. (Tehran, Iran).

<sup>&</sup>lt;sup>n</sup> NFE (nitrogen-free extract) = 100–(crude protein + crude lipid + ash + crude fiber + moisture).

# Experimental animals and design

A total of 800 kutum fish juveniles were obtained from a state-owned fish hatchery center (Siahkal Fish Stock Enhancement and Propagation Center, Guilan Province, Iran). After being moved into 100L aerated fish tanks, were transferred to the Mirzakochak khan Centre for Fisheries sciences & technology. The fish were initially transferred in 2000 L tank and acclimated to rearing conditions for 10 days.

The juveniles were introduced into 21 glass tanks (aquaria, 50×35×40 cm) and reared for 12 weeks. Thirty fish (initial mean body weight 0.59±0.05 g) were randomly distributed into each tank (Ahmadian *et al.*, 2015) with aeration supplied through an air pump (HAILEA, Aco-450, China).

Water quality, including temperature, dissolved oxygen, and pH was measured and recorded daily basis using digital multi meter (GENWAY 370, UK). During the rearing period, the average water temperature, dissolved oxygen, nitrite, nitrate and pH levels were estimated in different treatments at  $20.57\pm1.90^{\circ}$ C,  $6.76 \pm 0.65$ mg/L.  $0.16\pm0.02$ mg/L,  $3.80\pm0.35$ mg/L, and  $8.18 \pm 0.12$ , respectively. The daily photoperiod was set at 12-h light and 12h dark. The daily water recirculation rate was 20% of the whole volume (12 L) in each tank. In addition, during the experimental period each tank was completely washed and filled with fresh water every 2 weeks.

The fish were fed three times a day (at 08:00, 13:00 and 20:00 h) by a

feeding rate of 8% wet body weight. The weight and length of kutum juveniles were measured every two weeks for monitoring their growth and determining their daily feeding requirement per tank. After the feeding trial, fish were counted and weighed and the different indices such as weight gain rate (WGR), specific growth rate (SGR), condition factor (CF) and feed conversion ratio (FCR) were calculated.

At the end of the experiments, 5 juveniles were collected in each tank (from each treatment 15 juveniles) and kept at -20°C for carcass analysis. The proximate chemical analysis of fish carcasses were determined based on the standard international methods (AOAC, 1990). To determine the moisture, 5 g of the sample was placed in drier at 105°C until constant weight (approximately for 24 h). Protein content was measured according to the Kjeldahl method after Kjeldahltherm acid digestion by (KBL20S, Gerhardt, Germany). Lipid measurement was carried out through the petroleum ether extraction and the use of Soxhlet-Extraktion system (B-810, Büchi, Netherlands) with fiber content specified using by glass crucibles and crude fiber extraction system (Foss Fibertec 1020 Hot and 1021 Cold extractor, Denmark).

# Statistical analysis

Statistical analysis was done using a SPSS software package (18.0, SPSS Statistics, IBM, Chicago, IL, USA). After confirming the normality and homogeneity of variance, differences between means were tested for

significance (p < 0.05) by the one way **ANOVA** analyses. Tukey HSD's multiple-range test was used determined significant differences among means. **Optimal** protein requirement in the diet was calculated using broken-line regression model (Robbins et al., 2006) and SAS version 9 software package.

## Results

The survival rate of kutum fish juveniles fed experimental diets at different protein levels is presented in Figure 1. After 12 weeks of the feeding trial, the survival rate among kutum juveniles fed the 40% diet showed to be higher (97.78%)than that of in treatments, specifically those fed the 28% diet (88.33%) (p < 0.05). The growth performance of juvenile kutum fish is indicated in Table 2. Final weight and fish condition factor in all treatments were not significantly affected by the dietary protein level (p>0.05). Weight gain rate and special growth rate of kutum juveniles fed the 28, 32 and 52% crude protein were significantly lower than those of the fish fed the 36, 40 and 44% CP diets (*p*<0.05).




Figure 1: Survival of *Rutilus frisii kutum* juveniles fed diets containing various levels of protein after 12 weeks rearing period.

Parameters of nutrient utilization (FCR, PER and PPV) were significantly affected by the levels of dietary protein (Table 2, p<0.05). FCR among juveniles tended to improve in line with an increase in protein content of diets up to 40% protein level (p<0.05). PER significantly decreased with incremental

levels of dietary protein (p<0.05). PER of fish fed the 32% protein diet was significantly higher than those of fish fed the other experimental diets. Protein productive value (PPV) was progressed with increasing dietary protein level and was found to be the highest for fish fed 40% CP (p<0.05). No significant

difference were observed in feed intake among fish fed the experimental diets. The production rate in experimental treatments was affected by protein content of diets to the extent that a rise in protein level was associated with significant production increase (p<0.05). The lowest production (16.42±1.30g in tank) and the highest production level (23.86±1.05g in tank) were estimated in 28% and 40% diets respectively (Table 2).

Table 2: Growth performance and feed utilization (mean±SD) of *Rutilus frisii kutum* juveniles fed diets with different levels of crude protein for 12 weeks.

diets with unicidit tevels of crude protein for 12 weeks.							
Parameters	Dietary protein level (%, dry matter)						
	28	32	36	40	44	48	52
Initial weight (g)	$0.61\pm0.00$	$0.62\pm0.01$	$0.57\pm0.03$	$0.60\pm0.02$	$0.57\pm0.03$	$0.58\pm0.03$	0.61±0.03
Final weight (g)	$1.31\pm0.01$	$1.39\pm0.03$	$1.34\pm0.07$	$1.42\pm0.03$	$1.35\pm0.07$	$1.35\pm0.08$	$1.37\pm0.07$
Weight gain (g) <sup>1</sup>	$0.70\pm0.01^{a}$	$0.78\pm0.02^{ab}$	$0.77\pm0.03^{ab}$	$0.83\pm0.02^{b}$	$0.78\pm0.04^{ab}$	$0.77\pm0.05^{ab}$	$0.76\pm0.04^{ab}$
Weight gain rate <sup>2</sup>	$114.75\pm2.32^a$	$125.94\pm1.22^{b}$	134.53±1.19°	$138.57\pm2.18^{c}$	$138.29\pm2.34^{c}$	$133.28 \pm 2.56^{c}$	125.85±1.85 <sup>b</sup>
Specific growth	$0.91\pm0.02^{a}$	$0.97\pm0.01^{b}$	1.01±0.01°	$1.03\pm0.02^{c}$	1.03±0.01°	$1.01\pm0.02^{c}$	$0.97\pm0.01^{b}$
rate (%/day) <sup>3</sup>							
Feed intake	$2.30\pm0.03$	$2.01\pm0.02$	$2.07\pm0.08$	$2.04\pm0.13$	$2.00\pm0.18$	$2.11\pm0.21$	2.27±0.15
(g/fish) <sup>4</sup>							
Feed conversion	$3.28\pm0.03^{b}$	$2.58\pm0.06^{a}$	$2.70\pm0.10^{ab}$	$2.47\pm0.11^{a}$	$2.56\pm0.35^{a}$	$2.73\pm0.24^{ab}$	$2.98\pm0.24^{ab}$
ratio <sup>5</sup>							
Protein efficiency	$1.09\pm0.01^{de}$	$1.21\pm0.03^{e}$	$1.03\pm0.04^{cd}$	$1.01\pm0.04^{cd}$	$0.90\pm0.12^{bc}$	$0.77\pm0.08^{ab}$	$0.65\pm0.05^{a}$
Ratio <sup>6</sup>							
Protein	204.72±90.50ab	267.86±96.75ab	294.11±133.98ab	409.86±82.58 <sup>b</sup>	343.64±114.22ab	215.58±69.41ab	122.88±2.55a
productive value							
$(\%)^7$							
Condition factor	$0.82\pm0.02$	$0.88\pm0.05$	$0.88\pm0.05$	$0.81\pm0.06$	$0.89\pm0.11$	$0.82\pm0.10$	$0.79\pm0.04$
$(g/m^3)^8$							
Production	$16.42\pm1.30^{a}$	$20.04\pm0.32^{b}$	$20.33\pm0.80^{b}$	$23.86 \pm 1.05^{c}$	21.26±1.49bc	20.47±1.63b	20.12±0.58b
(g/tank) <sup>9</sup>							

Values within the same row with different letters are significantly (p<0.05) different.

According to the broken-line regression analysis of weight gain (%) and protein production value (%) against dietary protein content, the optimum protein requirement of the kutum diet was estimated at 43.23% and 42.85% respectively (Figs. 2 and 3).

The result of carcass composition analysis of experimental fish feeding on diets with different protein levels is indicated in Table 3. Initial carcass composition of fish contained more moisture than final carcass composition of fish but relatively lower protein and lipid contents, regardless of experimental diets.

<sup>&</sup>lt;sup>1</sup> Weight gain: final mean weight – initial mean weight.

<sup>&</sup>lt;sup>2</sup> Weight gain rate: 100×(final mean weight – initial mean weight) / initial mean weight.

<sup>&</sup>lt;sup>3</sup> Specific growth rate: 100×[Ln (final mean weight)–Ln (initial mean weight)] / days of the experiment.

<sup>&</sup>lt;sup>4</sup>Feed intake (dry matter, g/fish) = total feed fed (g)/fish.

<sup>&</sup>lt;sup>5</sup> Feed conversion ratio: g dry feed / g weight gain.

<sup>&</sup>lt;sup>6</sup> Protein efficiency ratio: g weight gain / g protein fed.

<sup>&</sup>lt;sup>7</sup> Protein productive value: 100×g protein gain / g protein fed.

<sup>&</sup>lt;sup>8</sup> Condition factor: 100×fish weight (g) / fish length (cm) <sup>3</sup>.

<sup>&</sup>lt;sup>9</sup> Production: final mean biomass (g) – initial mean biomass (g).




Figure 2: The quadratic broken-line model fitting weight gain (%) to dietary protein levels in *Rutilus frisii kutum* juveniles fed the experimental diets ( $R^2 = 0.9621$ ). Ymax, maximum growth rate; X, a protein level required for Ymax. Each graphic point represents the average value of each tank.




Figure 3: The quadratic broken-line model fitting protein productive value (%) to dietary protein levels in *Rutilus frisii kutum* juveniles fed the experimental diets ( $R^2 = 0.8163$ ). Ymax, maximum growth rate; X, a protein level required for Ymax. Each graphic point represents the average value of each tank.

Moisture level in the carcass of kutum juveniles declined with increasing of diet protein but no significant difference was observed in the dietary treatments (p>0.05). Crude protein content in the carcass of kutum juveniles was affected

by dietary protein level (p<0.05). The protein content in the carcass of fish (17.28±0.52) fed the 40% protein showed to be significantly higher than those in the fish receiving 28%, 32% and 52% dietary proteins (p<0.05). Although

lipid variation in the carcass of kutum juveniles was similar to that of protein, there were no significant differences in the amount of such index across treatments involving the use of diets of varying protein levels (p>0.05). The highest lipid content (12.6±1.20) was

detected in the carcasses of kutum juveniles fed the 44% diet. The result of the present study showed that the protein level of diet does not have any impact on the ash content of kutum fish juvenile's carcass (p>0.05).

Table 3: Whole-body compositions (fresh-wt. basis) (mean±SD) of *Rutilus frisii kutum* juveniles fed different dietary protein levels for 12 weeks.

Distant matrix land (0/)	Body composition						
Dietary protein level (%)	Moisture %	Protein %	Lipid %	Ash %			
Initial <sup>1</sup>	69.48±1.72 <sup>b</sup>	13.99±0.42a	8.71±0.99 a	4.05±0.51			
28	$67.61 \pm 0.58^{ab}$	$15.27 \pm 0.60^{ab}$	$10.39 \pm 0.21^{ab}$	$3.91\pm0.45$			
32	$67.27 \pm 0.42^{ab}$	$15.67 \pm 0.63^{ab}$	$10.49\pm0.89^{ab}$	$3.85 \pm 0.65$			
36	$67.21 \pm 0.29^{ab}$	$16.33\pm1.11^{bc}$	$10.52 \pm 1.46^{ab}$	$3.85\pm0.38$			
40	$65.76 \pm 0.30^{a}$	$17.28\pm0.52^{c}$	$11.42\pm0.66^{ab}$	$3.84\pm0.44$			
44	$65.55 \pm 1.15^{a}$	$16.94\pm0.94^{bc}$	$12.06\pm1.20^{b}$	4.12±0.36			
48	$66.51 \pm 0.54^{a}$	$16.10\pm0.53^{bc}$	$11.51 \pm 1.25^{ab}$	$3.93\pm0.39$			
52	$67.10\pm1.04^{ab}$	$15.40 \pm 0.13^{ab}$	$10.06\pm1.09^{ab}$	$3.95\pm0.36$			

Values within the same column with different letters are significantly different (p<0.05).

## Discussion

After 12 weeks of the feeding trial, the survival of experimental fish averaged 93.25%. The survival was significantly different between the diets containing 28 and 40% CP. Survival rate of fish under trial condition depends on many factors including water quality, fish density, sanitation of the rearing environment and more specifically the quality and quantity of fish feed. In the present study, increasing dietary protein level up to 40% resulted in relative improvement of survival rate in juvenile kutum fish whereas the lowest survival rate was noticed in fish fed the 28% diet. This might account that diets of lower than 30% protein level are not suitable for kutum juveniles. However, researchers reported that the survival of other fish such as catfish (Khan, 1993;

Deng, 2011) and cyprinids (Jin *et al.*, 2015; Suharmili *et al.*, 2015) have shown to be less affected by protein level.

In this study, growth parameters (i.e. WG, WGR, and SGR) tended to improve significantly in line with an increased diet protein and the highest growth performance was obtained at 40% protein level. A number of researches had already suggested the positive effects of protein levels on fish growth parameters (Steffens, 1981; Li et al., 2010; Jin et al., 2015), while other studies reported a reverse relationship between protein levels of diet and weight gain in common carp, Cyprinus carpio (Cho et al., 2001) and Indian carp, Cyprinus catla (Parveen and Sheri, 1994; Satpathy et al., 2003). Such a different result might have been due to

<sup>&</sup>lt;sup>1</sup>The initial fish stock before starting the experiment.

optimizing dietary protein level to growth and supplementation of energy requirements through carbohydrates and lipids (Caballero et al., 2002). The optimum protein content in a diet relies upon sufficient provision of energy since part of the protein is used as energy reserve when fish feed energy is insufficient (Lovell, 1991; Salhi et al., 2004). Therefore, there should be a suitable protein / energy ratio in the diet to effectively utilize the protein. Jin et al. (2015)reported that grass (Ctenopharyngodon idella) juveniles fed the 40% protein diet enjoyed the best growth performance which is similar to the result of the present study.

In this study, the highest FI and FCR were observed among kutum juveniles fed 28% protein diet. The greater feed intake in dietary treatment containing lower protein and their higher FCR level might have been due to the fact that the nutrients in diets should be commensurate with tissue formation and repair requirements and metabolic functions of the fish body (Winfree and Stickney, 1981). When fish fed diets with sub-optimal protein contents, they tend to consume higher amounts of feed so as to meet their protein and energy requirements for growth and development (Gaylord and Gatlin, 2001; Wang et al., 2016). Similar to the results reported from other studies (Steffens, 1981; Tuan and Williams, 2007), the findings of the present study indicated FCR improvement upon increasing diet protein up to 40%. This might be attributed to sufficient provision of energy and its effective use for inducing

maximum growth of juveniles in these treatments.

In this study, an increase in dietary protein level decreased PER and raised PPV. Such a phenomenon has also been in other fish species reported (Dabrowski, 1977; Lee et al., 2002; Suharmili et al., 2015; Jiang et al., 2016). Generally, in carnivorous fish, PER is inversely correlated linearly with protein (Yang et al., 2003; Salhi et al., 2004). A dietary protein content above of required optimum level is used as an energy source (Shyong et al., 1998; Yang et al., 2003) which might induce increased enzymatic activity that breaks down amino acids in hepatopancreas, triggering greater nitrogen diffusion and thus reducing PER/PPV (Shimeno et al., 1981). The increased PPV associated with a rise in protein level (up to 40%) and P/E (up to 27.4 g/Mj) might have been caused by the formation of a proper P/E ratio within the diet subsequently suitable protein retention in the body of experimental fish. As indicated in earlier studies, fish need a specific P/E ratio in their feed to gain an optimum growth (Tucker. Neverian et al., 2008b) and P/E ratio for optimum growth defined for fish species ranges in 19-27 g/Mj (NRC, 2011).

In the present study, kutum fish juveniles with better growth rates enjoyed also the highest CF as was reported in grass carps by Jin *et al.* (2015). Higher CF values represented the appropriate nutritional and health status of fish and their suitable growth rate which is an ideal in aquaculture (Ayode, 2011; Castro *et al.*, 2016).

Estimating of yield amount is considered as a foundation of economic production estimation in aquaculture operations (El-Saidy and Gaber, 2005). In this study, fish production, estimated as per biomass and survival rate, began to rise when protein level in the diet was increased. Since kutum fish juveniles survival and growth rate were influenced by protein diet, it was evident that production increase could, likewise be expected.

Protein and lipid contents in the body of kutum fish fed the diet with high protein levels were significantly higher than the initial fish and those fed by a diet containing lower protein, whereas moisture level declined with an increase in protein content. In addition, protein in the body of fish feeding on 40% dietary protein turned to be significantly greater than those in treatments using feeds of 28, 32 and 52% protein levels. In agreement with this study, the whole body protein in cyprinids such as lemon fin barb hybrid, Hypsibarbus wetmorei  $\triangle \times Barbodes gonionotus \bigcirc$  (Suharmili et al., 2015) and Mahseer, Tor putitora (Hossain et al., 2002) began to rise with an increase in dietary protein level. The highest whole body protein specified in the body of grass carp juveniles fed 40% diet (Jin et al., 2015) which is in line with the present research.

In the present study, the highest protein and lipid contents were observed among juveniles fed diets containing 40 and 44% protein, respectively. This might reflect adequate protein level and ultimately, suitable P/E ratio that

induces maximum protein retention in body. It seems that a high protein level in diet (more than 44% in diet) results in extra energy generation. It has been indicated in previous research that excess energy in diet might entail the production of fat fish, decreased feed intake and could deter other feed ingredients from being utilized (Shiau and Lan, 1996).

The whole-body moisture of the experimental fish was negatively affected by increased protein levels in the diet, giving results similar to other research (Kim and Lee, 2005; Alam et al., 2008). The moisture decline could have been due to increased lipid in juveniles' bodies. The overall body and lipids of juveniles moisture remained nearly constant (Deng, 2011), parallel with reports often suggesting a reverse relationship between these two items in fish (Marais and Kissil, 1979; Afzal Khan et al., 2003).

The Increased protein level is associated with elevated fish feed cost, so that diets containing high protein level is known as an important restraint by many fish farmers, diminishing fish farming productivity. On the other hand, insufficient provision of protein and energy supply in feed might, apart from increasing fish feed production costs, diminish water quality due to feeding waste (Lee et al., 2002). Therefore, determining the protein requirement in the fish diet is necessary for promoting efficiency and securing feed profitability in fish rearing operations. According to broken line analyzing, the optimum protein levels for maximum PWG and PPV obtained 43.23% and 42.85% respectively. These values stand higher than the optimum protein levels for the other cyprinids species such as nobilis *Hypophthalmichthys* (Santiago and Reyes, 1991), lemon fin barb hybrid (34.6%) (Suharmili et al., 2015), Catla catla (Dars et al., 2010) and Leptobarbus hoevenii (Farahiyah et al., 2017), but that is almost equal with protein requirement of grass carp (Dabrowski, 1977; Jin et al., 2015). However, the required protein level in diet might vary according to size and species of fish, the feed formulation, rearing condition, water temperature, and more specifically, the energy level of diet as well as protein quality (Elangovan and Shim, 1997).

Based on the results obtained here, the optimum protein requirement in the diet of the Caspian kutum fish juveniles stands higher than that of other cyprinid species, suggesting that diets with high protein contents need to be used for maximum growth of this species. Although the estimated optimum protein level of the diet varied from 42.85 to 43.23%, it would be safe to suggest a 40% protein in any practical diet for rearing the kutum juveniles since it has caused a similar growth performance and the best protein retention rate. However, more studies need to be done on quality and sources of diet protein, energy requirement and P/E ratio aimed to devise a practical cost-effective diet for such species.

## References

- Afzal Khan, M., Khalil Jafari, A., Kumar Chadha, N. and Usmani, 2003. Growth N.. and body composition of rohu (*Labeo rohita*) fed diets containing oilseed meals: partial or total replacement of fish meal with soybean meal. Aquaculture Nutrition. 391-396. 9. DOI: 10.1046/j.1365-2095.2003.00268.x.
- Ahmadian, E., Malekzadeh Viayeh, R. and Zahmatkesh, A., 2015. Caspian whitefish (Rutilus frisii kutum, 1901) potential Kamensky, a aquaculture candidate: study on the cumulative effects of salinity and temperature on culture performance. Iranian Journal of **Fisheries** Sciences, 14(3), 623-633.
- Akpinar, Z., Sevgili, H., Ozgen, T., Demir, A. and Emre Y., 2012. Dietary protein requirement of juvenile shi drum, Umbrina cirrosa (L). *Aquaculture Research*, 43(3), 421-429. DOI: 10.1111/j.1365-2109.2011.02845.x.
- Alam, M.S., Watanabe, W.O. and Carrol P.M., 2008. Dietary protein requirements of juvenile Black sea bass, *Centropristis striata*. *Journal of the World Aquaculture Society*, 39(5), 656-663. DOI:10.1111/j.1749-7345.2008.00204.x.
- AOAC (Association of Official Analytical Chemists), 1990. Official Methods of Analysis. Fifth edition. Arlington, Virginia, USA.
- Ayode, A.A., 2011. Length—weight relationship and diet of African carp *Labeo ogunensis* (Boulenger, 1910) in Asejire lake southwestern Nigeria. *Journal of Fisheries and Aquatic Science*, 6(4), 472-478. DOI: 10.3923/jfas.2011.427.478.

- **Bowen, S. H., 1987.** Dietary protein requirements of fishes a reassessment. *Canadian Journal of Fisheries and Aquatic Sciences,* 44(11), 1995-2001. DOI: org/10.1139/f87-244.
- Caballero, M.J., Obach, A., D., Rosenlund, G., Montero, Gisvold, M. and Izquierdo, M.S., 2002. Impact of different dietary lipid sources on growth, lipid digestibility, tissue fatty acid composition and of rainbow histology trout. Oncorhynchus mykiss. Aquaculture, 214(1-4), 253-271. DOI:10.1016/S0044-8486(01)00852-3.
- Castro, MG., Castro, M.J., De Lara, A.R., Monroy, D.M.C., Ocampo, C.J.A. and Davila, FS., 2016. Length, weight and condition factor comparison of *Carassius auratus* (Linnaeus, 1758) juveniles cultured in biofloc system. *International Journal of Fisheries and Aquatic Studies*, 4(6), 345-350.
- Chang-an, W., Qi-you, X., Zhi-gang, Z., Jin-nan, L., Lian-sheng, W. and Liang, L., 2014. Effects of dietary protein and temperature on growth and flesh quality of Songpu Mirror carp. *Journal of Northeast Agricultural University*, 21 (2), 53-61. DOI:10.1016/S1006-8104(14)60034-9.
- Cho, S.H., Jo, J.Y. and Kim, D.S., 2001. Effects of variable feed allowance with constant energy and ratio of energy to protein in a diet for constant protein input on the growth of common carp *Cyprinus carpio* L. *Aquaculture Research*, 32(5), 349–356. DOI: 10.1046/j.1365-2109.2001.00564.x.

- Craig, S. and Helfrich, L., 2009. Understanding fish nutrition, feeds, and feeding. publication 420-256. College of Agriculture and Life Sciences, Virginia Polytechnic Institute and State University.
- **Dabrowski, K., 1977.** Protein requirements of grass carp fry (*Ctenopharyngodon idella* Val.). *Aquaculture*, 12(1), 63–73. DOI: 10.1016/0044-8486(77)90047-3.
- Dars, B.A., Narejo, N.T., Dayo, A., Lashar, P.K., Laghari, M.Y. and Waryani, B., 2010. Effect of different protein on growth and survival of *Catla catla* (Hamilton) reared in glass aquaria. *Sindh University Research Journal*, 42(1), 65-68.
- Deng, B., Zhang, X., Bi, B., Kong, L. and Kang, B., 2011. Dietary protein requirement of juvenile Asian redtailed catfish *Hemibagrus wyckioides*. *Animal Feed Science and Technology*, 170(3), 231–238. DOI:10.1016/j.anifeedsci.2011.08.014.
- Elangovan, A. and Shim, K.F., 1997. Growth response of juvenile *Barbodes altus* fed isocaloric diets with variable protein levels. *Aquaculture*, 158(3), 321-329. DOI:10.1016/S0044-8486(97)00199-3.
- El-Saidy, D.M.S.D. and Gaber. M.M.A., 2005. Effect of dietary protein levels and feeding rates on performance, production growth traits and body composition of Nile tilapia, Oreochromis niloticus (L.) cultured in concrete tanks. Aquaculture Research, 36(2), 163-10.1111/j.1365-171. DOI: 2109.2004.01201.x.

- **Falahatkar, B., Mohammadi, H. and Noveirian, H., 2012.** Effects of different starter diets on growth indices of Caspian Kutum, Rutilus frisii kutum larvae. *Iranian Journal of Fisheries Sciences*, 11(1), 28-36.
- FAO., 2020. The State of World Fisheries and Aquaculture 2020. Sustainability in action. Rome. DOI:10.4060/ca9229en.
- Farahiyah, I.J., Zainal Abidin, A.R., Ahmad, A. and Wong, H.K., 2017. Optimum protein requirement for the growth of Jelawat fish (*Leptobarbus hoevenii*). *Malaysian Journal of Animal Science*, 20(2), 39-46.
- Gaylord, T.G. and Gatlin, D.M., 2001. Dietary protein and energy modifications maximize to compensatory growth of channel catfish (Ictalurus punctatus). 337-348. Aquaculture, 194(**3-4**), DOI:10.1016/S0044-8486(00)00523-8.
- Hossain, M.A., Hasan, N., Azad Shah, A. K. M. and Hussain, M. G., 2002. Optimum dietary protein requirement of Mahseer, *Tor putitora* (Hamilton) fingerlings. *Asian Fisheries Science*, 15(3), 203-214.
- Hosseini Shekarabi, S.P., Shamsaie Mehrgan, M., Banavreh, A. and F., 2021a. **Partial** Foroudi, replacement of fishmeal with corn protein concentrate in diets for rainbow trout (Oncorhynchus mykiss): **Effects** growth on performance, physiometabolic responses, fillet quality. and Aquaculture Research, 52(1), 249-259. DOI: 10.1111/are.14887.
- Hosseini Shekarabi, S.P., Shamsaie Mehrgan, M., Banavreh, A., 2021b. Feasibility of superworm, Zophobas

- morio, meal as a partial fishmeal replacer in fingerling rainbow trout, Oncorhynchus mykiss, diet: growth performance, amino acid profile, proteolytic enzymes activity and pigmentation. *Aquaculture Nutrition*, 27(4), 1077-1088. DOI:10.1111/anu.13249.
- Jiang, S., Wu, X., Luo, Y., Wu, M., Lu, S., Jin, Z. and Yao, W., 2016. Optimal dietary protein level and protein to energy ratio for hybrid grouper (*Epinephelus fuscoguttatus* ♀ × *Epinephelus lanceolatus* ♂) juveniles. *Aquaculture*, 465, 28–36. DOI:10.1016/j.aquaculture.2016.08. 030.
- Jin, Y., Tiana, L.X., Xie, S.W., Guo, D.G., Yang, H. J., Liang, G.Y. and Liu, Y.J., 2015. Interactions between protein dietary levels, growth performance, feed utilization, gene expression and metabolic products in juvenile grass carp (Ctenopharyngodon idella). Aquaculture, 437, 75-83. DOI:10.1016/j.aquaculture.2014.11. 031.
- Kaushik, S.J., Doudet, T., Médale, F., Aguirre, P. and Blanc, D., 1995. Protein and energy needs for maintenance and growth of Nile tilapia (*Oreochromis niloticus*). *Journal of Applied Ichthyology*, 11(3-4), 290-296. DOI: 10.1111/j.1439-0426.1995.tb00029.x.
- Khan, M.S., Ang, K.J., Ambak, M.A. and Saad, C.R., 1993. Optimum dietary protein requirement of a Malaysian freshwater catfish, *Mystus nemurus*. *Aquaculture*, 112(2), 227-235. DOI:10.1016/0044-8486(93)90448-8.

- **Kim, L.O. and Lee, S.M., 2005.** Effects of the dietary protein and lipid levels on growth and body composition of bagrid catfish, *Pseudobagrus fulvidraco*. *Aquaculture*, 243(1), 323-329.
  - DOI:10.1016/j.aquaculture.2004.11. 003.
- Lee, S.M., Jeon, I.G. and Lee, J.Y., 2002. Effects of digestible protein and lipid levels in practical diets on growth, protein utilization and body composition of juvenile rockfish (Sebastes schlegeli). Aquaculture, 211(1), 227–239.DOI:10.1016/S0044-8486(01)00880-8.
- Li, X., Liu, W., Jiang, Y., Zhu, H. and Ge, X., 2010. Effects of dietary protein and lipid levels in practical diets on growth performance and body composition of blunt snout bream (*Megalobrama amblycephala*) fingerlings. *Aquaculture*, 303(1), 65–70.DOI:10.1016/j.aquaculture.2010.03.014.
- **Lovell, R.T., 1991.** Nutrition of aquaculture species. *Journal of Animal Science*, 69(**10**), 4193-4200.
- Mahmoodi, Z., Alaf Noveirian, H., Falahatkar, B. and Khoshkholgh, M., 2013. The effect of different dietary protein and lipid levels on performance in growth Caspian Kutum (Rutilus frisii kutum. Kamensky, 1901). Iranian Scientific Fisheries Journal, 22(1), 101-116.
  - DOI:10.22092/ISFJ.2017.110106.
- Marais, J.F.K. and Kissil, G.Wm., 1979. The influence of energy level on the feed intake, growth, food conversion and body composition of *Sparus aurata*. *Aquaculture*, 17(3),

- 203-219. DOI:10.1016/0044-8486(79)90124-8.
- McGoogan, B.B. and Gatlin, D.M., 1999. Dietary manipulations affecting growth and nitrogenous waste production of red drum, *Sciaenops ocellatus*. Effects of dietary protein and energy levels. *Aquaculture*, 178 (3), 333–348. DOI:10.1016/S0044-8486(99)00137-4.
- Mirzajani, A., Hamidian, A.H., Abbasi, K.,and Karami, M., 2016. Distribution and abundance of fish in the southwest of Caspian Sea coastal waters. *Russian Journal of Marine Biology*, 42(2), 178–189.
- Neverian, H.A., Mostafazadeh, S. and Toluei, M.H., 2005. A study on various protein levels on growth indices (SR, WG, RGR, FCR and PER) of *Rutilus frisii kutum*, Kamenskii 1901(Advanced fry). *Pajouhesh & Sazandegi*, 68, 61-68(in Persian with an abstract in English).
- Neverian, H., Shabanipour, N., Khoushkholgh, M.R. and Hosseini, M.R., 2008a. Effect of dietary digestible energy level on growth indices of kutum (*Rutilus frisii kutum Kamenskii*, 1901). *Iranian Journal of Fisheries Sciences*, 7(2), 205-214. DOI: 10.22092/ijfs.2018.114778.
- Neverian, H., Zahmatkesh, A., Zamani, H.A., Ghanaatparast, A., 2008b. Determination of the optimum level of vitamin premix for the diet of Caspian frisii kutum (advanced fry). *Pajouhesh-va-Sazandegi*, 279(21), 166-174.
- NRC (National Research Council), 2011. Nutrient requirements of fish. National Academy Press, Washington, D.C.

- Parveen, S. and Sheri, A.N., 1994. Influence of different levels of protein on growth performance and body composition of major carps. *Pakistan Journal of Agricultural Sciences*, 31(4), 345-350.
- Robbins, K.R., Saxton, A.M. and Southern, L.L., 2006. Estimation of nutrient requirements using brokenline regression analysis. *Journal of Animal Science*, 84(E. Suppl.), 155–165.

DOI:10.2527/2006.8413\_supple155x

- Salhi, M., Bessonart, M., Chediak, G., Bellagamba, M. and Carnevia, D., 2004. Growth, feed utilization and body composition of black catfish, *Rhamdia quelen*, fry fed diets containing different protein and energy levels. *Aquaculture*, 231(1), 435-444.
  - DOI:10.1016/j.aquaculture.2003.08. 006.
- Samavat, Z., Shamsaie Mehrgan, M., Jamili, S., Soltani, M. and Hosseini Shekarabi, S. P., 2019. Determination of grapefruit (Citrus paradisi) peel extract bio-active substances and its application in Caspian white fish (Rutilus frisii diet: Growth, *kutum*) haematobiochemical parameters and intestinal morphology. Aquaculture Research, 2496-2504. 50(9), DOI: 10.1111/are.14204.
- Santiago, C.B. and Reyes, O.S., 1991.

  Optimum dietary protein level for growth of bighead carp (*Aristichthys nobilis*) fry in a static water system. *Aquaculture*, 93(2), 155-165.

  DOI:10.1016/0044-8486(91)90214-R.
- Satpathy, B.B., Mukherjee, D. and Ray, A.K., 2003. Effects of dietary

- protein and lipid levels on growth, feed conversion and body composition in rohu, *Labeo rohita* (Hamilton), fingerlings. *Aquaculture Nutrition*, 9(1), 17–24. https://doi: 10.1046/j.1365-2095.2003.00223.x.
- Shiau, S.Y. and Lan, C. W., 1996.

  Optimum dietary protein level and protein to energy ratio for growth of grouper (*Epinephelus malabaricus*).

  Aquaculture, 145(1), 259-266.

  DOI:10.1016/S00448486(96)01324-5.
- Shimeno, S., Takeda, M., Takayama, S. and H., Sasaki, H., 1981. Response of nitrogen excretion to change of dietary composition in carp. *Bulletin of the Japanese Society for the Science of Fish*, 47,191-195.
- Shyong, W.J., Huang, C.H. and Chen, H.C., 1998. Effects of dietary protein concentration on growth and muscle composition of juvenile *Zacco barbata*. *Aquaculture* 167(1), 35-42. DOI:10.1016/S0044-8486(98)00313-5.
- Siddigui, T.Q. and Khan, M.A., 2009. Effects of dietary protein levels on feed utilization, growth, protein efficiency retention and body composition of young Heteropneustes fossilis (Bloch). Fish Physiology and Biochemistry, 35(3), 479-488. DOI: 10.1007/s10695-008-9273-7.
- **Steffens, W., 1981.** Protein utilization by rainbow trout (*Salmo gairdneri*) and carp (*Cyprinus carpio*): A brief review. *Aquaculture*, 23(1), 337-345. DOI:10.1016/0044-8486(81)90026-0.
- Suharmili, R., Kamarudin, M.S., Saad, C.R., Ina-Salwany, M.Y., Ramezani-Fard, E. and Mahmud,

- M.H., 2015. Effects of varying dietary protein level on the growth, feed efficiency and body composition of lemon fin barb hybrid fingerlings. *Iranian Journal of Fisheries Sciences*, 14(2), 425-435.
- Tuan, L.A. and Williams, K.C., 2007. Optimum dietary protein and lipid specifications for juvenile Malabar grouper (Epinephelus malabaricus). *Aquaculture*, 267(1), 129–138. DOI:10.1016/j.aquaculture.2007.03. 007.
- **Tucker, J.W., 1992.** *Marine fish nutrition.* Proceedings from the Aquaculture Nutrition Workshop, Salamander Bay, Australia, pp. 25-40.
- Wang, J., Jiang, Y., Li, X., Han, T., Yang, Y., Hu, S. and Yang, M., 2016. Dietary protein requirement of juvenile red spotted grouper (*Epinephelus akaara*). Aquaculture, 450,289–294.
- Winfree, R.A. and Stickney, A.R., 1981. Effects of dietary protein and energy on growth, feed conversion efficiency and body composition of *Tilapia aurea*. The Journal of

- *Nutrition*, 111(**6**), 1001-1012. DOI: 10.1093/jn/111.6.1001.
- Yang, S.D., Lin, T. S., Liou, C.H. and Peng, H.K., 2003. Influence of dietary protein levels on growth performance, carcass composition lipid classes juvenile of Spinibarbus (Oshima). hollandi Aquaculture Research, 34(8), 661-666. DOI: 10.1046/j.1365-2109.2003.00880.x.
- Yun, B., Yu, X., Xue, M., Liu, Y., Wang, J., Wu, X., Han, F. and Liang, X., 2015. Effects of dietary protein levels on the long-term growth response and fitting growth models of gibel carp (*Carassius auratus gibelio*). Animal Nutrition, 1(2), 70 -76. DOI: 10.1016/j.aninu.2015.05.003.
- Zahmatkesh, A. and Karimzadeh, K., 2018. Study of growth performance, survival and carcass composition of Kutum (*Rutilus frisii kutum*) fed diets containin,g different levels of protein and zinc. *Journal of Applied Ichthyological Research* 6(2), 73-89(in Persian).