# **Research Article**



# Assessment of tidal and lunar influence on larval diversity and distribution along the CVCAs of tropical Kali estuary, Karwar, West coast of India

Rajeshree P.1\*; Shivakumar H.1

Received: July 2022 Accepted: December 2022

### **Abstract**

Larval ingress, diversity and distributions were studied for a period of 13 months along the critically vulnerable areas of the tropical Kali estuary, Karwar, West coast of India. Planktonic and hydrological samplings were carried out following the tidal regime and lunar periodicity at Zone (1) (categorized as euhaline <30 PSU to polyhaline 18.0 to 30 PSU in condition), subsequently, Zone (2) (mesohaline 5.0 to 18.0 PSU to oligohaline 0.5 to 5.0 PSU). A total of 26 different larval forms were recorded, representing 10 taxa, of which crustaceans are dominant. The predominant larval forms documented during the study period were Copepod nauplii, Barnacle nauplii, Eggs & fish larvae and Crab zoea, respectively. Larval forms, viz., Planula, Cercaria, Rhabditiform, Alima (Squilla), Bipinnaria, Ophiopluteus and Doliolaria distributions was restricted to euhaline zone due to tidal ingress. The distribution of Bivalve, Crab and Shrimp larvae along the mesohaline zone represents larval recruitment in their preferred habitat. The larval distribution showed typical seasonal variation governed by hydrological parameters and tidal regimes. Multivariate RDA analysis of larval distributions attributed to salinity and water temperature. The Shannon-Wiener diversity index was higher (2.92) during the post-monsoon season in particular to full moon of the flood tide. Overall, the Kali Estuary represents a suitable habitat for juvenile nurture and larval ingress enhances the estuarine productivity.

Keywords: Larvae, Tides, Diversity, Distribution, Kali Estuary

<sup>1-</sup>Karnatak University Dharwad's, Department of Studies in Marine Biology, Post Graduate Centre, Kodibag, Karwar, Karnataka, India

<sup>\*</sup>Corresponding author's Email:rajeshree7pednekar@gmail.com

### Introduction

Tropical semi-enclosed estuaries ranked among the world's most productive and dynamic ecosystems (Sreekanth et al., 2017). These brackish water wetland ecosystems are the most challenging habitat and are the gateway for migratory species to complete their life cvcle. Moreover, mangrove canopy along the mudflats ofestuaries productivity enhances biological essential subsequently, offers an complex structural habitat for juvenile nurture. Also. the estuarine environment provides ample food resources for the early life stages of organisms (Holt and Holt, 2000). Most of the marine faunal life cycle accomplishes in these shallow sheltered zones as the tidal cycle bridges their ingress and exit. In particular specific ecological-based larval migration evokes mainly through the flood tide transport, which includes olfactory cues, currents, salinity, temperature and turbulence associated with amplitude and phase between shelf region to estuarine waters (Forward and Tankersley, 2001). Similarly, limiting factors for spatial species recruitment and distribution across the estuary are salinity, temperature, pH, dissolved oxygen, turbidity nutrients (Darboe, 2002; Perumal et al., 2009; Badsi et al., 2010; Cloern et al., 2017; Tanmoy and Mandal, 2020). Moreover, Oueiroga and Blanton (2004), state that successful estuarine species recruitment depends upon the strategies employed by behavioral larvae. The estuarine-dependent robust resilient larvae grow to the adulthood along the favorable salinity gradient, significantly enhance the regional biodiversity and associated fishery.

Among the crustaceans, molluscan and pisces larval ingress holds a key position from the commercial perspective (Katsanevakis et al., 2008; Kunze et al., 2013; Dias et al., 2020; Tarimo et al., 2022). Thus, larval totally depends ingress on the hydrodynamics environmental and signals selectively tidal to meteorologically driven currents back into the estuaries and avoid being flushed back out when these currents reverse (Buskey, 2018). Variability in the delivery of larvae from offshore to the nursery ground up to the upstream freshwater zone will certainly impact dynamics. trophic population structure and associated fisheries.

Larvae are the post-embryonic stages of the life cycle, which morphologically differ from the adults and might possess independent locomotion. Many marine species represent a bipartite life cycle characterized by a dispersive larval phase and a sedentary adult phase. Thus, quantifying larval forms and their dispersal is a goal of ecology and conservation (Kritzer and Sale, 2010). The spatial and temporal distribution of planktonic larval studies carried out across the globe are restricted to species/group viz., penaeid prawn larval ingress (Staple and Vance, 1985; Goswami and Usha, 1992), brachyuran crab larval distribution (Garrison, 1999) and fish larvae (Rao, 1970; Hare et al., 2005).

In general, larval diversity and ingress documentation was lacking. Aganist this backdrop, an attempt has been made to document the overall larval ingress, distribution, seasonal abundance, and wetland utility to forecast the nursery role of the estuarine ecosystem along the Kali estuary, Karwar, West coast of India.

## Materials and methods

Study area

The Kali estuary is a typical, mesotidal well-mixed tropical estuarine ecosystem of the West coast of India (14° 50 19.19" N - 74° 07 13.80" E). The river originates at the plateau of the Western Ghats, flows westward for about 184km, and layouts into the Arabian Sea, Karwar. The estuary has a narrow opening followed by a width of 1.5km in the mid-region with an average water column of 10m. Kali estuary is a tropical monsoon-influenced ecosystem in which the southwest monsoon (June-September) average rainfall recorded was 2760mm (IMD-Indian Meteorological Department), of which June and July were the wettest months. Tides in this tropical coast are mixed semi-diurnal with a large variation in the tidal range of 1.5 to 2.5m. The mangrove canopy of 606.47ha offers a network of sheltered zones innumerable ecological services (Mesta et al., 2016) along the Kali estuary. Being a resilient habitat, this wetland ecosystem was declared as a critically vulnerable coastal area (CVCA) by Management Indian Coastal Zone Authorities (CRZ, 2011).

Sampling stations were set along the estuarine stream based on the salinity gradient and active fishing pockets. Based on the salinity gradient, Zone 1 (Lat 14° 50' 33" N; Long 074° 08' 11"E) was the main channel of the estuary which categorized as euhaline (<30 PSU) to polyhaline (18.0 to 30 PSU) in nature. Zone 2 (Lat: 14° 50' 18" N Long: 074° 10' 00" E) is a backwater wetland zone of the estuary categorized as mesohaline (5.0 to 18.0 PSU) to oligohaline (0.5 to 5.0 PSU), besides densely covered mangrove vegetation (Fig. 1). Kali estuary supports the livelihood of fishermen as traditional fishing activity prevails throughout the year.

Field sampling methods and laboratory analysis

Chemical and biological samples were collected at monthly intervals following the tidal (flood tide and ebb tide) and lunar phases (full moon and new moon) for a period of 13 months (January 2018 to January 2019) at two fixed study zones. Field samplings were carried out during the daylight of flood tide and ebb tide, wherein fortnightly samplings were carried out during the nocturnal rising lunar tide. Sampling regimes were followed on a specific duration of mid-flood tide and subsequent ebb tide. Larval samples were collected through a plankton net 50µM mesh size, (0.5m mouth diameter) hauled horizontally for a period of 10 min at uniform speed through a motorized boat from surface water up to the depth of 5m.




Figure: 1: Map showing study area and location of sampling zones along Kali Estuary, Karwar, West coast of India.

A flowmeter attached to the plankton net registered the volume of water filtered (Archambault and Bourget, 1999; Goncalves et al., 2010). Samples collected were transferred to labeled bottles and preserved in 5% neutralized formalin. The known volume of the sample was concentrated to the desired (10mL). volume Samples were homogenized by vigorous random stirring and a triplicate sub-sample of 1ml each was transferred the Sedgwick counting chamber for enumeration. The mean of sub-samples is calculated and multiplied by the known volume of the sample. Larval samples were identified based on relevant literature (Young et al., 2002; Suthers and Rissik, 2009; Conway, 2012a, 2012b; Weiss, 2017) through a compound microscope fitted with a camera at a magnification of 4X, 10X and 40X. In parallel to larval sampling, water parameters, viz., pH, salinity (PSU), water temperature (°C) and dissolved oxygen (mg/L) were recorded in-situ through a hand-held multiprobe water analyzing kit (SD 400 Oxi L-Lovibond).

### Statistical analysis

Environmental variables analyzed for tidal and lunar phases were grouped seasonally, Pre-monsoon (PRE), Monsoon (MON), and Post-monsoon seasonally (POM) seasons. The variables categorized environmental were subjected to analysis of variance (ANOVA) with IBM SPSS (Statistical package for the social science ver. 22) software to find the variance of means among seasons and studied zones

followed bv Pearson's correlation coefficient to assess the interrelationship. Prior to the data analysis, all data sets were checked for the normality using Kolmogorov-Smironov test and homogeneity of variances using the Levene test (Sokal and Rohlf, 1995). Further, to test the variance between seasonal distributions, a non-parametric Kruskal-Wallis test (two or more groups) was performed to derive the significance between the seasons, followed by a post-hoc Mann-Whitney's U-test (two groups). Univariate diversity indices for larval community viz., the Shannon-Wiener (H) diversity index (Shannon, 1948) was computed (Tidal and Lunar phase) through the PAST software (PAlaeontological **Statistics** Ver. 4.07b).

In order to elucidate the relationship of larval distribution with environmental parameters a Detrended correspondence analysis (DCA) was carried out. The DCA gradient plot did not exceed the deviation of 3.0, thus multivariate Redundancy Data Analysis (RDA) was employed to discover the relationship between larval distribution and environmental factors. RDA was plotted to determine the correlations between tidal-larval diversity environmental parameters, similarly, lunar-larval diversity and environmental parameters, followed by seasonal-larval diversity and environmental parameters. Only dominant larvae were considered omitted and rare larvae were subsequently, statistical significance was evaluated based on the Monte Carlo Permutation Test (MCPT) with CANOCO 4.5 software (Leps and Smilauer, 2003).

### **Results**

Environmental clauses

The Physicochemical parameters studied during the study period are represented in (Table 1 and Fig. 2).

Table 1: Zone wise environmental parameters with seasonal significance difference.

Water parameters	PRE	MON	POM	mean±SD	Seas signifi	
•					F	p 0.526 0.000 0.000
pH	$7.7 \pm .21$	$7.7 \pm .30$	$7.7 \pm .2$	$7.7 \pm .24$	0.647	0.526
Dissolved oxygen (mg/L)	$4.9 \pm .87$	$6.0 \pm .71$	$5.1 \pm .7$	$5.38\pm.90$	16.427	0.000
Salinity (PSU)	$27.1 \pm 6.8$	13.3±10.9	26.7±7.6	22.7±10.61	27.952	0.000
Water temperature (°C)	27.9±1.7	$25.5\pm2.03$	$27.4 \pm 2.25$	$27.03\pm2.26$	12.155	0.000

Significance level at 5%

The physicochemical characteristics studied for the tides and lunar phases are categorized primarily into three conventional seasons; southwest monsoon (June-September), postmonsoon (October-January), and premonsoon (February-May), respectively. Kali estuary is majorly influenced by

river run-off where salinity markedly varies seasonally. The mean salinity values were higher during pre and postmonsoon (27.17±6.8 and 26.7±7.6 PSU) seasons wherein, salinity sharply decreased during the southwest monsoon (13.3±10.9 PSU).




Figure: 2: Spatio-temporal variations of environmental parameters with significance difference.

The water temperature followed the cycle with a mean of seasonal 27.03±2.26°C, high-temperature values represented during the pre and post- $(27.9\pm1.79)$ monsoon season 27.4±2.25°C). The pH value ranged from 7.1 to 8.3 with a mean of 7.7±0.24, wherein, dissolved oxygen varied from 3.12 to 6.93 mg/L with a mean of 5.38±0.90 mg/L during the study period. Overall. seasonal significant differences (p<0.05) were observed in the water temperature and salinity, followed by dissolved oxygen (Table 1). Moreover, the physicochemical parameters of the tidal regime and lunar phase also represented seasonal significant variation between the studied zones (Fig. 2). A significant variation (p=0.016) of salinity and (p=0.030)dissolved oxygen was observed during flood tide at Zone (1), whereas. salinity (p=0.001)significant at Zone (2). The significant variation during the ebb tide was represented by pH (p=0.014), salinity (p=0.001), dissolved oxygen (p=0.001) and water temperature (p=0.001) at Zone (1), whereas, salinity (p=0.002) and water temperature (p=0.023) were significant at Zone (2). Similarly, the phase showed lunar seasonal significance for salinity (p=0.001) and dissolved oxygen (p=0.002) at Zone (1), whereas, рH (p=0.002),salinity (p=0.001)and dissolved oxygen (p=0.034) at Zone (2) during the full moon phase. During the new moon phase, salinity (p=0.008) and dissolved oxygen (p=0.045) were statistically significant at Zone (1), wherein, salinity

(p=0.001), dissolved oxygen (p=0.005), and water temperature (0.025) at Zone (2) (Fig. 2). The person's correlation coefficient showed a strong positive correlation between the water temperature and salinity (r=0.889, p=0.037) during the study period.

Percentage composition of larvae distribution

A total of 26 larval forms of 10 taxa. Platyhelminthes, namely Cnidaria. Nematoda, Annelida. Mollusca, Arthropoda, Echinodermata, Bryozoa, Brachiopoda and Ichthyoplankton, were recorded in this study. Arthropods and crustaceans were the dominant group, with Copepod nauplii (15.30%), Crab zoea (8.50%), Barnacle nauplii (8.30%) followed by Shrimp zoea (8.20%), respectively. The Egg & fish larvae (10.3%), Bivalve veliger (6.90%) and Gastropod veliger (6.10%) are the other major predominant groups (Table 4). During the study period, larval forms such as Polycheate larvae, Barnacle nauplii, Barnacle cypris, Shrimp nauplii, Shrimp zoea, Shrimp mysis, post-larvae, Shrimp Crab zoea, Copepod nauplii, Bivalve veliger, Gastropod veliger, Cyphonautes and & fish larvae were Egg the representatives of both the studied zones (Tables 2 to 4).

The ingress of larval forms was markedly influenced by the tidal regimes. During the flood tide, *Ephyra*, *Bipinnaria* and *Ophiopluteus* were recorded throughout the year along the euhaline zone, wherein *Planula*, *Cercaria*, *Rhabditiform*, *Alima* 

(Squilla), and *Doliolaria* larvae were recorded during the pre-monsoon and post-monsoon seasons.

Table 2: Tidal and lunar seasonal percentage (%) composition of larvae at zone 1.

		lunar seasonal percentage (%) composition of larvae at zone 1.  ZONE 2										
Larval forms	F	lood ti	de	]	Ebb tid			ull mod	n	N	ew mo	on
	P	M	PO	P	M	PO	P	M	PO	P	M	PO
Cnidaria												
Ephyra (Ep)	3.0	X	2.5	1.2	X	1.0	5.2	1.9	2.5	3.2	X	3.4
Planula (Pla)	X	X	0.1	X	X	X	0.2	X	0.4	0.2	X	0.2
Actinula (Act)	0.6	X	4.1	4.9	X	4.6	3.7	4.3	1.4	2.8	X	2.8
Platyhelminthes												
Cercaria (Cer)	X	X	X	X	X	X	0.3	X	0.3	X	X	X
Nematoda												
Rhabditiform	0.3	X	0.1	X	X	X	0.3	X	0.2	X	X	X
(Rha)												
Annelida												
Polychaete (Pol)	4.2	3.3	4.9	3.4	2.5	4.8	5.7	1.8	6.7	7.5	1.3	6.5
Arthropoda												
Alima (Al)	X	X	X	X	X	X	X	X	0.1	X	X	X
Barnacle nauplii	7.9	6.1	9.3	7.5	1.3	10.7	9.8	3.7	7.5	9.2	2.6	8.9
(BNau)	10.0	<b>5</b> 0	<i>5</i> 4	<i>5</i> 1	0.2	6.0	<i>c</i> 0	1.5	7.1	0.1	2.0	
Barnacle cypris	10.8	5.9	5.4	5.4	8.2	6.8	6.9	4.5	7.1	8.1	2.0	6.6
(BCyp)	2.1	0.6	1 1	0.4	1.0	0.8	1.1	1.0	2.0	1.3	1.0	1.4
Shrimp nauplii (NauSr)	2.1	0.6	1.1	0.4	1.0	0.8	1.1	1.0	2.0	1.5	1.0	1.4
Shrimp zoea	3.6	5.7	10.6	10.4	8.3	9.0	8.6	6.3	7.2	3.5	11.6	9.7
(ZoSr)	3.0	3.7	10.0	10.4	0.5	9.0	0.0	0.5	1.2	3.3	11.0	9.1
Shrimp mysis	5.3	4.6	2.5	4.7	13.5	5.8	2.1	9.6	3.2	5.2	3.6	2.3
(MySr)	3.3	4.0	2.5	7.7	13.3	3.0	2.1	7.0	3.2	3.2	5.0	2.3
Shrimp Post-larvae	0.8	1.0	2.1	3.1	3.3	1.2	4.3	6.1	4.9	4.0	2.9	4.6
(PlSr)												
Ephausiid nauplii	1.3	X	0.4	0.4	X	0.3	2.0	0.3	4.5	3.5	1.5	3.2
(EpNau)		<b>7</b> 0	0.0	6.0	2.2	1.0		2.4	6.0		6.0	<b>7</b> 0
Crab zoea (CrZo)	7.7	5.8	9.0	6.9	3.2	10	6.5	2.4	6.8	6.6	6.8	5.8
Megalopa (Meg)	0.1	0.8	0.6	0.2	X	0.5	0.1	0.2	0.6	0.3	0.7	0.5
Crab Post-larvae	0.4	0.2	0.5	0.5	0.7	0.6	0.5	X	1.2	1.2	X	0.4
(PlCr)	17.0	21.5	10.1	17.0	26.2	12.4	162	11.0	11.1	160	22.4	10.0
Copepod nauplii (NaCop)	17.9	31.5	12.1	17.9	36.2	13.4	16.3	11.9	11.1	16.9	32.4	10.8
Mollusca												
Bivalve veliger	5.3	9.4	7.1	7.5	3.6	7.3	2.9	14.1	5.6	3.7	13.7	5.9
(BiVvel)												
Gastropod veliger (GaVel)	7.8	8.4	4.5	6.8	3.1	9.2	1.6	13.6	4.3	4.1	12.8	5.0
Echinodermata	0.6		0.2				0.5	0.0	0.6	0.1		Λ 1
Bipinnaria (Bip)	0.6	X	0.3	X	X	X 0.2	0.5	0.8	0.6	0.1 1.2	X 0.5	0.1
Ophiopluteus	0.5	X	1.1	0.8	X	0.2	4.2	1.1	3.7	1.2	0.5	1.3
(Oph) Doliolaria (Dol)	0.6	X	0.5	0.3	X	0.3	0.8	X	0.9	0.5	X	0.5
Bryozoa												
Cyphonautes	3.9	3.4	4.8	0.7	0.6	0.4	4.8	0.6	2.5	1.7	1.5	3.0
(Cyp)	3.7	3.4	7.0	0.7	0.0	0.4	7.0	0.0	2.5	1.7	1.5	3.0
Brachiopoda												
Lingula (Lin)	2.6	5.8	3.3	2.5	X	2.7	6.2	8.5	5.2	6.2	0.8	6.8
0 \ /												

Table 2 continue	d:											
Ichthyoplankton Egg&Fish larvae (Eg&F)	12.7	7.4	12.9	14.7	14.5	10.4	5.3	7.3	9.5	8.9	4.2	10.3
TOTAL	23	16	24	21	14	21	25	20	26	23	17	23

Note: 'x' indicates absence of larvae, P-Pre-monsoon, M-monsoon, Po-Post-monsoon.

The seasonal higher larval diversity (24 larval forms) was recorded during the flood tide of post and pre-monsoon seasons at Zone (1). Similarly, along Zone (2) an average of 18 larval forms were documented, representing the flood tide of post and pre-monsoon seasons. The larval diversity was found to be minimal during the ebb tide

compared to the flood tide. In the lunar phase, larval ingress was higher during the full moon phase of post and premonsoon periods consisting of 26 larval forms at Zone (1) wherein, Zone (2) represented 17 larval representatives during the post-monsoon period (Tables 2 and 3).

Table 3: Tidal and lunar seasonal percentage (%) composition of larvae at zone 2.

ZONE 2

Larval forms/						ZON											
Abbreviation	F	lood tic	le	]	Ebb tid	e	F	ull moo	n	N	ew mo	on					
Abbieviation	P	M	PO	P	M	PO	P	M	PO	P	M	PO					
Cnidaria												<u>.</u>					
Ephyra (Ep)	X	X	X	X	X	X	X	X	X	X	X	X					
Planula (Pla)	X	X	X	X	X	X	X	X	X	X	X	X					
Actinula (Act)	1.1	X	1.7	1.5	X	2.0	1.0	X	1.4	1.1	5.4	0.8					
Platyhelminthes																	
Cercaria (Cer)	X	X	X	X	X	X	X	X	X	X	X	X					
Nematoda																	
Rhabditiform (Rha)	X	X	X	X	X	X	X	X	X	X	X	X					
Annelida																	
Polychaete (Pol)	3.7	11.8	5.3	3.2	2.5	5.7	6.1	1.0	7.2	5.0	3.7	9.1					
Arthropoda																	
Alima (Al)	X	X	X	X	X	X	X	X	X	X	X	X					
Barnacle nauplii (BNau)	8.5	2.7	10	5.0	1.0	7.7	8.4	3.7	9.0	9.6	11.6	8.0					
Barnacle cypris (BCyp)	13.5	7.1	5.5	6.2	3.6	4.6	6.0	4.5	7.1	9.9	2.7	6.6					
Shrimp nauplii (NauSr)	X	X	0.1	X	X	X	X	X	0.1	X	X	0.1					
Shrimp zoea (ZoSr)	7.1	12.6	9.9	16.1	7.6	6.4	8.3	1.8	6.8	8.6	X	7.9					
Shrimp mysis (MySr)	6.7	5.5	7.6	6.4	5.2	8.1	3.6	1.8	6.2	4.4	X	4.7					
Shrimp Post-larvae (PlSr)	2.9	1.4	2.8	5.9	4.1	0.9	9.3	2.7	4.2	9.6	X	5.0					
Ephausiid nauplii (EpNau)	0.1	1.1	0.4	X	X	X	4.3	X	1.7	2.0	X	1.8					
Crab zoea (CrZo)	9.1	11.5	11.1	10.4	8.3	14.6	11.0	10.0	9.0	7.1	10.7	12.6					
Megalopa (Meg)	0.5	0.5	0.2	0.3	X	0.3	0.2	X	0.1	X	X	0.1					
Crab Post-larvae (PlCr)	0.5	0.7	0.1	0.4	0.6	0.3	1.5	X	1.2	1.1	X	0.8					
Copepod nauplii (NaCop)	8.6	18.4	20.6	14.8	43.4	19.8	12.7	17.3	16	9.6	44.4	15.1					
Mollusca			- 0		40.4			40 =			• •	40.4					
Bivalve veliger (BiVvel)	9.0	5.4	6.0	6.2	10.4	6.3	3.3	40.7	6.9	7.1	3.0	10.4					
Gastropod veliger (GaVel)	7.1	5.3	7.1	5.1	2.4	7.1	3.6	3.2	8.3	8.1	X	6.2					
Echinodermata																	
Bipinnaria (Bip)	X	X	X	X	X	X	X	X	X	X	X	X					
Ophiopluteus (Oph)	X	X	X	X	X	X	X	X	X	X	X	X					

Table 3 continued:												
Doliolaria (Dol)	X	X	X	X	X	X	X	X	X	X	X	X
Bryozoa Cyphonautes (Cyp)	X	X	0.1	X	X	X	X	X	X	X	X	x
Brachiopoda Lingula (Lin)	6.9	1.4	2.0	1.0	0.8	3.4	8.0	2.5	5.3	3.9	X	3.0
<b>Ichthyoplankton</b> Egg&Fish larvae (Eg&F)	14.7	14.8	9.5	17.6	10.3	12.5	12.6	10.7	9.5	12.6	18.6	7.8
TOTAL	16	15	18	15	13	15	16	12	17	15	8	17

Note: 'x' indicates absence of larvae, P-Pre-monsoon, M-monsoon, Po-Post-monsoon.

Among the lunar phases, the new moon represented fewer larval forms than the full moon.

Larval abundance showed significant seasonal variation (*p*<0.001) between the studied zones through the Kruskal-Wallis test (Table 5). Furthermore, Mann-Whitney's U test signifies the significance between pre-monsoon and monsoon seasons (PRE and MON),

followed by monsoon and post-monsoon seasons (MON and POM), respectively (p<0.05). There were no significant variations (p>0.05) between the pre-monsoon and post-monsoon seasons (PRE and POM) for the tidal cycle. However, a significant difference was observed during the full moon phase at Zone (1) (Table 6).

Table 4: Mean, SD and larval percentage (%) composition (zone 1 and zone 2).

Larval forms	Mean (ind. m <sup>-3</sup> )	SD (ind. m <sup>-3</sup> )	Percentage (%)
Cnidaria			
Ep	492.9	921.99	1.5
Pla	28.6	124.09	0.1
Act	720.71	1121.59	2.2
Platyhelminthes			
Cer	21.09	139.2	0.1
Nematoda			
Rha	17.47	110.82	0.1
Annelida			
Pol	1904.8	2106.12	5.7
Arthropoda			
Al	2.08	21.18	0
BNau	2737.11	2631.58	8.3
BCyp	2243.87	2139.60	6.8
NauSr	258.81	530.69	0.8
ZoSr	2714.11	2632.42	8.20
MySr	1580.81	1700.77	4.80
PlSr	1325.51	1637.11	4.00
EpNau	606.85	1334.91	1.80
CrZo	2828.63	2604.32	8.50
Meg	110.26	247.22	0.30

Table 4 continued:			
PlCr	233.35	510.72	0.70
NaCop	5054.89	3728.3	15.30
Mollusca			
BiVvel	2271.77	2273.74	6.90
GaVel	2014.35	1997.96	6.10
Echinodermata			
Bip	61.63	210.50	0.20
Oph	344.02	999.40	1.00
Dol	100.44	251.32	0.30
Bryozoa			
Сур	511.66	1323.72	1.50
Brachiopoda			
Lin	1483.96	2042.28	4.5
Ichthyoplankton			
Eg&F	3463.13	2882.09	10.3

The total mean larval abundance during the full moon phase was  $(2059\pm2809 \text{ ind. m}^{-3})$ , of which post-monsoon season represented maximum mean

abundance of  $(3297\pm3502 \text{ ind. m}^{-3})$  (Table 5 and Fig. 3).

Table 5: Total mean larval abundance (mean±SD) between seasons (Kruskal-Wallis test).

	PRE	MON	POM	Total mean abundance (ind. m <sup>-3</sup> )	df	P
	mean±SD	mean±SD	mean±SD	mean±SD	-	
Zone1 FT	996±1402	557±1203	1880±2666	1201±2020	2	< 0.001
Zone2 FT	870±1241	327±649	1720±2816	1030±1995	2	< 0.001
Zone1 EbT	844±1313	337±809	$1704\pm2424$	1019±1819	2	< 0.001
Zone2 EbT	831±1380	203±555	1167±1883	767±1481	2	< 0.001
Zone1 FM	1761±2140	812±1514	3297±3502	2059±2809	2	< 0.001
Zone2 FM	1356±1729	239±957	2369±3023	1402±2339	2	< 0.001
Zone1 NM	1462±1732	411±950	2720±3060	1623±2388	2	< 0.001
Zone2 NM	1223±1572	113±389	1776±2522	1094±1928	2	< 0.001

Table 6: Total mean larval abundance (mean±SD) with significance difference between two seasons (Mann-Whitney's U test).

	PRE*MON	PRE*MON	PRE*POM		PF	PRE*MON				
ZONES	Z	Asymp. Sig.(2-tailed)	Z	Asymp. Sig.(2-tailed)	Z	Asymp. Sig.(2-tailed)				
Z1FT	3.496	0.001*	-2.048	0.122	-5.734	0.001*				
Z2FT	2.869	0.012*	-1.011	0.937	-4.035	0.001*				
Z1EbT	3.266	0.003*	-2.365	0.054	-5.808	0.001*				
Z2EbT	3.357	0.002*	-0.789	1.000	-4.327	0.001*				
Z1FM	3.861	0.001*	-2.874	0.012*	-6.944	0.001*				
Z2FM	4.549	0.001*	-1.325	0.555	-6.121	0.001*				
Z1NM	5.334	0.001*	-2.283	0.067	-7.905	0.001*				
Z2NM	6.040	0.001*	-0.234	1.000	-6.601	0.001*				

<sup>\*</sup>Implies the result is significant at p<0.001, Z is absolute average value, PRE-pre-monsoon; MON-monsoon; POM- post-monsoon.




Figure 3: Total mean abundance of larval distribution across zones according to seasonal, tidal and lunar cycle.

Larval diversity indices

The Shannon Wiener (*H*) diversity index (mean=2.44, SD= 0.31) ranged from 1.64 to 2.92. During the study period, the Shannon wiener index increased at Zone (1) of the full moon during post-monsoon season (26 larvae

and H=2.92) (Fig. 4). However, diversity index enormously varied from studied zones explaining Zone (1) with highest diversity (post-monsoon season) and Zone (2) with minimum diversity predominantly in monsoon season.




Figure 4: Shannon Wiener (H) diversity indices.

Redundancy data analysis (RDA) for larval distribution

The results of multivariate Redundancy Data **Analysis** (RDA) combining physicochemical variables with dominant larval communities were depicted in triplots (Figs. 5 to 7). The larval-environmental relations for the first two axes varied from 66.3 to 22.6% and 47.7 to 46.9% for the tide and lunar phases, respectively (Figs. 5 and 6). Among the environmental factors, the Monte Carlo permutations showed that salinity, test water temperature, dissolved oxygen and pH were significant in explaining the relationship between the larval and tidal cycle (F=13.521, P=0.002) contrarily, there was no statistically significant variation (F=3.93, P=0.68) observed for larvae and lunar phase.

The RDA (Fig. 5) explains a strong positive correlation of larvae like Rhabditiform. Planula. Bipinnaria. Cyphonautes, Megalopa, Ophiopluteus, Doliolaria, Ephyra, Shrimp nauplii and Lingula with salinity and pH during the flood tide at Zone (1). Further, the larval distributions. viz.. **Bivalve** Barnacle nauplii, Barnacle veliger, cypris, Actinula, Gastropod veliger, Egg & fish and Crab post-larvae were positively correlated to water temperature during the ebb tide.




Figure 5: The RDA ordination triplot for larvae with respect to environmental parameters and tidal cycle. Environmental parameters (continuous line) and larvae (dotted lines) are shown by arrows, tides with zones are displayed by filled dots. (Abbreviations: WT-Water temperature (°C), DO- Dissolved oxygen (mg/L), Sal- Salinity; Z1FT- Zone 1 Flood tide, Z1EbT – Zone 1 ebb tide, Z2FT- Zone 2 Flood tide, Z2EbT- Zone 2 ebb tide; Larval abbreviation presented in Table 2).

Similarly, the RDA plot for the lunar phase explains the positive correlation for larvae such as Megalopa, Rhabditiform, Bipinnaria, Ophiopluteus, Cercaria, Alima, Doliolaria, Cyphonautes, Planula, Shrimp nauplii, Actinula and Ephyra with salinity during the full moon and new moon phases at Zone (1) (Fig. 6).

The RDA eigenvalues for the larvalenvironmental relationship for the first two axes between seasons were found to vary from 54.5 to 14.2%, registering maximum larvae during the postmonsoon and pre-monsoon seasons (Fig. 7).




Figure 6: The RDA ordination triplot for larvae with respect to environmental parameters and the lunar cycle. Environmental parameters (continuous line) and larvae (dotted lines) are shown by arrows, lunar phase with zones is displayed by filled dots. (Abbreviations: WT-Water temperature (°C), DO-Dissolved oxygen (mg/L), Sal-Salinity; Z1FM-Zone1 Full moon, Z1NM- Zone1 new moon, Z2FM -Zone2 Full moon, Z2NM- Zone2 new moon; Larval abbreviation presented in Table 2).




Figure 7: The RDA ordination triplot for larvae with respect to environmental parameters and seasons. Environmental parameters (continuous line) and larvae (dotted lines) are displayed by arrows, filled dots represent seasons (Abbreviations: WT- Water temperature (°C), DO- Dissolved oxygen), Sal- Salinity; PRE- Pre-monsoon, MON-Monsoon, POM- Post-monsoon; Larval abbreviation presented in Table 2).

### Discussion

Tropical estuaries are biogeochemical hotspots with higher primary production, organic matter and nursery grounds (Baran, 2000). Being the transitional zone between freshwater and marine environment, hydrological parameters vary significantly. In the present study, significant seasonal variations of hydrological parameters observed, which of water were temperature, salinity and dissolved oxygen markedly fluctuated along the study zones. These changes are mainly attributed to regular tidal fluctuations and seasonal changes in the freshwater flow (Costa et al., 2009a). Further, water parameters are generally stable during non-rainy seasons due to wellmixed tidal salinity, pronounced solar radiation and reduced turbidity along the tropical estuary (Rajaneesh and Mitbavakar, 2013). In the present study, water pH was consistent with persistent alkaline conditions throughout the study at both zones. Dissolved oxygen is the most vital parameter of water quality for biological productivity, which was ambient during the study. Dissolved oxygen levels are affected by rising temperature and salinity during nonrainy seasons (Saravanakumar et al., 2008) and similar trends were observed during the study. The present study infers that salinity is one of the most important parameters influencing the distribution of larvae across the Kali estuary (p<0.05). Seasonal salinity variations were pronounced during the monsoon season compared to pre and post-monsoon, mainly due to the southwest monsoon and river runoff (Bhat, 1988; VishnuRadhan, 2015; Vineetha *et al.*, 2015; Kumbhar *et al.*, 2020).

The relationship between the distribution of larvae and hydrological parameters has been studied in several estuaries around the globe. Salinity, water temperature, pH, chlorophyll a, and turbidity are the most important parameters which influence distribution and abundance of larvae (Mouny and Dauvin, 2002). The geoclimatic larval distribution states that salinity mainly regulates larval abundance distribution and along tropical estuaries (Costa et al., 2009b; Atique et al., 2016), wherein, water temperature is responsible across temperate estuaries (Villate, 1994). Moreover, studies showed that the salinity gradient restricts the spatial distribution of planktonic communities across the estuaries due to the explicit tolerance range (Vezi et al., 2019; Rose et al., 2019).

In this study, 26 larval forms were recorded and their distributions were higher at Zone (1), particularly during the post-monsoon season due to the rising tides (Srichandan et al., 2021). Subsequently, Zone (2) is considered a mesohaline zone with 18 larval forms which may be considered as estuarine recruiters. The larval forms such as Ephyra, Planula, Alima (Squilla), Bipinnaria, Ophiopluteus, Cercaria, Rhabditiform and Doliolaria are

observed along Zone (1), indicating tidal ingress/salinity passive Similar studies carried out along the prominent estuaries of the east coast of India reported 17 to 21 larval forms with higher larval density during the and pre-monsoon post season. Moreover. their abundances are attributed to the organic matter. catchment area, and stable hydrological post-rainy parameters season (Santhanam and Perumal 2003: Baliarsingh et al., 2014). Such results are well-matched with the present study observations.

With reference to the percentage composition of the larval distribution, Copepod nauplii (15.30%), Egg & fish larvae (10.52%), Shrimp zoea (8.20%), Barnacle nauplii (8.30%) and Bivalve veliger (6.9%) were found to be dominant during the study period. The following larvae have prolonged larval duration and maximum larval forms were benthic in habitat. Copepod nauplii were numerically dominant throughout the studied period, which is attributed to continuous breeding and reproductive higher capacity subsequently, adaptation to variable environmental conditions (Ramaiah and Nair, 1997; An et al., 2012; Baliarsingh et al., 2014). Dominate larval forms recorded during monsoon season (Copepod nauplii, Egg & fish larvae, Bivalve veliger, and Gastropod veliger) can be attributed to their euryhaline adaptations; subsequently, the occurrence of Egg & fish larval coincides with the breeding pattern induced by the season (Santhanam and

2003). A study on Perumal. the barnacle larval population along the well-mixed estuarine gradient reported higher ingress during the post-monsoon period at Mondovi and Zuari estuarine complexes (Grinson et al., 2013), which represents geographical and hydrological similarities with the Kali estuary. Larvae follow the tidally driven migration pattern with higher abundance during the flood tide at both stations. Zone (1) mainly comprised marine habited larval diversity, which indicates intrusion from the adjacent sea, wherein larval compositions along Zone (2) represent marine-brackish with horizontal dispersion to their preferred habitat. During the study, crustacean larval ingress was higher during the lunar phase which demonstrates nocturnal tidal upward movement similar to the study of Srichandan et al. (2021).

ordination The of larval and environmental relationship bv Redundancy Data analysis showed that the increasing salinity influences the dispersal of larvae and other species due to incoming tidal contributes (Mohan and Rao, 1972; Vineetha et al., 2015; Srichandan et al., 2021) whereas, the lunar-tidal cycle is related to the spawning period due to the intensity of moonlight and plays an important role in structuring dispersion of larval forms (Pankhurst and Porter, 2003; Oliveira et al., 2009; Takemura et al., 2010). Crucially, the low diversity and abundance during the monsoon season are due to changes in the coastal current followed by monsoonal rainfall and river runoff (Rakesh *et al.*, 2008).

Hence, the present study surmises that the larval distributions are typically complemented by the tidal, lunar phases and seasonally fluctuating environmental parameters that shape the estuarine larval complex.

Kali estuary is a critically vulnerable coastal area (CVCA) and its effective ecosystem management and biodiversity accounts are valuable. The estuary supports traditional fishing activity throughout the year and the inner reach of Zone (1) and the meeting point of Zone (2) is considered a bivalve bed. Subsequently, Zone (2) is known as the shrimp fishing basket sustaining the associated fishermen's livelihood. The present study generates the baseline database on larval diversity and distributions, which are critical components of the estuarine food web. Hence, considering the Kali estuary a delicate larval pool and protecting its sensitive habitat is crucial in juvenile management and conservation.

## Acknowledgements

We thank to University Grants commission (UGC) and Karnatak University, Dharwad's for providing fellowship to carryout research work (No. KU/Sch/URS/2017-18/648).

### References

An, P.X., Du, H.Z., Zhang, H.J., Li, P.Y. and Qi, W.J., 2012. Structure of Zooplankton community in Hulunk lake, China. *Procedia* 

- Environmental Science, 13, 1099-1109.
- Archambault, P. and Bourget, E., Influence of shoreline configuration on spatial variation of meroplanktonic larvae, recruitment and diversity of benthic subtidal communities. Journal of **Experimental** Marine Biology, 238(2), 161-184. https://doi.org/10.1016/S0022-0981(98)00146-4
- Atique, P., Costa, D.K.G., Monteiro, M.C., Pereira, L.C. and Coasta, R.M., 2016. Copepod assemblages in a highly dynamic equatorial estuary on Brazilian Amazon coast. *Marine Ecology*, 38(1), e12385. https://doi.org/10.1111/maec.12385.
- Badsi, H., Ali, H.O., Loudiki, M., Hafa, E.M., Chakli, R. and Aamiri, A., 2010. Ecological factors affecting the distribution of zooplankton community in the Massa Lagoon (Southern Morocco). African Journal of Environmental Science and Technology, 4(11), 751-762.
- Baliarsingh, S.K., Srichandan, S., Biraja, K., Sahu, K.C., Lotliker, A.A. and Kumar, S. T., 2014. Zooplankton community distribution along offshore transects of some Indian estuaries of east coast: A taxonomic investigation during a summer cruise. *Indian Journal of Geo-Marine Science*, 43, 1781-1791.
- **Baran, E., 2000.** Biodiversity of estuarine fish faunas in West Africa. Naga, ICLARM Quarterly, 23, 4-9.
- Bhat, U.G., Neelakantan., Kusuma,

- N. and Naik, U.G., 1988. Environmental characteristics of the marine and estuarine habitats of Karwar: an overview. *Journal of Indian Fisheries Association*, 18, 401-412.
- Buskey, J.E., 2018. Recruitment of estuarine dependent species of commercial and recreational importance through the Aransas Ship channel. University of texas libraries, Austin. *Marine Science*.
- Cloern, J.E., Jassby, A.D., Schraga, T.S., Nejad, E. and Martin, C., 2017. Ecosystem variability along the estuarine salinity gradient: Examples from long-term study of San Francisco Bay. *Journal of Limnology and Oceanography*, 62, S272-S291.

https://doi.org/10.1002/Ino.10537

- Conway. **P.V.D..** 2012a. Marine zooplankton of southern Britain. Part Radiolaria. Heliozoa. Foraminifera, Ciliophora, Cnidaria, Ctenophora, Platyhelminthes, Nemertea, Rotifera and Mollusca. A. John Occasional G. (ed.). Publications, Marine **Biological** Association of the United Kingdom, No. 25, Plymouth, United Kingdom, 138.
  - http://dx.doi.org/10.13140/2.1.3274. 3369
- Conway, P.V.D., 2012b. Marine zooplankton of southern Britain Part 2: Arachnida, Pycnogonida, Cladocera, Facetotecta, Cirripedia and Copepoda. Occasional Publication of the Marine Biological Association No. 26, Plymouth (UK).

- Marine Biological Association of the United Kingdom, 163. http://dx.doi.org/10.13140/2.1.4704. 4800
- Costa, L.A., Huszar, V.L.M. and Ovalle, A.R., 2009a. Phytoplankton functional groups in a tropical estuary: Hydrological control and nutrient limitation. *Estuaries and Coasts*, 32(3), 508-52. http://dx.doi.org/10.1007/s12237-009-9142-3
- Costa, R.M., Leite, R.N. and Pereira, C.C.L., 2009b. Mesozooplankton of the Curuca estuary (Amazon coast, Brazil). Proceedings of the 10<sup>th</sup> International Coastal Symposium. *Journal of Coastal Research*, Special Issue, 56, 400-404.
- **CRZ** (Coastal regulation zone CRZ), **2011**. The Gazette of India part 2 section 3 sub-section (ii) published by authority. New Delhi
- Darboe, F.S., 2002. Fish species abundance and distribution in the Gambia Estuary. United Nations University, Fisheries training program. Final Project 2002. Reykjavik, Iceland: United Nations University.
- Dias, E., Barros, A.G., Hoffman, J.C., Antunes, C. and Morais, P., 2020. Habitat use and food sources of European flounder larvae (Platichthys flesus, L. 1758) across the Minho River estuary salinity gradient (NW Iberian Peninsula). Regional Studies in Marine Science, 34.
  - http://dx.doi.org/10.10.1016/j.rsma.2 020.101196

- **Forward, Jr R.B. and Tankersley, A.R., 2001**. Selective tidal-stream transport of marine animals. *Oceanography and Marine Biology*, 39, 305-353.
- Garrison, L.P., 1999. Vertical migration behavior and larval brachyuran transport in crabs. Marine Ecology Progress Series, 176. 103-113. http://dx.doi.org/10.3354/meps17610 3
- Goncalves, A.M., Pardal, M.A., Marques, S.C., De Troch, M. and Azeiteiro, U.M., 2010. Distribution and composition of small-sized zooplankton fraction in a temperate shallow estuary (western Protugal). Fresenius Environmental Bulletin, 19, 3160-3176.
- Goswami, S.C. and Usha, G., 1992. Lunar, diel and tidal variability in penaid prawn larval abundance in Mondovi estuary, Goa. *Indian Journal of Marine Science*, 21, 21-25.
- Grinson, G., Desai, D.V., Gaonkar, C.A., Aboobacker, V.M., Vethamon, P. and Anil, A.C., 2013.

  Barnacle larval transport in the Mondovi-Zuari estuarine system, Central west coast of India. *Journal of Oceanography*, 69(4), 451-466. https://doi.org/10.1007/s 10872-013-0186-4.
- Hare, J.A., Thorroid, S., Walsh, H.,
  Reiss, C., Valle- Levinson, A. and
  Jones, C., 2005. Biophysical mechanism of larval fish ingress into the Chesapeake Bay. *Marine Ecology Progress Series*, 303, 295-

310. https://doi.org/10.3354/meps303295

- Holt, G.J. and Holt, S.A., 2000. Vertical distribution and the role of physical processes in the feeding dynamics of two larval sciaendis Sciaenops ocellatus and Cynoscion nebulosus. Marine Ecological Progress Series, 193, 181-190. https://doi.org/10.1111/maec.12385.
- Katsanevakis, S., Lefkaditou, G.E., Galinou- Mitsoudi, S., Koutsoubas, D. and Zenetos, A., 2008. Molluscan species of minor commercial interest in Hellenic seas: Distribution, exploitation and conservation status. *Mediterranean Marine Science*, 9(1), 77-118. https://doi.org/10.12681/mms.145.
- Kritzer, J.P. and Sale, P.F., 2010. Marine metapopulations. Elsevier.
- Kumbhar, N., Thomas, S. and Loka, J., 2020. Seasonal abundance and diversity of finfish in a tropical estuary, Karwar, Southwest coast of India. *Journal of Marine Biological Association*, *India*, 62(2), 79-84. http://dx.doi.org/10.6024/jmbai.2020.62.2.2218-09
- **Kunze, H. B., Morgan, S. G. and Lwiza, K. M., 2013**. Field test of the behavioral regulation of larval transport. *Marine Ecology Progress Series*, 487(1), 71-87. https://doi.org/10.3354/meps1028.
- Leps, J. and Smilauer, P., 2003.

  Multivariate analysis of ecological data using CANOCO. Cambridge University Press, Cambridge. https://doi.org/10.1111/j.1442-993.2005.01433.x

- Mesta, P., Chandran, S.D.M. and Ramachandran, **V.T..** 2016. Mangrove mapping of Uttara Kannada District. Lake 2016: Conference on conservation sustainable management of ecologically sensitivity regions in Western Ghats (Ed.). Indo- German Environmental Program - GIZ.
- Mohan, C.P. and Rao, S.S.T., 1972. Tidal cycle studies in relation to zooplankton distribution in the Godavari estuary. *Proceedings. Indian Academy Sciences*, 75, 23-31. https://doi.org/10.1007/BF03045331
- Mouny, P. and Dauvin, J.C., 2002. Environmental control mesozooplankton community structure in the Seine estuary (English Channel). Oceanologica 25 Acta. (1).13-22. http://dx.doi.org/10.1016/S0399-1784(01)01177-X
- Oliveira, C., Vera, L.M., Lopez-Olmeda, J.F., Guzman, J.M., Manonos, E., Ramos, J. and Sanchez-Vazquez, F.J., 2009. Monthly day/night changes seasonal daily rhythms of sexual steroids in Senegal Sole (Solea senegalensis) under natural fluctuating controlled or environmental conditions. *Comparative Biochemistry* PhysiologyPart A: Molecular and Integrative Physiology, 152(2), 168-175.
  - https://doi.org/10.1016/j.cbpa.2008.0 9.012
- Pankhurst, N.W. and Porter, M.J.R., 2003. Cold and dark or warm and

- light: variations on the theme of environmental control of reproduction. *Fish Physiology and Biochemistry*, 28, 385-389.https://doi.org/10.1023/B:FISH. 0000030602.51939.50
- Perumal, N.V., Rajkumar, M., Perumal, P. and Rajasekar, T.K., 2009. Seasonal variation of plankton diversity in the Kaduviyar estuary, Nagapattinam, southeast coast of India. *Journal of Environmental Biology*, 30 (6), 1035-1046.
- Queiroga, H. and Blanton, J., 2004. Interaction between behavior and physical forcing in the control of horizontal transport of Decapoda crustacean larvae. *Advance in Marine Biology*, 47, 107-214. https://doi.org/10.1016/s0065-2881(04)47002-3
- Rajaneesh, K.M. and Mitbavkar, S., 2013. Factors controlling the temporal and spatial variation in Synechococcus abundance in monsoonal estuary. *Marine Environmental Research*, 92, 133-143.
  - https://doi.org/10.1016/j.marenvres.2 013.09.010
- Rakhesh, M., Raman, A.V., Kalavati, C., Subramanian, B.R., Sharma, V.S., Sunitha., Babu, E. and Sateesh, N., 2008. Zooplankton community structure across an eddy generated upwelling band close to a tropical bay-mangrove ecosystem. *Marine Biology*, 154, 953-972. https://doi.org/10.1007/s00227-008-0991-2

- Ramaiah, N. and Nair, V., 1997.

  Distribution and abundance of Copepods in the pollution gradient zones of Bombay harbor- Thane creek-Basin Creek, west coast of *Indian Journal of Marine Science*, 26, 20-25.
- Rao, A.V.P., 1970. Observation on the larval ingress of the Milk fish, *Chanoschanos* (Forskal) into the Pulicate Lake. *Indian Journal of Marine Biological Association*, 13(2), 249-257.
- Rose, T.H., Tweedley, J.R., Warwick, R.M. and Potter, I.C., 2019.
  Zooplankton dynamics in a highly eutrophic microtidal estuary. *Marine Pollution Bulletin*, 142, 433-451. http://dx.doi.org/10.1016/j.marpolbul .2019.03.047
- Santhanam, P. and Perumal, P. 2003.

  Diversity of zooplankton in Parangipettai coastal waters, southeast coast of India. *Journal of Marine Biological Association*, *India*, 45(2), 144-151.
- Saravanakumar, A., Rajkumar, M., Sesh., Serebiah, J. and Thivakaran, G.A., 2008. Seasonal variations physic-chemical in characteristics of water, sediment and soil texture in arid mangroves of Kachchh-Gujarat. Journal of Environmental Biology, 29(**5**), 725-732.
- Shannon, C.E., 1948. The mathematical theory of communication. *Bell System Technical Journal*, 27(3), 379 423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x

- **Sokal, R.R. and Rohlf, J.F., 1995**. Biometry. The principles and practice of Statistics in Biological Research. 3<sup>rd</sup> edition. W. H. Freeman and Co., New York.
- Sreekanth, G.B., Lekshmi, N.M. and Singh. N.P., 2017. **Temporal** in Fish community patterns structure: Environmental perturbation form a well- mixed Tropical Estuary. Proceedings of the National Academy of Sciences, India Section B: Biological Sciences, 87, 135-145. https://doi.org/10.1007/s4011- 015-
- Srichandan, S., Baliarsingh, K.S., Lotliker, A.A., Sahu, K.B., Roy, R. and Nair, B.T.M., 2021.

  Unravelling tidal effect on zooplankton community structure in tropical estuary. *Environmental Monitoring and Assessment*, 193, 362. https://doi.org/10.1007/s10661-021-09112-z

0581-2.

- Staples, D.J. and Vance, J.D., 1985. Shortterm and longterm influences on the immigration of post larval banana prawns Penaeus merguiensis, into a mangrove estuary of the Gulf of Carpentaria, Australia. Marine Ecology Progress Series. 23(1), 15-29. http://dx.doi.org/10.3354/meps02301 5
- Suthers, I.M. and Rissik, D., 2009. Plankton: A guide to their Ecology and Monitoring for Water Quality. *Journal of Plankton Research*, 32(2), 261-262.

- https://doi.org/10.1111/j.1442-9993.2012.02360.x
- **Takemura, A. and Rahman, M.S., Park, Y.J., 2010**. External and internal controls of lunar-related reproductive rhythms in fishes. *Journal of Fish Biology*, 76, 7-26. https://doi.org/10.1111/j.1095-8649.2009.02481.x
- Tanmoy, N. and Sumit M., 2020. Unravelling the spatio-temporal variation of zooplankton community from the river Matla in the Sundarbans Estuarine System, India. *Oceanologia*, 62 (3), 326-346. https://doi.org/10.1016/j.oceano.202 0.03.005
- Tarimo, B., Winder, M., Mtolera, M.S.P., Muhando, C.A. and **2022**. Gullstrom, M., Seasonal distribution of Fish larvae in mangrove-seagrass seascapes of Zanzibar (Tanzania). Scientific Report, 12. 4196. https://doi.org/10.1038/s41598-022-07931-9
- Vezi, M., Downs, T.C., Wepener, V. and O'Brien, G., 2019. Response of zooplankton communities to altered water quality and seasonal flow changes in selected river dominated estuaries in KwaZulu-Natal, South Africa. *Ecohydrology and Hydrobiology*, 19(3), 393- 406. http://dx.doi.org/10.1016/j.ecohyd.2 019.01.005
- Villate, F., 1994. Temporal variability of the spatial distribution of the zooplankton community in a coastal embayment of the Basque country in relation to physical phenomena.

*Hydrobiologia*, 288, 79-95. http://dx.doi.org/10.1007/BF000071 28

Vineetha, G., Jyothibabu, R., Madhu, N.V., Kusum, K.K., Sooria, P.M., Shivaprasad, A., Reny, P.D. and Deepak, M.P., 2015. Tidal influence on the diel vertical migration pattern of zooplankton in a tropical monsoonal estuary. *Wetlands*, 35, 597-610. http://doi.org/10.1007/s13157-015-0650-6

Vishnuradhan, R., 2015. Southwest monsoon influence the water quality and waste assimilative capacity in the Mondovi estuary (Goa state, India). *Chemistry and Ecology*, 31(3), 217-234. https://doi.org/10.1080/02757540.20 14.961435

Weiss, H., 2017. Key to the Larvae of Common Decapoda Crustaceans (Lobster, Crabs and Shrimps) in Long Island Sound. Project Oceanology.

Young, C.M., Sewell, M.A. and Rice, M.E., 2002. marine **Atlas** of invertebrate larvae. Academic Press, New York, N. Aquaculture 34, Research, 437-437 https://doi.org/10.1046/j.1365-2109.2003.00819.x