

Research Article

Dietary effect of chitosan on immunity, antioxidant status, and immune-related gene expression in juvenile starry sturgeon (*Acipenser stellatus*)

Zakeri D.¹, Pazooki J.^{1*}, Mohseni M.², Jamshidi S.²
¹Department of Animal Sciences and Marine Biology, Faculty of Life Sciences and Biotechnology, Shahid Beheshti University, Tehran, Iran

²International Sturgeon Research Institute, Iranian Fisheries Science Research Institute (IFSRI), Agricultural Research, Education and Extension Organization (AREEO), Rasht, Iran

*Correspondence: pazooki2001@yahoo.com

Keywords

Chitosan,
Aquaculture,
Stellate sturgeon,
Inflammatory genes,
Real-time PCR

Abstract

This study investigated the effects of the dietary supplement chitosan (CH) on various parameters including hematological indices, immune response, liver enzyme levels, antioxidant capacity, and the expression of immune-related genes in juvenile starry sturgeon (*Acipenser stellatus*). 180 juvenile starry sturgeons with an average weight of 31.90 ± 0.73 g were divided into five feeding groups for 60 days includes: control (CTR), 1.5 g (CH_{1.5}), 3.0 g (CH_{3.0}), 4.5 g (CH_{4.5}), and 6.0 g (CH_{6.0}) of CH powder per kg of a basal diet. Blood analyses conducted after the dietary trial demonstrated that CH significantly increased total red blood cells, white blood cells, hematocrit, and hemoglobin levels, with the highest values recorded in the CH_{6.0} group. CH enhanced immune responses as evidenced by elevated levels of total protein, albumin, globulin, lysozyme activity, and immunoglobulin, particularly in the CH_{6.0} group. Additionally, dietary CH reduced serum lipid levels, and the lowest liver enzyme activities were observed in the CH_{4.5} and CH_{6.0} groups. CH also stimulated the activity of antioxidant enzymes, with the most pronounced antioxidant activity noted in the CH_{6.0} group. Moreover, CH improved the regulation of the relative expression of interleukin-1 beta (*IL-1β*) and tumor necrosis factor alpha (*TNF-α*) genes in liver, intestine, spleen, and head kidney tissues, with the highest expression levels for both genes recorded in the CH_{6.0} group. These findings suggest that the inclusion of 6 g kg⁻¹ CH in the diet of starry sturgeon enhances hematological and biochemical parameters, boosts antioxidant capacity and immune response, and supports immune-related gene expression.

Article info

Received: July 2025

Accepted: February 2026

Published: July 2026



Copyright: © 2025 by the authors.
Licensee MDPI, Basel, Switzerland.
This article is an open access article
distributed under the terms and
conditions of the Creative Commons
Attribution (CC BY) license
(<https://creativecommons.org/licenses/by/4.0/>).

Introduction

The aquaculture industry is undergoing rapid development and has increasingly shifted toward intensive farming practices. A significant challenge in fish farming is the declining survival rates against disease and contamination, particularly during the early life stages, where stress from high fish density can negatively impact growth performance, immune response, and increase susceptibility to diseases. The use of antibiotics to treat diseases poses further issues, including environmental risks, the emergence of bacterial resistance, the accumulation of these substances in farmed fish, and potential transmission to humans, the ultimate consumers (Harikrishnan *et al.*, 2012). Consequently, the use of antibiotics in aquaculture is prohibited in many countries (Yu *et al.*, 2022). To address these concerns, it is crucial to enhance the immune systems of fish, especially in economically important species. Research has increasingly focused on environmentally friendly alternatives, such as dietary supplements that can bolster the aquatic immune system.

Recently, the use of immunostimulants in farmed fish has risen, positioning them as viable alternatives to antibiotics. Immunostimulants, either natural or synthetic substances, serve to stimulate and regulate the immune system, thereby enhancing immune responses in the body. They can increase phagocytic activity, antibody production, lysozyme activity, and antioxidant enzyme activity, among other effects (Anderson, 1992). Among these immunostimulants, biological materials, particularly by-products from

other industries, are recognized for their environmental friendliness.

Chitosan (CH), β -(1-4)-N-acetyl-D-glucosamine, a biopolymer derived from the deacetylation of chitin, is a biological immunostimulant. It is found in the exoskeletons of insects, the shells of crustaceans, and the cell walls of fungi (Perini *et al.*, 2020). CH is characterized by its non-toxic, biodegradable, and biocompatible properties, along with numerous physiological functions, which include antibacterial (Kong *et al.*, 2010), antioxidant (Chang *et al.*, 2018), antitumor (Qin *et al.*, 2002), hypolipidemic (Zhang *et al.*, 2012), growth-promoting (Imefon Udo, 2018), and immune-stimulating effects (Fadl *et al.*, 2020). Recently, CH has garnered considerable attention in aquaculture, with studies highlighting its positive impacts on growth performance, intestinal morphology, non-specific immunity, pathogen resistance, antioxidant activity, and lipid metabolism in various fish species (Chen and Chen, 2019; Dawood *et al.*, 2020; Younus *et al.*, 2020; Stanek *et al.*, 2023).

The antioxidant potential of CH has also been increasingly recognized. During metabolic processes, reactive oxygen species (ROS) are generated, capable of interacting with crucial biological macromolecules such as lipids, proteins, and nucleic acids. Under normal circumstances, a delicate balance exists between ROS production and elimination. However, any disruption of this balance can lead to oxidative stress (Yan *et al.*, 2017). The antioxidant role of CH has been established across various fish species (Dawood *et al.*, 2020; Abdel-Tawwab *et*

al., 2021), with antioxidant status serving as an indicator of an organism's ability to effectively remove ROS (Enomoto *et al.*, 2001).

Sturgeons are among the most valuable aquatic species, yet their populations face threats of extinction due to overfishing, habitat loss, and various forms of pollution (Cvijanović *et al.*, 2024). Accordingly, artificial breeding for these fish presents a promising approach to conservation and sustainability. The starry sturgeon (*Acipenser stellatus*) is one of the most valuable sturgeon, which is valuable for caviar production due to its early sexual maturity and is considered one of the most profitable aquaculture species. However, farmed sturgeon suffer from infectious diseases, with the development of sturgeon farming, it is necessary to use immunostimulants to strengthen the immune system of these fish.

Since chitosan (CH) exerts multifaceted effects on immune gene regulation, antioxidant activity, and lipid metabolism, its therapeutic efficacy in enhancing physiological functions is contingent upon the specific requirements of different fish species. Therefore, studying individual species is crucial for determining the most effective use of CH. While recent advances have illuminated the efficacy of CH across various fish species, there remains limited information regarding its effects in sturgeon. This study aimed to investigate the impact of dietary CH on the immune response, antioxidant status, liver enzymes, and the expression of immune-related genes, specifically interleukin-1 beta (*IL-1β*) and tumor necrosis factor alpha (*TNF-α*), in starry sturgeon.

Materials and methods

Ethical statement

All animal experiments performed in this study were ethically approved by the Shahid Beheshti University Ethics Committee for the Use of Animals (IR.SBU.REC.1402.171).

Chitosan extraction

Shrimp shells (*Penaeus semisulcatus*) were washed to remove residual tissues and impurities, then dried at 60°C for 24 h and ground to obtain a homogeneous powder. The extraction of chitosan followed the method outlined by Fernandez-Kim (2004). To eliminate CaCO₃, 7% hydrochloric acid was added to the shrimp shell powder at a ratio of 1:20 (w v⁻¹) for 4 h at 65°C. Following this, 9% sodium hydroxide (1:20 w v⁻¹) was introduced for an additional 4 h at 65°C. Decolorization was performed using 0.32% sodium hypochlorite (1:15 w v⁻¹) for 5 min at room temperature to yield chitin. Subsequently, deacetylation was executed by adding 55% NaOH for 1 h at 75°C at a 1:20 (w v⁻¹) ratio to extract chitosan. The degree of deacetylation was assessed using Fourier-transform infrared spectroscopy (FTIR) (TENSOR 27, Bruker Optics Inc, Billerica, MA, USA) through the KBr tablet method over a wavelength range of 400-4000 cm⁻¹, revealing a deacetylation degree of 80%.

Preparation of diets and study design

Five specific diets were formulated, as detailed in Table 1. A basal diet without CH served as the control (CTR), while four additional diets incorporated 1.5 g (CH_{1.5}), 3.0 g (CH_{3.0}), 4.5 g (CH_{4.5}), and 6.0 g (CH_{6.0}) of chitosan powder per kg of the

basal diet. The diets were prepared via the extrusion method, whereby all dry ingredients were ground and mixed thoroughly using an electric mixer (Raybony, Garma Electric, Amol, Iran) for 10 min. Subsequently, the mixture was combined with fish oil and water to achieve

a paste-like consistency. This dough was then processed through an extruder system (Polylab, Thermo Fisher Scientific Company, USA) and dried at 40°C for 24 h. The dried feed was stored at -20°C until required.

Table 1: Formulation and proximate composition of the experimental diets.

Ingredient (g kg⁻¹)	CTR	CH_{1.5}	CH_{3.0}	CH_{4.5}	CH_{6.0}
Fish meal	500	500	500	500	500
Corn meal	140	138.5	137	135.5	134
Soybean meal	210	210	210	210	210
Soybean oil	60	60	60	60	60
Fish oil	60	60	60	60	60
Vitamin premix [†]	20	20	20	20	20
Mineral premix [‡]	10	10	10	10	10
Chitosan	0.0	1.5	3.0	4.5	6.0

Proximate analysis (g kg⁻¹ as fed basis)					
Moisture	90	91	91	90	90
Crude protein	489	491	488	490	490
Crude lipid	136	135	135	136	137
Ash	70.1	70.5	70.7	70.4	70.5
Fiber	25.8	25.8	25.5	25.6	25.5
Gross energy (kJ g ⁻¹)	18.2	18.1	18.5	18.1	18.3

[†] Vitamin premix (as m kg⁻¹): Alpha tocopherol 60 IU; Calciferol 3000 IU; Thiamine 15; Riboflavin 30; Pyridoxine 15; B12 0.05; Nicotinic acid 175; Folic acid 5; Ascorbic acid 500; Inositol 1000; Biotin 2.5; Calcium pantothenate 50. [‡] Mineral premix (Unit kg⁻¹) (Science Laboratory): Cobalt sulphate 100 mg; Zinc sulphate 10 g; Iron sulphate 6 g; Choline chloride 6 g; Selenium 20 mg; Manganese sulphate 5 g; Copper sulphate 600 mg; Iodine 600 mg

Fish and rearing conditions

A 60-day feeding trial was conducted at the International Sturgeon Research Institute (Rasht, Gilan, Iran). Initial acclimatization for the fish lasted two weeks during which they were fed a basic diet. Following acclimatization, 180 starry sturgeon juveniles with a mean initial weight of 31.90±0.73 g were randomly distributed into fifteen fiberglass tanks (1000 L, 12 fish per tank) and fed the five formulated diets, with three replicates for each treatment. The water source consisted of a mixture of aerated groundwater and Sefidrud River water at a flow rate of 2 L min⁻¹. Water

quality parameters were monitored daily (temperature 17.51±0.2°C, dissolved oxygen 9.25±0.22 mg L⁻¹, and pH 7.50±0.10). Throughout the feeding trial, fish were fed manually four times daily (08:00, 12:00, 16:00, and 20:00) until visual satiation. Any uneaten food and waste were siphoned out before each feeding to maintain optimal water quality.

Sampling

At the end of the feeding trial, feeding was ceased for 24 h. Three fish from each replicate were then anesthetized using clove powder (300 mg L⁻¹). Blood samples

were collected from the caudal vessels using heparinized syringes for hematological tests. For serological tests, additional blood samples were collected without heparin, centrifuged (3000 rpm, 10 min, 4°C), and stored at -20°C until further analysis. For the gene expression study, spleen, head kidney, liver, and intestinal tissues were harvested from three anesthetized fish per tank, immediately frozen in liquid nitrogen, and subsequently stored at -80°C.

Hematological parameters measurements

The total counts of red blood cells (RBCs), white blood cells (WBCs), hematocrit (Hct), hemoglobin (Hb), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), and differential leukocyte cells (diff) were assessed following the method described by Feldman *et al.* (2000).

Blood serum immunological and biochemical parameters measurements

Serum lysozyme (LYZ) activity was measured using a turbidometric assay in accordance with the methodology outlined by Ellis (1990). The determination of total immunoglobulin (IgM) was executed using Siwicki and Anderson (1993) method. The total protein content was assessed both before and after IgM precipitation with a 12% polyethylene glycol solution (Sigma, USA). Respiratory burst activity was evaluated using the nitro blue tetrazolium (NBT) dye method based on the protocol established by Anderson and Zeeman (2020). The levels of total protein, albumin, triglycerides, cholesterol, glucose, calcium,

alkaline phosphatase (ALP), aspartate aminotransferase (AST), and alanine aminotransferase (ALT) were quantified using commercial kits (Pars Azmun, Karaj, Iran) in accordance with the manufacturer's instructions and analyzed using an automated analyzer (Fuji Photo Film Ltd., Tokyo, Japan). Additionally, the serum globulin concentration was calculated by subtracting albumin from the total protein content (Abdel-Wahab *et al.*, 2021).

The activity of serum antioxidant enzymes, including catalase (CAT), superoxide dismutase (SOD), and glutathione peroxidase (GPx), was evaluated spectrophotometrically using commercial kits (ZellBio GmbH, Lonsee, Germany).

Real-time quantitative PCR (RT-qPCR)

RNA extraction was conducted using an RNA extraction kit (QIAGEN, Hilden, Germany) following the manufacturer's protocol. The quantity and quality of RNA were assessed by measuring absorbance at 260/280 nm using a NanoDrop 1000 spectrophotometer (Thermo Scientific, Waltham, USA) and through electrophoresis on a 1% agarose gel. RNA samples were treated with a DNase I enzyme kit (Sinaclone, Tehran, Iran) in accordance with the manufacturer's instructions. cDNA synthesis was performed using a cDNA synthesis kit (Thermo Fisher Scientific, Massachusetts, USA). Primers for this study were designed based on sequences available for the sturgeon species in the NCBI database, with their specific sequences provided in Table 2. Beta-actin (β -actin) served as the reference gene for data normalization. Relative expression levels in tissue samples

were analyzed via RT-qPCR using the CFX-96 Real-Time PCR Thermocycler (Bio-Rad, Hercules, CA, USA) in a reaction volume of 20 μ L, which comprised 10 μ L of 2X SYBER Green (Ampliqon, Copenhagen, Denmark) and 100 nM of each primer. The amplification was carried out under the following conditions: initial

cooling of the products at 55°C for 90 s, followed by gradual heating from 55°C to 95°C at a rate of 0.5°C every 5 s. The $2^{-\Delta\Delta C_t}$ method for relative gene expression analysis, as described by Livak and Schmittgen (2001), was employed to calculate gene expression levels.

Table 2: Primer pair sequences of the genes used for quantitative real-time PCR.

Gene name	Gene abbreviation	Sequences of primers	Accession number
Interleukin-1 beta	<i>IL-1β</i>	F: TGGTGGAGAAGAGAACGCAAGATG R: GCGTCACTGCCTGTCTCATGG	MF818014.1
Tumor necrosis factor alpha	<i>TNF-α</i>	F: AAAATGTCCAGCCATCAAAGCA R: GTGAGGAATGATCCCGTTGGT	XM_034003960.3
Beta-actin	<i>β-actin</i>	F: TTGCCATCCAGGCTGTGCT R: TCTCGGCTGTGGTGAA	AY878120.1

Statistical analysis

All data are expressed as mean $\pm$ standard error (SE). SPSS software (Version 26) was utilized for all statistical analyses. The normality of the data distribution was verified using the Kolmogorov-Smirnov test, after which One-way ANOVA was performed to analyze the collected data. Significant differences between mean values were identified using Duncan's multiple range test at a p -value of 0.05.

Results

Hematological parameters

The results regarding hematological parameters are presented in Table 3. Significant increases in RBCs, Hb, and Hct were observed in the CH_{6.0} group compared to other groups ($p < 0.05$). While MCV increased in the CH-treated groups, no significant differences were noted among the CH_{3.0}, CH_{4.5}, and CH_{6.0} groups. MCH values in the treated groups did not show

significant differences when compared to the CTR group ($p > 0.05$). Additionally, there was no significant difference in MCHC among the groups ($p > 0.05$). WBC counts displayed significant differences between the groups ($p < 0.05$), with increases noted in the CH_{4.5} (6.03) and CH_{6.0} (6.11) groups. The diff results indicated that monocytes and eosinophils did not exhibit significant differences across the treated and CTR groups ($p > 0.05$). However, neutrophil levels rose in the CH-treated groups, and the CH_{6.0} group had the highest percentage of lymphocytes ($p < 0.05$).

Blood serum immunological and biochemical parameters

The results detailing blood serum immunological and biochemical parameters of juvenile starry sturgeon are shown in Table 4. The highest total protein levels were recorded in the CH_{6.0} group

(3.12 g dL⁻¹), while no significant differences were seen between the CTR, CH_{1.5}, and CH_{3.0} groups ($p>0.05$). In groups with higher CH levels (CH_{4.5} and CH_{6.0}), serum albumin levels increased, but no

significant differences were observed between the CTR group and the CH_{1.5} and CH_{3.0} groups ($p>0.05$). Globulin levels were highest in the CH_{6.0} group ($p<0.05$).

Table 3: Hematological parameters of juvenile starry sturgeon fed with experimental diets for 60 days.

Parameter	CH level (g kg ⁻¹)				
	CTR	CH _{1.5}	CH _{3.0}	CH _{4.5}	CH _{6.0}
RBC (×10 ⁶ mm ⁻³)	0.53 ± 0.01 ^a	0.53 ± 0.00 ^a	0.53 ± 0.00 ^a	0.54 ± 0.01 ^a	0.59 ± 0.01 ^b
Hb (g dL ⁻¹)	5.18 ± 0.10 ^a	5.12 ± 0.03 ^a	5.19 ± 0.14 ^a	5.14 ± 0.06 ^a	5.74 ± 0.09 ^b
Hct (%)	22.80 ± 0.20 ^a	22.60 ± 0.11 ^a	22.86 ± 0.49 ^a	23.20 ± 0.68 ^a	24.67 ± 0.32 ^b
MCV (fL)	415.00 ± 0.57 ^a	421.00 ± 0.57 ^b	422.66 ± 1.76 ^{bc}	424.00 ± 1.53 ^{bc}	426.33 ± 1.20 ^c
MCH (Pg)	97.13 ± 0.37	95.96 ± 0.28	95.73 ± 0.89	96.63 ± 0.26	95.93 ± 0.18
MCHC (g dL ⁻¹)	22.70 ± 0.25	22.70 ± 0.57	22.70 ± 0.20	23.17 ± 0.03	22.10 ± 0.15
WBC (×10 ³ mm ⁻³)	4.00 ± 0.05 ^a	4.43 ± 0.18 ^a	5.26 ± 0.39 ^b	6.03 ± 0.09 ^c	6.11 ± 0.05 ^c
Neutrophil (%)	9.33 ± 0.88 ^a	12.00 ± 1.52 ^{ab}	13.66 ± 0.33 ^b	13.33 ± 0.88 ^b	14.67 ± 0.33 ^b
Lymphocyte (%)	79.33 ± 0.33 ^a	79.66 ± 0.33 ^a	81.66 ± 1.33 ^a	80.67 ± 1.20 ^a	88.33 ± 0.67 ^b
Monocyte (%)	3.33 ± 0.33	5.00 ± 0.57	5.33 ± 0.66	3.67 ± 0.33	4.33 ± 0.33
Eosinophil (%)	0.66 ± 0.33	0.33 ± 0.33	0.00 ± 0.00	0.33 ± 0.33	0.00 ± 0.00

Mean±SE (n = 3); Different letters in horizontal rows indicate a significant difference between treatments ($p<0.05$). RBC: Red blood cells, Hb: Hemoglobin, Hct: Hematocrit, MCV: Mean corpuscular volume, MCH: Mean corpuscular hemoglobin, MCHC: Mean corpuscular hemoglobin concentration, WBC: White blood cells

Table 4: Serum immunological and biochemical parameters of juvenile starry sturgeon fed with experimental diets for 60 days.

Parameter	CH level (g kg ⁻¹)				
	CTR	CH _{1.5}	CH _{3.0}	CH _{4.5}	CH _{6.0}
Lysozyme Activity (u/ml/min)	27.87 ± 0.48 ^a	30.20 ± 0.25 ^b	32.17 ± 0.38 ^c	32.17 ± 0.38 ^c	33.70 ± 0.06 ^d
Total Immunoglobulin (mg mL ⁻¹)	13.80 ± 0.21 ^a	14.83 ± 0.03 ^{ab}	14.02 ± 0.51 ^{ab}	15.61 ± 0.24 ^{ab}	16.13 ± 0.02 ^b
NBT (mg mL ⁻¹)	1.15 ± 0.01	1.21 ± 0.04	1.28 ± 0.01	1.31 ± 0.02	1.16 ± 0.01
Total protein (g dL ⁻¹)	2.34 ± 0.01 ^a	2.48 ± 0.02 ^a	2.36 ± 0.11 ^a	2.66 ± 0.04 ^b	3.12 ± 0.04 ^c
Albumin (g dL ⁻¹)	0.97 ± 0.02 ^a	0.95 ± 0.01 ^a	0.97 ± 0.01 ^a	1.15 ± 0.02 ^b	1.19 ± 0.01 ^b
Globulin (g dL ⁻¹)	1.37 ± 0.01 ^a	1.52 ± 0.02 ^a	1.39 ± 0.10 ^a	1.50 ± 0.05 ^a	1.93 ± 0.03 ^b
Triglycerides (mg dL ⁻¹)	510.67 ± 10.81 ^a	452.00 ± 7.51 ^b	412.33 ± 8.37 ^c	332.33 ± 7.31 ^d	322.67 ± 15.50 ^d
Cholesterol (mg dL ⁻¹)	144.33 ± 3.71 ^a	123.00 ± 4.58 ^b	113.66 ± 1.45 ^b	114.66 ± 1.20 ^b	91.67 ± 2.40 ^c
ALT (U L ⁻¹)	46.17 ± 1.96 ^a	39.47 ± 1.10 ^b	38.60 ± 0.35 ^b	31.10 ± 0.56 ^c	31.13 ± 0.34 ^c
AST (U L ⁻¹)	410.67 ± 6.57 ^a	400.00 ± 3.79 ^{ab}	392.33 ± 4.67 ^b	351.33 ± 4.10 ^c	342.00 ± 3.51 ^c
ALP (U L ⁻¹)	309.00 ± 5.86 ^a	299.67 ± 2.40 ^a	260.67 ± 2.33 ^b	243.67 ± 4.63 ^c	233.33 ± 2.40 ^c
Glucose (mg dL ⁻¹)	48.73 ± 1.33 ^a	56.83 ± 0.83 ^b	55.50 ± 0.55 ^b	59.73 ± 0.26 ^c	61.80 ± 0.45 ^c
Calcium (mg dL ⁻¹)	9.54 ± 0.25 ^a	9.92 ± 0.09 ^a	9.85 ± 0.17 ^a	10.67 ± 0.04 ^b	10.81 ± 0.11 ^b

Mean±SE (n=3); Different letters in horizontal rows indicate a significant difference between treatments ($p<0.05$). ALT: Alanine aminotransferase, AST: Aspartate aminotransferase, ALP: Alkaline phosphatase

Serum LYZ activity showed significant differences across all groups ($p < 0.05$), with the highest values in the CH_{6.0} group (33.70) and the lowest in the CTR group (27.86). IgM levels significantly differed between the CTR group and CH_{6.0} ($p < 0.05$). No significant differences in the NBT parameter were detected between treated groups and the CTR group ($p > 0.05$).

Triglyceride levels decreased with increasing CH doses in the diet, with the highest triglyceride level in the CTR group at 510.67 mg dL⁻¹, demonstrating significant differences in the treated groups ($p < 0.05$). The lowest triglyceride levels were noted in the CH_{4.5} and CH_{6.0} groups ($p > 0.05$). Significant differences in cholesterol levels were also observed ($p < 0.05$), with the highest level in the CTR group (144.33 mg dL⁻¹) and the lowest in the CH_{6.0} group (91.67 mg dL⁻¹). The lowest serum glucose level was recorded in the CTR group (48.73 mg dL⁻¹), with higher levels noted in the CH_{4.5} and CH_{6.0} groups. Calcium levels were highest in the CH_{4.5} and CH_{6.0} groups, significantly differing from the other groups ($p < 0.05$).

Serum liver enzymes

Table 4 illustrates that the levels of liver enzymes ALT, AST, and ALP decreased with increasing CH levels in the diet. The highest ALT level was observed in the CTR group, showing significant differences from the treated groups ($p < 0.05$). The lowest AST levels were recorded in the CH_{4.5} and CH_{6.0} groups, while the highest AST levels were found in the CTR group. Both the CH_{4.5} and CH_{6.0} groups displayed the lowest ALP levels, whereas the highest ALP levels were seen in the CTR and CH_{1.5} groups.

Antioxidant enzyme activity

Table 5 presents the results of antioxidant enzyme activity. Dietary CH increased CAT activity, with the highest CAT activity found in the CH_{6.0} group and the lowest in the CTR group. SOD activity was not significantly different in the treated groups compared to the CTR group. GPx activity also increased in the CH_{6.0} group relative to the other groups.

Table 5: Antioxidant activity of juvenile starry sturgeon fed with experimental diets for 60 days.

CH level (g kg ⁻¹)	CAT (u mL ⁻¹)	SOD (u mL ⁻¹)	GPx (u mL ⁻¹)
CTR	111.00 ± 1.53 ^a	52.03 ± 0.95 ^a	204.00 ± 2.08 ^a
CH _{1.5}	125.33 ± 0.88 ^b	54.17 ± 0.91 ^a	207.67 ± 2.40 ^a
CH _{3.0}	127.00 ± 1.53 ^{bc}	54.53 ± 0.75 ^a	254.67 ± 1.76 ^b
CH _{4.5}	130.33 ± 0.88 ^c	52.43 ± 0.91 ^a	253.67 ± 2.03 ^b
CH _{6.0}	136.33 ± 0.88 ^d	58.32 ± 0.62 ^c	269.00 ± 2.31 ^c

Mean ± SE (n = 3); Different letters in vertical rows indicate a significant difference between treatments ($p < 0.05$). CAT: Catalase, SOD: Superoxide dismutase, GPx: Glutathione peroxidase.

Immune-related gene expression (*IL-1β* and *TNF-α*)

The *IL-1β* gene expression results are illustrated in Figure 1. The highest levels of

IL-1β gene expression in the liver and intestine were observed in the CH_{6.0} group, demonstrating significant differences from the other groups ($p < 0.05$).





Figure 1: Relative mRNA expression of *IL-1 β* gene in the liver (a), intestine (b) spleen (c) and head kidney (d) of juvenile starry sturgeon fed with experimental diets for 60 days. Mean $\pm$ SE (n=3 repeat); Different letters indicate a significant difference between treatments ($p < 0.05$).

The lowest expression was found in the CTR group for the spleen, while the CH_{6.0} group had the highest. In the head kidney, *IL-1 β* expression levels were highest in the CH_{6.0} group, with no significant difference between the CTR and CH_{1.5} groups ($p>0.05$).

Figure 2 presents the *TNF- α* gene

expression results. Statistical analysis revealed that *TNF- α* gene expression in the liver and spleen increased with higher CH doses, peaking in the CH_{6.0} group. In the intestine, the highest *TNF- α* expression was also seen in the CH_{6.0} group. The head kidney exhibited the highest *TNF- α* expression in the CH_{6.0} group and the lowest in the CTR group.






Figure 2: Relative mRNA expression of *TNF-α* gene in the liver (a), spleen (b), intestine (c) and head kidney (d) of juvenile starry sturgeon fed with experimental diets for 60 days. Mean±SE (n=3 repeat); Different letters indicate a significant difference between treatments ($p<0.05$).

Discussion

Strengthening the immune system is one of the most fundamental challenges in sturgeon farming, particularly during the early stages of life. Researchers are actively seeking compounds that can stimulate and modulate the immune systems of cultured animals. The present study investigates the impact of CH on the function of the immune system in juvenile starry sturgeon at a molecular level. The findings indicate that CH, functioning as an immunostimulant, enhances immune response and increases the expression of immune-related genes in juvenile starry sturgeon.

Hematological indices are frequently employed as biological indicators to monitor the overall health of fish (Fazio, 2019). The efficacy of dietary supplements for fish can also be evaluated through these indices. Variations in these values may arise from changes in the health status and nutritional behavior of the fish (Dawood *et al.*, 2020). In our study, we observed significant positive changes in hematological indices in response to the CH diet. Specifically, the RBC count, Hb, and

Hct levels increased in the CH_{6.0} group. Additionally, while MCV levels rose in the treated groups, dietary CH did not significantly affect MCH or MCHC. The increase in MCV values likely indicates larger RBCs with a greater capacity for gas exchange (Snyder and Sheafar, 1999). In vertebrates, MCV and MCH values serve as indicators for assessing anemia and iron deficiency. WBCs play a crucial role in the immune system of fish, contributing to both the cellular component of innate immunity and the release of humoral substances such as cationic antimicrobial peptides, components of the complement system, lectins, and cytokines. In our study, the highest WBC counts were observed in CH_{4.5} and CH_{6.0} groups.

Elevated levels of hemoglobin and RBCs suggest an increased metabolic rate for nutrient processing in the fish body, while higher Hct and WBC counts are typically associated with improved immune function (Fazio, 2019). The increase in WBCs, particularly lymphocytes, in our study can be attributed to the positive immunostimulant effect of CH on the health of starry sturgeon, potentially

bolstering their immunity and preventing diseases. Previous studies by Abd El-Naby *et al.* (2020) and Younus *et al.* (2020) demonstrated similar increases in WBCs, RBCs, Hb, and Hct in *Oreochromis niloticus* and *Hypophthalmichthys molitrix* that were fed chitosan nanoparticles (ChNPs), attributing these effects to the immunostimulant potential of ChNPs. Kamali Najafabad *et al.* (2016) reported that WBCs increased in *Rutilus frisii* fed diets containing 1 and 2 g kg⁻¹ CH.

Biochemical serum parameters serve as biomarkers for evaluating the health status of aquatic animals. Innate immunity is also linked to total protein, albumin, and globulin levels. Serum proteins, including albumin, are vital for maintaining osmotic blood pressure (Nya and Austin, 2009), while serum globulins, such as gamma globulins, serve as sources of immunoglobulins reflecting antibody concentration and immune status in fish (Goda, 2008). In our study, total protein and albumin levels increased in the CH_{4.5} and CH_{6.0} groups, with globulin also peaking in the CH_{6.0} group, indicating enhanced immunity in starry sturgeon. Studies involving *Epinephelus bruneus* fed with 1% and 2% CH have shown similar increases in total protein, albumin, and globulin levels in the treated groups, which aligns with our findings (Harikrishnan *et al.*, 2012). Research by Dawood *et al.* (2020) on *Liza ramada* also showed that total protein, albumin, and globulin levels rose in groups receiving 1.0 and 2.0 g kg⁻¹ of ChNPs compared to controls and lower doses.

LYZ activity directly indicates aquatic animals' health and safety status (Wu,

2020). In our study, LYZ activity increased with the dietary CH dosage. IgM is a critical immune protein that plays roles in bactericidal activity, immune regulation, and agglutination (Krogdahl *et al.*, 2000). The CH_{6.0} group exhibited significant differences in IgM levels compared to the control group, corroborating findings by Chen and Chen (2019), who reported increased LYZ and IgM in *Misgurnus anguillicadatus* when fed CH at doses of 1, 5, and 10 g kg⁻¹. Similarly, a study by Oushani *et al.* (2020) observed increases in LYZ, IgM, and total protein levels in *Oncorhynchus mykiss* fed with combinations of clinoptilolite and CH or ChNP-containing strains. Since monocytes and neutrophils are the primary sources of serum LYZ, the observed increases in serum LYZ in the CH-treated groups may be attributed to a higher percentage of neutrophils in the blood.

CH serves as a prebiotic that enhances the intestinal microbiome. Additionally, its role in immune responses may be linked to alterations in the intestinal microbiome and an increase in probiotics, which act as immunostimulants (Song *et al.*, 2014). Prebiotics can traverse the intestinal epithelium to stimulate leukocytes (Hoseinifar *et al.*, 2015). As a prebiotic, CH likely stimulates leukocytes, thereby contributing to enhanced immunity. Furthermore, it plays a significant role in the non-specific defense system of fish and other vertebrates, particularly within gut-associated lymphoid tissue (GALT), which is crucial for regulating the immune response to dietary supplements. The microbial flora of the digestive system may also influence this regulatory process

(Lazado and Caipang, 2014). The observed increase in IgM and total protein following CH administration may stem from changes in the gastrointestinal microbial flora, leading to enhancements in the non-specific immune system mediated by the GALT (Hoseinifar *et al.*, 2015). While the specific mechanisms by which CH stimulates immunity in fish are not yet fully understood, these findings suggest its potential as a modulator of the immune system, warranting further investigation.

CH has been noted to reduce cholesterol and triglyceride levels in species such as *Rutilus caspicus* (Zahmatkesh *et al.*, 2020) and *Cyprinus carpio* (Stanek *et al.*, 2023), corroborating our findings. This reduction in serum cholesterol and triglycerides in our study can be attributed to the hypolipidemic effects of CH. CH interacts with cholesterol and other sterols through hydrophobic interactions, forming hydrophobic aggregates in the small intestine. At a pH level of 7-8, these CH-lipid complexes solidify and are subsequently excreted in feces (Wydro *et al.*, 2007).

The dietary supplementation of CH resulted in increased calcium levels in the CH_{4.5} and CH_{6.0} groups. Several factors influence mineral absorption via CH, including environmental pH, intestinal morphology, the water solubility of CH, molecular weight, and the degree of polymerization (Schmitz *et al.*, 2019). Research by Zhao *et al.* (2022) confirmed that CH can inhibit the digestion of peptide-calcium chelate in the stomach, allowing more calcium to enter the intestine and enhancing calcium absorption. Additionally, elevated calcium levels may be associated with increased albumin

production, as calcium binds to albumin in the bloodstream (Stanek *et al.*, 2023).

In this study, we examined liver enzymes AST, ALT, and ALP, which are transaminase enzymes predominantly found in the mitochondria of liver cells. Any degeneration, inflammation, or necrosis of liver cells can lead to the release of these enzymes, resulting in elevated serum levels. ALP, ALT, and AST levels are generally considered indicators of liver damage influenced by nutritional status, stress, pollutants, or pathogenic infections (Taghavizadeh *et al.*, 2020).

Our findings indicated that serum levels of liver enzymes decreased with higher dietary CH levels, suggesting that CH does not adversely affect liver tissue. This protective effect may be due to CH's role in stabilizing cellular absorption and preventing the release of enzymes from liver cells into the bloodstream (Ramasamy *et al.*, 2014). Moreover, the liver-protective properties of CH might be linked to its stimulatory effects on the immune system in fish, given the liver's essential role in producing acute phase proteins and managing the destruction of pathogens, antigens, and inflammation-related molecules (Parker and Picut, 2012). The results of studies by Wu (2020) showed that diets containing 4, 6, and 8 g kg⁻¹ CH decreased the liver enzymes of *O. niloticus*. In the study by Abd El-Naby *et al.* (2020), ALT and AST activities in *O. niloticus* were reduced by the addition of ChNPs in the diet.

The current study demonstrated that CH increased the activities of antioxidant enzymes CAT, SOD, and GPx in starry sturgeon, with the highest levels observed

in the CH_{6.0} group. This suggests that CH can enhance the antioxidant capacity of sturgeon. Previous studies have also shown that CH increases antioxidant activity in various fish species, including *Trachinotus ovatus* (Yu *et al.*, 2022), *Dicentrarchus labrax* (Abdel-Tawwab *et al.*, 2021), *M. anguilladatus* (Chen and Chen, 2019), and hybrid sturgeon (*A. baerii*♀ × *A. schrenckii*♂) (Xu *et al.*, 2023). The antioxidant properties of CH can be attributed to its ability to inhibit and chelate free radicals, such as hydroxyl radicals, by donating a hydrogen atom or an electron pair (Ngo and Kim, 2014). The structural basis for its antioxidant function lies in its amino groups and primary and secondary hydroxyl groups (Feng *et al.*, 2007).

Studying immune-related gene expression can provide insights into preventing, controlling, and treating diseases, particularly in protected species (Tian *et al.*, 2021). *IL-1β*, a pro-inflammatory cytokine, plays a crucial role in responding to microbial attacks and tissue damage, as it enhances phagocyte activity, macrophage proliferation, and leukocyte migration while triggering the secretion of various cytokines, including *TNF-α* from immune cells (Giri *et al.*, 2015). *TNF-α* is vital for immunity, secreted by leukocytes in response to immune system stimulation (Grayfer *et al.*, 2011).

In the current study, we evaluated the expression of *IL-1β* and *TNF-α* genes in the liver, intestine, spleen, and head kidney tissues, revealing that chitosan (CH) significantly increased the relative expression of these genes. Notably, the highest levels of *IL-1β* and *TNF-α* gene

expression were observed in the CH_{6.0} group. These findings align with previous research, including the study by Nikapitiya *et al.* (2018), which reported that dietary chitosan nanoparticles (ChNPs) enhanced *IL-1β* gene expression in zebrafish larvae. Similarly, Zou *et al.* (2016) found that feeding *C. carpio* with the prebiotic *Agaricus bisporus* led to an increased expression of *IL-1β* and *TNF-α* genes. Additionally, Oushani *et al.* (2020) reported that a combination of CH and clinoptilolite, as well as ChNP and clinoptilolite, could elevate *IL-1β* gene expression in the head kidney of *O. mykiss*. Moreover, Panase *et al.* (2023) demonstrated that the inclusion of *Bacillus subtilis* and fructooligosaccharides stimulated *IL-1β* and *TNF-α* gene expression in *O. niloticus*.

Dietary supplements, such as probiotics and prebiotics, can directly activate primary defense mechanisms by influencing receptors and associated genes, thereby enhancing immunity and resistance to pathogens (Bricknell *et al.*, 2006). CH serves as an immune system stimulant, likely affecting these receptors and genes, and activating primary defense responses. As previously mentioned, the increase in *IL-1β* and *TNF-α* gene expression in response to CH is likely linked to alterations in the gastrointestinal bacterial flora, which impacts the gut-associated lymphoid tissue (GALT) system and strengthens the primary defense system. Overall, assessing the expression levels of cytokine system genes can serve as an indicator of immune system status (2007).

Further research is warranted to elucidate

the mechanisms underlying CH stimulation.

The results indicate that CH can bolster the innate immunity of starry sturgeon. Variations in hematological and safety parameters across different studies are influenced by factors such as species, age, nutrition, and sex of the fish (Ahmed *et al.*, 2019).

Conclusion

Our study demonstrates that CH, as an immunostimulant, effectively stimulates and modulates the immunity of juvenile starry sturgeon. Based on these findings, the optimal dose of CH is determined to be 6 g kg⁻¹, which enhances immune response, elevates the expression of immune-related genes, reduces serum lipid levels, improves liver health, and boosts antioxidant capacity. CH presents a viable alternative to antibiotics in the rearing of starry sturgeon.

Acknowledgments

The authors gratefully acknowledge the financial support and technical assistance provided by the International Sturgeon Research Institute, Agricultural Research, Education and Extension Organization.

Conflicts of interest

The authors declare no conflict of interest.

Reference

Abd El-Naby, A.S., Al-Sagheer, A.A., Negm, S.S. and Naiel, M.A., 2020. Dietary combination of chitosan nanoparticle and thymol affects feed utilization, digestive enzymes, antioxidant status, and intestinal morphology of *Oreochromis*

niloticus. *Aquaculture*, 515, 734577. DOI:10.1016/j.aquaculture.2019.734577

Abdel-Tawwab, M., Khalil, R.H., Diab, A.M., Khallaf, M.A., Abdel-Razek, N., Abdel-Latif, H.M. and Khalifa, E., 2021. Dietary garlic and chitosan enhanced the antioxidant capacity, immunity, and modulated the transcription of HSP70 and Cytokine genes in Zearalenone-intoxicated European seabass. *Fish & Shellfish Immunology*, 113, 35-41. DOI:10.1016/j.fsi.2021.03.012

Abdel-Wahab, M.M., Taha, N.M., Lebda, M.A., Elfeky, M.S. and Abdel-Latif, H.M., 2021. Effects of bovine lactoferrin and chitosan nanoparticles on serum biochemical indices, antioxidative enzymes, transcriptomic responses, and resistance of Nile tilapia against *Aeromonas hydrophila*. *Fish & Shellfish Immunology*, 111, 160-169. DOI:10.1016/j.fsi.2021.01.017

Ahmed, F., Soliman, F.M., Adly, M.A., Soliman, H.A., El-Matbouli, M. and Saleh, M., 2019. Recent progress in biomedical applications of chitosan and its nanocomposites in aquaculture: A review. *Research in Veterinary Science*, 126, 68-82. DOI:10.1016/j.rvsc.2019.08.005

Anderson, D.P., 1992. Immunostimulants, adjuvants, and vaccine carriers in fish: applications to aquaculture. *Annual Review of Fish Diseases*, 2, 281-307. DOI:10.1016/0959-8030(92)90067-8.

Anderson, D.P. and Zeeman, M.G., 2020. Immunotoxicology in fish. In: Rand, G.M. (ed) *Fundamentals of Aquatic*

- Toxicology. CRC Press, Boca Raton. pp 371-404.
- Bricknell, I.R., Bron, J.E. and Bowden, T.J., 2006.** Diseases of gadoid fish in cultivation: a review. *ICES Journal of Marine Science*, 63(2), 253-266. DOI:10.1016/j.icesjms.2005.10.009
- Chang, S.H., Wu, C.H. and Tsai, G.J., 2018.** Effects of chitosan molecular weight on its antioxidant and antimutagenic properties. *Carbohydrate Polymers*, 181, 1026-1032. DOI:10.1016/j.carbpol.2017.11.047
- Chen, J. and Chen, L., 2019.** Effects of chitosan-supplemented diets on the growth performance, nonspecific immunity and health of loach fish (*Misgurnus anguillicadatus*). *Carbohydrate Polymers*, 225, 115227. DOI:10.1016/j.carbpol.2019.115227
- Cvijanović, G., Đikanović, V., Galambos, L., Pengal, P. and Smederevac-Lalić, M., 2024.** Past and future of sturgeon species (Acipenseridae) in western balkans: Case for permanent conservation or sustainable management. In *Ecological Sustainability of Fish Resources of Inland Waters of The Western Balkans: Freshwater Fish Stocks, Sustainable Use and Conservation*, 411-452. Cham: Springer International Publishing.
- Dawood, M.A., Gewaily, M.S., Soliman, A.A., Shukry, M., Amer, A.A., Younis, E.M., Abdel-Warith, A.W.A., Van Doan, H., Saad, A.H., Aboubakr, M. and Abdel-Latif, H.M., 2020.** Marine-derived chitosan nanoparticles improved the intestinal histomorphometrical features in association with the health and immune response of grey mullet (*Liza ramada*). *Marine Drugs*, 18(12), 611. DOI:10.3390/md18120611
- Ellis, A.E., 1990.** Serum antiproteases in fish. *Techniques in fish immunology*, 95-99.
- Enomoto, A., Itoh, K., Nagayoshi, E., Haruta, J., Kimura, T., O'Connor, T., Harada, T. and Yamamoto, M., 2001.** High sensitivity of Nrf2 knockout mice to acetaminophen hepatotoxicity associated with decreased expression of ARE-regulated drug metabolizing enzymes and antioxidant genes. *Toxicological Sciences*, 59(1), 169-177. DOI:10.1093/toxsci/59.1.169
- Fadl, S.E., El-Gammal, G.A., Abdo, W.S., Barakat, M., Sakr, O.A., Nassef, E., Gad, D.M. and El-Sheshtawy, H.S., 2020.** Evaluation of dietary chitosan effects on growth performance, immunity, body composition and histopathology of Nile tilapia (*Oreochromis niloticus*) as well as the resistance to *Streptococcus agalactiae* infection. *Aquaculture Research*, 51(3), 1120-1132. DOI:10.1111/are.14458
- Fazio, F., 2019.** Fish hematology analysis as an important tool of aquaculture: A review. *Aquaculture*, 500, 237-242. DOI:10.1016/j.aquaculture.2018.10.030
- Feldman, B.F. and Zinkl, J.G., 2000.** e Jain, NC Schalm's Veterinary Hematology.
- Feng, T., Du, Y., Li, J., Wei, Y. and Yao, P., 2007.** Antioxidant activity of half N-acetylated water-soluble chitosan *in vitro*. *European Food Research and Technology*, 225(1), 133-138. DOI:10.1007/s00217-006-0391-0

- Fernandez-Kim, S.O., 2004.** Physicochemical and functional properties of crawfish chitosan as affected by different processing protocols. PhD Thesis. Louisiana State University and Agricultural & Mechanical College. USA.
- Giri, S.S., Sen, S.S., Chi, C., Kim, H.J., Yun, S., Park, S.C. and Sukumaran, V., 2015.** Effect of cellular products of potential probiotic bacteria on the immune response of *Labeo rohita* and susceptibility to *Aeromonas hydrophila* infection. *Fish & shellfish immunology*, 46(2), 716-722. DOI:10.1016/j.fsi.2015.08.012
- Goda, A.M.S., 2008.** Effect of dietary Ginseng herb (Ginsana® G115) supplementation on growth, feed utilization, and hematological indices of Nile Tilapia, *Oreochromis niloticus* (L.), fingerlings. *Journal of the World Aquaculture Society*, 39(2), 205-214. DOI:10.1111/j.1749-7345.2008.00153.x
- Grayfer, L., Hodgkinson, J.W., Hitchen, S.J. and Belosevic, M., 2011.** Characterization and functional analysis of goldfish (*Carassius auratus* L.) interleukin-10. *Molecular immunology*, 48(4), 563-571. DOI:10.1016/j.molimm.2010.10.013
- Harikrishnan, R., Kim, J.S., Balasundaram, C. and Heo, M.S., 2012.** Immunomodulatory effects of chitin and chitosan enriched diets in *Epinephelus bruneus* against *Vibrio alginolyticus* infection. *Aquaculture*, 326, 46-52. DOI:10.1016/j.aquaculture.2011.11.034
- Hoseinifar, S.H., Esteban, M.Á., Cuesta, A. and Sun, Y.Z., 2015.** Prebiotics and fish immune response: a review of current knowledge and future perspectives. *Reviews in Fisheries Science & Aquaculture*, 23(4), 315-328. DOI:10.1080/23308249.2015.1052365
- Kamali Najafabad, M., Imanpoor, M.R., Taghizadeh, V. and Alishahi, A., 2016.** Effect of dietary chitosan on growth performance, hematological parameters, intestinal histology and stress resistance of Caspian kutum (*Rutilus frisii kutum* Kamenskii, 1901) fingerlings. *Fish Physiology and Biochemistry*, 42(4), 1063-1071. DOI:10.1007/s10695-016-0197-3
- Kong, M., Chen, X.G., Xing, K. and Park, H.J., 2010.** Antimicrobial properties of chitosan and mode of action: a state of the art review. *International Journal of Food Microbiology*, 144(1), 51-63. DOI:10.1016/j.ijfoodmicro.2010.09.012
- Krogdahl, RØed and Baeverfjord, 2000.** Feeding Atlantic salmon *Salmo salar* L. soybean products: Effects on disease resistance (furunculosis), and lysozyme and IgM levels in the intestinal mucosa. *Aquaculture Nutrition*, 6(2), 77-84. DOI:10.1046/j.1365-2095.2000.00129.x
- Lazado, C.C. and Caipang, C.M.A., 2014.** Mucosal immunity and probiotics in fish. *Fish & Shellfish Immunology*, 39(1), 78-89. DOI:10.1016/j.fsi.2014.04.015
- Livak, K.J. and Schmittgen, T.D., 2001.** Analysis of relative gene expression data using real-time quantitative PCR and the

- $2^{-\Delta\Delta CT}$ method. *Methods*, 25(4), 402-408. DOI:10.1006/meth.2001.1262
- Ngo, D.H. and Kim, S.K., 2014.** Antioxidant effects of chitin, chitosan, and their derivatives. *Advances in Food and Nutrition Research*, 73, 15-31. DOI:10.1016/B978-0-12-800268-1.00002-0
- Nikapitiya, C., Dananjaya, S.H.S., De Silva, B.C.J., Heo, G.J., Oh, C., De Zoysa, M. and Lee, J., 2018.** Chitosan nanoparticles: A positive immune response modulator as display in zebrafish larvae against *Aeromonas hydrophila* infection. *Fish & Shellfish Immunology*, 76, 240-246. DOI:10.1016/j.fsi.2018.03.010
- Nya, E.J. and Austin, B., 2009.** Use of garlic, *Allium sativum*, to control *Aeromonas hydrophila* infection in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases*, 32(11), 963-970. DOI:10.1111/j.1365-2761.2009.01100.x
- Oushani, A.K., Soltani, M., Sheikhzadeh, N., Mehrgan, M.S. and Islami, H.R., 2020.** Effects of dietary chitosan and nano-chitosan loaded clinoptilolite on growth and immune responses of rainbow trout (*Oncorhynchus mykiss*). *Fish & Shellfish Immunology*, 98, 210-217. DOI:10.1016/j.fsi.2020.01.016
- Panase, A., Thirabunyanon, M., Promya, J. and Chitmanat, C., 2023.** Influences of *Bacillus subtilis* and fructooligosaccharide on growth performances, immune responses, and disease resistance of Nile tilapia, *Oreochromis niloticus*. *Frontiers in Veterinary Science*, 9, 1094681. DOI:10.3389/fvets.2022.1094681
- Parker, G.A. and Picut, C.A., 2012.** Immune functioning in non lymphoid organs: the liver. *Toxicologic Pathology*, 40(2), 237-247. DOI:10.1177/0192623311428475
- Perini, M., Nardin, T., Venturelli, M., Pianezze, S. and Larcher, R., 2020.** Stable isotope ratio analysis as a fast and simple method for identifying the origin of chitosan. *Food Hydrocolloids*, 101, 105516. DOI:10.1016/j.foodhyd.2019.105516
- Qin, C., Du, Y., Xiao, L., Li, Z. and Gao, X., 2002.** Enzymic preparation of water-soluble chitosan and their antitumor activity. *International Journal of Biological Macromolecules*, 31(1-3), 111-117. DOI:10.1016/s0141-8130(02)00064-8
- Ramasamy, P., Subhapradha, N., Shanmugam, V. and Shanmugam, A., 2014.** Protective effect of chitosan from *Sepia kobeensis* (Hoyle 1885) cuttlebone against CCl₄ induced hepatic injury. *International Journal of Biological Macromolecules*, 65, 559-563. DOI:10.1016/j.ijbiomac.2014.02.009
- Schmitz, C., González Auza, L., Koberidze, D., Rasche, S., Fischer, R. and Bortesi, L., 2019.** Conversion of chitin to defined chitosan oligomers: Current status and future prospects. *Marine Drugs*, 17(8), 452. DOI:10.3390/md17080452
- Siwicki, A.K. and Anderson, D.P., 1993.** Nonspecific defense mechanisms assay in fish. II. Potential killing activity of neutrophils and macrophages, lysozyme

- activity in serum and organs and total immunoglobulin level in serum. In: Siwicki, A.K., Anderson, D.P. and Waluga, J. (eds) Fish disease diagnosis and prevention methods., Olsztyn, 1993, pp. 105–112. Poland
- Snyder, G.K. and Sheafor, B.A., 1999.** Red blood cells: centerpiece in the evolution of the vertebrate circulatory system. *American Zoologist*, 39(2), 189–198. DOI:10.1093/icb/39.2.189
- Song, S.K., Beck, B.R., Kim, D., Park, J., Kim, J., Kim, H.D. and Ringø, E., 2014.** Prebiotics as immunostimulants in aquaculture: a review. *Fish & Shellfish Immunology*, 40(1), 40–48. DOI:10.1016/j.fsi.2014.06.016
- Stanek, M., Mazurkiewicz, J., Rawski, M., Bogucka, J., Ziółkowska, E., Dankowiakowska, A. and Kierończyk, B., 2023.** Effect of chitosan on common carp (*Cyprinus carpio*) fry growth performance, feed utilization and nutriphysiological status. *Aquaculture Reports*, 30, 101622. DOI:10.1016/j.aqrep.2023.101622
- Taghavizadeh, M., Shekarabi, S.P.H., Mehrgan, M.S. and Islami, H.R., 2020.** Efficacy of dietary lysophospholipids (Lipidol™) on growth performance, serum immuno-biochemical parameters, and the expression of immune and antioxidant-related genes in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 525, 735315. DOI:10.1016/j.aquaculture.2020.735315
- Tian, G., Xie, M., Wang, B., Gao, W., Yuan, H., Wei, H., Zhang, Y. and Xu, Q., 2021.** Identification and expression of IL-1 β in the endangered Dabry's sturgeon (*Acipenser dabryanus*). *Aquaculture Reports*, 20, 100698. DOI:10.1016/j.aqrep.2021.100698
- Udo, I.U., Etukudo, U. and Anwana, U.I.U., 2018.** Effects of chitosan and chitosan nanoparticles on water quality, growth performance, survival rate and meat quality of the African catfish, *Clarias gariepinus*. *Nanoscience*, 1(1), 12–25. DOI:10.31058/j.nano.2018.11002
- Wu, S., 2020.** The growth performance, body composition and nonspecific immunity of Tilapia (*Oreochromis niloticus*) affected by chitosan. *International Journal of Biological Macromolecules*, 145, 682–685. DOI:10.1016/j.ijbiomac.2019.12.235
- Wydro, P., Krajewska, B. and Haczydro, K., 2007.** Chitosan as a lipid binder: A Langmuir monolayer study of chitosan–lipid interactions. *Biomacromolecules*, 8(8), 2611–2617. DOI:10.1021/bm700453x
- Xu, H., Wang, X., Liang, Q., Xu, R., Liu, J. and Yu, D., 2023.** Dietary chitosan moderates the growth rate, antioxidant activity, immunity, intestinal morphology and resistance against *Aeromonas hydrophila* of juvenile hybrid sturgeon (*Acipenser baerii*♀ $\times$ *Acipenser schrenckii*♂). *International Journal of Biological Macromolecules*, 224, 1012–1024. DOI:10.1016/j.ijbiomac.2022.10.185
- Yan, J., Guo, C., Dawood, M.A.O. and Gao, J., 2017.** Effects of dietary

- chitosan on growth, lipid metabolism, immune response and antioxidant-related gene expression in *Misgurnus anguillicaudatus*. *Beneficial Microbes*, 8(3), 439-450. DOI:10.3920/BM2016.0177
- Younus, N., Zuberi, A., Mahmood, T., Akram, W. and Ahmad, M., 2020.** Comparative effects of dietary micro- and nano-scale chitosan on the growth performance, non-specific immunity, and resistance of silver carp *Hypophthalmichthys molitrix* against *Staphylococcus aureus* infection. *Aquaculture International*, 28(6), 2363-2378. DOI:10.1007/s10499-020-00595-0
- Yu, W., Yang, Y., Zhou, Q., Huang, X., Huang, Z., Li, T., Wu, Q., Zhou, C., Ma, Z. and Lin, H., 2022.** Effects of dietary Astragalus polysaccharides on growth, health and resistance to *Vibrio harveyi* of *Lates calcarifer*. *International Journal of Biological Macromolecules*, 207, 850-858. DOI:10.1016/j.ijbiomac.2022.03.176
- Zahmatkesh, A., Karimzadeh, K. and Faridnia, M., 2020.** Effect of dietary selenium nanoparticles and chitosan oligosaccharide on biochemical parameters of Caspian roach (*Rutilus caspicus*) under malathion stress. *Caspian Journal of Environmental Sciences*, 18(1), 59-71. DOI:10.22124/cjes.2020.3979
- Zhang, J., Zhang, W., Mamadoub, B. and Xia, W., 2012.** A comparative study on hypolipidemic activities of high and low molecular weight chitosan in rats. *International Journal of Biological Macromolecules*, 51(4), 504-508. DOI:10.1016/j.ijbiomac.2012.06.018
- Zhao, M., He, H., Guo, D., Zhang, X., Jia, L., Hou, T. and Ma, A., 2022.** Chitosan oligosaccharides-tripolyphosphate microcapsules as efficient vehicles for desalted duck egg white peptides-calcium: Fabrication, entrapment mechanism and *in vivo* calcium absorption studies. *LWT*, 154, 112869. DOI:10.1016/j.lwt.2021.112869
- Zou, H.K., Hoseinifar, S.H., Miandare, H.K. and Hajimoradloo, A., 2016.** Agaricus bisporus powder improved cutaneous mucosal and serum immune parameters and up-regulated intestinal cytokines gene expression in common carp (*Cyprinus carpio*) fingerlings. *Fish & Shellfish Immunology*, 58, 380-386. DOI:10.1016/j.fsi.2016.09.050