# Seasonal change of thyroid histomorphological structure and hormone production in yellowfin seabream (*Acanthopagrus latus*) in the Persian Gulf

Salamat N.\*1, Havasi M.1, Earfani Majd N.2, Savari A.1

Received: May 2012 Accepted: September 2012

## **Abstract**

Seasonal changes of the thyroid gland structure and hormones secretion was examined in yellowfin seabream (*Acanthopagrus latus*) in the northwest of Persian Gulf (Musa creek). Thyroid gland composed of follicles scattered around the ventral aorta, near the gills. Follicular cells varied according to secretion of the gland during warm and cold seasons. Thyroid hormones (Triidothyronine [T3] and Thyroxine [T4]) were detected in the fish serum in levels ranged from 4.09-1.30 ng/mL for T3 and from 1.10-0.21 ng/mL for (T4) in the warm and cold seasons, respectively. The results showed that the height of thyroid epithelium and plasma concentration of thyroid hormones (thyroid activity) in *A. latus* increased significantly during spring and summer. The peak of these factors occurred in midsummer (August). Then, the thyroid activity decreased significantly during autumn and early winter from October to December according to decrease of temperature. T3 and T4 increased significantly from January to April.

**Keywords**: Yellowfin seabream, *Acanthopagrus latus*, Thyroid gland, Triidothyronine, Thyroxine, Histology, Plasma

<sup>1-</sup> Department of Marine Biology, School of Marine Science, Khorramshahr Marine Science and Technology University, P.O. Box: 669, Khorramshahr, Iran.

<sup>2-</sup> Department of Histology, Faculty of Veterinary Medicine, Shahid Chamran University of Ahvaz, Iran.

<sup>\*</sup>Corresponding author's email: salamatnegin@yahoo.com

# Introduction

Thyroid hormones (THs) include triiodothyronine (T3) and thyroxine (T4), regulate growth, development, differentiation. metabolism. maintenance of homeostasis in vertebrates (Szischa et al., 2005). In all vertebrates embryogenesis, organogenesis and growth acutely depend on thyroid hormones (Power et al., 2001). Although there is an extensive diversity in teleosts. developmental stages in most of them include of larva, juvenile, and adult, which appear to regulate by THs (Wright and 2001). As it seems, Alves, thyroid (THs) involve hormones in many physiological processes in teleost fishes. It has been suggested that photoperiod, temperature, and food intake may play species specific role in regulation of seasonal thyroid cycles (Comeau et al., 2000). These seasonal changes may act to promote growth, migratory activity, and reproductive development (Leatherland, 1994). It has been found that the changes of thyroid gland depend on species or population and are sensitive to food intake and diet composition models (Mackenzie et al., 1998). Acanthopagrus latus is a valuable seabream, which has distributed off Indo-West Pacific included Persian Gulf to Australia. It is also recorded from the Gulf of Tadjourah, Djibouti (By et al., 2003). As no detailed study has been carried out on the thyroid patterns of A. latus, in the Persian Gulf (Zangi branch, Musa creek), the present study was conducted on changes of the morphometric structure of thyroid gland and the secretion of thyroid hormones, T3 and T4 in A. latus, in two cold and warm seasons.

# Materials and methods

Sampling conditions

The number of 60 male (all yellowfin seabreams are protandrous namely are male in the first age of their life and then will become females) with weight of 138.5±6.05 g were collected from the northwest Persian Gulf (Musa creek) during a year from July 2007 to June 2008. The water temperature and salinity were ranging from 11 to 34°C and 40 to 43 ppt during cold and warm seasons. respectively.

# Blood sampling

Fish were euthanized with MS222 (200 ppm/100 L). Blood samples (2 cc) were then collected from the caudal vasculature in heparinized syringes (to prevent blood coagulation and the separation of serum, although serum also can be used). The blood samples were kept in ice for up to 30 min and then, plasma was separated using centrifuge (1000 rpm in 10 minutes) and frozen at -20°C for further thyroid hormones analysis.

# Thyroid gland histology

The *A. latus* were dissected to expose the internal organs and the jaws were cut at the corners to expose pharyngeal region. All tissues between the gills were fixed in Bouin's fixative for 72 h and then stored in 70% ethanol. Tissues were dehydrated using an ethanol series and embedded in paraffin (Biswas et al., 2006). The samples were then sectioned at 5-6µm and were stained with hematoxylin and eosin (H&E) and PAS for further basic histological analyses. The cell height of the thyroid epithelium was measured in a total of 15 follicles per fish. Measurements were

made at four points within each follicle at 90° from one another.

Thyroid hormones analysis

The radioimmunoassays (RIAs) were performed using T3 and T4 RIA kits (Immunotech, Beckman Culture Company, France) and gamacounter to determine T3 and T4 levels in plasma, as previously described by Van der Geyten et al. (2001). Their concentrations were computed in ng mL<sup>-1</sup> plasma. The T3 RIA had an intraassay variability of 2.2% and an interassay variability of 9.6%. The T4 RIA had an intra-assay variability of 2.8% and an inter-assay variability of 11.0%. For the T3 RIA cross-reactivity with T4 was 0.1-0.5%, whereas for the T4 RIA crossreactivity with T3 was 3.5%. For both RIA systems, plasma dilution tests and loading tests showed good parallelism with the standard curve.

Statistical analysis

All values of thyroid hormone levels and the height of follicles epithelium were represented as means±SE. The significant difference between warm and cold season values was analyzed using t test. Also correlation was taken between thyroid epithelial cell height and water temperature (Confidence level of 0.05).

#### Results

Structure of thyroid tissue

The results showed that the thyroid gland of A. latus, such as other teleosts is not capsulated. It was composed of follicles, scattered throughout the pharyngeal region along with the dorsal surface of ventral aorta and bronchial arteries. The follicles were round and their walls were consisted of epithelial cells, include follicular cells and a few parafollicular cells, surrounding the central lumen full of colloid fluid. The epithelial cells were cuboidal and squamous during warm and cold seasons, respectively. The mean water temperature of Musa creek, Zangi branch and the mean epithelial cell height for fishes (A. latus) during a year is shown in Figure 1.




Figure 1: Changes of thyroid epithelial cell heights (mean  $\pm$  SE) according to water temperature changes during a year in *Acanthopagrus latus* 




Figure 2: Active thyroid follicle in August (A and B), Parafollicular cells (black circle), follicular epithelial height (two head white arrows); Inactive thyroid follicle in December (C). (H&E; ×40)

The results showed that there is 20% correlation between epithelial cell height and water temperature. Follicular epithelial cells had maximum height in August, then their height significantly decreased to January, after which it slowly increased throughout the winter (P<0.05). Fish thyroid gland was characterized predominance of macrofollicles rich in colloid material during warm months (especially July to August) (Fig. 2A and B), whereas in cold months (especially October to December) thyroid gland showed some microfollicles with less colloid content and more interstitial connective tissue (Fig. 2C). There was a significant increase in ratio of parenchyma to stroma in summer in comparison with winter (P<0.05).

Seasonal change of plasma T3 and T4

The results recorded with the RIA method are shown in Fig. 3. This method confirms that the plasma level of T3 and T4 increased significantly from January to April, and again from April to June. This level was maintained up in summer and the peak of them in plasma occurs during August  $(4.09 \pm 0.41 \text{ and } 1.10 \pm 0.11 \text{ ng})$  $mL^{-1}$ . respectively), then declining significantly during autumn and early from October winter to December (P<0.05) to reach their lowest level in November  $(1.30 \pm 0.36 \text{ and } 0.21 \pm 0.12 \text{ ng})$ mL<sup>-1</sup>, respectively). Both hormones varied similarly across seasons and there was 99% correlation (at the level of 0.01) between two hormones. The increasing of T3 and T4 were correlated with increase of temperature (98 and 42%, respectively) and with the height of thyroid epithelial cell (84 and 82%, respectively).

Table 1: Seasonal variations of the T3 and T4 plasma concentrations during the year in the yellowfin seabream (mean ± SE).

seabream (mean ± SE).		
	T3	T4
July	4.00±0.20	1.04±0.11
Aug.	4.09±0.41	1.10±0.11
Sep.	3.87±0.15	1.00±0.10
Oct.	1.33±0.21	0.24±0.13
Nov.	1.30±0.36	0.21±0.12
Dec.	1.56±0.34	0.31±0.11
Jan.	2.64±0.11	0.64±0.19
Feb.	2.75±0.16	0.72±0.14
Mar.	2.91±0.23	0.75±0.24
Apr.	3.43±0.14	0.82±0.11
May.	3.51±0.24	0.85±0.21
Jun.	3.55±0.2	0.86±0.15




Figure 3: Seasonal variations of the thyroid hormones plasma concentrations during the year in the yellowfin seabream. Bars represent mean  $\pm$  SE.

## **Discussion**

This study established information for two forms of thyroid hormone measured in yellowfin seabreamin warm and cold seasons. No literature indicated any of these hormones, had been previously investigated for fishes in the Persian Gulf. Thyroid hormones are one component of a large complex network of responses to a number of environmental and physiological factors, many of which also growth, development influence and metabolism (Myers et al., 2006).

It was shown that in Atlantic sting ray, *Dasyatis sabina*, follicular cells vary in size and shape, according to the activity of the gland (Volkoff et al., 1999). This was in agreement with the finding of the present study. The surrounding epithelial cells are flattened, cuboidal, or columnar, depending on their activity. Tall, columnar epithelial cells with basophilic colloid containing vacuole-like spaces, characteristics of an active thyroid gland, were seen in warm season. In *Solea senegalensis*, thyroid represented colloid-

filled follicles surrounded by a cuboidal epithelium during summer, suggesting a high activity state of this organ (Ortiz et al., 2006). Swift (1960) suggested that the seasonal changes in thyroidal activity in many teleosts are regulated primarily by water temperature. This relationship of glandular activity and water temperature is interpreted as further evidence that the basic function of the thyroid is concerned in the control of the animal's metabolism, compensate for changes in the environmental temperature. Thus the release of thyrotropic hormone from the pituitary would seem to be influenced by the environmental temperature. In the present study, mean plasma T3 and T4 showed similar seasonal changes patterns, which was in agreement with the finding of Pavlidis et al. (1991). They also reported that similar seasonal variations occurred in circulating thyroid hormones strains of rainbow two trout mykiss Shasta (Oncorhynchus and Kamloops). The results showed that both

hormones decreased significantly during autumn and early winter from October to December according to decrease of temperature. Loter et al. (2007) also reported minimum thyroid hormones in cold months.

In the present study, T3 and T4 increased significantly from January to April, and again from April to July. Duncan et al. (1989) reported a significant increase in the concentrations of both thyroxine and triiodothyronin in mature female channel catfish (Ictalurus punctatus) in February, which was coincident with the steroid hormone peak. They suggested that endocrine interactions between thyroid activity and reproduction may be occurring at this time. Thyroid activity increase in the winter corresponds with intermediate temperatures and feed consumption during rapid reproductive development and spawning period of A. latus which spawn during late winter and early spring (Abou-Seedo et al., 2003). In normal diploid catfish, Heteropneustes fossilis, a general inverse relationship between thyroid hormone levels and advanced reproductive state has been observed (Cyr et al., 1988). It seems that thyroid hormones involve in reproductive maturity. Increase of T3 and T4 plasma concentrations in spring coincides with increasing ambient temperature but the results of the present study showed that the peak activity occurs during midsummer when temperature increase precipitously from July to September. Loter et al. (2007) reported that both increased T4 substrate availability (higher plasma T4 levels) and increased temperature would lead to much greater T3 production in summer. In

summary, the activities of the thyroid hormones deiodination pathways appear to be regulated to provide a much greater availability of T3 in summer, when fish are eating and growing most actively, than in winter (Loter et al., 2007). Decreased food consumption during cold season may depress thyroid hormone cycles in many fishes. The seasonal trend is consistent with the hypothesis that thyroid hormone production is activated during periods of increased nutrient assimilation (MacKenzie et al., 1998).

high A11 together, magnitude seasonal changes of thyroid hormones in male A. latus suggest that this species provides an excellent opportunity to examine the relative contributions of the generation mechanisms of dynamic cycles in circulating thyroid hormone levels. This study was designed to determine basal concentrations of thyroid hormone in A. latus, utilizing assays which have been validated for this species. The relationships between these hormones and food deprivation. reproductive state. other circulating hormones, immunoglobulins and contaminants can now be identified by further analysis.

## References

Abou-Seedo, F.S., Dadzie, S. and Al-Kanaan, K.A., 2003. Sexuality, sex change and maturation patterns in the yellowfin seabream, *Acanthopagrus latus* (Teleostei: Sparidae) (Houttuyn, 1782). *Journal of Applied Ichthyology*, 19, 65–73.

Biswas, A., Kundu, S., Roy, S., De, J., Pramanik, M. and Ray, A.K., 2006.

Thyroid hormone profile during

- annual reproductive cycle of diploid and triploid catfish, *Heteropneustes fossilis* (Bloch). *General and Comparative Endocrinology*, **147**, 126–132.
- By, F., Abou-Seedo, S., Dadzie, S. and Al-Kanaan, K.A., 2003. Sexuality, sex change and maturation patterns in the yellowfin seabream, *Acanthopagrus latus* (Teleostei: Sparidae). *Journal of Applied Ichthyology*, 19, 65-73.
- Comeau, L.A., Campana, S.E., Hanson, J.M. and Chouinard, G.A., 2000. Seasonal changes of thyroid hormones in field-collected Atlantic cod in relation to condition indices, water temperature and photoperiod. *Journal of Fish Biology*, **57**, 571–588.
- Cyr, D.G., Brommage, N.R., Duston, J. and Eales, J.G., 1988. Seasonal patterns in plasma levels of thyroid hormones and sex steroids in relation to photoperiod-induced changes in spawning time in rainbow trout, *Salmo gairdneri*. *General and Comparative Endocrinology*, **69**, 217–225.
- **Leatherland, J.F., 1994.** Reflections on the thyroidology of fishes: from molecules to humankind. *Guelph Ichthyology Reviews*, **2**, 1–67.
- Loter, T.C., MacKenzie, D.S., McLeese, J. and Eales, J.G., 2007. Seasonal changes in channel catfish thyroid hormones reflect increased magnitude of daily thyroid hormone cycles. *Aquaculture*, 262, 451–460.
- MacKenzie, D.S., Thomas, P. and Farrar, S.M., 1989. Seasonal changes in thyroid and reproductive steroid hormones in female channel catfish

- (*Ictalurus punctatus*) in pond culture. *Aquaculture*, **78**, 63-80.
- MacKenzie, D.S., VanPutte, C.L.M. and Leiner, K., 1998. Nutrient regulation of endocrine function in fish. *Aquaculture*, 161, 3–25.
- Myers, M.J., Rea, L.D. and Atkinson, S., 2006. The effects of age, season and geographic region on thyroid hormones in Steller sea lions (Eumetopias jubatus). Comparative Biochemistry and Physiology, 145, 90–98.
- Ortiz, J.B., Ruane, N.M., Pousão-Ferreira, P., Dinis, M.T. and Sarasquete, C., 2006. Thyroid gland development in Senegalese sole (*Solea senegalensis* Kaup 1858) during early life stages: A histochemical and immunohistochemical approach. *Aquaculture*, 260, 346–356.
- Pavlidis. M., Dessypris. and Christofidis. I.. 1991. Seasonal fluctuations in plasma thyroid hormones, in two strains of rainbow trout (Oncorhynchus mykiss), during the first and second reproductive with cycle: relation their photoperiodically altered spawning time. Aquaculture, 99, 365-385.
- Power, D.M., Llewellyn, L., Faustino, M., Nowell, M.A., Bjornsson, B.T., Einarsdottir, I.E., Canario, A.V.M. and Sweeney, G.E., 2001. Thyroid hormones in growth and development of fish. *Comparative Biochemistry and Physiology*, 130, 447–459.
- Szischa, V., Papandroulakisb, N., Fanourakia, E. and Pavlidisa, M., 2005. Ontogeny of the thyroid hormones and cortisol in the gilthead

- sea bream, *Sparus aurata. General and Comparative Endocrinology*, **142**, 186–192.
- **Swift, D.R., 1960.** Cyclical activity of the thyroid gland of fish in relation to environmental changes. *Symposia of the Zoological Society of London*, **2**, 17–27.
- Van der Geyten, S., Toguyeni, A., Baroiller, J.F., Fauconneau, B., Fostier, A., Sanders, J.P., Visser, T.J., Kühn, E.R. and Darras, V.M., 2001. Hypothyroidism induces type I iodothyronine deiodinase expression in tilapia liver. *General and Comarative Endocrinology*, 124, 333–342.
- Volkoff, H., Wourms, J.P., Amesbury, E. and Snelson, F. F., 1999. Structure of the thyroid gland, serum thyroid hormones, and the reproductive cycle of the Atlantic stingray, *Dasyatis sabina*. *Journal of Experimental Biology*, 284(5), 505-516.
- Wright, M. L. and Alves, C. D., 2001. The decrease in plasma melatonin at metamorphic climax in *Rana catesbeiana* bullfrog tadpoles is induced by thyroxine. *Comparative Biochemistry and Physiology Part A*, 129, 653–663.